

Contents

Acknowledgments — VI

Prologue — 1

P.1 Southwestern Australia and its Functional Biodiversity — 1

P.2 Environmental and Biotic Constraints — 2

P.3 Our Book — 4

1 Evolution and Diversity of the Flora — 6

1.1 A Radiating and Hyperdiverse Flora — 6

1.2 Evolutionary Constraints and Responses — 11

1.3 The SouthWest Flora 45 Million Years Ago and Now — 13

1.4 A Relict Podocarp — 17

1.5 Fire and Evolution of the Proteaceae — 18

1.6 Floristics and Phytogeography — 22

1.7 The SouthWest as a Biodiversity Hotspot — 27

2 Fire Adaptations — 30

2.1 Fire-enhancement — 40

2.2 Fire-dependence — 40

2.3 Fire-resistance — 42

2.4 Fire-avoidance — 44

2.4.1 Post-fire Persistence *via* Resprouting — 45

2.4.2 Lignotubers — 46

2.4.3 Epicormic Growth — 48

2.4.4 Rhizomes — 48

2.5 Fire-sensitivity — 49

2.6 Flammability — 50

2.7 Evolution of Fire-adapted Traits — 52

3 Drought Responses — 58

3.1 Herbaceous Drought Avoiders and Resisters — 59

3.2 Drought Response of Woody Plants — 60

3.2.1 Diallagy — 62

3.3 Classifying Species into Functional Groups — 62

3.3.1 Functional Significance of Rooting Depth Patterns for Groundwater-Dependent Vegetation — 66

3.3.2 Woody Perennial Rooting Patterns — 67

3.4 Stomatal Location—a Drought Response? — 68

3.5 Case Study: Ecophysiology of Dimorphic Leaves in *Hakea trifurcata* — 70

4	Carnivorous Plants — 72
4.1	<i>Drosera</i> — 72
4.2	<i>Byblis</i> — 75
4.3	<i>Cephalotus</i> — 76
4.4	<i>Utricularia</i> — 77
5	Parasitic Plants — 79
5.1	Stem Hemiparasites - Mistletoes — 79
5.1.1	Host Mimicry — 80
5.1.2	Seed Dispersal — 82
5.2	Root Hemiparasites — 84
5.2.1	<i>Nuytsia floribunda</i> (Loranthaceae) — 84
5.2.2	<i>Olex phyllanthi</i> (Olacaceae) — 86
5.2.3	<i>Santalum</i> (Santalaceae) — 86
5.3	Stem Holoparasites — 87
5.3.1	<i>Pilostyles</i> (Apodanthaceae) — 87
5.3.2	<i>Cuscuta</i> (Convolvulaceae) — 89
5.4	Stem Hemi or Holoparasite? <i>Cassytha</i> (Lauraceae) — 89
6	Specialised Nutrient Uptake Mechanisms — 91
6.1	Nitrogen — 91
6.1.1	Cyanobacterial Symbionts — 91
6.1.2	Rhizobial Symbionts — 93
6.1.3	Actinorhizal Symbionts — 93
6.2	Phosphorus — 95
6.2.1	Mycorrhizal Associations — 95
6.2.2	Arbuscular Mycorrhizas — 96
6.2.3	Ectomycorrhizas — 97
6.2.4	Orchid and Ericoid Mycorrhizas — 98
6.3	Case Study: Dispersal of Ectomycorrhizal Fungi by Marsupials — 103
6.4	Root Clusters — 104
6.4.1	Proteoid Roots — 104
6.4.2	Dauciform Roots — 106
6.4.3	Capillaroid Roots — 106
6.5	The Bizarre Root System of <i>Kingia australis</i> — 106
6.6	Nutrient-impoveryished Soils Promote Species Richness — 108
7	Pollination Strategies — 109
7.1	Outbreeding Strategies — 110
7.1.1	Protandry — 111
7.1.1.1	<i>Eucalyptus</i> (Myrtaceae) — 111
7.1.1.2	<i>Pimelea</i> (Thymelaeaceae) — 112

- 7.1.1.3 Goodeniaceae — **112**
- 7.1.1.4 *Banksia* (Proteaceae) — **113**
- 7.1.1.5 *Petrophile* and *Isopogon* (Proteaceae) — **113**
- 7.1.1.6 *Grevillea* and *Hakea* (Proteaceae) — **115**
- 7.1.1.7 *Persoonia* (Proteaceae) — **116**
- 7.1.1.8 *Stylidium* (Stylidiaceae) — **116**
- 7.1.1.9 *Eremophila* (Scrophulariaceae) — **117**
- 7.1.2 Protogyny — **118**
- 7.2 Autogamy — **118**
- 7.3 Attracting Pollinators — **119**
- 7.4 Flower Colour Change — **120**
- 7.5 What Role Do Extrafloral Nectaries Play? — **122**

8 Pollination Syndromes — 126

- 8.1 Insect-pollinated Species — **127**
- 8.1.1 Case study: *Grevillea leucopteris* — **127**
- 8.1.2 Bee Pollination — **130**
- 8.1.3 Orchid Pollination — **135**
- 8.1.3.1 Food Rewards — **137**
- 8.1.3.2 Pollinator Deception — **138**
- 8.1.3.3 Gnat Pollination — **140**
- 8.1.3.4 Sexual Deception — **142**
- 8.2 Honeyeater-pollinated Species — **145**
- 8.2.1 Case Study: *Gastrolobium* — **148**
- 8.2.2 Pollen Dispersal — **149**
- 8.3 Mini-possum Pollination — **150**

9 Leaf Properties — 153

- 9.1 Leaf Size — **153**
- 9.2 Leaf Design — **156**
- 9.2.1 Reducing Summer Leaf Heat Loads — **156**
- 9.2.2 Heterophylly — **157**
- 9.3 Sclerophylly — **158**
- 9.3.1 Case Study: Two Banksias With Long-lived Leaves — **161**
- 9.4 Leaf-like Structures — **162**
- 9.5 Alternatives to Sclerophylly for Protecting Leaves — **164**
- 9.6 Leaves as Defensive Mechanisms Against Herbivores, Florivores and Granivores — **166**
- 9.6.1 Cryptic Mimicry — **169**
- 9.6.2 Fruits Hidden by Leaves — **169**
- 9.6.3 Herbivore Defence Among Restios — **171**

10	Seed Release and Dispersal Mechanisms — 172
10.1	Pyriscence of Serotinous Fruiting Structures — 173
10.2	Wind Dispersal — 175
10.2.1	Secondary Dispersal — 177
10.3	Ant Dispersal — 177
10.3.1	Fate of Ant-dispersed Seeds — 180
10.3.2	Case Study: <i>Adenanthos cygnorum</i> — 182
10.4	Vertebrate Dispersal — 183
10.4.1	Emus and Large Macropods as Long-Distance Seed Dispersal Agents — 184
10.5	Unassisted Dispersal — 187
10.5.1	Kangaroo Paws — 187
10.5.2	Orchid Seeds — 188
11	Seed Storage, Germination and Establishment — 189
11.1	Canopy-stored Seeds — 190
11.1.1	Adaptive Advantages of Canopy-stored Seeds — 193
11.1.2	Protective Properties of Woody Fruits — 193
11.1.3	Post-fire Germination and Seedling Establishment — 197
11.2	Soil-stored Seeds — 198
11.2.1	Advantages of Storing Seeds in the Soil — 202
11.2.2	Case Study: <i>Austrostipa compressa</i> and <i>Ehrharta calycina</i> — 203
11.3	Seed Size and Seedling Survival — 204
11.3.1	Small Seeds Produce Small Seedlings — 206
Bibliography — 209	
About the Authors — 234	
List of Figures — 235	
List of Tables — 252	
Index — 254	