



REVIEW

Open Access

Comparative energetics and kinetics of autotrophic lipid and starch metabolism in chlorophytic microalgae: implications for biomass and biofuel production

Sowmya Subramanian¹, Amanda N Barry², Shayani Pieris³ and Richard T Sayre^{1,2*}

Abstract

Due to the growing need to provide alternatives to fossil fuels as efficiently, economically, and sustainably as possible there has been growing interest in improved biofuel production systems. Biofuels produced from microalgae are a particularly attractive option since microalgae have production potentials that exceed the best terrestrial crops by 2 to 10-fold. In addition, autotrophically grown microalgae can capture CO₂ from point sources reducing direct atmospheric greenhouse gas emissions. The enhanced biomass production potential of algae is attributed in part to the fact that every cell is photosynthetic. Regardless, overall biological energy capture, conversion, and storage in microalgae are inefficient with less than 8% conversion of solar into chemical energy achieved. In this review, we examine the thermodynamic and kinetic constraints associated with the autotrophic conversion of inorganic carbon into storage carbohydrate and oil, the dominant energy storage products in Chlorophytic microalgae. We discuss how thermodynamic restrictions including the loss of fixed carbon during acetyl CoA synthesis reduce the efficiency of carbon accumulation in lipids. In addition, kinetic limitations, such as the coupling of proton to electron transfer during plastoquinone reduction and oxidation and the slow rates of CO₂ fixation by Rubisco reduce photosynthetic efficiency. In some cases, these kinetic limitations have been overcome by massive increases in the numbers of effective catalytic sites, e.g. the high Rubisco levels (mM) in chloroplasts. But in other cases, including the slow rate of plastoquinol oxidation, there has been no compensatory increase in the abundance of catalytically limiting protein complexes. Significantly, we show that the energetic requirements for producing oil and starch relative to the recoverable energy stored in these molecules are very similar on a per carbon basis. Presently, the overall rates of starch and lipid synthesis in microalgae are very poorly characterized. Increased understanding of the kinetic constraints of lipid and starch synthesis, accumulation and turnover would facilitate the design of improved biomass production systems.

Keywords: Biofuel, Algae, Plant, Storage carbohydrate, Starch, Oil, Energetics, Photosynthesis, Biomass, Metabolism, Cultivation

* Correspondence: rsayre@newmexicoconsortium.org

¹New Mexico Consortium, 100 Entrada Rd., Los Alamos, NM 87544, USA

²Bioscience Division, Los Alamos National Lab, M888, Los Alamos, NM 87545, USA

Full list of author information is available at the end of the article

Introduction

Concerns about higher energy prices, finite fossil fuel reserves, rising atmospheric CO₂ levels, and energy security have led to a growing interest in developing domestically produced renewable sources of energy using biomass production systems [1,2]. Given the low solar energy density on the earth's surface and the inefficiencies of photosynthesis, robust biomass production systems that are optimized for the local environment and which have the greatest energy-return-on-investment are (EROI) desirable for producing biofuels. Recently, there has been growing recognition that microalgae have among the highest potential to produce the greatest biomass per unit area in the shortest period of time. Furthermore, microalgae have enhanced environmental sustainability characteristics since they can utilize inorganic carbon (bicarbonate) sequestered in ponds that can be captured from CO₂ production point sources (power plants), unlike terrestrial plants. In addition, microalgae produce feedstocks (oils) for conversion to fuels that are compatible with the existing liquid transportation infrastructure [3]. Compared to crop plants, microalgae have 2–10 times the biomass yield potential. This enhanced productivity is in part attributed to the fact that all cells are photosynthetic unlike plants and that algae have active carbon concentrating mechanisms (CCMs) to enhance photosynthetic efficiency (Figure 1). Furthermore, some microalgae are capable of accumulating large amounts of lipids (up to 70% w/w) [4-7], can recycle water and nutrients from effluent streams, and do not directly compete with food production [2,8]. In addition, hydrocarbons produced by algae represent a potential means to store and sequester carbon [2]. Finally, from the

energy security perspective a production failure in an algal pond can generally be brought back on line in a matter of days whereas a crop failure in a terrestrial crop production system may take up to a year before another harvest can occur.

Similar to all biomass production systems, however, microalgae have restricted cultivation areas. Cost-effective algal biomass production is constrained by topography (slopes less than 1%), water availability, temperature, nutrients (CO₂, N and P), and the competing requirements for agricultural food production [5,9]. Production methods include open ponds, closed reactors, matrix-immobilized algae, and algal biofilms (for an in-depth review on production and harvesting techniques currently in use, see [10]). Several open pond systems have been evaluated at the pilot scale for microalgal growth, harvesting and biofuel production [11]. In contrast to open ponds, closed photobioreactors (PBRs) can be more productive on an aerial basis and allow stricter control over light, temperature, pH and nutrient inputs. However, recent comparative life cycle analyses of open pond versus closed PBR systems indicate that the higher capex and opex costs of closed PBRs make them prohibitively expensive to produce biofuels [12].

Unlike crop systems, many microalgae are also capable of heterotrophic growth and can use a variety of reduced carbon sources for biomass or oil accumulation. For example *Chlamydomonas reinhardtii* can assimilate acetate. Other species can be grown heterotrophically on sugars such as glucose which can enhance oil accumulation by as much as 900% [13]. Glycerol, a byproduct of the production of biodiesel production, can also be utilized for heterotrophic or mixotrophic growth by some

Biofuels from Microalgae




Figure 1 The advantages of algal biomass production systems. CCM; carbon concentrating mechanism.

microalgal species to achieve higher lipid productivity [14-16]. Although heterotrophic growth has been shown to produce high levels of desirable neutral lipids (for review, see [17]), addition of a reduced carbon source to algal ponds is not always feasible due to higher susceptibility to microbial contamination and cost. Finally, due to their low culture density (0.1% of mass) in ponds and small cell size (1–10 μm) considerable energy, accounting for up to 40% of production costs, may be required to harvest microalgae [8,18]. Substantial progress, however, is being made in the development of new technologies to reduce cultivation and harvesting costs (NAABB Consortium web site: <http://www.naabb.org>). Ultimately, for algal biofuel production systems to be financially, energetically and temporally successful, the financial (FROI) and EROI for algal biofuel production systems must approach or be greater than that of competing fuel technologies. Economic analyses indicate that improvement of lipid content and growth rate are the predominant cost drivers for economical algal biofuel production [18].

In this review, we consider the kinetic and thermodynamic constraints of the cellular aspects of microalgal (Chlorophytic) biomass production systems. We demonstrate that the metabolic and physiological constraints impacting biomass accumulation are poorly characterized and need targeted research investment to facilitate the development of targeted strategies to improve biomass production and accumulation in microalgae. Much of our understanding of photosynthesis and carbon fixation stems from studies on cyanobacteria, Chlorophyta and higher plants. Because microalgae are a very diverse group of organisms in their entirety, we will restrict the bulk of our discussions to the Chlorophyta.

Kinetics of biomass accumulation

Microalgae have a broad range of growth rates and energy densities depending on the species and growth conditions. Biomass productivity rates range from 15–30 g dry weight $\text{m}^{-2} \text{day}^{-1}$ [19,20] with an average energy content of 4.3 to 7.0 kcal/gdw [21,22]. The higher energy densities are typically associated with higher lipid contents. In general there is often a trade off, however, between cell division rates and the accumulation of energy storage products including storage carbohydrate (such as starch and glycogen) and neutral lipids or triacylglycerol (TAGs). Many studies have demonstrated that a variety of stress conditions that reduce growth rates lead to enhanced storage carbohydrate and/or oil accumulation [23-25]. Oil accumulation induced by stress may increase the energy content per dry weight by 50% or more but also may involve substantial lipid remodeling [24,25]. There are a number of factors, however, that determine overall rates of biomass accumulation at the cellular level. Some of these factors include: the absorption

spectrum of the photosynthetic pigments, the optical cross-section of the light-harvesting apparatus, energy transfer efficiency, enzyme rate kinetics, the numbers and concentrations of energy transfer complexes (electron transfer chains) and enzyme catalytic sites, substrate concentrations, metabolite flux rates, metabolic compartmentalization and feedback control, respiration rates, and the partitioning of reduced carbon between new cell growth and division and energy storage [26-28].

Kinetics of light capture and energy conversion

The earliest event in biological solar energy conversion is light capture. Energy capture and transfer by the light harvesting pigment-protein complexes is among the most energy efficient and fastest processes known in any biological system. Green algae (Chlorophyta) and higher plants initially capture photons via their light harvesting antenna complexes (LHC). The pigments associated with the LHC complexes in green algae account for about 80% of the total chlorophyll content with the remaining 20% of pigments associated with the proximal antenna and reaction center (RC) complexes where charge separation occurs [29]. In eukaryotic algae and plants the chlorophylls and carotenoids of the LHC complexes are bound to thylakoid membrane proteins in close association with the RCs. Energy transfer between pigments of the LHC complexes occurs on the femtosecond time scale [30]. Recent experimental and theoretical studies of electronic energy transfer processes in LHCs have revealed that energy transfer between pigments in the peripheral LHCs occurs at nearly 100% efficiency mediated by long-lived quantum coherence energy transfer processes [31-33]. Advances in techniques used to probe quantum coherence (2D- and multi-dimensional electronic spectroscopy) effects in energy transfer have provided a wealth of information on the mechanism and nature of energy transfer, charge separation and connectivity between the donor-acceptor states [32,34].

The first kinetic constraints in energy transfer following light capture occur at the interfaces between the peripheral and proximal antenna as well as between the proximal antenna and the RC. For example, energy transfer between the proximal antenna complexes, CP43/CP47 and the Photosystem II (PSII) RC occurs at a time scale of 20–30 ps, while the subsequent primary charge separation in the PSII-RC occurs within 1–3 ps [29,35]. These kinetic bottlenecks result in some energy losses at higher ($\geq 25\%$ of full sunlight) than saturating light intensities. However, the greatest kinetic bottleneck in photosynthetic electron transfer is the coupled transfer of protons and electrons associated with the diffusion and oxidation of plastoquinol (PQH_2) mediated by cytochrome b_6/f ($\text{Cyt}b_6f$) complex [36,37]. This has significant implications for the efficiency of photon utilization at various light intensities.

Further complicating the picture is the fact that each independent electron transfer complex in the photosynthetic apparatus has substantially different rate-limiting kinetics. The overall rate-limiting steps for charge separation or electron transfer in the photosystem I (PS I), PS II and *cytb₆f* complex are 1 ns, 1 μ s and 10 ms, respectively (Figure 2) [38–40]. Additional proton coupled steps also kinetically constrain energy transduction including the slow diffusion rates of plastoquinone [37] as well as the relatively slow turnover ($100\text{--}200\text{s}^{-1}$) of the ATP synthase complex [41,42]. It is noteworthy that a variety of metabolic control mechanisms may also impact rate kinetics. For example, the ATP synthase can function in the reverse direction *viz.*, hydrolysis of ATP depending on the trans-thylakoidal pH as well as its redox control by thioredoxins [43] leading to reduction in ATP synthesis rates. However, in a recent paper by Yamori *et al.*, in which the authors describe the effects of reduced ATP synthase and *Cytb₆f* contents on overall electron transfer rates and CO_2 assimilation rates, they observed that reductions in *Cytb₆f* content had the greatest impact on overall electron transfer rates, and not alterations in ATP synthase levels [44]. To further complicate matters the activity of the ATP synthase complex has been shown to be modulated by CO_2 concentrations [43,45–47].

Significantly, rate-limiting steps in electron transfer indirectly impact the means by which the energy of chlorophyll excited states in the LHC complex are dissipated either by photochemistry or by re-emission of trapped energy as heat or fluorescence. The relative proportion of captured energy that is converted into charge separated states or dissipated by non-photochemical quenching (NPQ) mechanisms is determined by both the light intensity, the optical cross-section of the light harvesting apparatus and the kinetics of electron transfer processes. In *Chlamydomonas* the apparent optical cross section of the photosystem II (PS II) antenna complex is about 70 \AA^2 [48]. At a photon flux density (400–

700 nm) of $2,000 \mu\text{mole photons/m}^2/\text{sec}$, equivalent to full sunlight intensity, the rate of photon capture approaches 1.2 photons per ms per PSII complex. This rate of photon capture is five to eight times faster than the rate-limiting step (PQH_2 oxidation) in photosynthetic electron transfer (discussed below). Thus, at full sunlight intensity about 75% of the energy that is captured by the LHC antenna complex does not drive photochemistry but is dissipated as heat or fluorescence by non-photochemical quenching (NPQ) mechanisms [49,50]. Similar inefficiencies in light capture and conversion have been observed in whole plants. Björkman *et al.*, [51] determined from a study of 42 species of plants that the average photochemical quantum yield for oxygen evolution was 0.108. Similarly, Long *et al.*, [52] measured a maximum, average photochemical quantum yield of 0.093 from CO_2 exchange measurements from 11 species of C3 plants. These studies indicate that most plants and algae have light harvesting systems that are larger than necessary for efficient light capture and energy conversion. Consistent with this observation, it has been demonstrated that reductions in the size of the LHC complex associated with the inhibition of chlorophyll b synthesis in green algae leads to higher rates of photosynthesis in algal cultures at high light intensities [33,50,53,54]. Recent studies using algae engineered to have a range of peripheral light-harvesting antenna sizes have demonstrated that the optimal antenna size for efficient conversion of light into chemical energy in green plants and in *Chlamydomonas* mutants is approximately 30% smaller than wild type, a size that is consistent with efficient coupling of the rate kinetics of light capture with the limitations of downstream electron transfer steps [33].

Kinetics of carbon fixation

The overall rate-limiting step in photosynthesis, however, is the initial fixation of CO_2 by the enzyme






Figure 2 Model of the photosynthetic electron transfer system of plants and Chlorophyta demonstrating the lateral membrane heterogeneity of the major protein complexes. Abbreviations: Fd, ferredoxin; PC, plastocyanin; PQ, plastoquinone; PSI, photosystem I; PSII, photosystem II

times the number of PSII and PSI RCs (Figures 3 and 4). Due to the very large number of total Rubisco catalytic sites the overall rates of carbon dioxide fixation approach maximum rates of electron transfer. The efficiency of Rubisco is reduced, however, by the





competitive fixation of oxygen versus carbon dioxide and the fact that much of the enzyme may be in an inactive state. Rubisco oxygenase activity reduces the efficiency of CO_2 fixation, the regeneration of the substrate RuBP, and results in the loss of previously fixed carbon during the decarboxylation of serine in the photorespiratory pathway (Figure 3) [56]. The net reduction in CO_2 fixation by the photorespiratory pathway has been estimated to be 25% of the maximum carboxylation efficiency [56]. Many microalgae facultatively (when grown in air) reduce photorespiratory losses by actively pumping bicarbonate into the cells where it is converted back to CO_2 . This can occur in specialized sub-organelle structures known as pyrenoids in eukaryotic algae where carbonic anhydrase is closely associated with Rubisco to facilitate the conversion of bicarbonate to carbon dioxide [57]. This type of algal CCM is best characterized in the green alga, *Chlamydomonas reinhardtii* [57,58]. In *Chlamydomonas*, bicarbonate is actively transported across the plasmamembrane by an ABC-type transporter (HLA3) that presumably requires at least one additional ATP/ CO_2 fixed, as shown by gene knockout studies [59]. This additional ATP demand to pump carbon dioxide into the cell is met by cyclic photophosphorylation [60]. Subsequently, bicarbonate is transported into the plastid by the LCIA transporter and dehydrated by carbonic anhydrase to generate CO_2 . Similarly, cyanobacteria employ a number of bicarbonate transporters and carbonic anhydrases to increase the CO_2 levels in specialized compartment called carboxysomes, where carbon fixation occurs. In diatoms, CCM occurs by both actively

increasing CO_2 levels near the site of carbon fixation (Rubisco), and C4 fixation.

It has been demonstrated in plants that there are additional kinetic constraints in the Calvin-Benson Cycle associated with the limiting abundance of enzymes that have very high equilibrium constants including; Seduheptulose biphosphatase (SBPase), fructose biphosphatase (FBPase) and aldolase [38]. Overexpression of the enzymes, FBPase and SBPase in tobacco lead to increased sugar and storage carbohydrate content, improving biomass yield [61]. Similar results have been achieved in rapidly dividing algal cells (Sayre, unpublished results). Metabolic flux rates through the Calvin-Benson Cycle have been described by a number of labs [62,63], however carbon flux through carbon storage and sequestration pathways leading to storage carbohydrate and oil synthesis have not been well characterized in microalgae. Thus, it is not possible to account for potential differences in rate kinetics for overall storage carbohydrate and oil synthesis.

Thermodynamics of energy capture and conversion in algae

The thermodynamic efficiency of light capture and energy conversion by the photosynthetic electron transfer system as well as the efficiency of carbon fixation, metabolism, respiration and carbon sequestration determine the overall energy efficiency of solar energy conversion into chemical energy present in biomass [64,65]. Of the total solar radiation that reaches the earth's surface, photosynthetically active radiation (PAR) is limited to 400–700 nm for most eukaryotes, and accounts for less than 50 % of the total solar photon flux [64].

In plants and the Chlorophytes, the light harvesting complexes (LHCs) or antenna protein-pigment complexes are comprised of a superfamily of proteins. The LHC proteins bind chlorophylls and carotenoids (Car) that capture light and transfer energy to the reaction centers (RCs). Chlorophyll *a* (Chl *a*), is present in all eukaryotic photosynthetic organisms. Additional species of chlorophyll including Chl *b*, Chl *c* or Chl *d* along with carotenoids (Car) expand the light absorption spectrum of the light harvesting antenna complexes [66]. In green algae (Chlorophyta) and higher plants, the peripheral antenna include Chl *a/b*-binding proteins whose pigments absorb predominantly in the blue (430–455 nm) and red (650–690 nm) region of the solar spectra.

In plants and green algae, the LHC proteins associated with photosystem II (LHCII/peripheral antenna) form a trimeric complex (major LHC) in association with three other proteins (minor LHC). Typically, the PSII RC and proximal antenna proteins of green algae and higher plants bind about 6 Chl *a*/2Car and 28 Chl *a*, respectively [67,68]. The PSII crystal structure of spinach demonstrates that the major [67,68] LHCII binds about 8

Chl *a*/6 Chl *b*/4 Car per monomer [69], while the minor LHCII bind about 5–6 Chl *a*/2–5 Chl *b*/2 Car. Structural studies of PSI revealed that the PSI associated LHC (LHCI) is a tetrameric protein with 14 Chl *a*+*b* molecules bound per monomer [70], and the PSI core complex binds about 200 molecules of Chl *a* and about 20 carotenoids.

Members of the LHC family also bind carotenoids [71]. The carotenoids serve two roles in the LHC: 1) They expand the spectra of light captured by the photosystems, and 2) dissipate the excess excitation energy from Chl through non-photochemical quenching (NPQ). As previously discussed, under high light conditions electron transfer is saturated, and this results in the accumulation of Chl singlet excited states ($^1\text{Chl}^*$). When photochemistry is saturated, $^1\text{Chl}^*$ relax to the ground state by emitting energy as heat and/or fluorescence or by singlet-singlet exciton annihilation. During the latter process, spin inversion results in the formation of $^3\text{Chl}^*$ which can interact with $^3\text{O}_2$ to produce singlet $^1\text{O}_2$ species, a very potent oxidant that causes damage to the photosynthetic apparatus or photoinhibition [72–74]. In addition, excited state Chls can be quenched by zeaxanthin [75] resulting in the emission of heat. This process is induced by increased acidification of the lumen, which triggers the xanthophyll cycle, wherein de-epoxidases convert violaxanthin to zeaxanthin. However, the exact mechanism by which zeaxanthin quenches the excess energy from Chl excited states remains a subject of controversy [49,76,77].

Thermodynamics of carbon reduction

The quantum efficiency of primary charge separation in photosystem I and II is quite high. Energy transfer from the PSII antenna pigments to the RC primary electron donor chlorophyll (P680) in PSII leads to charge transfer. The primary electron acceptor, pheophytin (Phe) accepts an electron from P680* to form a radical pair state (P680*Phe $^-$). The radical pair in turn reduces the secondary electron acceptor Q_A, a single electron-accepting plastoquinone molecule, with lower energy than Phe $^-$ resulting in the stabilization of the charge separated state. The transfer of electrons in an energetically downhill fashion contribute to the thermodynamic efficiency and the high quantum efficiency (0.8–0.9) of the charge separation process [78,79]. A similar process of charge separation and thermodynamically downhill energy transfer occurs in PSI-RC (P700). The photoexcited P700* reduces the primary acceptor Chl molecules (A₀ & A₁) which stabilizes primary charge separation with a quantum efficiency of ~ 1 [47,80].

The overall apparent quantum or thermodynamic efficiency of photosynthesis is generally normalized, however, to the number of photons required to evolve one

molecule of O₂ or fix one molecule of CO₂ (Figure 2 and 3). A theoretical maximum quantum efficiency of 8 red (680 nm) photons/CO₂ fixed was proposed by Emerson *et al.*, in the late 1950s [81]. Evidence based on the Emerson enhancement effect, which identified the role of the two photosystems working in series, and determination of the quantum requirement through calorimetric studies by Arnold *et al.*, [82] supported a quantum requirement of 10–12 photons per CO₂ fixed. While, the difference in the quantum yield owing to the difference in the measurement technique (O₂ evolution vs. CO₂ assimilation) can be appreciated, recent reports indicate that the quantum requirement for CO₂ fixation in plants in the absence and presence of photorespiration is ~ 9 and ~ 10 photons, respectively [83,84]. However, the experimental photosynthetic quantum efficiency ranges as high as 20 for plants under stress [85]. Given that the stoichiometric ratio of ATP and NADPH per CO₂ fixed is 3:2 (from Calvin cycle) and that 4 reducing equivalents are required to generate 2 NADPH or one O₂, the magnitude of the proton motive force necessary to drive synthesis of sufficient ATP to fix CO₂ then becomes the critical factor to balance the energy requirements for CO₂ fixation. Recent experimental evidence suggests that the H⁺/ATP ratio is approximately 4 [86–88]. These results are consistent with structural inferences from the ATP synthase complex. The CF₀ rotor ring of the CF₀-CF₁ complex of the ATP synthase has 14 C-subunits, each of which must be sequentially protonated to drive ATP synthesis. The fact that the ATPase has 3 catalytic sites that must complete a rotation cycle to generate 3 ATPs, indicates that the H⁺/ATP ratio is approximately 4.67 or 14 protons/3ATP [89,90]. Since the proton requirement/ATP synthesized is 4 and 4 ATPs are required for each CO₂ pumped (1 ATP) and fixed (3 ATP) the proton requirement (16 H⁺) to produce sufficient ATP to fix CO₂ cannot be met by linear electron transfer of 4 electrons to produce 2 NADPH. Additional proton motive force can be generated by a variety of mechanisms. On average, each light-driven charge separated state generates the stoichiometric equivalent of 1.25 protons/electron. This occurs via a combination of multiple proton generating or transducing steps including water splitting (1 H⁺/e $^-$), PQH₂ oxidation (1 H⁺/e $^-$), proton transfer via the Q cycle (1.5 H⁺/e $^-$), and cyclic photophosphorylation (1–1.5 H⁺/e $^-$). Considering water splitting and the Q-cycle alone, twelve proton equivalents are expected to be generated from 8 photochemical events associated with the production of 2 NADPH molecules [82]. Since the proton requirement/ATP synthesized is > 4 and 4 ATPs are required for each CO₂ pumped and reduced by the Calvin-Benson Cycle the proton requirement (~ 16 H⁺) to produce sufficient ATP to fix CO₂ cannot be met by

linear electron transfer. The additional proton requirement must be provided by cyclic photophosphorylation [91]. In cyclic photophosphorylation, electrons derived from ferredoxin reduce plastoquinone in the cytochrome b_6f complex. Reduced ferredoxin is generated by photosystem I. Reduced PQ molecules are protonated on the stromal side (Q_I site) of the $Cytb_6f$ complex and are re-oxidized at the Q_O site on the lumenal side of the thylakoid, thus increasing the proton gradient at the expense of NADPH synthesis.

As stated previously, cyclic photophosphorylation is also essential for active bicarbonate uptake in green microalgae [92,93]. In air-grown chlorophytic microalgae such as *Chlamydomonas* the biosynthesis of any reduced carbon storage product begins with the active uptake of bicarbonate and the subsequent reduction of the imported inorganic carbon or CO_2 via the Calvin-Benson Cycle. For *Chlamydomonas*, one additional ATP is required to support its active transport into the cell by a plasmamembrane localized ABC-type bicarbonate transporter, the HLA3 protein (David Kramer, personal communication). Recent evidence indicates that bicarbonate uptake by the HLA3 transporter requires one additional ATP/ CO_2 fixed. Bicarbonate is then transported into the chloroplast stroma by the LCIA transporter and eventually to the thylakoid lumen by another putative bicarbonate transporter. Carbonic anhydrases localized in the stroma and the lumen, catalyze the conversion of bicarbonate to CO_2 , effectively elevating the internal CO_2 concentration by 10-fold and competitively inhibiting the oxygenase reaction of Rubisco [57]. This inorganic carbon pumping system effectively reduces photorespiration by more than 90% improving overall photosynthetic efficiency [94,95].

Storage carbohydrate (starch) and oil production during autotrophic growth

During active photoautotrophic growth carbon in excess of that required for metabolism, respiration and growth is stored typically as carbohydrate or TAGs. As shown in Table 1, the storage carbohydrate (polysaccharides) or oil (lipid) content of algae can range anywhere from 6 to 64% of the total biomass. Typically, only carbohydrates and oils are utilized for the production of commercial fuels, however. For this reason, as well as to simplify the thermodynamic considerations for biofuels production, we only consider the energetic costs for carbohydrate and oil production. These calculations are made under standard physical conditions since we do not have a comprehensive understanding of the steady state pool size of products and reactants typically found in cells (Table 2). For stoichiometric considerations, we compare the energetic requirements for the synthesis of 6 TAG molecules containing fatty acids with the following chain

Table 1 Protein, carbohydrate and lipid fractions in several algal species

Algal species	Protein	Carbohydrates	Lipids
<i>Anabaena cylindrica</i> [96]	43–56	25–30	4–7
<i>Aphanizomenon flos-aquae</i> [96]	62	23	3
<i>Chlamydomonas reinhardtii</i> [96]	48	17	21
<i>Chlorella minutissima</i> [21]	9–24	14–42	31–57
<i>Auxenochlorella protothecoides</i> [97]	11–26	17–24	25–54
<i>Auxenochlorella protothecoides</i> [21]	36–38	41–52	11–23
<i>Chlorella pyrenoidosa</i> [96]	57	26	2
<i>Chlorella emersonii</i> [21]	28–32	11–41	29–63
<i>Chlorella sorokiniana</i> [21]	42–45	32–38	20–22
<i>Chlorella vulgaris</i> [96]	51–58	12–17	14–22
<i>Chlorella vulgaris</i> [21]	7–29	51–55	18–40
<i>Dunaliella salina</i> [96]	57	32	6
<i>Euglena gracilis</i> [96]	39–61	14–18	14–20
<i>Nannochloropsis</i> sp. [98]	23–59	5–17	9–62
<i>Porphyridium cruentum</i> [96]	28–39	40–57	9–14
<i>Scenedesmus obliquus</i> [96]	50–56	10–17	12–14
<i>Spirogyra</i> sp. [96]	6–20	33–64	11–21
<i>Arthrospira maxima</i> [96]	60–71	13–16	6–7
<i>Spirulina platensis</i> [96]	46–63	8–14	4–9
<i>Synechococcus</i> sp. [96]	63	15	11

Presented as a percentage of total dry weight. Related references are given in brackets.

lengths and degree of saturation, C16:0, C18:1, and C18:3, and having the molecular formula $C_{55}H_{98}O_6$. We compare these energetic requirements to the energetic required to synthesize starch (55 glucose) having the same number of carbon atoms (330) as 6 TAGs. We begin the energy accounting with the active import of inorganic carbon by HLA3 followed by the synthesis of triose-phosphate produced in the Calvin-Benson Cycle. Triose-phosphate is the substrate for fatty acid and TAG synthesis as well as glucose and starch synthesis.

Overall, the synthesis of a typical TAG (C16:0, C18:1 and C18:3; $C_{55}H_{98}O_6$) includes the energy required for acetyl CoA synthesis, fatty acid synthesis, fatty acid desaturation, and glycerol synthesis. As summarized in Table 2, TAG ($C_{55}H_{98}O_6$) synthesis requires 6.3 ATP/carbon and 2.9 NAD(P)H/carbon.

Glucose is derived from the condensation of two molecules of 3-phosphoglyceraldehyde. Activation of glucose units for starch polymerization requires the consumption of one additional ATP for the synthesis of ADP glucose. Overall, however, the energy required for storage carbohydrate synthesis is dominated by the energy required for carbon reduction by the Calvin-Benson Cycle. In total, the energy required for starch synthesis is 4.2 ATP/carbon and 2 NAD(P)H/carbon. Thus on a per

Table 2 Energetic (ATP and NAD(P)H) requirements for the synthesis of TAG and storage carbohydrate per unit carbon

	ATP/NAD(P)H
TAG synthesis partial reactions:	
3 CO ₂ → 3-phosphoglyceraldehyde→ acetylCoA	12 ATP, 4 NAD(P)H + CO ₂
AcetylCoA _(n-1used) mediated fatty acid elongation (e.g., C16:0)	7 ATP + 14 NADH
Fatty acid desaturation	1 NADH equivalent/bond desaturated
3 CO ₂ → Glycerol	9 ATP + 7 NAD(P)H
TAG (C₅₅H₉₈O₆) synthesis summary:	
26 acetyl CoA =	312 ATP, 104 NAD(P)H
Fatty acid elongation C16:0 + 2 C18:0 =	23 ATP, 46 NAD(P)H
4 desaturations	4 NAD(P)H equivalents
1 glycerol	9 ATP, 7 NAD(P)H
Total	344 ATP, 160 NAD(P)H
Per carbon	6.25 ATP, 2.93 NAD(P)H
Starch /glycogen synthesis partial reactions:	
6 CO ₂ → 2 3-phosphoglyceraldehyde → glucose	24 ATP, 12 NAD(P)H
Glucose → storage carbohydrate	1 ATP/glucose
Starch synthesis (55 units, 330 C) summary:	
Total for 55 glucose	1375 ATP, 660 NAD(P)H
Per Carbon	4.16 ATP, 2 NAD(P)H

carbon basis, the ATP and NAD(P)H requirements for starch synthesis are 50% and 45% lower, respectively than those required for TAG synthesis. If we assume the free energy of ATP hydrolysis and NADH oxidation (with molecular oxygen) are -50kJ/mole and -220 kJ/mole, respectively, then the energy cost per carbon for TAG synthesis (993 kJ/C) is 53% greater than for storage carbohydrate (650 kJ/C) synthesis under standard conditions. These energetic values are compromised, however, by limited knowledge of the *in vivo* concentrations of substrates and products and the energy of hydration.

Cells metabolize lipids and carbohydrates both for carbon skeletons to build other molecules as well as for energy metabolism. The efficiency of energy recovery from lipid and storage carbohydrate metabolism has been well studied in plants and animals. When saturated fatty acids are oxidized via the β -oxidation pathway and the citric acid cycle, the energy recovery is approximately 6.6 ATP equivalents/C. For glucose oxidation via glycolysis and the citric acid cycle the energy recovery is 5 ATP equivalents/carbon [99]. Thus, the energy recovery from fatty acid (lipid) oxidation is 32% greater than glucose (storage carbohydrate)

oxidation. Similar results are obtained if we directly combust TAGs or storage carbohydrate to produce energy (Table 3). Since TAGs have an average energy density (heat of combustion) of 38 kJ/g and storage carbohydrate has an average energy density of 15.5 kJ/g, then for an equivalent number of carbons, the energy content per carbon of TAG is 41% greater than that for starch (Table 3).

Interestingly, the relative biological energy input/carbon for TAG synthesis versus starch synthesis is substantially greater (53%) than the relative energy return from TAGs versus starch obtained either from respiration (32% greater for TAG than storage carbohydrate) or from direct combustion (41% greater for TAG than storage carbohydrate). Much of the increased energy costs for TAG versus storage carbohydrate synthesis can be attributed to the loss of reduced carbon that occurs during the conversion of pyruvate to acetyl CoA. Twenty six reduced carbon equivalents are lost during the decarboxylation of sufficient pyruvate to synthesize a typical TAG (C₅₅H₉₈O₆) molecule. Overall, these results beg the question: Does it make sense to engineer algae with enhanced oil levels versus engineering algae with high carbohydrate levels if the EROI for TAGs is 10-20% lower than that for starch? If the biofuel feedstock is to be used at a refinery to make diesel, Jet Propellant 8 (jet fuel JP8) or gasoline, however, then oil has a lower downstream energy cost to produce reduced fuels than does glucose due to reduced hydrogen requirements.

Biological implications of storage carbohydrate and oil accumulation in microalgae

What is missing from this discourse on the energetics and kinetics of starch and oil accumulation in algae is a greater understanding of overall cellular metabolism, flux rates, and cellular compartmentalization. Parameters that can impact the yields of starch and oil metabolism include; the kinetics, levels, and allosteric regulation (including feedback control) of the enzymes involved in

Table 3 Relative energy content (heat of combustion)/carbon for TAG and starch

	Energy
Mass for an equivalent number of moles of carbon (330) in TAG and starch	
6 moles TAG (C ₅₅ H ₉₈ O ₆) =	5124 g
55 moles glucose in starch (C ₆ H ₁₂ O ₆ , monomer) =	8928 g
Energy content of an equivalent number of moles of carbon (330) in TAG and starch	
330 C = 5124 g TAG x 38 kJ/g TAG =	194,712 kJ/330C = 590 kJ/C
330 C = 8928 g starch x 15.5 kJ/g storage carbohydrate =	138,384 kJ/330C = 419 kJ/C

their biosynthesis as well as the channeling of substrates and products through these and competing pathways. For example, starch accumulation in plastids can exert a feedback inhibition on photosynthesis, and so adequate sink strength is often necessary to achieve the greatest impact from improvements in photosynthetic rates [79,100,101]. Similarly, it has recently been shown that the substrates for TAG synthesis, acyl CoAs, can regulate the early dedicated events in fatty acid synthesis including acetyl CoA carboxylase activity [102]. There is also the role that these energy storage molecules play as substrates for the synthesis of other biologically important molecules.

Summary

It is argued that kinetic and thermodynamic constraints that determine the biomass or energy yield of biological systems are currently very poorly characterized. We have described kinetic constraints within the photosynthetic electron transfer apparatus and Calvin-Benson cycle that impact the efficiency of light utilization. Some of these constraints include; the coupling of proton to electron transfer associated with plastoquinone reduction and oxidation, the slow turnover numbers of Rubisco, and limiting concentrations of enzymes or effective enzyme catalytic turnover numbers. In some cases, these constraints can be overcome by increases in the numbers of catalytic sites or enzyme concentrations. In other cases, evolution has not provided an apparent resolution to the slow kinetics. This opens opportunities for metabolic engineering strategies to increase photosynthetic efficiency. For example, increasing the plastoquinone pool size could buffer transients in fast PSII and slow Cytb₆f electron transfer rates. Alternatively, increasing the activity or levels of the cytochrome b₆f complexes may accelerate electron transfer rates. In addition, the importance of optimizing Rubisco efficiency by concentrating CO₂ at the active site requires greater cyclic photophosphorylation activity to produce sufficient ATP for active bicarbonate uptake.

Lastly, by comparing the energy requirements for TAG and starch synthesis, we find that storage carbohydrates such as starch are energetically cheaper to make than TAG on a per carbon basis. This differential energy cost is largely attributed to the loss of fixed carbon occurring during the decarboxylation of pyruvate to synthesize acetyl CoA, an obvious target for metabolic engineering. Moreover, the higher energy density of TAG does not compensate for the energy investment. Thus, storage carbohydrate accumulation may be an effective strategy for efficient chemical energy accumulation in algae.

Abbreviations

ACCase: Acetyl CoA carboxylase; AGP: Adenosyl glucose phosphotransferase; CA: Carbonic anhydrase; CCM: Carbon concentrating mechanism; DGAT: Diacylglycerol acyltransferase; DOE: US Department of Energy;

EROI: Energy return on investment; FAS: Fatty acid synthase complex; Fd: Ferredoxin; FBpase: Fructose biphosphatase; FROI: Financial return on investment; G3PDH: Glycerol-3-phosphate dehydrogenase; GPAT: Glycerol-3-phosphate acyltransferase; GBSS: Granular bound starch synthase; HPI: Hexose phosphate isomerase; JP8: Jet propellant 8 jet fuel; LHC: Light harvesting complex; LPAT: Lysophosphatidate acyltransferase; NAABB: National Alliance for Advanced Biofuels and Bioproducts (DOE algal biomass program); PAP: Phosphatidic acid phosphatase; PAR: Photosynthetically active radiation (400–700 nm); PC: Plastocyanin; PQ: Plastoquinone; PSI: Photosystem I; PSII: Photosystem II; pSFBA: Plastidial Sedulose/Fructose-Bisphosphatase Aldolase; RC: Reaction centers of photosystem I or II; SBE: Starch branching enzyme; TAG: Triacylglycerol.

Competing interests

Dr. Richard Sayre is Chief Technology Officer for Phycal Inc.

Authors' contributions

All authors contributed equally to this publication. More specifically, SS contributed to writing sections on the energetics and kinetics of photosynthesis, organizing references and Figure 4. AB contributed to writing the introduction and summary, organizing the draft and references and Table 1 and Figure 2. SP contributed to writing on storage carbohydrate and oil metabolism and Figure 3. RS contributed to overall organization and thematic focus, energy calculations, Tables 2 and 3, Figure 1, and final editing. All authors read and approved the final manuscript.

Authors' information

Dr. Sowmya Subramanian is an associate research scientist at the New Mexico Consortium working on the metabolic engineering of electron transfer processes and carbon fixation to enhance photosynthetic efficiency. Dr. Amanda N. Barry is a postdoc at LANL working on engineering algal bioflocculation and assessment of algal biomass productivity using controlled photobioreactors. Dr. Shayani Pieris is an algal biotechnologist formerly at the New Mexico Consortium working on the enhancing of oil accumulation and photosynthesis in transgenic algae. Dr. Richard Sayre has a joint appointment with the New Mexico Consortium and Los Alamos National Laboratory and is Senior Research Scientist and Scientific Director for the Center for Advanced Biofuel Systems and the National Alliance for Advanced Biofuels and Bioproducts.

Acknowledgements

Sowmya Subramanian was supported by US-DOE Center for Advanced Biofuels contract, DE-SC0001295. Amanda N. Barry was supported by the Los Alamos Laboratory LDRD program. Shayani Pieris was supported by the US Air Force Office of Scientific Research, contract FA9550-08-1-0451. Richard Sayre was supported by US DOE National Alliance for Advanced Biofuels and Bioproducts, contract, DE-EE0003046, and by the US-DOE-EFRC, Center for Advanced Biofuels, contract, DE-SC0001295.

Author details

¹New Mexico Consortium, 100 Entrada Rd., Los Alamos, NM 87544, USA. ²Bioscience Division, Los Alamos National Lab, M888, Los Alamos, NM 87545, USA. ³Natural Sciences Division, Missouri Baptist University, One College Park Drive, St. Louis, MO 63141, USA.

Received: 26 April 2013 Accepted: 12 September 2013
Published: 19 October 2013

References

1. Khan SA, Rashmi J, Hussain MZ, Prasad S, Banerjee UC: **Prospects of biodiesel production from microalgae in India.** *Renew Sustain Energy Rev* 2009, **13**(9):2361–2372.
2. Sayre R: **Microalgae: the potential for carbon capture.** *Bioscience* 2010, **60**(9):722–727.
3. National Research Council: *Sustainable Development of Algal Biofuels in the United States.* Washington, DC: The National Academies Press; 2012.
4. Bondioli P, Della Bella L, Rivolta G, Chini Zittelli G, Bassi N, Rodolfi L, Casini D, Prussi M, Chiamonti D, Tedrici MR: **Oil production by the marine**

- microalgae *Nannochloropsis* sp. F&M-M24 and *Tetraselmis suecica* F&M-M33. *Bioresource technol* 2012, **114**:567–572.
5. Mata TM, Martins AA, Caetano NS: **Microalgae for biodiesel production and other applications: a review.** *Renew Sustain Energy Rev* 2010, **14**(1):217–232.
6. Rodolfi L, Chini Zittelli G, Bassi N, Padovani G, Biondi N, Bonini G, Tredici MR: **Microalgae for oil: strain selection, induction of lipid synthesis and outdoor mass cultivation in a low-cost photobioreactor.** *Biotechnol Bioeng* 2009, **102**(1):100–112.
7. Xu H, Miao X, Wu Q: **High quality biodiesel production from a microalga *Chlorella protothecoides* by heterotrophic growth in fermenters.** *J Biotechnol* 2006, **126**(4):499–507.
8. Chisti Y: **Biodiesel from microalgae beats bioethanol.** *Trends Biotechnol* 2008, **26**(3):126–131.
9. Wigmosta MS, Coleman AM, Skaggs RJ, Huesemann MH, Lane LJ: **National microalgae biofuel production potential and resource demand.** *Water Resour Res* 2011, **47**(3):W00H04.
10. Christenson L, Sims R: **Production and harvesting of microalgae for wastewater treatment, biofuels, and bioproducts.** *Biotechnol Adv* 2011, **29**(6):686–702.
11. Broekhuizen N, Park JBK, McBride GB, Craggs RJ: **Modification, calibration and verification of the IWA river water quality model to simulate a pilot-scale high rate algal pond.** *Water Res* 2012, **46**(9):2911–2926.
12. Richardson JW, Johnson MD, Outlaw JL: **Economic comparison of open pond raceways to photo bio-reactors for profitable production of algae for transportation fuels in the Southwest.** *Algal Res* 2012, **1**(1):93–100.
13. Liu J, Huang J, Sun Z, Zhong Y, Jiang Y, Chen F: **Differential lipid and fatty acid profiles of photoautotrophic and heterotrophic *Chlorella zofingiensis*: assessment of algal oils for biodiesel production.** *Bioresource Technol* 2011, **102**(1):106–110.
14. Chen YH, Walker TH: **Biomass and lipid production of heterotrophic microalgae *Chlorella protothecoides* by using biodiesel-derived crude glycerol.** *Biotechnol Lett* 2011, **33**(10):1973–1983.
15. Heredia-Arroyo T, Wei W, Hu B: **Oil accumulation via heterotrophic/mixotrophic *Chlorella protothecoides*.** *Appl Biochem Biotechnol* 2010, **162**(7):1978–1995.
16. O'Grady J, Morgan J: **Heterotrophic growth and lipid production of *Chlorella protothecoides* on glycerol.** *Bioproc Biosyst Eng* 2011, **34**(1):121–125.
17. Kosa M, Ragauskas AJ: **Lipids from heterotrophic microbes: advances in metabolism research.** *Trends Biotechnol* 2011, **29**(2):53–61.
18. Chen CY, Yeh KL, Aisyah R, Lee DJ, Chang JS: **Cultivation, photobioreactor design and harvesting of microalgae for biodiesel production: a critical review.** *Bioresource Technol* 2011, **102**(1):71–81.
19. Batan L, Quinn J, Willson B, Bradley T: **Net energy and greenhouse gas emission evaluation of biodiesel derived from microalgae.** *Environ Sci Technol* 2010, **44**(20):7975–7980.
20. Campbell PK, Beer T, Batten D: **Life cycle assessment of biodiesel production from microalgae in ponds.** *Bioresource Technol* 2011, **102**(1):50–56.
21. Illman AM, Scragg AH, Shales SW: **Increase in *Chlorella* strains calorific values when grown in low nitrogen medium.** *Enzyme Microbial Technol* 2000, **27**(8):631–635.
22. Williams PJB, Laurens LML: **Microalgae as biodiesel & biomass feedstocks: review & analysis of the biochemistry, energetics & economics.** *Energy Environ Sci* 2010, **3**(5):554–590.
23. Sharma KK, Schuhmann H, Schenk PM: **High lipid induction in microalgae for biodiesel production.** *Energies* 2012, **5**(5):1532–1553.
24. Metting FB Jr: **Biodiversity and application of microalgae.** *J Ind Microbiol* 1996, **17**(5–6):477–489.
25. Spolaore P, Joannis-Cassan C, Duran E, Isambert A: **Commercial applications of microalgae.** *J Biosci Bioeng* 2006, **101**(2):87–96.
26. Cardol P, Forti G, Finazzi G: **Regulation of electron transport in microalgae.** *Biochim Biophys Acta - Bioenergetics* 2011, **1807**(8):912–918.
27. Radakovits R, Jinkerson RE, Darzins A, Posewitz MC: **Genetic engineering of algae for enhanced biofuel production.** *Eukaryot Cell* 2010, **9**(4):486–501.
28. Work VH, D'Adamo S, Radakovits R, Jinkerson RE, Posewitz MC: **Improving photosynthesis and metabolic networks for the competitive production of phototroph-derived biofuels.** *Curr Opin Biotechnol* 2012, **23**(3):290–297.
29. Gokhale Z, Sayre RT: **Photosystem II, A Structural Perspective.** In *The Chlamydomonas Sourcebook*. Secondth edition. Edited by Stern DB. San Diego: Academic Press; 2009.
30. Croce R, Muller MG, Bassi R, Holzwarth AR: **Carotenoid-to-chlorophyll energy transfer in recombinant major light-harvesting complex (LHCII) of higher plants. I. Femtosecond transient absorption measurements.** *Biophys J* 2001, **80**(2):901–915.
31. Ishizaki A, Fleming GR: **Quantum coherence in photosynthetic light harvesting.** *Ann Rev Cond Matter Phys* 2012, **3**(1):333–361.
32. Merkli M, Berman G, Sayre R: **Electron transfer reactions: generalized spin-boson approach.** *J Math Chem* 2013, **51**:890–913.
33. Perrine Z, Negi S, Sayre RT: **Optimization of photosynthetic light energy utilization by microalgae.** *Algal Res* 2012, **1**:134–142.
34. Lewis KLM, Ogilvie JP: **Probing photosynthetic energy and charge transfer with Two-dimensional electronic spectroscopy.** *J Phys Chem Lett* 2012, **3**(4):503–510.
35. Pawlowicz NP, Groot ML, van Stokkum IH, Breton J, van Grondelle R: **Charge separation and energy transfer in the photosystem II core complex studied by femtosecond midinfrared spectroscopy.** *Biophys J* 2007, **93**(8):2732–2742.
36. Stiehl HH, Witt HT: **Quantitative treatment of the function of plastoquinone in photosynthesis.** *Z Naturforsch B* 1969, **24**(12):1588–1598.
37. Kirchhoff H, Mukherjee U, Galla HJ: **Molecular architecture of the thylakoid membrane: lipid diffusion space for plastoquinone.** *Biochem* 2002, **41**(15):4872–4882.
38. Harris GC, Königer M: **The 'high' concentrations of enzymes within the chloroplast.** *Photosynth Res* 1997, **54**(1):5–23.
39. Govindjee G, Kern JF, Messinger J, Whitmarsh J: *Photosystem II*. Hoboken, NJ: John Wiley & Sons, Ltd; 2001.
40. Johnson X, Alric J: **Interaction between starch breakdown, acetate assimilation, and photosynthetic cyclic electron flow in *Chlamydomonas reinhardtii*.** *JBiol Chem* 2012, **287**(31):26445–26452.
41. Junesch U, Gräber P: **The rate of ATP synthesis as a function of ΔpH in normal and dithiothreitol-modified chloroplasts.** *BBA - Bioenerg* 1985, **809**(3):429–434.
42. Etzold C, Deckers-Hebestreit G, Altendorf K: **Turnover number of *Escherichia coli* FOF1 ATP synthase for ATP synthesis in membrane vesicles.** *Eur J Biochem / FEBS* 1997, **243**(1–2):336–343.
43. Kramer DM, Avenson TJ, Edwards GE: **Dynamic flexibility in the light reactions of photosynthesis governed by both electron and proton transfer reactions.** *Trends Plant Sci* 2004, **9**(7):349–357.
44. Yamori W, Takahashi S, Makino A, Price GD, Badger MR, von Caemmerer S: **The roles of ATP synthase and the cytochrome b6/f complexes in limiting chloroplast electron transport and determining photosynthetic capacity.** *Plant Physiol* 2011, **155**(2):956–962.
45. Baker NR, Harbinson J, Kramer DM: **Determining the limitations and regulation of photosynthetic energy transduction in leaves.** *Plant Cell Environ* 2007, **30**(9):1107–1125.
46. Kanazawa A, Kramer DM: **In vivo modulation of nonphotochemical exciton quenching (NPQ) by regulation of the chloroplast ATP synthase.** *Proc Natl Acad Sci U S A* 2002, **99**(20):12789–12794.
47. Owens TG, Webb SP, Mets L, Alberte RS, Fleming GR: **Antenna size dependence of fluorescence decay in the core antenna of photosystem I: estimates of charge separation and energy transfer rates.** *Proc Natl Acad Sci USA* 1987, **84**(6):1532–1536.
48. Ley AC, Mauzerall DC: **Absolute absorption cross-sections for photosystem II and the minimum quantum requirement for photosynthesis in *Chlorella vulgaris*.** *BBA - Bioenerg* 1982, **680**(1):95–106.
49. Muller P, Li XP, Niyogi KK: **Non-photochemical quenching. A response to excess light energy.** *Plant Physiol* 2001, **125**(4):1558–1566.
50. Melis A: **Solar energy conversion efficiencies in photosynthesis: minimizing the chlorophyll antenna to maximize efficiency.** *Plant Sci* 2009, **177**(4):272–280.
51. Björkman O, Demmig B: **Photon yield of O₂ evolution and chlorophyll fluorescence characteristics at 77 K among vascular plants of diverse origins.** *Planta* 1987, **170**(4):489–504.
52. Long SP, Postl WF, Bolhár-Nordenkamp HR: **Quantum yields for uptake of carbon dioxide in C3 vascular plants of contrasting habitats and taxonomic groupings.** *Planta* 1993, **189**(2):226–234.
53. Mussgnug JH, Thomas-Hall S, Rupprecht J, Foo A, Klassen V, McDowall A, Schenk PM, Kruse O, Hankamer B: **Engineering photosynthetic light capture: impacts on improved solar energy to biomass conversion.** *Plant Biotechnol J* 2007, **5**(6):802–814.

54. Polle J, Kanakagiri S-D, Melis A: **tla1**, a DNA insertional transformant of the green alga *Chlamydomonas reinhardtii* with a truncated light-harvesting chlorophyll antenna size. *Planta* 2003, **217**(1):49–59.
55. Spreitzer RJ, Salvucci ME: RUBISCO: structure, regulatory interactions, and possibilities for a better enzyme. *Ann Rev Plant Biol* 2002, **53**(1):449–475.
56. Wingler A, Lea PJ, Quick WP, Leegood RC: Photorespiration: metabolic pathways and their role in stress protection. *Philos Trans R Soc Lond B Biol Sci* 2000, **355**(1402):1517–1529.
57. Moroney JV, Ynalvez RA: Proposed carbon dioxide concentrating mechanism in *Chlamydomonas reinhardtii*. *Eukaryot Cell* 2007, **6**(8):1251–1259.
58. Spalding MH: Microalgal carbon-dioxide-concentrating mechanisms: *Chlamydomonas* inorganic carbon transporters. *J Exp Bot* 2008, **59**(7):1463–1473.
59. Duanmu D, Miller AR, Horken KM, Weeks DP, Spalding MH: Knockdown of limiting-CO₂-induced gene HLA3 decreases HCO₃⁻ transport and photosynthetic C_i affinity in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* 2009, **106**(14):5990–5995.
60. Cardol P, Alric J, Girard-Bascou J, Franck F, Wollman F-A, Finazzi G: Impaired respiration discloses the physiological significance of state transitions in *Chlamydomonas*. *Proc Natl Acad Sci USA* 2009, **106**(37):15979–15984.
61. Miyagawa Y, Tamoi M, Shigeoka S: Overexpression of a cyanobacterial fructose-1,6-sedoheptulose-1,7-bisphosphatase in tobacco enhances photosynthesis and growth. *Nature Biotechnol* 2001, **19**(10):965–969.
62. Schwender J, Ohlrogge J, Shachar-Hill Y: Understanding flux in plant metabolic networks. *Curr Opin Plant Biol* 2004, **7**(3):309–317.
63. Szczotka M, Heise R, Tohge T, Nunes-Nesi A, Vosloh D, Huege J, Feil R, Lunn J, Nikoloski Z, Stitt M, *et al*: Metabolic fluxes in an illuminated *Arabidopsis rosette*. *Plant Cell* 2013, **25**(2):694–714.
64. Zhu XG, Long SP, Ort DR: Improving photosynthetic efficiency for greater yield. *Ann Rev Plant Biol* 2010, **61**:235–261.
65. Hohmann-Marriott MF, Blankenship RE: Evolution of photosynthesis. *Ann Rev Plant Biol* 2011, **62**:515–548.
66. Larkum AWD: Light Harvesting Systems in Algae. In *Photosynthesis in Algae*. Edited by Larkum AWD, Douglas SE, Raven JA. Netherlands: Kluwer Academic Publishers; 2003:277–282.
67. Zouni A, Witt HT, Kern J, Fromme P, Krauss N, Saenger W, Orth P: Crystal structure of photosystem II from *Synechococcus elongatus* at 3.8 Å resolution. *Nature* 2001, **409**(6821):739–743.
68. Kamiya N, Shen JR: Crystal structure of oxygen-evolving photosystem II from *Thermosynechococcus vulcanus* at 3.7-Å resolution. *Proc Natl Acad Sci USA* 2003, **100**(1):98–103.
69. Liu Z, Yan H, Wang K, Kuang T, Zhang J, Gui L, An X, Chang W: Crystal structure of spinach major light-harvesting complex at 2.72 Å resolution. *Nature* 2004, **428**(6980):287–292.
70. Ben-Shem A, Frolov F, Nelson N: Crystal structure of plant photosystem I. *Nature* 2003, **426**(6967):630–635.
71. Green BR, Durnford DG: The chlorophyll-carotenoid proteins of oxygenic photosynthesis. *Ann Rev Plant Physiol Plant Mol Biol* 1996, **47**:685–714.
72. Niyogi KK: Photoprotection revisited: genetic and molecular approaches. *Ann Rev Plant Physiol Plant Mol Biol* 1999, **50**:333–359.
73. Aro EM, Virgin I, Andersson B: Photoinhibition of photosystem II. Inactivation, protein damage and turnover. *Biochim Biophys Acta* 1993, **1143**(2):113–134.
74. Baker N: Photoinhibition of Photosynthesis. In *Light as an Energy Source and Information Carrier in Plant Physiology*, Volume 287. Edited by Jennings R, Zucchini G, Ghetti F, Colombetti G. New York, NY: Springer US; 1996:89–97.
75. Crimi M, Dorra D, Bosinger CS, Giuffra E, Holzwarth AR, Bassi R: Time-resolved fluorescence analysis of the recombinant photosystem II antenna complex CP29. Effects of zeaxanthin, pH and phosphorylation. *Eur J Biochem / FEBS* 2001, **268**(2):260–267.
76. Holt NE, Zigmantas D, Valkunas L, Li XP, Niyogi KK, Fleming GR: Carotenoid cation formation and the regulation of photosynthetic light harvesting. *Science* 2005, **307**(5708):433–436.
77. Avenso TJ, Ahn TK, Zigmantas D, Niyogi KK, Li Z, Ballottari M, Bassi R, Fleming GR: Zeaxanthin radical cation formation in minor light-harvesting complexes of higher plant antenna. *J Biol Chem* 2008, **283**(6):3550–3558.
78. Wiederrecht GP, Seibert M, Govindjee, Wasielewski MR: Femtosecond photodichroism studies of isolated photosystem II reaction centers. *Proc Natl Acad Sci USA* 1994, **91**(19):8999–9003.
79. Hillier W, Babcock GT: Photosynthetic reaction centers. *Plant Physiol* 2001, **125**(1):33–37.
80. Golbeck JH: *Photosystem I: The Light-Driven Plastocyanin: Ferredoxin Oxidoreductase*. Dordrecht: Springer; 2006.
81. Emerson R: The quantum yield of photosynthesis. *Annu Rev Plant Physiol* 1958, **9**(1):1–24.
82. Govindjee: On the requirement of minimum number of four versus eight quanta of light for the evolution of one molecule of oxygen in photosynthesis: a historical note. *Photosynth Res* 1999, **59**(2–3):249–254.
83. Skillman JB: Quantum yield variation across the three pathways of photosynthesis: not yet out of the dark. *J Exp Bot* 2008, **59**(7):1647–1661.
84. Hogewoning SW, Wientjes E, Douwstra P, Trouwborst G, van Ieperen W, Croce R, Harbinson J: Photosynthetic quantum yield dynamics: from photosystems to leaves. *Plant Cell* 2012, **24**(5):1921–1935.
85. Singaas EL, Ort DR, DeLucia EH: Variation in measured values of photosynthetic quantum yield in ecophysiological studies. *Oecologia* 2001, **128**:15–23.
86. Van Walraven HS, Strotmann H, Schwarz O, Rumberg B: The H⁺/ATP coupling ratio of the ATP synthase from thiol-modulated chloroplasts and two cyanobacterial strains is four. *FEBS Lett* 1996, **379**(3):309–313.
87. Turina P, Samoray D, Graber P: H⁺/ATP ratio of proton transport-coupled ATP synthesis and hydrolysis catalysed by CF₀F₁-liposomes. *EMBO J* 2003, **22**(3):418–426.
88. Steigmiller S, Turina P, Graber P: The thermodynamic H⁺/ATP ratios of the H⁺—ATP synthases from chloroplasts and *Escherichia coli*. *Proc Natl Acad Sci U S A* 2008, **105**(10):3745–3750.
89. Seelert H, Poetsch A, Dencher NA, Engel A, Stahlberg H, Muller DJ: Structural biology. Proton-powered turbine of a plant motor. *Nature* 2000, **405**(6785):418–419.
90. Vollmar M, Schlieper D, Winn M, Buchner C, Groth G: Structure of the c14 rotor ring of the proton translocating chloroplast ATP synthase. *J Biol Chem* 2009, **284**(27):18228–18235.
91. Sacksteder CA, Kanazawa A, Jacoby ME, Kramer DM: The proton to electron stoichiometry of steady-state photosynthesis in living plants: a proton-pumping Q cycle is continuously engaged. *Proc Natl Acad Sci USA* 2000, **97**(26):14283–14288.
92. Jia H, Oguchi R, Hope AB, Barber J, Chow WS: Differential effects of severe water stress on linear and cyclic electron fluxes through photosystem I in spinach leaf discs in CO₂-enriched air. *Planta* 2008, **228**(5):803–812.
93. Baker NR, Ort DR: Light and crop photosynthetic performance. In *Crop Photosynthesis: Spatial and Temporal Determinants*. Edited by Baker NR, Thomas H. Amsterdam: Elsevier Science Publishers; 1992:289–312.
94. Spalding MH: Photosynthesis and photorespiration in freshwater green algae. *Aquat Bot* 1989, **34**(1–3):181–209.
95. Birmingham BC, Coleman JR, Colman B: Measurement of photorespiration in algae. *Plant Physiol* 1982, **69**(1):259–262.
96. Becker EW: Micro-algae as a source of protein. *Biotechnol Adv* 2007, **25**(2):207–210.
97. Xiong W, Liu L, Wu C, Yang C, Wu Q: 13C-tracer and gas chromatography-mass spectrometry analyses reveal metabolic flux distribution in the oleaginous microalga *Chlorella protothecoides*. *Plant Physiol* 2010, **154**(2):1001–1011.
98. Hu H, Gao K: Response of growth and fatty acid compositions of *Nannochloropsis* sp. to environmental factors under elevated CO₂ concentration. *Biotechnology Lett* 2006, **28**(13):987–992.
99. Buchanan B, Gruissem W, Jones RL: *Biochemistry and Molecular Biology of Plants*. Rockville, MD: American Society of Plant Biologists; 2000.
100. Wunsche JN, Greer DH, Laing WA, Palmer JW: Physiological and biochemical leaf and tree responses to crop load in apple. *Tree Physiol* 2005, **25**(10):1253–1263.
101. Ihemere U, Arias-Garzon D, Lawrence S, Sayre R: Genetic modification of cassava for enhanced starch production. *Plant Biotechnol J* 2006, **4**(4):453–465.
102. Andre C, Haslam RP, Shanklin J: Feedback regulation of plastidic acetyl-CoA carboxylase by 18:1-acyl carrier protein in *Brassica napus*. *Proc Natl Acad Sci USA* 2012, **109**(25):10107–10112.

doi:10.1186/1754-6834-6-150

Cite this article as: Subramanian *et al.*: Comparative energetics and kinetics of autotrophic lipid and starch metabolism in chlorophytic microalgae: implications for biomass and biofuel production. *Biotechnology for Biofuels* 2013 **6**:150.