

RESEARCH ARTICLE

Open Access

Characterisation of *bla*_{TEM} genes and types of β -lactamase plasmids in *Neisseria gonorrhoeae* – the prevalent and conserved *bla*_{TEM-135} has not recently evolved and existed in the Toronto plasmid from the origin

Ibrahim Muhammad¹, Daniel Golparian¹, Jo-Anne R Dillon², Åsa Johansson³, Makoto Ohnishi⁴, Sunil Sethi⁵, Shao-chun Chen⁶, Shu-ichi Nakayama⁴, Martin Sundqvist¹, Manju Bala⁷ and Magnus Unemo^{1*}

Abstract

Background: Antimicrobial resistance (AMR) in *Neisseria gonorrhoeae* is a major concern worldwide. It has been recently feared that the *bla*_{TEM-1} gene is, via *bla*_{TEM-135}, evolving into an extended-spectrum β -lactamase (ESBL), which could degrade all cephalosporins including ceftriaxone. The aims of the present study were to characterize the *bla*_{TEM} genes, types of β -lactamase plasmids, the degradation of ampicillin by TEM-135 compared to TEM-1, and to perform molecular epidemiological typing of β -lactamase-producing *N. gonorrhoeae* strains internationally.

Methods: β -lactamase producing *N. gonorrhoeae* isolates (n = 139) cultured from 2000 to 2011 in 15 countries were examined using antibiograms, *bla*_{TEM} gene sequencing, β -lactamase plasmid typing, and *N. gonorrhoeae* multiantigen sequence typing (NG-MAST). Furthermore, the *bla*_{TEM} gene was sequenced in the first described Toronto plasmid (pJD7), one of the first Asian plasmids (pJD4) and African plasmids (pJD5) isolated in Canada. The degradation of ampicillin by TEM-135 compared to TEM-1 was examined using a MALDI-TOF MS hydrolysis assay.

Results: Six different *bla*_{TEM} sequences were identified (among isolates with 125 different NG-MAST STs), i.e. *bla*_{TEM-1} (in 104 isolates), *bla*_{TEM-135} (in 30 isolates), and four novel *bla*_{TEM} sequences (in 5 isolates). The *bla*_{TEM-1} allele was only found in the African and Asian plasmids, while all Rio/Toronto plasmids possessed the *bla*_{TEM-135} allele. Most interesting, the first described gonococcal Toronto plasmid (pJD7), identified in 1984, also possessed the highly conserved *bla*_{TEM-135} allele. The degradation of ampicillin by TEM-135 compared to TEM-1 was indistinguishable in the MALDI-TOF MS hydrolysis assay.

(Continued on next page)

* Correspondence: magnus.unemo@orebroll.se

¹Department of Laboratory Medicine, WHO Collaborating Centre for Gonorrhoea and other Sexually Transmitted Infections, National Reference Laboratory for Pathogenic Neisseria, Microbiology, Örebro University Hospital, SE-701 85, Örebro, Sweden

Full list of author information is available at the end of the article

(Continued from previous page)

Conclusions: *bla*_{TEM-135}, encoding TEM-135, is predominantly and originally associated with the Rio/Toronto plasmid and prevalent among the β -lactamase producing gonococcal strains circulating globally. *bla*_{TEM-135} does not appear, as previously hypothesized, to have recently evolved due to some evolutionary selective pressure, for example, by the extensive use of extended-spectrum cephalosporins worldwide. On the contrary, the present study shows that *bla*_{TEM-135} existed in the Toronto plasmid from its discovery and that *bla*_{TEM-135} is highly conserved (not further evolved in the past >30 years). Nevertheless, international studies for monitoring the presence of different *bla*_{TEM} alleles, the possible evolution of the *bla*_{TEM-135} allele, and the types of β -lactamase producing plasmids, remain imperative.

Keywords: Gonorrhoea, Antimicrobial resistance, *bla*_{TEM-1}, *bla*_{TEM-135}, TEM-1, TEM-135, Rio/Toronto plasmid, Extended-spectrum β -lactamase (ESBL)

Background

Gonorrhoea is the most prevalent bacterial sexually transmitted infection (STI) globally, according to the latest estimates by the World Health Organization (WHO) [1]. No vaccine is available and, accordingly, appropriate prevention, and particularly effective diagnosis and antimicrobial treatment are the cornerstones for control of gonorrhoea. *Neisseria gonorrhoeae* has developed resistance to all antimicrobials previously recommended for first-line empiric monotherapy, such as penicillins, tetracyclines, macrolides and fluoroquinolones [2-5]. Clinical resistance to the extended-spectrum cephalosporins (ESCs) has also been verified in the latest years in many countries [6-19]. Furthermore, the first three extensively-drug resistant (XDR [3]) gonococcal strains were recently described; all displayed a high-level of resistance to ceftriaxone, the last remaining option for empiric first-line antimicrobial monotherapy in most countries globally [9,18,20]. All the reported decreased susceptibility and resistance to ESCs has been due to an accumulation of chromosomal resistance determinants [2,4,5].

In 1976 the first β -lactamase producing *N. gonorrhoeae* strains, resulting in high-level resistance to penicillins but not affecting the MICs of ESCs, were reported [21,22]. These strains produced the traditional TEM-1 β -lactamase that hydrolyses the cyclic amide bond in the β -lactam ring [23]. β -lactamase producing gonococcal strains are currently widespread internationally [2,4]. The *bla*_{TEM} genes are located on a family of related β -lactamase plasmids, of which the most frequently described have been the Asian, African, and the mainly indistinguishable Rio and Toronto plasmids (named based on their epidemiological origin) [4,24-26]. However, other types of β -lactamase producing plasmids have also been described in gonococci, e.g. Nimes, New Zealand, Australian and Johannesburg [4,26-28]. The Asian plasmid has been considered to be the ancestral plasmid from which the other plasmids evolved through deletions and/or insertions. Accordingly, these β -lactamase producing plasmids may be characterised as either deletion derivatives of the Asian plasmid (Africa,

Rio/Toronto and Johannesburg) or insertion derivatives of either the Asian (New Zealand) or African (Nimes) plasmids [4,24-27].

Worryingly, the *bla*_{TEM-1} gene, encoding the TEM-1 β -lactamase, needs only a few specific single nucleotide polymorphisms (SNPs) to evolve into a gene encoding an extended-spectrum β -lactamase (ESBL), which could degrade all ESCs including ceftriaxone [29-32]. The spread of a potent ESBL, which also degrades ceftriaxone, in the gonococcal population might rapidly result in untreatable gonorrhoea in most settings worldwide. It has been hypothesized that the gonococcal *bla*_{TEM-1} has recently evolved into *bla*_{TEM-135}, which originally was identified in *Salmonella enterica* subsp. *enterica* serovar Typhimurium [33] and might be a precursor in the evolution into an ESBL gene [29,34,35]. TEM-135 producing gonococcal isolates have been described in 2004 and 2008 in Japan [29], from 2005 to 2007 in Thailand [34,36], and in 2007 and 2012 in China [35]. TEM-135, which only differs from TEM-1 by one SNP (T→C at position 539) resulting in the amino acid alteration M182T, requires solely one additional specific SNP to evolve into an ESBL such as TEM-20 [29-32,34]. In those ESBLs, the M182T alteration presumably stabilizes the active site topology reorganized by other mutations, which collaboratively results in the emergence of a stable ESBL [34,35,37]. However, there have been few reports, and none outside Asia, regarding surveillance of the different types of *bla*_{TEM} genes and β -lactamase producing plasmids in the gonococcal strains circulating worldwide.

The aims of this study were to characterize the *bla*_{TEM} genes, the types of β -lactamase plasmids, the degradation of ampicillin by TEM-135 compared to TEM-1, and to perform molecular epidemiological typing of β -lactamase-producing *N. gonorrhoeae* isolates cultured in 2000–2011 in 15 countries.

Methods

β -lactamase producing *N. gonorrhoeae* isolates

In total, 139 β -lactamase producing *N. gonorrhoeae* isolates were examined, including 136 *N. gonorrhoeae* clinical

isolates (collected from 2000 to 2011 in 15 WHO European (n = 40), African (n = 22), American (North and Latin America) (n = 10), Southeast Asian (n = 33) or Western Pacific (n = 31) countries) and three of the 2008 WHO *N. gonorrhoeae* reference strains, i.e. WHO M (isolated in the Philippines, 1992), WHO N (Australia, 2001), and WHO O (Canada, 1991) [38]. All isolates were cultured on selective culture media for 16–18 h at 37°C, in a 5% CO₂-enriched atmosphere. *N. gonorrhoeae* isolates were identified using characteristic colony morphology, Gram staining, positive oxidase test, a rapid carbohydrate utilization test, and the Phadebact GC Monoclonal Test (Boule Diagnostics AB, Huddinge, Sweden). All examined gonococcal isolates were cultured and stored as part of the routine diagnostics (standard care) and no patient identification information was used. Furthermore, to elucidate when *bla*_{TEM-135} emerged DNA from three previously published strains was also investigated. Those represented the first Toronto plasmid described (pJD7 [39,40], identified in 1984), one of the first Asian plasmids (pJD4 [39–41], identified in the late-1970s) and one of the first African plasmids (pJD5 [39–41], identified in the late-1970s) isolated in Canada.

Antimicrobial susceptibility testing

MICs (mg/L) of the isolates for ampicillin, cefixime, ceftriaxone, azithromycin, ciprofloxacin, and spectinomycin were determined using the Etest method (AB bioMérieux, Solna, Sweden), according to the instructions from the manufacturer. Breakpoints for susceptibility, intermediate susceptibility and resistance in accordance to the European Committee on Antimicrobial Susceptibility Testing (EUCAST; www.eucast.org) were used. β -lactamase production was identified using nitrocefin solution (Oxoid, Basingstoke, Hants, England). The 2008 *N. gonorrhoeae* WHO reference strains [38] were used for quality control in all antimicrobial susceptibility testing.

β -lactamase hydrolysis assay

To examine the degradation of ampicillin by TEM-135 compared to TEM-1, a selection of six β -lactamase producing gonococcal isolates (three *bla*_{TEM-135} and three *bla*_{TEM-1} isolates) were investigated using a hydrolysis assay. The non- β -lactamase producing *N. gonorrhoeae* reference strain WHO F [38] was included as negative control. All isolates were cultured on New York City agar media plates and incubated for 16–18 h at 37°C in a 5% CO₂-enriched atmosphere. The hydrolysis assay was performed as previously described [42] on a Microflex (Bruker Daltonics, GmbH, Germany), by recording spectra in the mass range of 0–1000 Da after incubation for 15 minutes, 1 h, 2 h and 3 h. For calibration, the Peptide calibration standard II (Bruker Daltonics, GmbH, Germany) was used. The peaks utilized for calibration was

CCA [M+H]⁺ at 190.05 Da, CCA [2 M+H]⁺ at 379.09 Da and Bradykinin (1–7) peak [M+H]⁺ at 757.40 Da. Generated spectra were manually examined using the Flex Analysis 3.1 software (Bruker Daltonics, GmbH, Germany), and peaks correlating to hydrolyzed or intact ampicillin [42] were identified.

Plasmid DNA extraction

The QIAGEN Spin Miniprep Kit (QIAGEN, Hilden, Germany) was used to extract and purify plasmid DNA, according to the instructions from the manufacturer. DNA was stored at 4°C prior to subsequent analysis.

Genomic DNA extraction

Genomic DNA was extracted using the robotized NorDiag Bullet (NorDiag ASA, Oslo, Norway) and the BUGS'n BEADS STI-fast kit (NorDiag ASA, Oslo, Norway), according to the instructions from the manufacturer. DNA was stored at 4°C prior to subsequent analysis.

Plasmid typing

Multiplex PCR was performed for β -lactamase plasmid typing on all isolates as previously described using the primers BL1, BL2, BL3, and BL4 [25].

Sequencing of the *bla*_{TEM} gene

The entire coding region, including the signal peptide of 23 amino acids (GenBank accession number AAR25033), of *bla*_{TEM} was PCR amplified in a LightCycler real-time PCR System (Roche Molecular Biochemicals, Mannheim, Germany) and subsequently sequenced as previously described [29]. Multiple-sequence alignments of nucleotide and amino acid sequences were performed using the BioEdit (version 5.0.9) software. For comparison and numbering of the amino acid positions, all amino acid sequences identified in the present study were compared to sequences at the Lactamase Engineering Database (<http://www.laced.uni-stuttgart.de/>) as well as at the β -Lactamase Classification and Amino Acid Sequences for TEM, SHV and OXA Extended-Spectrum and Inhibitor Resistant Enzymes database (<http://www.lahey.org/Studies/>). The scheme proposed by Ambler et al. [43] was used for numbering of amino acids.

Molecular epidemiological characterisation using *Neisseria gonorrhoeae* multiantigen sequence typing (NG-MAST)

Amplification and sequencing of the more variable segments of the *porB* and *thpB* genes examined in NG-MAST [44,45] was performed as previously described [46]. For assignment of *porB* and *thpB* allele numbers as well as NG-MAST STs, the NG-MAST website (<http://www.ng-mast.net/>) was used.

Results

Antimicrobial susceptibility of β -lactamase producing *N. gonorrhoeae* isolates (n = 139) from 2000 to 2011

The majority (83.5%, n = 116/139) of isolates were resistant to ampicillin and the remaining 16.5% (n = 23) had an intermediate susceptibility to ampicillin. Furthermore, 80.6% (n = 112) and 6.5% (n = 9) of the isolates were resistant to ciprofloxacin and azithromycin, respectively. Only one isolate (0.7%) was resistant to cefixime (MIC = 0.25 mg/L) and no isolates were resistant to ceftriaxone or spectinomycin.

β -lactamase plasmid types and bla_{TEM} alleles

The African-type β -lactamase plasmid was the most common (67.6% [94/139] of isolates), followed by the Rio/Toronto-type plasmid (18.7% [26/139] of isolates) and the Asian-type plasmid in 13.7% (19/139) of isolates (Table 1). No other β -lactamase plasmid types were found.

The amino acid sequences of all gonococcal TEM sequences identified in the present study are displayed in Figure 1. Briefly, one-hundred and four isolates (74.8%) possessed the bla_{TEM-1} allele, which was identical to the bla_{TEM-1} allele in *E. coli* (GenBank accession number AAR25033), and 30 (21.6%) of isolates had an identical $bla_{TEM-135}$ allele. Interestingly, all (n = 26) Rio/Toronto plasmids contained the $bla_{TEM-135}$ allele. However, the African plasmid and Asian plasmid in one (1.1%) and three (15.8%) isolates also contained the $bla_{TEM-135}$ allele. Furthermore, four novel amino acid substitutions in TEM were identified, that is, alterations in the signal peptide (P14T (n = 1) and P14S (n = 2)), and in the TEM coding sequence: E110K (n = 1) and G228S (n = 1), which is close to the substrate binding site. All these new bla_{TEM} alleles were possessed by African (n = 4) or Asian (n = 1) plasmids (Table 1).

Most interesting, the first described Toronto plasmid (pJD7 [39,40]), isolated in 1984, also possessed the $bla_{TEM-135}$ allele, while the Asian plasmid (pJD4 [39-41]) and African plasmid (pJD5 [39-41]), isolated in the late-1970s, had the bla_{TEM-1} allele.

The mean MIC of ampicillin in TEM-135 and TEM-1 producing isolates was 16 mg/L and 12 mg/L, respectively. Furthermore, the mean MIC of ceftriaxone in the TEM-135 and TEM-1 producing isolates was identical, that is, 0.016 mg/L. The isolates containing the novel bla_{TEM} alleles had also low MICs of ceftriaxone (MIC = 0.003-0.006 mg/L).

Degradation of ampicillin by TEM-135 and TEM-1

In the hydrolysis assay, the MALDI-TOF MS spectra showed that the degradation of ampicillin by TEM-135 and TEM-1 was indistinguishable and both β -lactamases degraded ampicillin in 15 minutes. For the non- β -lactamase producing *N. gonorrhoeae* reference strain WHO F [38], included as negative control, no hydrolysis was observed even after three hours of incubation (Additional file 1: Figure S1).

Genotyping using *N. gonorrhoeae* multiantigen sequence typing (NG-MAST)

In total, 125 different NG-MAST STs were identified, of which 87 had not been previously described. The most frequent STs were ST6058 (3.5% of isolates), ST6057 (2.9%), and ST1288 (2.2%). The majority of STs (93.6% of the STs) were represented by one isolate, whereas five ST were represented by two isolates. Due to the heterogeneity of the isolates and the many STs identified, it was not possible to correlate any β -lactamase plasmid type to any specific NG-MAST ST. Notably, the 30


Table 1 Type of bla_{TEM} allele, β -lactamase producing plasmid, number of NG-MAST STs, and year and WHO Region of isolation of *N. gonorrhoeae* isolates cultured in 2000–2011 in 15 countries

β -lactamase encoding gene (No. of isolates)	β -lactamase producing plasmid types (%)	Number different NG-MAST STs	Year (WHO Region) of isolation
bla_{TEM-1} (104)	African (85.6), Asian (14.4)	93	2000-2011 (All Regions ^a)
$bla_{TEM-135}$ (30)	Rio/Toronto (86.7), Asian (10), African (3.3)	29	2000-2009 (All Regions ^a)
$bla_{TEM-P14S^b}$ (2)	African (100)	2	2008 (European)
$bla_{TEM-P14T^b}$ (1)	Asian (100)	1	2003 (European)
$bla_{TEM-E110K}$ (1)	African (100)	1	2003 (African)
$bla_{TEM-G228S}$ (1)	African (100)	1	2009 (Western Pacific)
$bla_{TEM-135}^c$	Rio/Toronto		1984 (American)
bla_{TEM-1}^c	African		Late-1970s (American)
bla_{TEM-1}^c	Asian		Late-1970s (American)

^aWHO European, African, American (North and Latin America), Southeast Asian and Western Pacific Region.

^bAmino acid alteration in the 23 amino acids long signal peptide.

^cPreviously published strains possessing the first Toronto plasmid described (pJD7 [39,40]), one of the first Asian plasmids (pJD4 [39-41]) and African plasmids (pJD5 [39-41]) isolated in Canada.



gonococcal isolates possessing the *bla*_{TEM-135} allele were assigned to 29 different NG-MAST STs.

Discussion

The present study investigated the characteristics of *bla*_{TEM} genes and their association with β -lactamase plasmid type, the degradation of ampicillin by TEM-135 compared to TEM-1, and the molecular epidemiology of international β -lactamase-producing *N. gonorrhoeae* isolates. The isolates comprised a large collection of temporally (cultured from 2000 to 2011), geographically (from 15 WHO European, African, American (North and Latin America), Southeast Asian or Western Pacific countries) and genetically diverse (125 NG-MAST STs) gonococcal strains. Clearly, *N. gonorrhoeae* strains producing TEM-1 are widespread and all these strains carried the *bla*_{TEM-1} allele on an African or Asian plasmid. The highly conserved *bla*_{TEM-135} allele was predominantly found on the Rio/Toronto plasmids, i.e. only four of the 30 identified *bla*_{TEM-135} alleles were carried on African (n = 1) or Asian (n = 3) plasmids. In fact, all the identified Rio/Toronto plasmids in 26 genetically highly diverse isolates (26 different NG-MAST STs) contained a *bla*_{TEM-135} allele and according to our best knowledge *bla*_{TEM-1} has only been described in one Rio/Toronto plasmid ever [34]. These results strongly indicate that the Rio/Toronto-type plasmid is the origin of the *bla*_{TEM-135} allele. Interestingly, in the present study the *bla*_{TEM-135} allele was also identified on the first reported Toronto plasmid (pJD7 [39,40]), found in a gonococcal isolate from 1984. This shows that

the *bla*_{TEM-135} allele has not recently evolved and, on the contrary, was present in the Toronto plasmid from its discovery. This might also question the evolutionary origin of the Rio/Toronto plasmids, i.e. that these plasmids emerged directly through a deletion of 2273 bp in the Asian plasmid [25,27], in which the *bla*_{TEM-135} allele is rare internationally. Nevertheless, as shown in the present study the *bla*_{TEM-135} allele can also be found on African-type and Asian-type plasmids. Thus, it cannot be excluded that those *bla*_{TEM-135} alleles might have evolved from *bla*_{TEM-1} alleles by a SNP in those plasmids. However, considering how conserved the gonococcal *bla*_{TEM} alleles appear to be, a more plausible explanation might be that these *bla*_{TEM-135} alleles were acquired by horizontal transfer from strains possessing the *bla*_{TEM-135} allele on a Rio/Toronto plasmid. This hypothesis is further supported by the lack of antimicrobial selective pressure for the evolution of *bla*_{TEM-1} to *bla*_{TEM-135}. Accordingly, similar MICs of ampicillin and ceftriaxone, and of other antimicrobials, were displayed by the TEM-135 and TEM-1 producing isolates, and the two different TEM enzymes showed an indistinguishable degradation of ampicillin in the MALDI-TOF MS hydrolysis assay. Nevertheless, for detailed measurement of the kinetics of the ampicillin hydrolysis appropriate kinetic experiments (Kcat/Km) with a purified protein would be required. All the novel *bla*_{TEM} alleles found in the present study were carried by African (four strains) or Asian (one strain) plasmids, and none of these five isolates had any enhanced MIC of ceftriaxone.

Conclusions

*bla*_{TEM-135}, encoding TEM-135, is predominantly and originally associated with the Rio/Toronto plasmid and prevalent among the β -lactamase producing gonococcal strains circulating globally. *bla*_{TEM-135} does not appear, as previously hypothesized, to have recently evolved due to, for example, some evolutionary selective pressure by the extended-spectrum cephalosporins. On the contrary, *bla*_{TEM-135} existed in the Toronto plasmid from its discovery. The present study indicates that *bla*_{TEM-135} instead is highly conserved (not further evolved in the past >30 years). Accordingly, despite the extensive use of extended-spectrum cephalosporins globally no additional SNP has evolved in *bla*_{TEM-135}. The reasons for this remain unknown, however, both the lack of evolutionary positive selection for such SNPs and a decreased biological fitness of gonococcal strains with the resulting TEM alleles might be involved. Even so, only one specific SNP added to the *bla*_{TEM-135} allele would produce an ESBL (e.g. an additional G238S alteration to evolve into TEM-20), which would be able to degrade all extended-spectrum cephalosporins, possibly rendering gonorrhoea an untreatable infection. Accordingly, international studies for monitoring and enhancing our understanding of, in addition to the chromosomal ESC resistance determinants, the presence of different *bla*_{TEM} alleles, the possible evolution of the *bla*_{TEM-135} allele, and the β -lactamase producing plasmids, remain imperative.

Additional file

Additional file 1: Figure S1. β -lactamase hydrolysis assay. MALDI-TOF MS spectra of ampicillin after incubation with *Neisseria gonorrhoeae* (β -lactamase negative, TEM-1 producing and TEM-135 producing) and *Escherichia coli* ATCC 35218 (TEM-1 producing). Ampicillin alone (top) and *N. gonorrhoeae* incubated in water (second from top) are also shown in the figure. Intact ampicillin displayed the peaks of 350.4 Da, 372.4 Da and 394.4 Da while hydrolysed ampicillin displayed the peaks of 324.4 Da, 368.4 Da, 390.4 Da and 412.4 Da. The peak of 324.4 Da was observed with low intensity also in the top-spectrum indicating a slight spontaneous hydrolysis.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MO, SS, MB, JAD and MU designed, initiated and coordinated the study. IM, DG, AS, SC, SN, and MS coordinated and performed all the laboratory analyses. IM, DG and MU analysed and interpreted all the data, and wrote a first draft of the paper. All authors read, commented on and approved the final manuscript.

Acknowledgments

This study was supported by grants from the Research Committee of Örebro County and the Örebro University Hospital Research Foundation, Örebro, Sweden.

Author details

¹Department of Laboratory Medicine, WHO Collaborating Centre for Gonorrhoea and other Sexually Transmitted Infections, National Reference Laboratory for Pathogenic *Neisseria*, Microbiology, Örebro University Hospital,

SE-701 85, Örebro, Sweden. ²Department of Microbiology and Immunology, College of Medicine, University of Saskatchewan, Saskatoon, Saskatchewan, Canada. ³Department of Clinical Microbiology, Central Hospital, Växjö, Sweden. ⁴National Institute of Infectious Diseases, Tokyo, Japan. ⁵Department of Medical Microbiology, Post Graduate Institute of Medical Education and Research (PGIMER), Chandigarh, India. ⁶National Center for STD Control, Chinese CDC, Nanjing, China. ⁷Apex Regional STD Teaching, Training and Research Centre, VMMC and Safdarjang Hospital, New Delhi, India.

Received: 16 June 2014 Accepted: 18 August 2014

Published: 22 August 2014

References

- World Health Organization: *Global Incidence and Prevalence of Selected Curable Sexually Transmitted Infections* - 2008. Geneva: World Health Organization; 2012. Available at: http://www.who.int/reproductivehealth/publications/rtis/2008_STI_estimates.pdf; (Accessed: July 24, 2014).
- Lewis DA: **The gonococcus fights back: is this time a knock out?** *Sex Transm Infect* 2010, **86**:415–421.
- Tapsall JW, Ndowa F, Lewis DA, Unemo M: **Meeting the public health challenge of multidrug- and extensively drug-resistant *Neisseria gonorrhoeae*.** *Expert Rev Anti Infect Ther* 2009, **7**:821–834.
- Unemo M, Shafer WM: **Antimicrobial resistance in *Neisseria gonorrhoeae* in the 21st century: past, evolution and future.** *Clin Microbiol Rev* 2014, **27**:587–613.
- Unemo M, Nicholas RA: **Emergence of multi-drug resistant, extensively drug-resistant and untreatable gonorrhea.** *Future Microbiol* 2012, **7**:1401–1422.
- Deguchi T, Yasuda M, Yokoi S, Ishida K, Ito M, Ishihara S, Minamide T, Harada Y, Tei K, Kojima K, Tamaki M, Maeda S: **Treatment of uncomplicated gonococcal urethritis by double-dosing of 200 mg cefixime at a 6-h interval.** *J Infect Chemother* 2003, **9**:35–39.
- Yokoi S, Deguchi T, Ozawa T, Yasuda M, Ito S, Kubota Y, Tamaki M, Maeda S: **Threat to cefixime treatment for gonorrhea.** *Emerg Infect Dis* 2007, **13**:1275–1277.
- Ison CA, Hussey J, Sankar KN, Evans J, Alexander S: **Gonorrhoea treatment failures to cefixime and azithromycin in England, 2012.** *Euro Surveill* 2011, **16**(14):pii=19833.
- Unemo M, Golparian D, Nicholas R, Ohnishi M, Galloway A, Sednaoui P: **High-level cefixime- and ceftriaxone-resistant *Neisseria gonorrhoeae* in France: novel *penA* mosaic allele in a successful international clone causes treatment failure.** *Antimicrob Agents Chemother* 2012, **56**:1273–1280.
- Unemo M, Golparian D, Stry A, Eigentler A: **First *Neisseria gonorrhoeae* strain with resistance to cefixime causing gonorrhoea treatment failure in Austria, 2011.** *Euro Surveill* 2011, **16**(43):pii=19998.
- Unemo M, Golparian D, Syversen G, Vestheim DF, Moi H: **Two cases of verified clinical failures using internationally recommended first-line cefixime for gonorrhoea treatment, Norway, 2010.** *Euro Surveill* 2010, **15**(47):pii=19721.
- Lewis DA, Sriruttan C, Müller EE, Golparian D, Gumedde L, Fick D, de Wet J, Maseko V, Coetzee J, Unemo M: **Phenotypic and genetic characterization of the first two cases of extended-spectrum-cephalosporin-resistant *Neisseria gonorrhoeae* infection in South Africa and association with cefixime treatment failure.** *J Antimicrob Chemother* 2013, **68**:1267–1270.
- Allen VG, Mitterni L, Seah C, Rebbapragada A, Martin IE, Lee C, Siebert H, Towns L, Melano RG, Low DE: ***Neisseria gonorrhoeae* treatment failure and susceptibility to cefixime in Toronto, Canada.** *JAMA* 2013, **309**:163–170.
- Unemo M, Golparian D, Potočnik M, Jeverica S: **Treatment failure of pharyngeal gonorrhoea with internationally recommended first-line ceftriaxone verified in Slovenia, September 2011.** *Euro Surveill* 2012, **17**:1–4.
- Unemo M, Golparian D, Hestner A: **Ceftriaxone treatment failure of pharyngeal gonorrhoea verified by international recommendations, Sweden, July 2010.** *Euro Surveill* 2011, **16**:1–3.
- Tapsall J, Read P, Carmody C, Bourne C, Ray S, Limnios A, Sloots T, Whitley D: **Two cases of failed ceftriaxone treatment in pharyngeal gonorrhoea verified by molecular microbiological methods.** *J Med Microbiol* 2009, **58**:683–687.
- Chen YM, Stevens K, Tideman R, Zaia A, Tomita T, Fairley CK, Lahra M, Whitley D, Hogg G: **Failure of ceftriaxone 500 mg to eradicate pharyngeal gonorrhoea, Australia.** *J Antimicrob Chemother* 2013, **68**:1445–1447.
- Ohnishi M, Golparian D, Shimuta K, Saika T, Hoshina S, Iwasaku K, Nakayama S, Kitawaki J, Unemo M: **Is *Neisseria gonorrhoeae* initiating a future era of**

- untreatable gonorrhea?: Detailed characterization of the first strain with high-level resistance to ceftriaxone. *Antimicrob Agents Chemother* 2011, **55**:3538–3545.
19. Read PJ, Limnios EA, McNulty A, Whitley D, Lahra LM: One confirmed and one suspected case of pharyngeal gonorrhoea treatment failure following 500 mg ceftriaxone in Sydney, Australia. *Sex Health* 2013, **10**:460–462.
 20. Camara J, Serra J, Ayats J, Bastida T, Carnicer-Pont D, Andreu A, Ardanuy C: Molecular characterization of two high-level ceftriaxone-resistant *Neisseria gonorrhoeae* isolates detected in Catalonia, Spain. *J Antimicrob Chemother* 2012, **67**:1858–1860.
 21. Ashford WA, Golash RG, Hemming VG: Penicillinase-producing *Neisseria gonorrhoeae*. *Lancet* 1976, **2**:657–658.
 22. Phillips I: Beta-lactamase-producing, penicillin-resistant gonococcus. *Lancet* 1976, **2**:656–657.
 23. Bergström S, Norlander L, Norqvist A, Normark S: Contribution of a TEM-1-like beta-lactamase to penicillin resistance in *Neisseria gonorrhoeae*. *Antimicrob Agents Chemother* 1978, **13**:618–623.
 24. Pagotto F, Aman AT, Ng L-K, Yeung KH, Brett M, Dillon JA: Sequence analysis of the family of penicillinase-producing plasmids of *Neisseria gonorrhoeae* based on DNA sequencing. *Plasmid* 2000, **43**:24–34.
 25. Palmer HM, Leeming JP, Turner A: A multiplex polymerase chain reaction to differentiate β -lactamase plasmids of *Neisseria gonorrhoeae*. *J Antimicrob Chemother* 2000, **45**:777–782.
 26. Dillon JR, Yeung KH: β -lactamase plasmids and chromosomally mediated antibiotic resistance in pathogenic *Neisseria* species. *Clin Microbiol Rev* 1989, **2**:S125–S133.
 27. Müller EE, Fayemiwo SA, Lewis DA: Characterization of a novel β -lactamase-producing plasmid in *Neisseria gonorrhoeae*: sequence analysis and molecular typing of host gonococci. *J Antimicrob Chemother* 2011, **66**:1514–1517.
 28. Trembizki E, Buckley C, Lawrence A, Lahra M, Whitley D: On behalf of the GRAND study investigators: Characterisation of a novel *Neisseria gonorrhoeae* penicillinase-producing plasmid, Australia 2012. *Antimicrob Agents Chemother* 2014, **58**:4984–4985.
 29. Ohnishi M, Ono E, Shimuta K, Watanabe H, Okamura N: Identification of TEM-135 β -lactamase in penicillinase-producing *Neisseria gonorrhoeae* strains in Japan. *Antimicrob Agents Chemother* 2010, **54**:3021–3023.
 30. Arlet G, Goussard S, Courvalin P, Philippon A: Sequences of the genes for the TEM-20, TEM-21, TEM-22, and TEM-29 extended-spectrum β -lactamases. *Antimicrob Agents Chemother* 1999, **43**:969–971.
 31. Michael GB, Butaye P, Cloeckaert A, Schwarz S: Genes and mutations conferring antimicrobial resistance in *Salmonella*: An update. *Microb Infect* 2006, **8**:1898–1914.
 32. Huang W, Palzkill T: A natural polymorphism in beta-lactamase is a global suppressor. *Proc Natl Acad Sci U S A* 1997, **94**:8801–8806.
 33. Pasquali F, Kehrenberg C, Manfreda G, Schwarz S: Physical linkage of Tn3 and part of Tn1721 in a tetracycline and ampicillin resistance plasmid from *Salmonella Typhimurium*. *J Antimicrob Chemother* 2005, **55**:562–565.
 34. Nakayama S, Tribuddharat C, Prombhul S, Shimuta K, Srifueungfung S, Unemo M, Ohnishi M: Molecular analyses of TEM genes and their corresponding penicillinase-producing *Neisseria gonorrhoeae* isolates in Bangkok, Thailand. *Antimicrob Agents Chemother* 2012, **56**:916–920.
 35. Chen SC, Yin YP, Dai XQ, Yu RX, Han Y, Sun HH, Ohnishi M, Unemo M, Chen XS: Prevalence and molecular epidemiological typing of penicillinase-producing *Neisseria gonorrhoeae* and their bla(TEM-135) gene variants in Nanjing, China. *Sex Transm Dis* 2013, **40**:872–876.
 36. Srifueungfung S, Roongpisuthipong A, Asavapiriyonont S, Lolekha R, Tribuddharat C, Lokpichart S, Sungthong P, Tongtep P: Prevalence of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in HIV-seropositive patients and gonococcal antimicrobial susceptibility: an update in Thailand. *J Infect Dis* 2009, **62**:467–470.
 37. Orenica MC, Yoon JS, Ness JE, Stemmer WP, Stevens RC: Predicting the emergence of antibiotic resistance by directed evolution and structural analysis. *Nat Struct Biol* 2001, **8**:238–242.
 38. Unemo M, Fasth O, Fredlund H, Limnios A, Tapsall JW: Phenotypic and genetic characterization of the 2008 WHO *Neisseria gonorrhoeae* reference strain panel intended for global quality assurance and quality control of gonococcal antimicrobial resistance surveillance for public health purposes. *J Antimicrob Chemother* 2009, **63**:1142–1151.
 39. Yeung KH, Dillon JR, Pauzé M, Wallace E: A novel 4.9-kilobase plasmid associated with an outbreak of penicillinase-producing *Neisseria gonorrhoeae*. *J Infect Dis* 1986, **153**:1162–1165.
 40. Yeung KH, Dillon JR: In vitro transcription/translation products and molecular characterization of naturally occurring and in vitro deletion of the 7.2 kb plasmid of *Neisseria gonorrhoeae*. p. 209 215. In *The Pathogenic Neisseriae. Proceedings of the 4th international symposium, Asilomar, California, 21 25 October 1984*. Edited by Schoolnik GK. Washington, D.C: American Society for Microbiology (ASM); 1985.
 41. Pagotto F, Dillon JA: Multiple origins and replication proteins influence biological properties of beta-lactamase-producing plasmids from *Neisseria gonorrhoeae*. *J Bacteriol* 2001, **183**:5472–5481.
 42. Sparbier K, Schubert S, Weller U, Boogen C, Kostrzewa M: Matrix-assisted laser desorption ionization-time of flight mass spectrometry-based functional assay for rapid detection of resistance against β -lactam antibiotics. *J Clin Microbiol* 2012, **50**:927–937.
 43. Ambler RP, Coulson AF, Frère JM, Ghysen JM, Joris B, Forsman M, Levesque RC, Tiraby G, Waley SG: A standard numbering scheme for the class A beta-lactamases. *Biochem J* 1991, **276**:269–270.
 44. Martin IM, Ison CA, Aanensen DM, Fenton KA, Spratt BG: Rapid sequence-based identification of gonococcal transmission clusters in a large metropolitan area. *J Infect Dis* 2004, **189**:1497–1505.
 45. Unemo M, Dillon JA: Review and international recommendation of methods for typing *Neisseria gonorrhoeae* isolates and their implications for improved knowledge of gonococcal epidemiology, treatment, and biology. *Clin Microbiol Rev* 2011, **24**:447–458.
 46. Unemo M, Sjöstrand A, Akhras M, Gharizadeh B, Lindbäck E, Pourmand N, Wretling B, Fredlund H: Molecular characterization of *Neisseria gonorrhoeae* identifies transmission and resistance of one ciprofloxacin-resistant strain. *APMIS* 2007, **115**:231–241.

doi:10.1186/1471-2334-14-454

Cite this article as: Muhammad et al.: Characterisation of bla_{TEM} genes and types of β -lactamase plasmids in *Neisseria gonorrhoeae* – the prevalent and conserved bla_{TEM-135} has not recently evolved and existed in the Toronto plasmid from the origin. *BMC Infectious Diseases* 2014 **14**:454.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

