

STUDY PROTOCOL

Open Access

A randomized control trial to assess the impact of vitamin D supplementation compared to placebo on vascular stiffness in chronic kidney disease patients

Adeera Levin^{1*}, Taylor Perry², Prathibha De Zoysa², Mhairi K Sigrist³, Karin Humphries⁴, Mila Tang⁵ and Ognjenka Djurdjev²

Abstract

Background: Vitamin D deficiency is associated with cardiovascular (CV) risk in multiple populations, including those with chronic kidney disease (CKD). The active form of the hormone (1,25 OH₂D₃) binds to receptors in multiple organs. CKD patients are deficient in both 25 Vitamin D and 1,25 OH₂D₃. Clinical trial data demonstrating the benefits of vitamin D formulations are limited, and fail to show significant benefits on CV outcomes, and have compared different compounds, in various populations, and focused on a variety of outcomes. A understanding of the mechanism by which different vitamin D compounds confer CV protection in CKD is important for the design of future studies.

Methods/Design: This 3 arm randomized prospective double-blinded placebo-controlled study examining the impact of calcitriol (1,25 OH₂D₃) and 25-hydroxyvitamin D₃ supplementation compared to placebo on vascular stiffness, as measured by pulse wave velocity (PWV). Patients are enrolled from 2 tertiary care institutions if they meet inclusion criteria (stable estimated glomerular filtration rate (eGFR) between 15-45ml/min, <±5ml/min change in previous 6 months), on stable doses of renin-angiotensin aldosterone system blockade. For those already receiving vitamin D therapies, a 3 month washout period before randomization is mandatory. Treatment duration is 6 months; medications are given thrice weekly in fixed doses. The primary outcome measure is Vascular stiffness, measured non-invasively by pulse wave velocity (PWV). Other measurements include BP, kidney function and serial blood levels of biomarkers. The primary analysis will compare any vitamin D therapy versus placebo for the primary outcome defined as the change of PWV from baseline to 6 months. Analysis of covariance will be used to detect differences between vitamin D preparations in the magnitude of reduction in PWV.

Discussion: This study is novel in that we are using a robust study design in CKD patients (not on dialysis) comparing placebo to different forms of vitamin D supplementation in fixed doses, irrespective of baseline values. We hope to demonstrate the biological mechanistic effect of vitamin D supplementation on vascular function in order for this information to be used in designing larger randomized controlled trials.

Trial registration: Current Controlled Trials NCT01247311. Date of Registration: November 12, 2010.

Keywords: Chronic kidney disease, Vitamin D, Vascular stiffness, PWV, Randomized protocol study

* Correspondence: alevin@providencehealth.bc.ca

¹University of British Columbia, 1081 Burrard Street, Room 6010A, Vancouver, BC V6Z 1Y6, Canada

Full list of author information is available at the end of the article

Background

Cardiovascular (CV) disease in chronic kidney disease (CKD) is highly prevalent and is associated with a significant increase in morbidity and mortality [1]. Vascular dysfunction and CV disease are caused by a combination of atherosclerotic processes (mediated by dyslipidemia and smoking) and arteriosclerotic processes such as aging, diabetes, vitamin D deficiencies, hyperphosphatemia and hyperparathyroidism (HPTH). The latter is associated with vascular stiffness, loss of collagen and elastin fibers [2]. CKD patients have both 'conventional' and non-conventional risk factors for CV disease; and high prevalence of left ventricular hypertrophy (LVH) has been noted. To date, conventional CV disease risk reduction agents and strategies (renal angiotensin aldosterone system (RAAS) antagonists, anti-platelet agents, lipid lowering agents) have either not been systematically evaluated in all CKD populations, or do not appear to mitigate CV disease risk to the extent that they are effective in non-CKD populations. Furthermore, strategies to address non-conventional risk factors (anemia, hyperphosphatemia) on outcomes have either shown no impact, harm, conflicting results, or have not been evaluated in a rigorous randomized control trial with clinically meaningful endpoints [3].

Vitamin D deficiency, (25 OHD₃) is common in both CKD and general populations. It has been associated in animal and human studies with hypertension (HTN), diabetes, obesity and inflammation, and proteinuria. It is these associations that has led some to regard it as an important CV disease risk factor [4-6]. CKD patients have low levels of 25-hydroxyvitamin D (25 vitamin D), but also lack the physiological capacity to hydroxylate 25 vitamin D into its active form 1,25-dihydroxyvitamin D (1,25 vitamin D), which binds to the vitamin D receptors (VDR) found in blood vessels, the heart, muscles and kidneys. Animal studies describe the impact of VDR activators on inflammation in both kidneys and hearts. In addition, LVH has been noted, and reversed with supplementation [7]. [Human studies have demonstrated reduction in proteinuria [8,9] and some impact on changes in eGFR. These are small studies of limited duration. Thadhani et al. have found that treatment with paricalcitol (a VDR activator) in CKD patients reduces parathyroid hormone (PTH) levels within 4 weeks of commencing treatment versus a placebo [10], however, did not impact the primary outcome (LV mass), but did reduce the number of CV disease-related hospitalizations over the trial period. Large cohort studies have also demonstrated reductions in 1,25 vitamin D and 25 vitamin D in CKD populations in association with rises in serum PTH values [11], at eGFR levels commencing around 50mL/min/1.73m². While CKD patients may be able to hydroxylate 25 vitamin D outside the kidney, the

relative effectiveness of these mechanisms have not been studied [2]. The prevailing theory is that the combined 25 OHD₃ and 1,25 OH₂D₃ deficiencies seen in CKD do impact outcomes.

The association between 25 vitamin D deficiency and CV disease has been demonstrated in numerous populations [5,12]. The active hormone modulates a variety of processes directly associated with arteriosclerotic changes via its interactions with specific biomarkers and inflammatory processes [13-17]. As a recent Cochrane review asserts, no prospective randomized controlled study to date have demonstrated the effect of vitamin D supplementation to reduce CV disease events [18]. Given the multiple reported effects of vitamin D on PTH, phosphate (PO₄), blood pressure (BP), inflammation, insulin sensitivity and other factors, it is imperative to demonstrate a biological impact of vitamin D on vascular function, which could be directly in the causal pathway for CV disease [11,15,19,20].

There are non invasive methods of studying vascular stiffness, the most used of which is Pulse wave velocity (PWV) using standardized validated equipment (aSphygmoCor). This technique utilizes applanation tonometry to measure the velocity of the pulse wave between two points on the arterial tree [21]. Increased vascular stiffness (higher PWV values) is associated with LVH, poor CV disease outcomes, and death in dialysis populations [22,23]. The reproducibility and validity of this measurement has been demonstrated both in the general population and in those with kidney failure [22,24,25]. Utilizing PWV measurements, we will attempt to demonstrate changes in vascular stiffness with various forms of vitamin D supplementation in a non dialysis CKD population.

Methods/Design

Overarching objectives

The overarching objective of this study is to describe the impact of currently used formulations of vitamin D (25 vitamin D₃ and calcitriol) on vascular stiffness in patients with CKD, eGFR levels between 15 and 45 mL/min/1.73m². We hypothesize that supplementation with vitamin D analogues will result in greater reductions in PWV measurements than placebo when compared to baseline measurements.

Study cohort

Subjects will be drawn from a cohort of CKD patients treated according to best practices at university based tertiary care centers in Vancouver. Following written informed consent, subjects who fulfill eligibility criteria at screening (Table 1) will be randomized (1:1:1) to receive treatment of 25 vitamin D₃ at 5000IU three times a week, 1,25 vitamin D (calcitriol) at 0.5ug three times a week, or placebo.

Table 1 Inclusion/exclusion criteria

Inclusion criteria	Exclusion criteria
- Signed informed consent	- Refused informed consent
- eGFR between 15-45 ml/min	- Change of >5ml/min eGFR over the past 6 months
- Change of <5ml/min in eGFR over the past 6 months	- Planned transplant within 6 months
- Treated with maximal conventional CV disease risk reduction protocolized medications (ACEi or ARB)	- Likely to commence renal replacement therapy within 6 months after enrolment
- Currently receiving vitamin D therapies, but agreeing to a washout period of 3 months	- Terminal malignancies
	- Active infections
	- Active inflammatory diseases (SLE, vasculitis)

Funding sources

The study is funded from a peer reviewed grant from Kidney foundation of Canada (KFoC) in collaboration with Pfizer. The study addresses specific interests of the KFoC, focusing on Canadians with CKD at risk for CVD, and identifying potential benefits of adjunct therapy in a complex condition.

Statement of purpose

This physiological study will describe the impact of vitamin D therapies on a physiological measure related to CVD in CKD: vascular stiffness. Describing this mechanism would enhance the viability of testing longer term outcomes using these interventions.

Primary questions:

1. Does treatment with vitamin D analogues for a period of 6 months reduce PWV in CKD patients as compared to placebo?

Secondary questions:

1. Is there a difference between 1,25 vitamin D and 25 vitamin D formulations in the magnitude of reduction in PWV?
2. Does 6 month provision of 1,25 or 25 vitamin D lead to demonstrable reductions in BP (without changes in antihypertensive medication), or proteinuria (without change in ACEi/ARB therapy) relative to placebo?
3. What is the impact of 1,25 vitamin D and 25 vitamin D on blood levels of FGF-23, serum PTH, PO₄ Ca, and CRP in this population? Is there a relationship between changes in vascular stiffness and changes in these biochemical/hormonal parameters?

Rationale for the study

There is increasing evidence that vitamin D acting on multiple aspects of the physiological processes involved in the pathogenesis of CVD may be an important therapeutic strategy in CKD. The intellectual appeal of studying vitamin D supplementation in CKD has led to a number of proposals for studies of CV outcome in CKD, and some randomized control trials [3,10,26,27]. While these studies have shown promising signals (reduction in proteinuria and reduction in CV hospitalization and events), no studies have addressed the mechanism of how vitamin D supplementation might impact these outcomes in humans, other than via reduction in LV mass (which was not shown). While both 1,25 vitamin D and 25 vitamin D formulations have been used in clinical studies to date, controversy continues as to the form of vitamin D best used in clinical practice to supplement deficiencies [16,28,29].

Recent meta-analyses and systematic reviews [15,17, 26,30-32] have described epidemiological associations with 25 vitamin D deficiencies and outcomes, and supplementation as impacting intermediate outcomes (proteinuria), but not mortality or time to end stage kidney failure. Quantification of the impact of vitamin D on vascular stiffness will permit estimates of effect size and stimulate further research into combining medication strategies.

Interventions

We will prescribe fixed doses of 5000IU 25 vitamin D and 0.5µg of 1,25 vitamin D, three times a week, (compatible with doses given in other studies, and at intervals used in many clinical practices) in order to maximally stimulate the circadian impact of the vitamin D therapy. A 6 month treatment period is chosen to ensure "biologically meaningful exposure" which should permit measurable changes in blood levels and vascular compliance, as well as mitigate the effects of sun exposure. Medications are supplied in oral suspension with identical look and taste (in almond oil), so that patients and providers are blinded.

Duration of the study

Each patient will be followed for a period of 6 months post-randomization. Patients on existing vitamin D supplementation will have been followed for 9 months due to the additional 3 months of washout prior to randomization.

Study design

This will be a 3 arm prospective randomized double blind placebo controlled study of 128 stable CKD subjects examining the impact of vitamin D supplementation (1,25 vitamin D or 25 vitamin D formulations) compared to placebo on arterial stiffness and other parameters of vascular health. The study plan shown in Figure 1 will combine physiological experiments in a well characterized cohort




Figure 1 The study plan depicts the timeline for the completion of the study. Screening/recruitment includes the 3 month washout period. Vitamin D doses will be administered for 6 months allowing for the inclusion criteria. PWV will be measured at baseline (-3 months), post washout period (0 months) and post treatment (6 months) for both treatment arms and their respective placebos.

of patients with CKD, and utilize observational study methods to test and explore additional related hypotheses.

Sample size considerations

The study is powered to demonstrate differences in primary outcome between individual study subjects; the calculated sample size will be adequate to detect a clinically important effect of vitamin D on PWV (reduction of 1m/sec). The mean difference of 1m/s is clinically meaningful in that it has been demonstrated to confer a 34% increased risk of CV mortality in a cohort of CKD stage 5 patients on dialysis [33,34]. A sample size of 105 (35 placebo arm and 70 treatment arm) without attrition will provide 85% power to demonstrate the mean PWV difference of 1m/s between arms, with 0.05 target alpha for a two sided test. Thus, recruitment and enrolment of 128 subjects will allow for ~15% attrition.

Trial management

To ensure safety and data integrity, the single research office centrally located is responsible for the conduct at two sites within the city, both University teaching hospitals. The study will be overseen by independent Data Safety and Monitoring Board (DSMB) and also will review subject lab results and concomitant medications on a monthly basis. The DSMB will screen for the adverse events, including the possibility of hypervitaminosis, hypercalcaemia and hyperphosphatemia. In addition, random audits will be performed to assess data quality.

Variables measured

PWV measurements, clinical and biological samples will be obtained at pre-washout, baseline, and post treatment (3 time points). Blood and urine samples will be obtained at each of the 3 time points to test for serum PTH, 1,25 vitamin D, 25 vitamin D, fibroblast growth factor-23 (FGF-23), PO_4 , calcium (Ca), C-reactive protein (CRP), urinalysis and random urine albumin-to-creatinine ratio (UACR). All samples will be run in an accredited

laboratory using validated assays. BP will be done in duplicate and PWV will be done in triplicate at each visit using standardized BP cuffs and SphygmoCor, respectively [35]. Clinical and medication histories as well as rationale for specific biomarkers are listed in Additional file 1.

Statistical analysis

The primary analysis will compare vitamin D (25 vitamin and 1,25 vitamin D) therapies versus placebo for the primary outcome defined as the change of PWV from baseline to 6 months by covariance analysis (ANCOVA); where the 6 month measurement of PWV will be the dependent variable, allocated treatment the independent variable, and baseline PWV the baseline covariate [29,36,37]. Analyses will be considered statistically significant if the p-value is <0.05 for allocated treatment. In case of violation of ANCOVA assumptions we will use a mixed model approach.

Strengths and weaknesses

It is conceivable that the therapies as administered may not have an impact on our chosen outcomes of interest: vascular stiffness, BP or proteinuria (primary or secondary). The 6 month time period may not be sufficient exposure to demonstrate the effect of the vitamin D treatment, or the expected difference in PWV changes/reductions between the groups is different and thus the study may be underpowered. However, given that this study uniquely examines non dialysis advance stage CKD patients, using fixed dose drug administration schedules, physiological measurements and biochemical measurements serially over time, the study will add to the growing body of literature regarding the use of vitamin D supplementation in CKD population. If positive, it will inform future larger studies looking at additional measures such as LV mass, or function over time in relation to PWV, and will add to our understanding of the physiology of vascular changes in CKD. Given the current state of knowledge,

and the plethora of studies, reviews, and meta-analysis that fail to consistently demonstrate a benefit to Vitamin D supplementation, there are no ethical issues conducting a placebo-controlled trial.

Discussion

There is an accumulating body of evidence that vitamin D acting on multiple aspects of physiological processes involved in the pathogenesis of CVD and thus may be an important therapeutic strategy in CKD. Studies conducted to date have failed to demonstrate significant impact of hormone supplementation on outcomes. Cholecalciferol has been studied in 60 patients on hemodialysis, who were known to be 25 OHD₃ deficient and did not show any change in PWV [38]. Other studies conducted in both dialysis and non dialysis CKD patients, also included were those with demonstrable deficiencies for limited time periods (8 weeks) showed changes in serum levels of 25 and 1, 25 Vitamin D levels [39-42]. Although more pronounced in non dialysis patients, other outcomes were not measured. Another study demonstrated that 25 OHD₃ supplementation in post menopausal women did not reduce CV risk, when studied in a randomized control trial [43]. These studies were all relatively short term and focused on different questions than the ones we are attempting to answer in this study. We believe that attempting to establish a biological mechanistic effect of vitamin D supplementation on vascular function is necessary before further larger randomized controlled trials are undertaken. Given the purported differential ability of CKD patients to achieve appropriate concentrations of the active compound, the question of which supplementation is more appropriate, 25 vitamin D or 1,25 vitamin D in CKD remains controversial. The relationship of serum levels to disease, type of therapeutic supplementation (fixed dose or based on targeting serum levels) is unknown in CKD populations. Moreover, in various nephrology studies, attempts at targeting deficiencies or excesses of measured serum values in CKD patients in large randomized control trials has not met with success [44-47]. Thus, we have developed and designed a robust randomized control trial to attempt to address deficiencies in the literature, and to better understand mechanisms and complex biological interactions in this patient group.

We expect to find vitamin D to have an impact on vascular stiffness and BP irrespective of serum vitamin D levels. The impact will be greatest on those with highest PWV. The data generated herein will permit the development of a set of phenotypes using clinical and biomarker data. Such that characterization of patients according to PWV, BP, FGF-23 and vitamin D levels will be possible, and results of treatment strategies will be examined within the phenotype category. This information will be significant for future enrolment in clinical trials. The biomarker

analyses will aid in explaining any dissociation found between BP and vascular stiffness.

Before large randomized trials can be developed, these unanswered questions regarding the mechanism of disease require resolution to ensure pertinent study design. This is an 'intermediate' study which fully examines the physiological consequences of different vitamin D therapies in CKD populations. The findings will be used to understand physiological mechanisms, generate further hypotheses for testing, and ultimately contribute to evidence base studies which will inform clinical practice.

Ethics

The study has been approved by the University of British Columbia and Providence Health Care Research Ethics Committee (Registration number: H10-01689).

Additional file

Additional file 1: Timetable of study visits.

Abbreviations

CV: Cardiovascular; CKD: Chronic kidney disease; BP: Blood pressure; eGFR: Glomerular filtration rate; PWV: Pulse wave velocity; HPTH: Hyperparathyroidism; HTN: Hypertension; PTH: Parathyroid hormone; PO4: Phosphate; LVH: Left ventricular hypertrophy; DSMB: Data safety and monitoring board; FGF-23: Fibroblast growth factor-23; Ca: Calcium; CRP: C-reactive protein; UACR: Urine albumin-to-creatinine ratio; ANCOVA: Analysis of covariance.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

This article was written with large contributions from the original grant application and protocol. The concept, rationale, design logistics and analyses were determined by the investigators. All analyses will be conducted by OD, and all study materials remain property of the investigators. The sponsors (KFOC and Pfizer) have no role aside from funding the study. AL, MKS and KH developed and wrote the protocol; AL was the key author on the manuscript, provided constructive criticism and edits of the drafts of the manuscripts, but all authors contributed to the final product. All are- or have been- responsible for the integrity and running of the study from inception to current state. All authors read and approved the final manuscript.

Acknowledgements

We thank all of the patients and their physicians who are participating in this study. We gratefully acknowledge the financial support obtained from KFOC.

Contributions of investigators and coordinators

We thank all the vitamin D investigators and coordinators for data and sample collections: Lina Siosin, Jeffrey Gardner, Matthew Paquette, Rossitta Yung, Naama Rozen and Nancy Ferguson.

Author details

¹University of British Columbia, 1081 Burrard Street, Room 6010A, Vancouver, BC V6Z 1Y6, Canada. ²Providence Health Care Research Institute (PHCRI), St. Paul's Hospital, 1081 Burrard Street, Room 302, Vancouver, BC V6Z 1Y6, Canada. ³Division of Nephrology Research - University of British Columbia/ Providence Health Care, Vancouver, BC V6Z 1Y6, Canada. ⁴Providence Health Care (PHC), Division of Cardiology, Vancouver, BC V6Z 1Y6, Canada. ⁵BC Provincial Renal Agency (BCPRA), 700-1380 Burrard Street, Vancouver, BC V6Z 2H3, Canada.

Received: 6 March 2014 Accepted: 27 October 2014
Published: 7 November 2014

References

- Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY: **Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization.** *N Engl J Med* 2004, **351**(13):1296–1305. Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer.
- Levin A, Le Barbier M, Er L, Andress D, Sigrist MK, Djurdjev O: **Incident isolated 1,25(OH)(2)D(3) deficiency is more common than 25(OH)D deficiency in CKD.** *J Nephrol* 2012, **25**(2):204–210.
- Jun M, Lv J, Perkovic V, Jardine MJ: **Managing cardiovascular risk in people with chronic kidney disease: a review of the evidence from randomized controlled trials.** *Ther Adv Chronic Dis* 2011, **2**(4):265–278.
- Johal M, Levin A: **Vitamin D and parathyroid hormone in general populations: understandings in 2009 and applications to chronic kidney disease.** *Clin J Am Soc Nephrol* 2009, **4**(9):1508–1514.
- Holick MF: **Vitamin D deficiency.** *N Engl J Med* 2007, **357**(3):266–281.
- Autier P, Gandini S: **Vitamin D supplementation and total mortality: a meta-analysis of randomized controlled trials.** *Arch Intern Med* 2007, **167**(16):1730–1737.
- Bodyak N, Ayus JC, Achinger S, Shivalingappa V, Ke Q, Chen YS, Rigor DL, Stillman I, Tamez H, Kroeger PE, Wu-Wong RR, Karumanchi SA, Thadhani R, Kang PM: **Activated vitamin D attenuates left ventricular abnormalities induced by dietary sodium in Dahl salt-sensitive animals.** *Proc Natl Acad Sci U S A* 2007, **104**(43):16810–16815.
- Manson JE, Bassuk SS, Lee IM, Cook NR, Albert MA, Gordon D, Zaharris E, Macfadyen JG, Danielson E, Lin J, Zhang SM, Buring JE: **The Vitamin D and Omega-3 Trial (VITAL): rationale and design of a large randomized controlled trial of vitamin D and marine omega-3 fatty acid supplements for the primary prevention of cancer and cardiovascular disease.** *Contemp Clin Trials* 2012, **33**(1):159–171.
- de Zeeuw D, Agarwal R, Amdahl M, Audhya P, Coyne D, Garimella T, Parving HH, Pritchett Y, Remuzzi G, Ritz E, Andress D: **Selective vitamin D receptor activation with paricalcitol for reduction of albuminuria in patients with type 2 diabetes (VITAL study): a randomised controlled trial.** *Lancet* 2010, **376**(9752):1543–1551.
- Thadhani R, Appelbaum E, Pritchett Y, Chang Y, Wenger J, Tamez H, Bhan I, Agarwal R, Zoccali C, Wanner C, Lloyd-Jones D, Cannata J, Thompson BT, Andress D, Zhang W, Packham D, Singh B, Zehnder D, Shah A, Pachika A, Manning WJ, Solomon SD: **Vitamin D therapy and cardiac structure and function in patients with chronic kidney disease: the PRIMO randomized controlled trial.** *JAMA* 2012, **307**(7):674–684.
- Levin A, Bakris GL, Molitch M, Smulders M, Tian J, Williams LA, Andress DL: **Prevalence of abnormal serum vitamin D, PTH, calcium, and phosphorus in patients with chronic kidney disease: results of the study to evaluate early kidney disease.** *Kidney Int* 2007, **71**(1):31–38.
- Pilz S, Tomaschitz A, Ritz E, Pieber TR: **Vitamin D status and arterial hypertension: a systematic review.** *Nat Rev Cardiol* 2009, **6**(10):621–630.
- Levin A: **Kidneys, hearts, hormones and immunomodulators: integrated understandings.** *Blood Purif* 2006, **24**(1):46–50.
- Andrade J, Er L, Ignaszewski A, Levin A: **Exploration of association of 1,25-OH2D3 with augmentation index, a composite measure of arterial stiffness.** *Clin J Am Soc Nephrol* 2008, **3**(6):1800–1806.
- Pfeifer M, Begerow B, Minne HW, Nachtigall D, Hansen C: **Effects of a short-term vitamin D(3) and calcium supplementation on blood pressure and parathyroid hormone levels in elderly women.** *J Clin Endocrinol Metab* 2001, **86**(4):1633–1637.
- Nagpal J, Pande JN, Bhartia A: **A double-blind, randomized, placebo-controlled trial of the short-term effect of vitamin D3 supplementation on insulin sensitivity in apparently healthy, middle-aged, centrally obese men.** *Diabet Med* 2009, **26**(1):19–27.
- Alborzi P, Patel NA, Peterson C, Bills JE, Bekele DM, Bunaye Z, Light RP, Agarwal R: **Paricalcitol reduces albuminuria and inflammation in chronic kidney disease: a randomized double-blind pilot trial.** *Hypertension* 2008, **52**(2):249–255.
- Palmer SC, McGregor DO, Craig JC, Elder G, Macaskill P, Strippoli GF: **Vitamin D compounds for people with chronic kidney disease not requiring dialysis.** *Cochrane Database Syst Rev* 2009, **1**(4):Cd008175.
- Andress DL: **Vitamin D in chronic kidney disease: a systemic role for selective vitamin D receptor activation.** *Kidney Int* 2006, **69**(1):33–43.
- Gunta SS, Thadhani RI, Mak RH: **The effect of vitamin D status on risk factors for cardiovascular disease.** *Nat Rev Nephrol* 2013, **9**(6):337–347.
- Sigrist MK, Chiarelli G, Levin A, Romann A, Weber C: **Pulse wave velocity measurements are reproducible in multiple trained observers: a short report.** *Nephron Clin Pract* 2010, **116**(1):c60–c64.
- Sigrist MK, Taal MW, Bungay P, McIntyre CW: **Progressive vascular calcification over 2 years is associated with arterial stiffening and increased mortality in patients with stages 4 and 5 chronic kidney disease.** *Clin J Am Soc Nephrol* 2007, **2**(6):1241–1248.
- Meng X, Qie L, Wang Y, Zhong M, Li L: **Assessment of arterial stiffness affected by atorvastatin in coronary artery disease using pulse wave velocity.** *Clin Invest Med* 2009, **32**(6):E238.
- Frimodt-Moller M, Nielsen AH, Kamper AL, Strandgaard S: **Pulse-wave morphology and pulse-wave velocity in healthy human volunteers: examination conditions.** *Scand J Clin Lab Invest* 2006, **66**(5):385–394.
- Frimodt-Moller M, Nielsen AH, Kamper AL, Strandgaard S: **Reproducibility of pulse-wave analysis and pulse-wave velocity determination in chronic kidney disease.** *Nephrol Dial Transplant* 2008, **23**(2):594–600.
- Xu L, Wan X, Huang Z, Zeng F, Wei G, Fang D, Deng W, Li Y: **Impact of vitamin D on chronic kidney diseases in non-dialysis patients: a meta-analysis of randomized controlled trials.** *PLoS One* 2013, **8**(4):e61387.
- Agarwal R, Acharya M, Tian J, Hippensteel RL, Melnick JZ, Qiu P, Williams L, Battle D: **Antiproteinuric effect of oral paricalcitol in chronic kidney disease.** *Kidney Int* 2005, **68**(6):2823–2828.
- Hathcock JN, Shao A, Vieth R, Heaney R: **Risk assessment for vitamin D.** *Am J Clin Nutr* 2007, **85**(1):6–18.
- Senn S: *Statistical Issues in Drug Development.* 1st edition. Chichester: Wiley; 1997.
- de Borst MH, Hajhosseiny R, Tamez H, Wenger J, Thadhani R, Goldsmith DJ: **Active vitamin D treatment for reduction of residual proteinuria: a systematic review.** *J Am Soc Nephrol* 2013, **24**(11):1863–1871.
- Cheng J, Zhang W, Zhang X, Li X, Chen J: **Efficacy and safety of paricalcitol therapy for chronic kidney disease: a meta-analysis.** *Clin J Am Soc Nephrol* 2012, **7**(3):391–400.
- Agarwal R, Siva S, Dunn SR, Sharma K: **Add-on angiotensin II receptor blockade lowers urinary transforming growth factor-beta levels.** *Am J Kidney Dis* 2002, **39**(3):486–492.
- Sigrist M, Bungay P, Taal MW, McIntyre CW: **Vascular calcification and cardiovascular function in chronic kidney disease.** *Nephrol Dial Transplant* 2006, **21**(3):707–714. Epub 2005 Nov 2001.
- Blacher J, Safar ME, Guerin AP, Pannier B, Marchais SJ, London GM: **Aortic pulse wave velocity index and mortality in end-stage renal disease.** *Kidney Int* 2003, **63**(5):1852–1860.
- Terra SG, Blum RA, Wei GC, Lew RA, Digenio AG, Rajman I, Kazierad DJ: **Evaluation of methods for improving precision of blood pressure measurements in phase I clinical trials.** *J Clin Pharmacol* 2004, **44**(5):457–463.
- Van Breukelen GJ: **ANCOVA versus change from baseline: more power in randomized studies, more bias in nonrandomized studies [corrected].** *J Clin Epidemiol* 2006, **59**(9):920–925.
- Senn S: **Change from baseline and analysis of covariance revisited.** *Stat Med* 2006, **25**(24):4334–4344.
- Hewitt NA, O'Connor AA, O'Shaughnessy DV, Elder GJ: **Effects of cholecalciferol on functional, biochemical, vascular, and quality of life outcomes in hemodialysis patients.** *Clin J Am Soc Nephrol* 2013, **8**(7):1143–1149.
- Shoben AB, Rudser KD, de Boer IH, Young B, Kestenbaum B: **Association of oral calcitriol with improved survival in nondialyzed CKD.** *J Am Soc Nephrol* 2008, **19**(8):1613–1619.
- Navaneethan SD, Schold JD, Arrigain S, Jolly SE, Jain A, Schreiber MJ, Simon JF, Srinivas TR, Nally JV: **Low 25-hydroxyvitamin D levels and mortality in non-dialysis-dependent CKD.** *Am J Kidney Dis* 2011, **58**(4):536–543.
- Teng M, Wolf M, Lowrie E, Ofsthun N, Lazarus JM, Thadhani R: **Survival of patients undergoing hemodialysis with paricalcitol or calcitriol therapy.** *N Engl J Med* 2003, **349**(5):446–456.
- Wolf M, Shah A, Gutierrez O, Ankers E, Monroy M, Tamez H, Steele D, Chang Y, Camargo CA, Tonelli M, Thadhani R: **Vitamin D levels and early mortality among incident hemodialysis patients.** *Kidney Int* 2007, **72**(8):1004–1013.
- Hsia J, Heiss G, Ren H, Allison M, Dolan NC, Greenland P, Heckbert SR, Johnson KC, Manson JE, Sidney S, Trevisan M, Women Health Initiative Investigators: **Calcium/vitamin D supplementation and cardiovascular events.** *Circulation* 2007, **115**(7):846–854.

44. St Peter WL, Liu J, Weinhandl E, Fan Q: **A comparison of sevelamer and calcium-based phosphate binders on mortality, hospitalization, and morbidity in hemodialysis: a secondary analysis of the Dialysis Clinical Outcomes Revisited (DCOR) randomized trial using claims data.** *Am J Kidney Dis* 2008, **51**(3):445–454.
45. Pfeffer MA, Burdmann EA, Chen CY, Cooper ME, de Zeeuw D, Eckardt KU, Feyzi JM, Ivanovich P, Kewalramani R, Levey AS, Lewis EF, McGill JB, McMurray JJ, Parfrey P, Parving HH, Remuzzi G, Singh AK, Solomon SD, Toto R: **A trial of darbepoetin alfa in type 2 diabetes and chronic kidney disease.** *N Engl J Med* 2009, **361**(21):2019–2032.
46. Fellstrom BC, Jardine AG, Schmieder RE, Holdaas H, Bannister K, Beutler J, Chae DW, Chevaile A, Cobbe SM, Gronhagen-Riska C, De Lima JJ, Lins R, Mayer G, McMahon AW, Parving HH, Remuzzi G, Samuelsson O, Sonkodi S, Sci D, Suleymanlar G, Tsakiris D, Tesar V, Todorov V, Wiecek A, Wuthrich RP, Gottlow M, Johnsson E, Zannad F: **Rosuvastatin and cardiovascular events in patients undergoing hemodialysis.** *N Engl J Med* 2009, **360**(14):1395–1407.
47. Levin A, Lancashire W, Fassett RG: **Targets, trends, excesses, and deficiencies: refocusing clinical investigation to improve patient outcomes.** *Kidney Int* 2013, **83**(6):1001–1009.

doi:10.1186/1471-2261-14-156

Cite this article as: Levin *et al.*: A randomized control trial to assess the impact of vitamin D supplementation compared to placebo on vascular stiffness in chronic kidney disease patients. *BMC Cardiovascular Disorders* 2014 **14**:156.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

