

RESEARCH ARTICLE

Open Access



Improvement of blood inflammatory marker levels in patients with hypothyroidism under levothyroxine treatment

Roseane C. Marchiori¹, Luiz A. F. Pereira¹, Alexandre A. Naujorks², Diego L. Rovaris³, Daiane F. Meinerz⁴, Marta M. M. F. Duarte⁵ and João B. T. Rocha^{4*}

Abstract

Background: There are several specific inflammatory and oxidative correlates among patients with hypothyroidism, but most studies are cross-sectional and do not evaluate the change in parameters during the treatment. The aim of this study was to investigate the effect of levothyroxine replacement therapy on biomarkers of oxidative stress (OS) and systemic inflammation in patients with hypothyroidism.

Methods: In this prospective open-label study, 17 patients with recently diagnosed primary hypothyroidism due to Hashimoto's thyroiditis who were not taking levothyroxine were included. The following parameters were measured before and at 6 and 12 months of levothyroxine treatment with an average dose of 1.5 to 1.7 $\mu\text{g}/\text{kg}/\text{day}$: thyroid-stimulating hormone (TSH), free thyroxine (FT4), high-sensitivity C-reactive protein (hs-CRP), interleukin 1 (IL-1), IL-6, IL-10, interferon gamma (INF- γ), tumor necrosis factor alpha (TNF- α), thiobarbituric acid-reactive substances (TBARS), activity of aminolevulinic acid dehydratase (δ -ALA-D), nonprotein and total thiol (NP-SH and T-SH) groups, total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C) and triglycerides (TG). Generalized estimating equation (GEE) modeling was used to analyze the effects of LRT (at pre-treatment, 6 months and 12 months) on those variables. The hypothyroidism status (i.e., overt or subclinical hypothyroidism) was included as a confounder in all analyses. An additional GEE post hoc analysis was made to compare time points.

Results: There was a significant decrease in TSH over time ($P < 0.0001$), (initial levels were on average 32.4 $\mu\text{U}/\text{mL}$ and 10.5 $\mu\text{U}/\text{mL}$ at 12 months). There was a significant increase in FT4 ($P < 0.0001$) (initial levels were on average 0.8 ng/dL and 2.7 ng/dL at 12 months). There were significant changes in interleukin levels over time, with a significant increase in IL-10 ($P < 0.0001$) and significant decreases in IL-1 ($P < 0.0001$), IL-6 ($P < 0.0001$), INF- γ ($P < 0.0001$) and TNF- α ($P < 0.0001$). No significant difference in hs-CRP over time was observed ($P < 0.284$). There was a significant reduction in NP-SH ($P < 0.0001$).

Conclusions: This study observed significant changes in the inflammatory profile in hypothyroid patients under treatment, with reduction of pro-inflammatory cytokines and elevation of anti-inflammatory cytokine. In these patients, a decrease in low-grade chronic inflammation may have clinical relevance due to the known connection between chronic inflammation, atherosclerosis and cardiovascular events.

Keywords: Inflammation, Oxidative stress, Overt hypothyroidism, Subclinical hypothyroidism, Hashimoto's thyroiditis, Levothyroxine, Atherosclerosis

* Correspondence: jbtrocha@yahoo.com.br

⁴Departamento de Bioquímica e Biologia Molecular, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria, Av. Roraima n° 1000, Cidade Universitaria, Camobi, Santa Maria, RS, Brazil

Full list of author information is available at the end of the article

Background

The full or partial deficiency of thyroid hormone action is called hypothyroidism, which can be either overt (OH) or subclinical (SCH). SCH is characterized by a serum thyrotropin (TSH) level above the upper reference limit in combination with a normal level of free thyroxine (FT4), while OH is characterized by elevated TSH, in combination with subnormal FT4 [1]. The most common cause of primary hypothyroidism is chronic autoimmune thyroiditis, a disease also known as Hashimoto's thyroiditis [1]. It is characterized by diffuse infiltration of the gland with sensitized T lymphocytes, with gradual destruction and fibrous replacement of the thyroid parenchymal tissue, elevated serum antithyroid antibodies, evidence of goiter or thyroid glandular atrophy and dysfunction to varying degrees [2].

Hypothyroidism is a prevalent disorder [3–5], and both OH and SCH seem to exert deleterious effects on the cardiovascular system [6]. Several mechanisms may be involved in this interaction, and the increased risks of atherosclerosis and coronary heart disease are some of them [4, 7]. Atherosclerosis develops over a period of years; inflammation is implicated in all of its stages (from the initial leukocyte recruitment to eventual rupture of the unstable atherosclerotic plaque) and has also been considered the link between the traditional risk factors and evident modifications in the artery wall [8].

Numerous circulating inflammatory biomarkers are associated with increased acute coronary event risk. These biomarkers may reflect pathways involved in disease progression and may thus be potential tools for predicting atherosclerosis and cardiovascular events [9]. C-reactive protein (CRP) is one of the most widely studied biomarkers in the general population and has been used to assess cardiovascular risk in both healthy subjects and people with various disorders [10]. Most epidemiological evidence on the relevance of cytokines as inflammatory markers has been obtained for interleukin (IL)-6 [11, 12]. Long-term circulating IL-6 levels may be associated with coronary heart disease, such as the main risk factors already established [11]. A causal role for IL-6 signaling in coronary heart disease is still elusive, but has been considered [12]. IL-1, interferon gamma (IFN- γ) and tumor necrosis factor alpha (TNF- α) are also important both the development of coronary heart disease and plaque destabilization [13–15]. TNF- α levels are significantly higher in patients with myocardial infarction than in controls, and patients with persistently higher levels in the post-infarction period are at a three fold higher risk of developing new coronary episodes [15]. There is substantial evidence that IL-1 inhibition can reduce recurrent vascular events in a high-risk secondary prevention population [13]. High levels of systemic markers of IFN- γ activity predict long-term adverse

prognosis in patients with stable angina pectoris, demonstrating the capacity to identify individuals with vulnerable lesions, despite their stable clinical conditions [14]. Circulating levels of several different pro-inflammatory cytokines in initially healthy people are associated with risk of coronary heart disease outcomes in an approximately log-linear manner [16].

Oxidative stress (OS) can also serve as both a monitor of inflammation and a pathophysiologic mediator of atherosclerosis [8, 17]. The association between hypothyroidism and increased OS is controversial, but greater OS has been observed in patients with SCH and OH in comparison with euthyroid controls [18–23]. Inflammation observed in patients with Hashimoto's thyroiditis and hypothyroidism is considered the link between OS and increased risk of atherosclerosis and underlying cardiovascular disease in these patients [24–26].

Hypothyroidism is treated with substitutive doses of levothyroxine. Few studies have evaluated the effects of hypothyroidism treatment on either OS, with conflicting results [18–22], or systemic inflammation [19, 27–31], also with conflicting results. Thus, this study aims to assess inflammatory biomarkers and OS in patients with primary hypothyroidism at baseline and after 6 and 12 months of levothyroxine replacement therapy (LRT).

Methods

Subjects

Patients were recruited prospectively, in the outpatient endocrinology clinic, University Hospital of Santa Maria, Santa Maria, RS, Brazil. The study protocol was approved by the Human Ethics Committee of the Federal University of Santa Maria (number 0177.0.243.000.07). Before performing any procedure, all of the patients were extensively informed about the study and signed an informed consent form. Inclusion criteria were age between 18 and 69 years, recent OH defined as high thyrotropin (TSH greater than 4.2 μ IU/mL) and low FT4 or SCH, defined as a high level of TSH with FT4 level within the reference range (0.93 to 1.71 ng/dl). Patients with SCH had been referred to the clinic because of presentation of goiter and its symptoms (dry skin, muscle weakness, impaired memory and thinking, fatigue, cold intolerance, constipation), with stable thyroid dysfunction for at least 6 months, without evidence of other recent or ongoing diseases.

Stringent exclusion criteria were defined to limit sources of confounding factors that could influence the clinical or laboratory parameters under investigation. We did not include patients with current treatment for thyroid dysfunction; diabetes mellitus; pregnancy; malignancy; active smoking; alcohol consumption; history of previous or recent respiratory disease; previous or recent cardiovascular disease; history of connective tissue diseases;

or exposure to drugs such as interferon, amiodarone, lipid-lowering drugs, appetite suppressants, antioxidant vitamin supplementation, or beta-blockers.

A medical history, physical examination, thorax radiogram, spirometry, electrocardiogram, transthoracic Doppler echocardiogram and laboratory investigations were performed in all patients. Body mass index (BMI) was calculated by dividing the body weight by the square of the height (kg/m^2). The levothyroxine (FT4) replacement therapy was performed with an average dose of 1.5 to 1.7 $\mu\text{g}/\text{kg}/\text{day}$. SCH required a lower dose. Patients were instructed to take the medication orally, with water, always in a single dose in the morning, after fasting (between 30 and 60 min before the first meal). Inflammatory and oxidative stress profiles were performed simultaneously with the collection of the thyroid profile, at 6 and 12 months follow-up.

Laboratory assessments

At baseline and after 6 and 12 months of LRT, blood samples were collected to assess thyroid profile (TSH, FT4), lipid profile (total cholesterol [TC], high-density lipoprotein cholesterol [HDL-C], low-density lipoprotein cholesterol [LDL-C], triglycerides [TG]) and inflammatory biomarkers (high-sensitivity CRP [hs-CRP], IL-1, IL-6, IL-10, INF- γ and TNF- α). In addition, we measured markers of OS (aminolevulinic acid dehydratase [δ -ALA-D], total thiol group [T-SH] and nonprotein thiol group [NP-SH] and thiobarbituric acid-reactive substances [TBARS]). Dosages of antithyroglobulin antibody (antiTG Ab) and antiperoxidase antibody (antiTPO Ab) were performed only at pre-treatment.

Blood samples were collected by venipuncture after fasting for 12 h. The samples were centrifuged for 15 min at 2500 $\times g$ and serum aliquots were stored at -20 degrees centigrade to study IL-1, IL-6, IL-10, and INF- γ , TNF- α . An aliquot of whole blood was sent to the laboratory of Biochemistry of the Natural and Exact Sciences Center for further measures of δ -ALA-D activity and NP-SH, T-SH and TBARS levels.

The thyroid profile (TSH, FT4, antibodies) was measured by electrochemiluminescence (Immulite 2000, Siemens Healthcare Diagnostics, USA). We considered the following reference values: TSH: 0.27 to 4.2 uIU/mL, FT4: 0.93 to 1.71 ng/dl; antiTPO Ab: up to 35 IU/ml and antiTG Ab: up to 115 IU/ml.

The lipid profile was measured by the homogeneous enzyme method (Dimension 2008, Siemens Healthcare Diagnostics, USA). TC and TG were measured in serum according to the technique recommended by the manufacturer. HDL-C was measured in plasma supernatant after precipitation of lipoproteins containing apolipoprotein B with dextran sulfate and magnesium chloride as previously described by Bachorik and Albers [32].

LDL-C was estimated from the Friedewald equation [33]. Baseline values were: TC: up to 200 mg/dl, TG: 150 mg/dl, HDL-C: above 40 mg/dl for men and above 50 mg/dl for women; LDL-C: up to 130 mg/dl.

Hemoglobin was measured using the Sysmex automated system.

CRP was measured in serum by the immunoturbidimetric method, which measures hs-CRP (Dimension 2008, Siemens Healthcare Diagnostics, USA). The reference value was less than 0.5 mg/dl.

Cytokines were measured by enzyme-linked immunosorbent assay (ELISA) using commercial kits for human IL-1, IL-6, IL-10, INF- γ , and TNF- α (eBioscience, San Diego, USA) according to the manufacturer's instructions.

Lipid peroxidation was determined by TBARS quantification using the method of Ohkawa et al. [34] with modifications. Red blood cells were isolated and proteins precipitated with 5 % trichloroacetic acid on ice for 30 min. The precipitate was removed by centrifugation at 3,000 rpm and the supernatant were then incubated at 100 °C for 60 min in an acidic medium containing 0.35 % phosphoric acid, and 0.28 % thiobarbituric acid. After centrifugation, the reaction product was determined at 532 nm using malondialdehyde as a standard. The results are expressed as nmol MDA/ml of erythrocytes.

Thiol oxidation was assessed by determination of T-SH and NP-SH, as described by Ellman [35]. For NP-SH level 300 microliters of isolated red blood cells were hemolyzed with 100 μL of 10 % Triton solution for 10 min. Following by protein fraction precipitation with 200 μL of 20 % trichloroacetic acid, the free SH groups (NP-SH) were determined in the supernatant using the colorimetric method performed in a potassium phosphate buffer (1 M, pH 7.4) with the presence of 10 mM of 5,5'-dithio(bis-nitrobenzoic acid) (DTNB). A standard curve was constructed using glutathione. The NP-SH level was measured at 412 nm and is expressed as nmol NP-SH/ml erythrocytes.

The T-SH level was determined in plasma and the colorimetric assay was performed in 0.85 ml of potassium phosphate buffer (0.3 M, pH 7.0), using 0.05 ml of 5,5'-dithio(bis-nitrobenzoic acid). A standard curve was constructed using glutathione to determine the total thiol groups. The T-SH level was measured at 412 nm and is expressed as nmol T-SH/ml of plasma.

δ -ALA-D was determined in whole blood according to the method of Sassa [36]. Enzyme activity was measured in the presence or absence of the reducing agent dithiothreitol (DTT). The enzymatic reaction in hemolyzed blood (after mixing with Triton 0.05 %) was started after 10 min of sample pre-incubation, by adding the substrate aminolevulinic acid (ALA), followed by incubation at 37 °C for 60 min. The reaction product (porphobilinogen) was determined using Ehrlich's reagent, which

reacted and formed a pink-colored product measured at 555 nm. The activity is expressed as nmol PBG/hour/ml of blood.

Statistical analysis

Data were analyzed with SPSS 18.0 (SPSS Inc. Chicago, IL, USA). Generalized estimating equation (GEE) modeling was used to analyze the effects of LRT (at pre-treatment, 6 months, and 12 months) on thyroid and lipid profiles, and on systemic inflammation and OS. The type of model and the link function were defined according to dependent variable distribution. Each parameter was analyzed as to its nature to choose the most suitable working correlation matrix. The model was chosen as the criterion of lowest quasi-likelihood under the independence model criteria (QIC). The hypothyroidism status (i.e., OH or SCH) was included as a confounder in all analyses.

Because 17 models were tested, we considered *P*-values less than 0.0025 ($\alpha_{Bonf} = 0.05/17$) to be significant. Eight models were significant at the 0.0025 level and underwent an additional GEE post hoc analysis to compare time points. In these analyses, corrected *P*-values were calculated, and alpha was set at 0.05. Data are expressed as means and standard deviations, means and 95 % confidence intervals (95 % CIs) or numbers (*n*) and percentages (%), as indicated in the table footnotes.

Results

Clinical, anthropometric, and baseline thyroid profiles of the study participants are given in Table 1. The group included 17 patients with high TSH, 11 of whom had diminished FT4 (OH) and 6 of whom had normal FT4 (SCH). The initial 17 patients, 16 were reassessed at 6 and 12 months of treatment with levothyroxine. Thyroid, lipid, hemoglobin, inflammatory and OS profiles of hypothyroid patients at pre-treatment and 6 and 12 months after LRT are shown in Table 2. All patients had increased anti-thyroglobulin antibodies and/or antibody antiperoxidase levels.

GEE evaluation (Table 2) revealed a significant decrease in TSH over time ($P < 0.0001$). The initial TSH

Table 1 Clinical, anthropometric and pre-treatment thyroid profile of the study

Variables	Hypothyroidism (TSH ↑) (<i>n</i> = 17)
Gender (males)	6 (35.3)
Age (years)	45.8 (15.2)
BMI (Kg/m ²)	28.4 (5.8)
TSH (uIU/ml)	32.1 (30.2)
FT4 (ng/dl)	0.8 (0.4)
AntiTG Ab (IU/mL)	979.9 (1386.0)
AntiTPO Ab (IU/mL)	398.0 (309.9)

Data are mean and (standard deviation) or *n* and (%) for gender

levels were on average 32.4 uIU/mL and the levels achieved at 12 months were 10.5 uIU/mL. In the post hoc analysis, significance was observed only at 12 months of LRT compared with pre-treatment (Fig. 1a). There was also significant modification in FT4 over time ($P < 0.0001$). The initial FT4 levels were on average 0.8 ng/dl and the levels achieved at 12 months were on average 2.7 ng/dl. In the post hoc analysis, a significant difference was noted between 6-months FT4 and pre-treatment (Fig. 1b).

Regarding inflammatory profile, there were significant changes in interleukin levels over time, with a significant increase in IL-10 ($P < 0.0001$) and significant decreases of IL-1 ($P < 0.0001$), IL-6 ($P < 0.0001$), INF- γ ($P < 0.0001$) and TNF- α ($P < 0.0001$). The GEE post hoc analyses revealed that the 6- and 12-month values differed significantly from pre-treatment (Fig. 1c to 1g). No significant difference in hs-CRP over time was noted ($P = 0.284$).

The evaluation of OS during follow-up showed significant reduction of NP-SH ($P < 0.0001$). In the post hoc analysis, 6 and 12-month NP-SH levels differed from pre-treatment (Fig. 1h). No significant change were observed in T-SH ($P = 0.068$), δ -ALA-D ($P = 0.725$) or TBARS ($P = 0.239$) levels over time.

An increasing trend was detected for hemoglobin levels ($P = 0.016$). There were no significant changes in TC ($P = 0.457$), HDL-C ($P = 0.715$), LDL-C ($P = 0.342$) or TG ($P = 0.581$) during follow-up.

Discussion

The main finding of this open label study was that patients with hypothyroidism using levothyroxine presented significant changes in the inflammatory profile, with reduction of pro-inflammatory cytokines and increase of one anti-inflammatory cytokine. It was also observed reduction of one OS biomarker, although there were no significant changes in CRP or lipid profile.

Christ-Crain et al. [27] first reported increased CRP in patients with OH or SCH citing it as an additional risk factor for development of coronary heart disease in those patients, but further studies have shown conflicting results. Some studies have detected elevated CRP in individuals with hypothyroidism [5, 19, 24–26, 29, 37–39], while others have not [30, 40, 41]. Regarding the effect of LRT on CRP, in the current study, in accordance with the first report of Christ-Crain et al. [27] and others [30, 31, 37, 39, 41], the CRP level was unaffected. However, other studies have shown that normalization of thyroid state by LRT seems to effectively reduce serum CRP [19, 29, 38]. Marfella et al. [26] observed a reduction in CRP in treated SCH patients, but CRP was still significantly increased compared to the control group.

While pro-inflammatory cytokines have a detrimental role in atherosclerosis [8], the role of cytokines in

Table 2 Generalized estimating equation (GEE) models to evaluate the effects of Levothyroxine replacement therapy (LRT) on inflammatory, oxidative stress, hemoglobin, and lipid profile parameters of hypothyroid patients

Parameters	Pre-treatment	Six months	Twelve months	Wald Chi-square	df	P-value
	Mean (95 % CI)	Mean (95 % CI)	Mean (95 % CI)			
TSH (μ IU/mL)	32.4 (18.0–30.4)	20.7 (8.6–49.8)	10.5 (6.3–17.4)	18.496	2	<0.0001
FT4 (ng/dL)	0.8 (0.7–0.9)	3.4 (1.8–6.6)	2.7 (1.3–5.8)	33.684	2	<0.0001
IL-10 (pg/mL)	132.7 (124.5–141.0)	160.7 (150.7–170.6)	184.6 (174.8–194.4)	181.076	2	<0.0001
IL-1 (pg/mL)	73.0 (68.5–77.4)	57.9 (52.6–63.1)	45.0 (40.4–49.6)	321.214	2	<0.0001
IL-6 (pg/mL)	90.1 (83.7–96.4)	70.5 (63.9–77.1)	49.9 (44.1–55.6)	512.104	2	<0.0001
INF- γ (pg/mL)	144.8 (141.0–148.6)	98.6 (93.1–104.5)	80.0 (75.8–84.3)	386.001	2	<0.0001
TNF- α (pg/mL)	122.8 (120.5–125.1)	90.4 (86.8–94.0)	66.2 (61.6–71.1)	341.402	2	<0.0001
hs-CRP (mg/dl)	0.6 (0.3–1.1)	0.5 (0.3–0.8)	0.9 (0.5–1.7)	2.519	2	0.284
NP-SH (nmol NP - SH/mL erit)	1522.4 (1374.9–1670.0)	1098.1 (929.3–1267.0)	1004.6 (757.5–1251.7)	23.528	2	<0.0001
Total-SH (nmol T -SH/ml plasma)	823.3 (683.5–963.2)	624.0 (537.1–710.8)	660.5 (584.2–772.8)	5.388	2	0.068
δ -ALA-D (nmol PBG/hour/ml of blood)	3.6 (2.7–4.4)	3.1 (1.9–4.3)	3.0 (2.1–4.0)	0.644	2	0.725
TBARS (nmol MDA/mL erit)	5.3 (4.0–6.7)	7.2 (4.8–8.5)	6.7 (4.8–8.5)	2.862	2	0.239
Hemoglobin (g/dL)	13.1 (12.6–13.6)	13.4 (13.0–13.9)	13.6 (13.2–14.0)	8.252	2	0.016
TC (mg/dL)	201.5 (176.7–244.2)	198.4 (168.5–228.4)	198.4 (171.9–225.0)	1.567	2	0.457
HDL- C (mg/dL)	54.3 (48.8–59.7)	50.9 (41.8–60.0)	52.1 (45.8–58.4)	0.670	2	0.715
LDL- C (mg/dL)	124.2 (102.6–150.5)	115.5 (99.5–139.7)	115.5 (99.0–134.7)	2.146	2	0.342
TG (mg/dL)	117.5 (88.1–156.9)	139.2 (99.2–195.4)	119.6 (85.3–167.7)	1.085	2	0.581

df = degrees of freedom; adjusted for overt hypothyroidism and subclinical hypothyroidism

Hashimoto's thyroiditis is complex and often contradictory: a regulatory cytokine may either favor induction of tolerance against thyroid autoimmune disease or favor activation and/or exacerbation of autoimmune responses [2]. Drugarin et al. [42] observed increased serum IL-2, TNF- α and IFN- γ in subjects with OH due to autoimmune thyroiditis compared with healthy controls, in whom these cytokines were barely detectable. Recently, in patients with autoimmune thyroiditis, increased IL-6 and IL-15 (another pro-inflammatory cytokine) were detected, likely due to increased proliferation and increased pro-inflammatory cytokine synthesis in T-helper 17 cells [43]. Díez et al. [44] confirmed the relevance of activation of the TNF- α system in patients with hypothyroidism, showing that serum TNF- α and receptors of TNF- α were significantly higher than those detected in the control group. Thyroid hormone levels have been suggested to influence cytokine production. IL-6 level might be correlated with hypothyroidism severity because serum IL-6 level has been positively correlated with LRT dose and negatively correlated with FT4 level in hypothyroidism due to autoimmune thyroiditis [45]. Díez et al. [44] found no differences in TNF- α or receptors of TNF- α between patients with autoimmune and non-autoimmune hypothyroidism, implicating thyroid hormone deficiency per se in systemic cytokine production. Karanikas et al. [46] demonstrated no influence of thyroid

hormone on cytokine production patterns by T cells, as they reported that patients with high titers of anti-TPO antibodies had significantly higher percentages of cells producing INF- γ and TNF- α than healthy controls.

Based on previous studies, an association between hypothyroidism and low-grade inflammation may be suggested. Taddei et al. [24] demonstrated a higher plasma IL-6 level, impaired endothelium dysfunction vasodilatation and reduced nitric oxide availability in patients with SCH compared with euthyroid controls. Türemen et al. [25] observed endothelial dysfunction and significantly higher serum IL-6 and TNF- α levels in SCH patients with autoimmune thyroiditis compared to controls. Patients with untreated SCH who underwent thromboendarterectomy due to asymptomatic severe internal carotid artery stenosis presented higher plasma TNF- α , IL-6, and interleukin-18 levels, and their atherosclerotic plaques presented an active inflammatory reaction with higher TNF- α levels and other characteristics of instability compared with non-SCH controls [26].

Regarding potential changes in cytokine levels with LRT, we found that one year of LRT produced significant reductions in pro-inflammatory cytokines (IL-1, IL-6, INF- γ , TNF- α) and an increase in IL-10. There are few data in the literature about the effect of levothyroxine on serum cytokines. Díez et al. [44] observed no reduction of the high levels of TNF- α and TNF- α receptors after




Fig. 1 Post Hoc comparisons from significant generalized estimating equation. (GEE) models: Panel **A** (TSH); Panel **B** (FT4); Panel **C** (IL-10); Panel **D** (IL-1); Panel **E** (IL-6); Panel **F** (INF- γ); Panel **G** (TNF- α) and Panel **H** (NP-SH), as described in Table 2. * $P \leq 0.050$; ** $P \leq 0.010$; *** $P \leq 0.001$

normalization of thyroid function in hypothyroid patients. Guclu et al. [47] observed a significant decrease in serum IL-12 (one of the most important cytokines responsible for Th1-type cytokine responses), a statistically non-significant decrease in IFN- γ and no change in serum IL-2 or IL-4 in patients with hypothyroidism due to

Hashimoto's thyroiditis who underwent 12 weeks of LRT. Marfella et al. [26] observed significantly lower plasma TNF- α , IL-6 and IL-18 levels, low pro-inflammatory cytokine levels and macrophage infiltration in atherosclerotic lesions of treated SCH patients compared to untreated SCH patients.

The association between hypothyroidism and increased OS is controversial, as is the effect of LRT on oxidative stress biomarkers. In the current study, NP-SH was significantly reduced by 12 months of LRT, but no significant changes were observed in T-SH, δ -ALA-D or TBARS. Some studies have shown reduction of oxidative stress after LRT, [18, 23, 29], while other studies have reported no modification [19, 20]. Marfella et al. [26] detected elevated OS biomarkers in atherosclerotic plaques of SCH subjects compared to control plaques, and higher levels of OS in plaques of untreated SCH patients in comparison with plaques of treated SCH patients. The restoration of endothelial function by systemic administration of indomethacin (cyclooxygenase inhibitor) or local infusion of vitamin C (an antioxidant) reinforces that OS may be a link between the inflammation and endothelial dysfunction observed in SCH patients [24]. According to Marfella et al. [26], OS reduction can play an important role in reducing the inflammatory activity in atherosclerotic plaques observed in patients with SCH treated with thyroxine.

In our sample, which included both OH and SCH patients, there were no significant changes in TC, HDL, LDL or TG over 12 months of LRT. Dyslipidemia is a common finding in patients with OH, with elevated TC and LDL [3, 27, 48], which usually improve with levothyroxine use [48]. Less clear is the relationship between SCH and lipid profile. Some studies have shown no alterations in lipid profile in SCH patients compared with controls [4, 27, 40, 41], while others have observed abnormalities in serum cholesterol or TG in SCH [5, 38]. Teixeira et al. [49] have suggested that SCH presents an intermediary lipid profile between that observed in normal individuals and that usually observed in OH. The role of levothyroxine in changing the lipid profile in SCH is also not clear. Iqbal et al. [50], in a large epidemiological study, and Razvi et al. [51], in a study of 100 patients with SCH, reported TC and LDL-C reductions with LRT. Teixeira et al. [49] observed a significant lipid profile improvement after 1 year of LRT. However, other studies did not report modifications of lipid levels with LRT [19, 29, 31].

Three older meta-analyses concluded that thyroxine substitution has no effect on TG and HDL-C [48, 52, 53], whereas TC and LDL-C levels were reduced, with the most pronounced effect in those with the highest pretreatment serum TSH [53] and serum TC [48, 52]. However, a more recent meta-analysis based on the Cochrane methodology, including 12 randomized controlled trials of 6–14 months' duration, reported that the beneficial effects of treatment were slight and that only TC was affected [54].

Knowledge about the effects of levothyroxine on inflammatory, OS and lipid profiles has been increasing. The results are conflicting, but there are growing indications of favorable effects of levothyroxine on adhesion molecule levels in SCH [55], impaired endothelial function

and hemodynamic profile both in OH and SCH [28, 37, 41, 51, 56]. There are also indications that young patients with treated SCH, in comparison with untreated SCH patients, present a significantly lower risk of developing heart failure [57], a significantly lower all-cause mortality [58] and fewer events of ischemic heart disease [59]. The decreased immune overactivity observed in our study may be a mechanism contributing to these previous results.

Our study has strengths and limitations. The strengths are its prospective nature, the number of investigated inflammatory biomarkers, the rigorous patient selection and the careful statistical analysis with correction for confounding factors. The main limitations are the rather small sample size, heterogenous sample (composed of both OH and SCH), lack of a control group and lack of a placebo-controlled group.

Conclusion

In conclusion, this study observed a significant reduction of pro-inflammatory cytokines and an increase of one anti-inflammatory cytokine in hypothyroid patients using levothyroxine. In these patients, the reduction of low-grade chronic inflammation may have clinical relevance because of the known connection between chronic inflammation, atherosclerosis and cardiovascular events.

Abbreviations

δ -ALA-D: Aminolevulinic acid dehydratase; antiTG Ab: antithyroglobulin antibody; antiTPO Ab: antiperoxidase antibody; BMI: Body mass index; CRP: C-reactive protein; FT4: Free thyroxine; HDL-C: High-density lipoprotein cholesterol; hs-CRP: high-sensitivity C-reactive protein; IL-1: Interleukin 1; IL-6: Interleukin 6; IL-10: Interleukin 10; INF- γ : Interferon gamma; LRT: Levothyroxine replacement therapy; LDL-C: Low-density lipoprotein cholesterol; NP-SH: Nonprotein thiol group; OH: Overt hypothyroidism; OS: Oxidative stress; SCH: Subclinical hypothyroidism; TBARS: Thiobarbituric acid-reactive substances; TSH: Thyroid-stimulating hormone, TC, Total cholesterol; T-SH: Total thiol group; TG: Triglycerides; TNF- α : Tumor necrosis factor alpha.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed to the protocol and development of the rationale of the study as well as to the development of the final manuscript. RCM conceived of the study and participated in its design, collect data, coordination and wrote the manuscript. LAFP participated in its design, collect data and conduction of the study. AAN, DFM and MMMFD. carried out laboratory tests and helped to draft the manuscript. DLR performed statistical analysis and helped to draft the manuscript. JBTR conceived of the study and participated in its design, statistical analysis and helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors thank all the patients who participated in this study. The study was supported by agencies that promote and develop the post graduate in Brazil: CNPq (Conselho Nacional de desenvolvimento Científico e Tecnológico, National Council for Scientific and Technological Development), CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Coordination for the Improvement of Higher Education Personnel), FINEP (Financiadora de Estudos e Projetos, Studies and Projects Financing Agency), PRONEX – FAPERGS (Programa de apoio a núcleos de excelência – Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul, Support Program for Excellence Nuclei – Research Support Foundation of Rio Grande do Sul).

Author details

¹Departamento de Clínica Médica, Centro de Ciências da Saúde, Universidade Federal de Santa Maria, Av. Roraima n° 1000, Cidade Universitária, Camobi, Santa Maria, RS, Brazil. ²Serviço de Métodos Gráficos, Hospital Universitário de Santa Maria, Universidade Federal de Santa Maria, Av. Roraima n° 1000, Cidade Universitária, Camobi, Santa Maria, RS, Brazil. ³Departamento de Genética, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves, 9500, Porto Alegre, Brazil. ⁴Departamento de Bioquímica e Biologia Molecular, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria, Av. Roraima n° 1000, Cidade Universitária, Camobi, Santa Maria, RS, Brazil. ⁵Ciências da Saúde, Universidade Luterana do Brasil (ULBRA), campus Santa Maria, BR 287, Km 252, Trevo Maneco Pedroso, Boca do Monte, Santa Maria, RS, Brazil/Cx. Postal 21834.

Received: 21 February 2015 Accepted: 12 June 2015

Published online: 23 June 2015

References

- Garber JR, Cobin RH, Guarib H, Hennessey JV, Klein I, Mechanik JJ, et al. Clinical practice guidelines for hypothyroidism in adults: cosponsored by the American Association of Clinical Endocrinologists and the American Thyroid Association. *Endocr Pract.* 2012;18:989–1028.
- Ganesh BB, Bhattacharya P, Gopisetty A, Prabhakar BS. Role of cytokines in the pathogenesis and suppression of thyroid autoimmunity. *J Interferon Cytokine Res.* 2011;31:721–31.
- Canaris GJ, Manowitz NR, Mayor G, Ridgway C. The Colorado thyroid disease prevalence study. *Arch Intern Med.* 2000;160:526–34.
- Hak AE, Pols HAP, Visser TJ, Drexhage HA, Hofman A, Witteman JCM. Subclinical hypothyroidism is an independent risk factor for atherosclerosis and myocardial infarction in elderly women: the Rotterdam study. *Ann Intern Med.* 2000;132:270–8.
- Kvetny J, Heldgaard PE, Bladbjerg EM, Gram J. Subclinical hypothyroidism is associated with a low grade inflammation, increased triglyceride levels and predicts cardiovascular disease in males below 50 years. *Clin Endocrinol (Oxf).* 2004;61:232–8.
- Klein I, Danzi S. Thyroid disease and the heart. *Circulation.* 2007;116:1725–35.
- Cappola AR, Ladenson PW. Hypothyroidism and atherosclerosis. *J Clin Endocrinol Metab.* 2003;88:2438–44.
- Libby P. Inflammation in atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2012;32:2045–51.
- Welsh P, Packard CJ, Sattar N. Novel antecedent plasma biomarkers of cardiovascular disease: improved evaluation methods and comparator benchmarks raise the bar. *Curr Opin Lipidol.* 2008;19:563–71.
- Buckley DJ, Fu R, Freeman M, Rogers K, Helfand M. C-reactive protein as a risk factor for coronary heart disease: a systematic review and meta-analysis for the U.S. Preventive Services Task Force. *Ann Intern Med.* 2009;151:483–95.
- Danesh J, Kaptoge S, Mann AG, Sarwar N, Wood A, Angleman SB, et al. Long-term interleukin-6 levels and subsequent risk of coronary heart disease: two new prospective studies and a systematic review. *PLoS Med.* 2008;5:e78.
- Interleukin-6 Receptor Mendelian Randomisation Analysis (IL6R MR) Consortium. The interleukin-6 receptor as a target for prevention of coronary heart disease: a mendelian randomisation analysis. *Lancet.* 2012;379:1214–24.
- Ridker PM, Howard CP, Walter V, Everett B, Libby P, Hensen J, et al. Effects of Interleukin-1 Beta inhibition with Canakinumab on hemoglobin A1c, lipids, C-reactive protein, interleukin-6, and fibrinogen. A phase IIb randomized, placebo-controlled trial. *Circulation.* 2012;126:2739–48.
- Pedersen ER, Midttun O, Ueland PM, Schartum-Hansen H, Seifert R, Iglund J, et al. Systemic markers of Interferon-gamma-mediated immune activation and long-term prognosis in patients with stable coronary artery disease. *Arterioscler Thromb Vasc Biol.* 2011;31:698–704.
- Ridker PM, Rifai N, Pfeffer M, Sacks F, Lepage S, Braunwald E. Elevation of Tumor Necrosis Factor-alpha and increased risk of recurrent coronary events after myocardial infarction. *Circulation.* 2000;101:2149–53.
- Kaptoge S, Seshasai SRK, Gao P, Freitag DF, Butterworth AS, Borglykke A, et al. Inflammatory cytokines and risk of coronary heart disease: new prospective study and updated meta-analysis. *Eur Heart J.* 2014;35:587–9.
- Sugiyama S, Kugiyama K, Aikawa M, Nakamura S, Ogawa H, Libby P. Hypochlorous acid, a macrophage product, induces endothelial apoptosis and tissue factor expression: involvement of myeloperoxidase-mediated oxidant in plaque erosion and thrombogenesis. *Arterioscler Thromb Vasc Biol.* 2004;24:1309–14.
- Baskol G, Atmaca H, Tanriverdi F, Baskol M, Kocer D, Bayram F. Oxidative stress and enzymatic antioxidant status in patient with hypothyroidism before and after treatment. *Exp Clin Endocrinol Diabetes.* 2007;115:522–6.
- Kebapçılar L, Akinci B, Bayraktar F, Comlekci A, Solak A, Demir T, et al. Plasma thiobarbituric acid-reactive substance levels in subclinical hypothyroidism. *Med Princ Pract.* 2007;16:432–6.
- Mehmetçik G, Becer E, Akbey A. Serum total antioxidant status, lipid profile, malondialdehyde and erythrocyte superoxide dismutase levels in Hashimoto thyroiditis patients treated with levothyroxine. *Türkiye Kinikleri J Med Sci.* 2012;32:1241–6.
- Bhimte B, Agrawal BK, Sharma VK, Chauhan SS. Oxidative stress status in hypothyroid patients. *Biomed Res.* 2012;23:286–8.
- Geranova J, Gadjeva V. Oxidative stress and antioxidant enzyme activities in patients with Hashimoto's thyroiditis. *Comp Clin Pathol.* 2007;16:259–64.
- Erdamar H, Demirci H, Yaman H, Erbil MK, Yakar T, Sancak B, et al. The effect of hypothyroidism, hyperthyroidism, and their treatment on parameters of oxidative stress and antioxidant status. *Clin Chem Lab Med.* 2008;46:1004–10.
- Taddei S, Caraccio N, Virdis A, Dardano A, Versari D, Ghiadoni L, et al. Low-grade systemic inflammation causes endothelial dysfunction in patients with Hashimoto's thyroiditis. *J Clin Endocrinol Metab.* 2006;91:5076–82.
- Türemen EE, Çetinarslan B, Sahin T, Cantürk Z, Tarkun I. Endothelial dysfunction and low grade chronic inflammation in subclinical hypothyroidism due to autoimmune thyroiditis. *Endocr J.* 2011;58:349–54.
- Marfella R, Ferraraccio F, Rizzo MR, Portoghese M, Barbieri M, Basilio C, et al. Innate immune activity in plaque of patients with untreated and L-thyroxine-treated subclinical hypothyroidism. *J Clin Endocrinol Metab.* 2011;96:1015–20.
- Christ-Crain M, Meier C, Guglielmetti M, Huber PR, Riesen W, Staub JJ, et al. Elevated C-reactive protein and homocysteine values: cardiovascular risk factors in hypothyroidism? A cross-sectional and a double-blind, placebo-controlled trial. *Atherosclerosis.* 2003;166:379–86.
- Taddei S, Caraccio N, Virdis A, Dardano A, Versari D, Ghiadoni L, et al. Impaired endothelium-dependent vasodilatation in subclinical hypothyroidism: beneficial effect of levothyroxine therapy. *J Clin Endocrinol Metab.* 2003;88:3731–7.
- Ozcan O, Cakir E, Yaman H, Akgul EO, Erturk K, Beyhan Z, et al. The effects of thyroxine replacement on the levels of serum asymmetric dimethylarginine (ADMA) and other biochemical cardiovascular risk markers in patients with subclinical hypothyroidism. *Clin Endocrinol (Oxf).* 2005;63:203–6.
- Aksoy DY, Cinar N, Harmanci A, Karakaya J, Yildiz BO, Usman A, et al. Serum resistin and high sensitive CRP levels in patients with subclinical hypothyroidism before and after L-thyroxine therapy. *Med Sci Monit.* 2013;19:210–5.
- Anagnostis P, Efstathiadou ZA, Slavakis A, Selamatzidou D, Poulasouchidou M, Katergari S, et al. The effect of L-tiroxine substitution on lipid profile, glucose homeostasis, inflammation and coagulation in patients with subclinical hypothyroidism. *Int J Clin Pract.* 2014. doi:10.1111:1–7.
- Bachorik PS, Albers JJ. Precipitation methods for quantification of lipoproteins. *Methods Enzymol.* 1986;129:78–100.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the low-density lipoprotein cholesterol in plasma without the use of preparative ultracentrifuge. *Clin Chem.* 1972;18:499–502.
- Ohkawa H, Ohishi H, Yagi K. Assay for lipid peroxide in animal tissues by thiobarbituric acid reaction. *Anal Biochem.* 1979;95:351–8.
- Ellman GL. Tissue sulfhydryl groups. *Arch Biochem Biophys.* 1959;82:70–7.
- Sassa S. Delta-aminolevulinic acid dehydratase assay. *Enzyme.* 1982;28:133–45.
- Nagasaki T, Inaba M, Shirakawa K, Hiura Y, Tahara H, Hameda Y, et al. Increased levels of C-reactive protein in hypothyroid patients and its correlation with arterial stiffness in the common carotid artery. *Biomed Pharmacother.* 2007;61:167–72.
- Poyrazoglu OK, Ozkan Y, Ozden M, Colak R, Ozalp G, Dönder E. L-thyroxine treatment of patients with subclinical hypothyroidism reduce inflammation. *Open Endocrinol J.* 2009;3:34–7.
- Bilgicir O, Bilgicir F, Calam M, Calam OG, Yuksel A. Comparison of pre- and post-levothyroxine high-sensitivity c-reactive protein and fetuin-a levels in subclinical hypothyroidism. *Clinics.* 2015;70:97–101.

40. Luboshitzky R, Herer P. Cardiovascular risk factors in middle-aged women with subclinical hypothyroidism. *Neuroendocrinol Lett.* 2004;25:262–6.
41. Peleg RK, Efrati S, Benbassat C, Fygenzo M, Golik A. The effect of levothyroxine on arterial stiffness and lipid profile in patients with subclinical hypothyroidism. *Thyroid.* 2008;18:825–30.
42. Drugarin D, Negru S, Korech A, Zosin I, Cristea C. The pattern of a Th1 cytokine in autoimmune thyroiditis. *Immunol Lett.* 2000;71:73–7.
43. Figueroa-Vega N, Alfonso-Pérez M, Benedicto I, Sánchez-Madrid F, González-Amaro R, Marazuela M. Increased circulating pro-inflammatory cytokines and Th17 lymphocytes in Hashimoto's thyroiditis. *J Clin Endocrinol Metab.* 2010;95:953–62.
44. Díez JJ, Hernanz A, Medina S, Bayon C, Iglesias P. Serum concentrations of tumor necrosis factor- α and soluble TNF- α receptor p55 in patients with hypothyroidism and hyperthyroidism before and after treatment. *Clin Endocrinol (Oxf).* 2002;57:515–21.
45. Papanas N, Papazoglou D, Papatheodorou K, Antonoglou C, Kotsiou S, Maltezos E. Thyroxine replacement dose in patients with Hashimoto disease: a potential role for interleukin-6. *Cytokine.* 2006;35:166–70.
46. Karanikas G, John P, Wahl K, Schuetz M, Dudczak R, Wilhelm M. T-Lymphocyte cytokine production patterns in nonimmune severe hypothyroid state and after thyroid hormone replacement therapy. *Thyroid.* 2004;14:488–92.
47. Guclu F, Ozmen B, Kirmaz C, Kafesciler SO, Degirmenci PB, Taneli F, et al. Down-regulation of the auto-aggressive processes in patients with hypothyroidism Hashimoto's thyroiditis following substitutive treatment with L-thyroxine. *Eur Cytokine Netw.* 2009;20:27–32.
48. Tanis BC, Westendorp RGJ, Smelt AHM. Effect of thyroid substitution on hypercholesterolaemia in patients with subclinical hypothyroidism: a reanalysis of intervention studies. *Clin Endocrinol (Oxf).* 1996;44:643–9.
49. Teixeira PF, Reuters VS, Ferreira MM, Almeida CP, Reis FA, Buescu A, et al. Lipid profile in different degrees of hypothyroidism and effects of levothyroxine replacement in mild thyroid failure. *Transl Res.* 2008;151:224–31.
50. Iqbal A, Jorde R, Figenschau Y. Serum lipid levels in relation to serum thyroid-stimulating hormone and the effect of thyroxine treatment on serum lipid levels in subjects with subclinical hypothyroidism: the Tromsø Study. *J Intern Med.* 2006;260:53–61.
51. Razvi S, Ingole L, Keeka G, Oates C, McMillan C, Weaver JU. The beneficial effect of L-thyroxine on cardiovascular risk factors, endothelial function, and quality of life in subclinical hypothyroidism: randomized, crossover trial. *J Clin Endocrinol Metab.* 2007;92:1715–23.
52. Danese MD, Ladenson PW, Meinert CL, Powe NR. Effect of thyroxine therapy on serum lipoproteins in patients with mild thyroid failure: a quantitative review of the literature. *J Clin Endocrinol Metab.* 2000;85:2993–3001.
53. Ineck BA, Ng TMH. Effect of subclinical hypothyroidism and its treatment on serum lipids. *Ann Pharmacoter.* 2003;37:725–30.
54. Villar HC, Saconato H, Valente O, Atallah AN. Thyroid hormone replacement for subclinical hypothyroidism. *Cochrane Database Syst Rev.* 2007;18:CD003419.
55. Bilgir F, Bilgir O, Calan M, Calan O, Isikyakar T. Subclinical hypothyroidism: comparison of adhesion molecule levels before and after levothyroxine therapy. *J Intern Med Res.* 2014;42:806–14.
56. Monzani F, Caraccio N, Kozakowa M, Dardano A, Vittone F, Virdis A, et al. Effect of levothyroxine replacement on lipid profile and intima-media thickness in subclinical hypothyroidism: a double-blind, placebo-controlled study. *J Clin Endocrinol Metab.* 2004;89:2099–106.
57. Rodondi N, Bauer DC, Cappola AR, Cornuz J, Robbins J, Fried LP, et al. Subclinical thyroid dysfunction, cardiac function and the risk of heart failure: the cardiovascular health study. *J Am Coll Cardiol.* 2008;52:1152–9.
58. Razvi S, Weaver JU, Vanderpump MP, Pearce SHS. The incidence of ischemic heart disease and mortality in people with subclinical hypothyroidism: reanalysis of the Wickham Survey Cohort. *J Clin Endocrinol Metab.* 2010;95:1734–40.
59. Razvi S, Weaver JU, Butler TJ, Pearce SHS. Levothyroxine treatment of subclinical hypothyroidism, fatal and nonfatal cardiovascular events, and mortality. *Arch Intern Med.* 2012;172:811–7.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

