



EDITORIAL

Open Access

Spotlight on prostate cancer: the latest evidence and current controversies

Sigrid Carlsson^{1,2} and Andrew Vickers^{3*}

Abstract

Recent decades have seen dramatic changes in the management of prostate cancer based on novel research findings. Prostate-specific antigen (PSA) screening has been introduced, and then recently modified to include new strategies and biomarkers. Management of advanced disease has been transformed by the rapid introduction of new agents. We have moved from a “one-size-fits-all” approach in prostate cancer management to multidisciplinary strategies tailored to the individual patient and his specific cancer. This editorial marks the launch of the article collection *Spotlight on prostate cancer* (<http://www.biomedcentral.com/bmcmed/series/SPR>), and here, guest editors Sigrid Carlsson and Andrew Vickers give an overview of the past, present, and future of prostate cancer research and management.

Keywords: Prostate cancer, Prevention, Screening, Treatment, Risk stratification

Editorial

Introduction

In the nineteenth century, prostate cancer was described as a “very rare disease” [1]. It is now the most common male cancer, and the second or third most common cause of cancer-related death in men in the US and in Europe [2,3]. Our new series, *Spotlight on prostate cancer*, will address a broad range of research and clinical topics related to this common and important public health challenge. We welcome submissions of research articles covering prostate cancer epidemiology and prevention, screening and risk stratification, disease management and new therapies, biomarkers, molecular genomics, and translational studies.

During the past three decades, we have witnessed tremendous shifts in the way that we approach prostate cancer. The introduction of prostate-specific antigen (PSA) screening dramatically altered the presentation of the disease, but we are now moving away from a “one-size-fits-all” approach to new ways of individualizing screening. Better understanding of prostate cancer biology has led to the introduction of several new agents for advanced disease, transforming what had previously

been a relatively barren field. Marker research has rapidly developed, with novel genomic markers now identifying the molecular drivers of aggressive disease. We are less aggressive in our approach to low-risk disease, moving away from radical treatment toward an increased use of active surveillance, while becoming more aggressive in attempting to control high-risk tumors. Instead of administering hormonal treatment alone, we also attack the tumor-invaded organs with surgery or kill the cancer cells with radiation.

Are we on the way to solving the quandary of prostate cancer, eloquently expressed by Dr. Willet Whitmore as: “Is cure necessary in those for whom it is possible, and is cure possible in those for whom it is necessary?”. Every approach involves harms and benefits. Every strategy can be debated. The intricacies and challenges of trying our best to understand and manage this disease continue to puzzle and to fascinate us. This new series, *Spotlight on prostate cancer*, will include commentaries as well as debate and opinion pieces.

Although knowledge of the etiology of prostate cancer is still an area of investigation, evidence of nutritional and dietary targets for primary prevention is accumulating. In the review *Nutrition, dietary interventions and prostate cancer: the latest evidence*, Lin and colleagues provide an overview of recent literature on the possible influences of diet and nutrients on prostate cancer outcomes [4].

* Correspondence: vickersa@mskcc.org

³Department of Epidemiology & Biostatistics, Memorial Sloan Kettering Cancer Center, 485 Lexington Avenue, New York, NY 10017, USA
Full list of author information is available at the end of the article

The controversy surrounding whom to screen, and how best to screen, continues. We have witnessed the rise of screening with PSA. We have observed the rapid and dramatic increased incidence of the disease in the US, followed by a decrease in prostate cancer mortality [2], after the widespread implementation of PSA testing, together with improvements in treatment [5]. We have followed the success of large-scale European screening trials that demonstrate a benefit of screening on prostate cancer mortality [6,7].

Every other year or so we read updated and conflicting guidelines [8], with the United States Preventive Services Task Force (USPSTF), as the outlier, now recommending against the use of routine screening [9].

In the commentary *Prostate-specific antigen based screening: Controversy and guidelines*, Kim and Andriole review the four largest randomized trials of screening and treatment and provide some reasons why these studies yielded apparently conflicting results [10]. Some readers may interpret the results of the screening trials as conflicting; the European ERSPC trial [6] indicating “yes, screening works” and the US-led PLCO trial [11] indicating “no, it does not”. Some could make the same claim about the trials of radical prostatectomy versus watchful waiting; the Scandinavian SPCG-4 trial [12] indicating “yes, radical prostatectomy works” and the US PIVOT trial [13] indicating “no, it does not”.

In our view, however, it is not a question of whether screening and treatment work or do not work. The question is: For which men, and in which situations, might they work, when used appropriately and in a risk-stratified manner?

The settings in which these four studies took place played important roles. While the American population was being heavily screened - essentially changing the question of the PLCO away from an evaluation of the effects of screening versus no screening - the European control population remained largely unscreened, and concomitantly, screening was shown to reduce prostate cancer mortality. Men in the Scandinavian population in SPCG-4 had clinically palpable, higher-risk disease, and surgery was abandoned for men with positive nodes; men in the American population in the PIVOT trial had mainly PSA-detected, lower-risk disease, were older, and surgery was more aggressive.

Owing to the heterogeneity of the disease, risk stratification has become an important aspect of screening, diagnosis, management, and treatment of prostate cancer. With personalized medicine, we attempt to tailor the right treatment, for the right patient, at the right time. Modern management of prostate cancer now includes active surveillance, that is, monitoring the disease and delaying curative intervention until signs of disease progression. Robot-assisted radical prostatectomy is now

more commonly performed than open procedures. Our approach to high-risk disease is multidisciplinary, and we continue to make progress in the treatment of advanced prostate cancer with the advent of new drugs for castration-resistant disease, and finding the most effective sequence in which to administer them. Reports on prostate cancer management and treatment are encouraged and will be welcome in *Spotlight on prostate cancer*.

Competing interests

AV is named on a patent application for a statistical method to detect prostate cancer. The method has been commercialized by OPKO Health. AV receives royalties from sales of the test and has stock options in OPKO Health. SC has no conflicts of interest to disclose.

Authors' contributions

SC and AV wrote the manuscript and approved the final manuscript submitted for publication.

Authors' information

SC and AV are the guest editors of our article collection *Spotlight on prostate cancer*. SC is an associate professor of experimental urology with a Ph.D. in medicine (urology). She has an M.P.H. from Harvard School of Public Health. AV holds a D. Phil. in clinical medicine and specializes in research methodology and statistics. SC and AV have more than 10 years of experience in the study of screening, early detection, and treatment of prostate cancer. Both are on staff at Memorial Sloan Kettering Cancer Center, New York, NY, USA.

Author details

¹Department of Surgery (Urology Service), Memorial Sloan Kettering Cancer Center, New York, NY, USA. ²Department of Urology, Institute of Clinical Sciences, Sahlgrenska Academy at University of Göteborg, Göteborg, Sweden. ³Department of Epidemiology & Biostatistics, Memorial Sloan Kettering Cancer Center, 485 Lexington Avenue, New York, NY 10017, USA.

Received: 4 March 2015 Accepted: 4 March 2015

Published online: 24 March 2015

References

- Adams J. The case of scirrhus of the prostate gland with corresponding affection of the lymphatic glands in the lumbar region and in the pelvis. *Lancet*. 1853;1:393.
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin*. 2015;65:5–29.
- Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JW, Comber H, et al. Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer*. 2013;49:1374–403.
- Lin PH, Aronson W, Freedland SJ. Nutrition, dietary interventions and prostate cancer: the latest evidence. *BMC Medicine*. 2015;13:3. <http://www.biomedcentral.com/1741-7015/13/3>.
- Etzioni R, Tsodikov A, Mariotto A, Szabo A, Falcon S, Wegelin J, et al. Quantifying the role of PSA screening in the US prostate cancer mortality decline. *Cancer Causes Control*. 2008;19:175–81.
- Schröder FH, Hugosson J, Roobol MJ, Tammela TL, Zappa M, Nelen V, et al. Screening and prostate cancer mortality: results of the European Randomised Study of Screening for Prostate Cancer (ERSPC) at 13 years of follow-up. *Lancet*. 2014;384:2027–35.
- Hugosson J, Carlsson S, Aus G, Bergdahl S, Khatami A, Lodding P, et al. Mortality results from the Göteborg randomised population-based prostate-cancer screening trial. *Lancet Oncol*. 2010;11:725–32.
- Loeb S. Guideline of guidelines: prostate cancer screening. *BJU Int*. 2014;114:323–5.
- Moyer VA. U.S. Preventive Services Task Force. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2012;157:120–34.
- Kim EH, Andriole GL. Prostate-specific antigen based screening: controversy and guidelines. *BMC Medicine*. 2015. DOI: 10.1186/s12916-015-0296-5.

11. Andriole GL, Holmberg L, Garmo H, Rider JR, Taari K, Busch C, et al. Prostate cancer screening in the randomized prostate, lung, colorectal, and ovarian cancer screening trial: mortality results after 13 years of follow-up. *J Natl Cancer Inst.* 2012;104:1–8.
12. Bill-Axelson A, Brawer MK, Jones KM, Barry MJ, Aronson WJ, Fox S, et al. Radical prostatectomy or watchful waiting in early prostate cancer. *N Engl J Med.* 2014;370:932–42.
13. Wilt TJ et al. Radical prostatectomy versus observation for localized prostate cancer. *N Engl J Med.* 2012;367:203–13.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

