

CASE REPORT

Open Access

Tocilizumab treatment safety in rheumatoid arthritis in a patient with multiple sclerosis: a case report

Hiroe Sato^{1,2*}, Daisuke Kobayashi¹, Asami Abe¹, Satoshi Ito¹, Hajime Ishikawa¹, Kiyoshi Nakazono¹, Akira Murasawa¹, Takeshi Kuroda², Masaaki Nakano³ and Ichiei Narita²

Abstract

Background: Multiple sclerosis is a relatively rare disease, and complications of multiple sclerosis and rheumatoid arthritis are much rarer. Since anti-tumor necrosis factor therapy increases exacerbations of multiple sclerosis, complications of demyelinating diseases contraindicate anti-tumor necrosis factor therapy. There have been few reports of anti-interleukin-6 receptor therapy for patients with rheumatoid arthritis complicated with multiple sclerosis.

Case presentation: A 53-year-old Japanese woman with multiple sclerosis and rheumatoid arthritis was admitted to our hospital because her rheumatoid arthritis was uncontrolled with oral methotrexate, tacrolimus, and prednisolone. She had developed multiple sclerosis when she was 25 years old and was treated with glucocorticoid therapy. Her multiple sclerosis was in remission for more than 9 years. Because anti-tumour necrosis factor therapy can exacerbate demyelinating disease, the anti-interleukin-6 receptor antibody tocilizumab was started at 8 mg/kg every 4 weeks. At the second administration of tocilizumab, complete remission was achieved. She has remained in remission with tocilizumab without recurrence of multiple sclerosis for more than 5 years.

Conclusion: Anti-interleukin-6 therapy was safely used in this patient with rheumatoid arthritis without exacerbations of multiple sclerosis.

Keywords: Rheumatoid arthritis, Multiple sclerosis, Tocilizumab, Interleukin-6, Tumour necrosis factor

Background

Multiple sclerosis (MS) is an autoimmune, inflammatory, demyelinating disease of the central nervous system characterised by repeated relapses and remissions. The prevalence rate of MS in Japan is reportedly 8 to 9 per 100,000 persons. High-dose glucocorticoid therapy is used for initial and relapsed progression of MS with or without immunosuppressive agents. Although MS is an autoimmune inflammatory disease, as is rheumatoid arthritis (RA), and although the level of tumour necrosis factor (TNF) in cerebrospinal fluid is correlated with the

severity and progression of the disease [1], anti-TNF therapy fails and actually increases exacerbations [2,3]. Therefore, complications of demyelinating diseases contraindicate anti-TNF therapy. There have been few reports of anti-interleukin (IL)-6 receptor therapy for patients with RA complicated with MS.

We herein describe a patient with RA and MS treated with anti-IL-6 receptor therapy.

Case presentation

A 53-year-old Japanese woman was admitted to Niigata Rheumatic Centre, Shibata city, Japan. She had been diagnosed with MS associated with right optic neuritis and thoracic myelitis when she was 25 years old and treated with high-dose prednisolone (PSL). The myelitis had relapsed three times when she was 36, 37 and 40 years old and treated with high-dose PSL. Oligoclonal

* Correspondence: hiroe-212@umin.ac.jp

¹Department of Rheumatology, Niigata Rheumatic Center, 1-2-8 Honcho, Shibata City, Niigata 957-0054, Japan

²Division of Clinical Nephrology and Rheumatology, Niigata University Graduate School of Medical and Dental Sciences, 1-757 Asahimachi-Dori, Chuoku, Niigata City, Niigata 951-8510, Japan

Full list of author information is available at the end of the article

IgG band was found in cerebral spinal fluid (CSF) and IgG and myelin basic protein in CSF were elevated (4.9 mg/dL and 1.2 mg/dL, respectively). Brain T2 weighted magnetic resonance imaging (MRI) showed high intensity area beside left lateral ventricle indicating asymptomatic plaque lesion due to MS. High intensity area was also shown in T2 weighted MRI of cervical spinal cord. Anti-aquaporin 4 antibody was negative. Slight right hemiparesis remained, and she needed a cane to walk outside. The MS achieved remission and PSL was stopped for 9 years. When she was 50 years old, polyarthritis developed, and rheumatoid factor and C-reactive protein (CRP) levels were high. She was diagnosed with RA. The PSL was restarted at 7.5 mg daily and methotrexate (MTX) was begun. Because the MTX could not be increased over 8 mg/week because of mild elevation of transaminases, tacrolimus (3 mg daily; TAC) was added to MTX and leukocyte apheresis was performed. However, the RA activity remained high: the CRP was 2.3 mg/dL and the disease activity score (DAS28ESR) was 4.94 (moderate disease activity). Furthermore, joint space narrowing of both knees and ankles had progressed obviously over 1 year. Because anti-TNF therapy can exacerbate demyelinating disease, the anti-IL-6 receptor antibody tocilizumab (TCZ) was started at 8 mg/kg every 4 weeks. At the second administration of TCZ, the CRP was <0.1 mg/dL and the DAS28ESR was 2.0 (complete remission). The MTX and TAC were tapered and stopped in 6 months, and the PSL was tapered to 0.5 mg daily in 1 year. The health assessment questionnaire disability index (HAQ DI) in 1 year was 1.88 and functional disability was remained. At the 5-year follow-up, she remained in remission with TCZ.

Serum interferon (IFN) γ was negative (≤ 0.1 IU/mL) and serum high sensitivity TNF- α was within normal range (1.6 pg/mL) before starting TCZ therapy. Both of them kept the same levels for a year. Serum IL-6 level was elevated, 51.2 pg/mL (normal range; ≤ 4.0 pg/mL) before starting TCZ therapy and it was 57.1 pg/mL a year later.

Discussion

Complications of MS and RA are rare, and only a few cases have been reported [4,5]. In two case reports, the duration between the onset of each disease was long (8–20 years), and RA or MS was preceded by the other and the two diseases did not flare at the same time [4,5]. Our patient with MS also developed RA 25 years after the onset of MS, which was well controlled without medication for 9 years. Thus, the onset mechanisms of MS and RA are expected to fundamentally differ despite the fact that both are autoimmune inflammatory diseases.

On the other hand, there are several reports of patients with RA who developed demyelinating diseases during anti-TNF therapy [6–8] and patients with MS who developed inflammatory arthritis during IFN- β therapy [9,10]. The mechanisms of demyelinating disease induced by anti-TNF therapy are not clear, but inhibition of TNF leads to IFN- γ production, which is associated with MS [11]. Moreover, TNF polymorphism may be associated with anti-TNF therapy-induced demyelinating diseases [12]. IFN- β therapy for preventing recurrence of relapsing-remitting MS and secondary progressive MS is now widely used. Arthritis reportedly develops during IFN- β therapy in patients with MS, and the HLA phenotype may be involved in its pathogenesis [9]. Elevated levels of IL-6 in the serum in response to IFN- β therapy may also be associated with arthralgia [10].

In comparison, few studies have reported anti-IL-6 therapy in demyelinating disorders. A case report of a 72-year-old woman with leukoencephalopathy that developed in a Phase 3 clinical trial of TCZ for treating RA has been published [13]. However, the relationship between TCZ and leukoencephalopathy was not clear because the possibility of infection had not been completely excluded and the symptoms did not improve after discontinuing TCZ.

Studies involving mouse models of MS (experimental autoimmune encephalomyelitis, EAE) and RA (collagen-induced arthritis, CIA) have revealed that Th-17 plays an important role in the development of both diseases [14,15]. Furthermore, IL-6 mainly affects the differentiation of Th-17 in the mouse, and anti-IL-6 therapy inhibits the onset of EAE and CIA. Meanwhile, TNF affects local inflammation [14,15]. Thus, anti-IL-6 therapy may control EAE and CIA earlier and more radically than anti-TNF therapy. In humans, IL-6 mainly affects the differentiation of Th-17, as in the mouse, and anti-IL-6 therapy should theoretically inhibit both RA and MS. However, the cytokine pathways differ in humans and mice, and further studies are needed to determine whether anti-IL-6 therapy can be used safely in patients with MS.

Conclusion

Here, we reported a patient with RA complicated by MS who was treated with anti-IL-6 therapy for more than 5 years without an exacerbation of the MS. Because anti-TNF therapy can induce and worsen demyelinating diseases, anti-IL-6 therapy is a potential treatment for patients with RA complicated by MS.

Consent

Written informed consent was obtained from the patient for publication of this Case Report. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Abbreviations

RA: Rheumatoid arthritis; MS: Multiple sclerosis; TNF: Tumour necrosis factor; IL: Interleukin; PSL: Prednisolone; CSF: Cerebral spinal fluid; MRI: Magnetic resonance imaging; CRP: C-reactive protein; MTX: Methotrexate; TAC: Tacrolimus; TCZ: Tocilizumab; IFN: Interferon; EAE: Experimental autoimmune encephalomyelitis; CIA: Collagen-induced arthritis.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HS, DK, AA, SI, HI, KN and AM made substantial contributions to conception and acquisition of data and analysis and interpretation of data. TK, MN and IN helped to draft the manuscript. All authors read and approved the final manuscript.

Author details

¹Department of Rheumatology, Niigata Rheumatic Center, 1-2-8 Honcho, Shibata City, Niigata 957-0054, Japan. ²Division of Clinical Nephrology and Rheumatology, Niigata University Graduate School of Medical and Dental Sciences, 1-757 Asahimachi-Dori, Chuoku, Niigata City, Niigata 951-8510, Japan. ³Department of Medical Technology, School of Health Sciences, Faculty of Medicine, Niigata University, 2-746 Asahimachi-Dori, Chuoku, Niigata City 951-8518, Japan.

Received: 30 January 2014 Accepted: 2 September 2014

Published: 12 September 2014

References

- Sharief MK, Hentges R: Association between tumor necrosis factor- α and disease progression in patients with multiple sclerosis. *N Engl J Med* 1991, **325**:467-472.
- The Lenercept Multiple Sclerosis Study Group and The University of British Columbia MS/MRI Analysis Group: TNF neutralization in MS: results of a randomized, placebo-controlled multicenter study. *Neurology* 1999, **53**:457-465.
- van Oosten BW, Barkhof F, Truyen L, Boringa JB, Bertelsmann FW, von Blomberg BM, Woody JN, Hartung HP, Polman CH: Increased MRI activity and immune activation in two multiple sclerosis patients treated with the monoclonal anti-tumor necrosis factor antibody cA2. *Neurology* 1996, **47**:1531-1534.
- Attout H, Toussiot E, Auge B, Chavot D, Wendling D: Rheumatoid arthritis and multiple sclerosis in the same patient. Two case-reports. *Rev Rhum Engl Ed* 1999, **66**:169-172.
- Mpofu S, Moots RJ: A case of multiple sclerosis associated with rheumatoid arthritis and positive anticardiolipin antibodies. *Ann Rheum Dis* 2003, **62**:376.
- Fromont A, De Seze J, Fleury MC, Maillefert JF, Moreau T: Inflammatory demyelinating events following treatment with anti-tumor necrosis factor. *Cytokine* 2009, **45**:55-57.
- Lozeron P, Denier C, Lacroix C, Adams D: Long-term course of demyelinating neuropathies occurring during tumor necrosis factor- α -blocker therapy. *Arch Neurol* 2009, **66**:490-497.
- Tristano AG: Neurological adverse events associated with anti-tumor necrosis factor alpha treatment. *J Neurol* 2010, **257**:1421-1431.
- Levesque MC, Ward FE, Jeffery DR, Weinberg JB: Interferon-beta1A-induced polyarthritis in a patient with the HLA-DRB1*0404 allele. *Arthritis Rheum* 1999, **42**:569-573.
- Nakatsuji Y, Nakano M, Moriya M, Kishigami H, Tatsumi C, Tada S, Sadahiro S, Naka T, Mitani K, Funauchi M, Azuma T, Watanabe S, Kinoshita M, Kajiyama K, Yuasa Y, Kaide M, Takahashi MP, Naba I, Hazama T, Sakoda S, Osaka Neurological Research Consortium: Beneficial effect of interferon-beta treatment in patients with multiple sclerosis is associated with transient increase in serum IL-6 level in response to interferon-beta injection. *Cytokine* 2006, **36**:69-74.
- Mitoma H, Horiuchi T, Hatta N, Tsukamoto H, Harashima S, Kikuchi Y, Otsuka J, Okamura S, Fujita S, Harada M: Infliximab induces potent anti-inflammatory responses by outside-to-inside signals through transmembrane TNF- α . *Gastroenterology* 2005, **128**:376-392.

- Kirk CW, Droogan AG, Hawkins SA, McMillan SA, Nevin NC, Graham CA: Tumour necrosis factor microsattelites show association with multiple sclerosis. *J Neurol Sci* 1997, **147**:21-25.
- Kobayashi K, Okamoto Y, Inoue H, Usui T, Ihara M, Kawamata J, Miki Y, Mimori T, Tomimoto H, Takahashi R: Leukoencephalopathy with cognitive impairment following tocilizumab for the treatment of rheumatoid arthritis (RA). *Intern Med* 2009, **48**:1307-1309.
- Fujimoto M, Serada S, Mihara M, Uchiyama Y, Yoshida H, Koike N, Ohsugi Y, Nishikawa T, Ripley B, Kimura A, Kishimoto T, Naka T: Interleukin-6 blockade suppresses autoimmune arthritis in mice by the inhibition of inflammatory Th17 responses. *Arthritis Rheum* 2008, **58**:3710-3719.
- Serada S, Fujimoto M, Mihara M, Koike N, Ohsugi Y, Nomura S, Yoshida H, Nishikawa T, Terabe F, Ohkawara T, Takahashi T, Ripley B, Kimura A, Kishimoto T, Naka T: IL-6 blockade inhibits the induction of myelin antigen-specific Th17 cells and Th1 cells in experimental autoimmune encephalomyelitis. *Proc Natl Acad Sci U S A* 2008, **105**:9041-9046.

doi:10.1186/1756-0500-7-641

Cite this article as: Sato et al.: Tocilizumab treatment safety in rheumatoid arthritis in a patient with multiple sclerosis: a case report. *BMC Research Notes* 2014 **7**:641.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

