

RESEARCH ARTICLE

Open Access

Deciphering the genetic basis of microcystin tolerance

Anke Schwarzenberger^{1*}, Thomas Sadler¹, Susanne Motameny², Kamel Ben-Khalifa³, Peter Frommolt⁴, Janine Altmüller², Kathryn Konrad² and Eric von Elert¹

Abstract

Background: Cyanobacteria constitute a serious threat to freshwater ecosystems by producing toxic secondary metabolites, e.g. microcystins. These microcystins have been shown to harm livestock, pets and humans and to affect ecosystem service and functioning. Cyanobacterial blooms are increasing worldwide in intensity and frequency due to eutrophication and global warming. However, *Daphnia*, the main grazer of planktonic algae and cyanobacteria, has been shown to be able to suppress bloom-forming cyanobacteria and to adapt to cyanobacteria that produce microcystins. Since *Daphnia*'s genome was published only recently, it is now possible to elucidate the underlying molecular mechanisms of microcystin tolerance of *Daphnia*.

Results: *Daphnia magna* was fed with either a cyanobacterial strain that produces microcystins or its genetically engineered microcystin knock-out mutant. Thus, it was possible to distinguish between effects due to the ingestion of cyanobacteria and effects caused specifically by microcystins. By using RNAseq the differentially expressed genes between the different treatments were analyzed and affected KOG-categories were calculated. Here we show that the expression of transporter genes in *Daphnia* was regulated as a specific response to microcystins. Subsequent qPCR and dietary supplementation with pure microcystin confirmed that the regulation of transporter gene expression was correlated with the tolerance of several *Daphnia* clones.

Conclusions: Here, we were able to identify new candidate genes that specifically respond to microcystins by separating cyanobacterial effects from microcystin effects. The involvement of these candidate genes in tolerance to microcystins was validated by correlating the difference in transporter gene expression with clonal tolerance. Thus, the prevention of microcystin uptake most probably constitutes a key mechanism in the development of tolerance and adaptation of *Daphnia*. With the availability of clear candidate genes, future investigations examining the process of local adaptation of *Daphnia* populations to microcystins are now possible.

Keywords: *Daphnia*, Microcystin, Tolerance, Transcriptome, Molecular basis

Background

One fundamental topic in modern evolutionary ecology is the understanding of the genetic mechanisms underlying adaptation of organisms to changes in their environments. Several attempts to identify the molecular basis of adaptation have already been made for a variety of organisms: e.g. (i) Xia et al. [1] have found differences in nucleotide diversity patterns at drought-related candidate genes in two species of tomatoes indicating local adaptation. (ii) Concerning animals, Feldman et al. [2] found that adaptive

evolution of the garter snake to toxic prey has occurred independently via the de novo acquisition of beneficial mutations in the skeletal muscle sodium channel. (iii) In the British peppered moth *Biston betularia*, a single core sequence has been identified to carry a signature of strong selection that is responsible for industrial melanism [3].

With the increasing availability of next-generation sequencing (NGS) technologies, it is possible to investigate the burning questions regarding mechanisms underlying adaptations also for non-model organisms with a well-understood ecology [4]. The first ecological model organism whose genome has only recently been released is *Daphnia* [5], a globally distributed grazer of algae and cyanobacteria. Its ecoresponsive genome [6,7] and

* Correspondence: Anke.Schwarzenberger@gmx.de

¹University of Cologne, Cologne Biocenter, Aquatic Chemical Ecology, Zuelpicher Str. 47b, 50674 Cologne, Germany

Full list of author information is available at the end of the article

well-studied ability to adapt to many biotic and abiotic factors [8-10] makes it the perfect system with which to study evolutionary and adaptive processes on the genetic level [11].

Especially *Daphnia*'s ability to adapt to cyanobacteria and their toxins has been extensively studied during the last years [8,10,12]. Cyanobacteria negatively affect *Daphnia* by reducing somatic growth [13,14] and inhibit feeding [15]. Also a decline in *Daphnia* biomass due to cyanobacteria has been observed in several field studies [16-18]. However, the generality of this negative correlation between cyanobacterial and *Daphnia* biomass has recently been questioned in an experiment [19] and in several field studies [20,21], demonstrating that *Daphnia* have the potential to adapt to increasingly tolerate dietary cyanobacteria. Cyanobacteria and their toxins are becoming more and more of an ecological threat due to global warming and eutrophication [22], and new solutions for the management of freshwater ecosystems are needed. Therefore, it is a key issue to elucidate the underlying molecular mechanisms of *Daphnia*'s ability to tolerate cyanobacterial toxins and to therefore possibly suppress cyanobacterial blooms [19]. Toxic cyanobacterial secondary metabolites that frequently occur in cyanobacterial blooms are the well-studied microcystins and serine protease inhibitors [23,24]. Both toxin types have been shown to negatively affect *Daphnia* [13,14].

Identifying candidate genes is a major issue in genetics of adaptation. For cyanobacterial protease inhibitors, these candidate genes are easy to determine, as digestive proteases are the direct targets of protease inhibitors. Schwarzenberger et al. [14] found that the tolerance of different *Daphnia* genotypes depended on the residual activity of proteases; increased gene expression and enhanced activity of the non-inhibited protease type also seemed to play a role. Von Elert et al. [25] demonstrated that tolerance to cyanobacterial protease inhibitors was acquired by remodelling the affected digestive protease type.

However, in the case of microcystins, candidate genes are not as easily identified. From *in vitro* studies it has been proposed that protein phosphatases 1 and 2a are direct targets of microcystins in *Daphnia* [26]. However, it remains unclear which major physiological pathways are affected by a putative binding of microcystins to these protein phosphatases, and which of the associated elements cause the difference in tolerance among different *Daphnia* genotypes. Pflugmacher et al. [27] proposed that a glutathione-microcystin conjugate formed *in vitro* by glutathione S-transferase (GST) might be the first step in detoxification of microcystins in *Daphnia*. However, only one of twelve GST genes was found to be up-regulated in response to dietary microcystins in a recently published *D. pulex* microarray study [12], calling the role of GST as a mechanism of tolerance into question. In the same microarray study, several oxidative stress

genes were up-regulated in *D. pulex* after ingestion of a microcystin producing cyanobacterium [12]. Oxidative stress responses have been observed in different aquatic organisms after exposure to microcystins [28]. However, it remains unclear whether the regulation of these genes in *Daphnia* was due to the microcystins or rather a general response to the ingestion of cyanobacteria. It is also not known whether these genes might explain tolerance to microcystins. The ingested cyanobacterium also contained other secondary metabolites; therefore the effects on gene expression could not exclusively be attributed to microcystins in this study [12]. Due to the importance of *Daphnia* as a global grazer of cyanobacteria and its capability to control cyanobacterial blooms, it is now essential to identify the candidate genes that are regulated after ingestion of microcystins. It is also essential to investigate the involvement of these candidate genes in microcystin tolerance by separating the effects of the pure microcystins from general cyanobacterial responses.

Different from Asselman [12], who only used a microcystin-producing cyanobacterium in a *Daphnia* microarray study, we here measured gene expression in transcriptomes of one tolerant clone of *D. magna*, which was fed with either 100% of a green alga or 90% of this alga with 10% of the wild-type strain of *M. aeruginosa* PCC7806 that produces both microcystins and protease inhibitors, or with 10% of its microcystin-free mutant. This cyanobacterial system, which only differed in microcystin-production, allowed disentangling effects on gene expression of *D. magna* due to microcystins from gene expression effects caused by dietary protease inhibitors. We pair-wise determined differentially expressed genes due to the different food sources and calculated significantly affected KOG-categories. From these KOG-categories we chose several candidate genes for microcystin tolerance from the comparison of green algal food/ food mixture with the wild-type strain and green alga/ mutant. By measuring differences in gene expression via qPCR in four *D. magna* clones from two ponds with or without cyanobacteria we were investigating the underlying molecular mechanisms of microcystin tolerance and local adaptation.

Results and discussion

For the transcriptome analyses we chose a clone of *Daphnia magna* (clone A [29]) that has been shown to be relatively tolerant to dietary microcystins and was therefore assumed to have distinct mechanisms for coping with dietary microcystins. In order to avoid differences in juvenile development, growth and reproduction between the cyanobacterial treatments, (which would probably obscure specific effects of microcystins on gene expression) we only added 10% of cyanobacterial carbon to the food mixtures. A significant reduction in somatic

growth rate (one-way ANOVA: $F_{2,9} = 53.46$, $p < 0.05$) was observed in comparison to high quality food, while no differences in *D. magna* growth between microcystin-containing and microcystin-free cyanobacterial food was found (Tukey HSD after one-way ANOVA, $p = 0.84$). This indicates that *D. magna* was still able to cope with such a low concentration of microcystins without additional detectable costs besides costs associated with feeding on cyanobacteria in comparison to control food. For each pair-wise comparison between food treatments, more than 1000 differentially expressed (DE) genes were found: CM: 1178; CW: 1651; MW: 1060 (Figure 1). Due to a very strict database search of *D. magna* against the *D. pulex* genome ($e\text{-value} \leq e^{-20}$, score ≥ 300 , similarity between sequences $\geq 40\%$) only around 20% of these DE genes were assigned to functional genes. This very low number probably also resulted from the fact that a high number of genes might be lineage specific for *D. magna* (comparisons with other species also resulted in less than 25% of DE genes with known functions (Additional file 1: Table S1)). Additionally, some of the DE genes might be improperly predicted by Cufflinks. However, the strictness of the database search ensured a high probability for the correct prediction of gene names and functions of the DE genes. Additionally, this strictness also enables to generalize our results for other *Daphnia* species, since these genes are probably more conserved than DE genes that were excluded. In all comparisons with WT, more DE genes were down-regulated than up-regulated

(binomial tests; $p < 0.05$). Several serine proteases (trypsin and chymotrypsins) were regulated in response to dietary trypsin inhibitors produced both by the wild-type and the mutant of *M. aeruginosa* PCC 7806. This finding corroborates earlier findings with the same *D. magna* clone [14] obtained by qPCR.

The DE genes which could be assigned to *D. pulex* genes were assigned to KOGs and KOG-categories [30]. KOG categories, which were highly significantly affected by dietary microcystins (Table 1), contained genes which were involved in transport processes: intracellular trafficking, secretion and vesicular transport (cat. U) and secondary metabolites biosynthesis transport and catabolism (cat. Q). A total of 64 ATP-binding cassette (ABC) proteins were identified in the *D. pulex* ABC-transporter gene family, and many of them were assigned to drug efflux transporters homologous to other species [31]. It has been suggested that the increased expression of ABC transporters constitutes an adaptive mechanism enabling organisms to be resistant to drugs and toxicants [32]. Here, in *D. magna*, many transporter genes were down-regulated in response to dietary microcystins, which suggests that these transporters are involved in microcystin up-take rather than in excretion. Gene regulation of these transporter genes was probably partly responsible for the observed higher number of down-regulated in comparison to up-regulated genes in the treatments with the microcystin-producing cyanobacterium. However, a permease was up-regulated in *Daphnia*, which is in




Figure 1 Venn diagrams of all DE genes (CM: 1178 genes; CW: 1651; MW: 1060) and genes that could be assigned to gene IDs from the wfleabase (dappu ID; CM: 182 genes; CW: 308; MW: 130) of the three comparisons. CW: all DE genes in the comparison between *D. magna* grown on 100% *C. klinobasis* and *D. magna* grown on 10% wild-type *M. aeruginosa* PCC7806. CM: all DE genes in the comparison between *D. magna* grown on 100% *C. klinobasis* and *D. magna* grown on 10% mutant *M. aeruginosa* PCC7806. MW: all DE genes in the comparison between *D. magna* grown on 10% wild-type *M. aeruginosa* PCC7806 and *D. magna* grown on 10% wild-type *M. aeruginosa* PCC7806. Overlapping planes share DE genes; i.e. these DE genes appear in more than one comparison. The unequal diameters of the circles in the Venn diagrams result from different numbers of DE genes.

Table 1 Gene names, accession numbers from wfleabase.org (for *D. pulex* and *D. magna*) and primer sequences used in qPCR analyses

Gene	<i>D.pulex</i>	<i>D. magna</i>	Primer forward (5'-3')	Primer reverse (5'-3')
transporter of the ABC superfamily	DappuDraft_99518	m8AUGep24bs03373g21t1_ext	TCTCGGGTGGTGAAGGAAA	ATGCCTGGACGATGTTCTGA
multidrug/pheromone exporter	DappuDraft_127589	m8AUGapi5_contig28666g180t1	CGCCATACCAATGTCGCTT	CTGCCTGCCAACATTTGT
permease of the major facilitator superfamily	DappuDraft_303762	m8AUGep24bs00868g315t1_ext	CGGTTCGACTATAACGCTGC	TTCTTTGCCAGGCTTGACAC

accordance with the finding that permeases have been shown to extrude cytotoxic compounds out of yeast cells [33]. Whereas transporter genes were clearly identified as novel candidate genes for microcystin tolerance, no previously suggested candidates from *in vitro* studies (i.e. glutathione synthetase and glutathione S-transferase [27]) were differentially expressed in response to microcystins. The finding that no GST gene was differentially expressed in our transcriptome is in line with Asselman et al. [12] who have found only one GST to be up-regulated. Both findings indicate a minor relevance of conjugation to glutathione for inducible tolerance to microcystins.

Three differentially expressed transporter genes were chosen for candidate gene expression analyses with qPCR. These genes were a transporter of the ABC superfamily (subfamily ABCG), a multidrug/pheromone exporter (subfamily ABCB/MDR) and a permease of the major facilitator

superfamily. Four *D. magna* clones were chosen for this gene expression study: Two clones showed a low tolerance to dietary microcystins (T1 and T2, Figure 2), while the other two clones had a high tolerance to the wild-type strain of *M. aeruginosa* that produces microcystins (Additional file 1: Table S1 and S2, Figure 2). In the qPCR analyses, the ABC superfamily and the multidrug/pheromone exporter genes were significantly down-regulated in three of four clones in response to supplemented pure microcystin LR (Figure 3A and B), which confirmed the results from the transcriptome study. For both genes, one of the clones with low tolerance did not regulate gene expression at all, whereas the two tolerant clones showed the strongest down-regulation of gene expression. Additionally, an expected significant increase in permease gene expression was observed only in one of the tolerant clones (Figure 3C). It is quite likely




Figure 2 Juvenile somatic growth rates of four clones of *D. magna* from two different populations with (T1 and T2; tolerant population) or without (S1 and S2; sensitive population) naturally occurring cyanobacteria. *D. magna* were either grown on 100% *Chlamydomonas klinobasis* or on 90% *C. klinobasis* and 10% of either the wild-type strain of *Microcystis aeruginosa* PCC7806 or its microcystin-free mutant. Different letters indicate significant difference between treatments for each clone separately (Tukey HSD after one-way ANOVA; $p < 0.05$).




Figure 3 Relative gene expression of three transporter genes (A: a transporter of the ABC superfamily, B: a multidrug/pheromone exporter, and C: a permease of the major facilitator superfamily) of four *D. magna* clones from two different populations with (T1 and T2; tolerant population) or without (S1 and S2; sensitive population) naturally occurring cyanobacteria. *D. magna* were grown on 100% *C. klinobasis* with either control liposomes or with liposomes supplemented with microcystin LR. Asterisks indicate significant differences ($p < 0.05$; t-test) in gene expression in relation to gene expression on control liposomes (set to one; horizontal line).

that the decrease in ABC-transporter and multidrug-transporter gene expression leads to lower production of transporter protein, which in turn results in a lower up-take of microcystins into the cells. A higher production

of permeases might on the other hand cause a stronger excretion of microcystins from the cells.

Since the two tolerant clones showed a higher transporter gene regulation in response to supplemented microcystin LR than the other clones, we concluded that the regulation of transporter genes actually plays a role in tolerance to dietary microcystins. Interestingly, both tolerant clones and the clone used in the transcriptome stemmed from the same population with frequent cyanobacterial blooms [24]. Therefore, local adaptation of this population to microcystins due to higher transporter-gene regulation is plausible. Another strong indication of the involvement of transporter genes in adaptation to microcystins is a particularly high number of gene duplications observed in several ABC-transporter subfamilies [31]. For example, gene family expansions in phenotypically important genes have been shown to be responsible for adaptation of insects to insecticides [34]. Future work should therefore address the role of transporter genes in local adaptation of *Daphnia* to microcystins in natural systems, as well as how these adapted populations might be used for the management of lakes with frequent cyanobacterial blooms.

Conclusions

Until recently it has been impossible to identify the genetic mechanisms underlying adaptive traits in non-model organisms. Whereas adaptations to natural environments have been demonstrated in ecologically relevant organisms, the genetic basis of adaptation has mainly been investigated in genetic model organisms whose adaptations to environmental variability are of minor interest. In the case of *Daphnia*, studies identifying genes that account for adaptive traits are scarce, and are based on qPCR analyses of candidate genes which were identified on the basis of *a-priori* knowledge of their function [35,36]. By using a transcriptome-wide approach, we have been able to answer the question as to which genes are involved in microcystin tolerance. Not only did we find new candidate genes, but also demonstrated that the regulation of these genes clearly accounts for the tolerance of *Daphnia* to microcystins. In most adaptation studies, changes in DNA sequences (e.g. mutations [2], SNPs [37], etc.) were identified to constitute the genetic mechanism. Here, we suggest that in some cases adaptation of individuals or populations might arise from the adapted organism's ability to differential express few important genes.

We here confirm that with the knowledge of the ecological background of an organism and a smart experimental design the huge amount of sequences produced in transcriptome approaches can be minimized to few genes with high potential. By releasing new ecoresponsive genomes, with *Daphnia* as a first example, and with the ever-increasing application of NGS approaches, it is

now possible to link well-studied adaptive traits to the discovery of the underlying genetic mechanisms.

Methods

Animal cultivation and food experiment

A clone of *Daphnia magna* (clone A [29]) originating from Lake Bysjön, Sweden (where it coexisted with cyanobacteria), was cultured at 20°C in membrane-filtered tap water. Fifteen animals per litre were kept under non-limiting food concentrations (2 mg C_{part} / l) with the green alga *Chlamydomonas klinobasis* as food alga. Only third clutch neonates which had been born within 12 h were used for the experiment.

The green alga *Chlamydomonas klinobasis* was cultivated semi-continuously at 20°C in a five litre batch culture in cyanophycean medium [38] with one litre exchanged every other day. The cyanobacterium *Microcystis aeruginosa* (UTEX LB 2063 and PCC 7806) and its genetically engineered microcystin synthetase knock-out mutant (PCC 7806 mcy⁻ [39]) were cultivated in chemostat cultures (dilution rate 0.045/ d) in cyanophycean medium at 20°C. The wild-type strain PCC 7806 contains two microcystin variants (RR and LR) as well as strong trypsin inhibitors (cyanopeptolins [40,41]). The two cyanobacterial strains only differed in the content of microcystins, while the amount of trypsin inhibitors was comparable (HPLC data not shown). All food organisms were grown under constant light (cyanobacteria: 95 µE/ m²/ s; *C. klinobasis* 130 µE/ m²/ s). Carbon concentrations (mg C/ ml) of the autotrophic food suspensions were measured for several dilutions per alga and cyanobacterium with an elemental analyser (ThermoFisher Scientific, Inc., Waltham, MA USA), and regression lines were drawn between these concentrations and the photometric light extinction at 470 nm for each dilution. These regression lines served to calculate the volume of each autotrophic food suspension needed for the carbon concentration applied in the experiments.

From a cohort of newborn *D. magna*, 15 animals each were transferred to one litre of membrane-filtered tap water and fed with 2 mg C/ l of either 100% *C. klinobasis* or 90% *C. klinobasis* and 10% of either of the two cyanobacteria. Food and medium were exchanged daily. The experiment was run in triplicate and lasted until all animals reached the size at first reproduction: Somatic growth rates of *D. magna* were determined from the dry weight of animals collected at the start and of five animals per replicate at the day when the first clutch (egg stage one) of the animals was released in the brood chamber (day seven: 100% of *C. klinobasis*; day eleven: 10% *M. aeruginosa*) as according to [42]. Levene's tests were conducted to ensure homogeneity of variances, and growth rates were analyzed with ANOVAs and Tukey's HSD for post-hoc comparisons. The three food treatments will be referred to as C

(for 100% *C. klinobasis*), WT (for 90% *C. klinobasis* and 10% PCC 7806) and Mut (for 90% *C. klinobasis* and 10% PCC 7806 mcy⁻).

Transcriptome: RNA extraction, sequencing and data analyses

At the end of the food experiment, the RNA of ten animals of each of three replicates was extracted using the RNeasy Mini Kit (Qiagen). Only animals bearing eggs at stage one [43] were chosen for RNA extraction. In order to remove any traces of genomic DNA, the RNA was treated with Desoxyribonuclease I (Fermentas) following the manufacturer's instructions. The integrity of the RNA was verified with a 2100 bioanalyzer (Agilent). After reverse transcription and library construction, the cDNA was sequenced in paired-end mode with 36 bp read length on an Illumina Genome Analyzer IIx.

The reads of each sample were aligned to the *Daphnia magna* v. 2.4 reference assembly using TopHat version 1.2.0. By default, TopHat reports up to 40 multiple hits per read if it maps at different positions in the reference genome. Additional file 1: Table S2 provides alignment statistics for every sample. Afterwards, Cufflinks version 1.0.3 was used to identify potential transcripts and quantify their expression. Cufflinks counts reads that have multiple hits in the alignment with reduced but equal weight at each mapping position. Expression values were compared between the three groups of samples using Cuffdiff. Differentially expressed genes and isoforms were identified and FPKM values calculated. The resulting potential transcripts were blasted against the *Daphnia pulex* genome database (wfleabase.org [44]) to assign gene names and functions to the differentially expressed genes. All differentially expressed genes were classified according to eukaryotic orthologous groups [45] and KEGG metabolic pathways [46]. The number of KOGs of each of the three comparisons (either C/WT, or C/Mut, or Mut/WT) were counted and assigned to their specific category. From the whole number of KOGs from each category and the number of differentially expressed KOGs from each comparison, the significantly affected KOG categories between the comparisons were calculated with exact binomial tests (chi²-tests and Fisher's exact tests) using the program Statistica 6.0. Here, we considered highly significant p-values from the chi²-tests to be p < 0.05, medium p-values to be p < 0.1 and low but still significant p-values to be p < 0.2.

Liposome production

Control liposomes were produced according to [25] and subsequently enriched tenfold through centrifugation and subsequent resuspension in 1 ml liposome buffer [47]. For passive loading of the liposomes with microcystin, 500 µl

of this suspension were added to 3 µg MC-LR and incubated for four h at 60°C.

Validation of transcriptome results with qPCR

Four *D. magna* clones (T1, T2, S1 and S2) were chosen for validation of transcriptome results by qPCR. These clones were pre-cultured in the same way as clone A. In single clone experiments, five newborn *D. magna* per each of the three replicates were kept for five days on 100% *C. klinobasis* in 250 ml water. Afterwards, three replicates were fed with 100% *C. klinobasis* and 20 µl concentrated control liposomes [47] for two additional days, while three other replicates were fed with 100% *C. klinobasis* and 20 µl liposomes supplemented with microcystin LR. 20 µl of liposomes supplemented with microcystin should contain 60 ng of MC-LR. Even if 50% were lost in the experiment, this should be similar to the measured concentration used in the transcriptome study with 10% of the wild-type strain of *M. aeruginosa* PCC7806 (120 ng microcystin per litre). Food and medium were exchanged daily. After 48 h, when RNA was extracted, all animals were in the same developmental phase in all treatments.

For qPCR analyses of relative protease gene expression, RNA was extracted using the RNeasy Mini Kit (Qiagen) following the manufacturer's instructions. RNA was purified with DNase I (Fermentas) and immediately reverse transcribed with the High-capacity cDNA Reverse Transcription Kit (ABI). The integrity of the RNA was verified with a NanoDrop. RNA concentrations were determined with a Qubit fluorometer (Invitrogen) as per the manufacturers' instructions. QPCR and data analyses were performed as according to [48] and were close to the MIQE guidelines [49].

QPCR was conducted on a 7300 real time PCR system (Applied Biosystems). Each reaction contained 2.5 ng of cDNA template, 10 µl Power SYBR® Green PCR Master Mix (Applied Biosystems) and 2.5 µM of each primer (Table 1) in a final volume of 20 µl. Each reaction was conducted in biological triplicates. The cycling parameters were 95°C for 10 min (to activate the DNA polymerase) followed by 40 cycles at 95°C for 15 s and at 60°C for 1 min. After the actual analysis, dissociation curves were performed to verify that no primer-dimers had been amplified. For normalization, six different endogenous controls (actin, alpha-tubulin, 18S ribosomal RNA (18S), succinate dehydrogenase (SucDH), TATA-box binding protein (TBP), and ubiquitin-conjugating enzyme (UBC)) were used in qPCR analysis. These endogenous controls were chosen from a given set of ten reference genes [50].

Availability of supporting data

NCBI Sequence Read Archive (SRA) accession numbers for the transcriptome data: Study: SRP045518;

BioProject: PRJNA258118; BioSamples: SAMN02988312, SAMN02988313, SAMN02988314; Runs: SRR1552196, SRR1552197, SRR1552198, SRR1552199, SRR1552200, SRR1552201, SRR1552219, SRR1552220, SRR1552221.

Additional file

Additional file 1: Table S1. Database search on DE genes from the transcriptome. DE genes from the transcriptome were blasted against several databases implemented in NCBI and OrthoDB. The organisms used in these databases were either Arthropoda, Crustacea in general or a set of selected arthropod species (*Daphnia pulex*, *Drosophila melanogaster*, *Tribolium castaneum*, *Ixodes scapularis* and *Apis mellifera*). **Table S2.** TopHat Alignment Statistics. Chlamy: *D. magna* fed with 100% *Chlamydomonas klinobasis*; Mut: *D. magna* fed with 90% *C. klinobasis* and 10% of the microcystin-free mutant strain of *M. aeruginosa* PCC7806; WT: *D. magna* fed with 90% *C. klinobasis* and 10% of the WT strain of *M. aeruginosa* PCC7806.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AS planned and performed all experiments under the supervision of EVE. AS also analyzed the transcriptome data after Illumina sequencing by JA and KK. Bioinformatic calculations were performed by KB-K, PF and SM. TS supported AS and EVE in the interpretation of the results. All authors discussed the results and commented on the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank Dörthe Becker for helpful comments on the data analysis, Lutz Becks for proofreading the manuscript, and Lea von Ganski for help in conducting the experiments. We would also like to thank Frederic Bartlett for English correction. This study was supported by grants from the German Research Foundation (DFG) to E.V.E. (E1179/9-1 and another grant within the Collaborative Research Centre SFB 680 Molecular Basis for Evolutionary Innovations). We are grateful for the access to preliminary sequence data on *Daphnia magna* that were produced by The Center for Genomics and Bioinformatics at Indiana University and distributed via wFleaBase in collaboration with the *Daphnia* Genomics Consortium. The *Daphnia magna* genome project is supported in part by NIH award 5R24GM078274 "Daphnia Functional Genomics Resources" and by the METACyt Initiative of Indiana University, and funded in part through a major grant from the Lilly Endowment, Inc.

Author details

¹University of Cologne, Cologne Biocenter, Aquatic Chemical Ecology, Zuelpicher Str. 47b, 50674 Cologne, Germany. ²University of Cologne, Cologne Center for Genomics, Weyertal 115b, 50931 Cologne, Germany.

³University of Cologne, RRZK, Weyertal 121, 50931 Cologne, Germany.

⁴University of Cologne, CECAD Cologne, Robert-Koch-Str. 21, 50931 Cologne, Germany.

Received: 15 May 2014 Accepted: 29 August 2014

Published: 9 September 2014

References

1. Xia H, Camus-Kulandaivelu L, Stephan W, Tellier A, Zhang Z: Nucleotide diversity patterns of local adaptation at drought-related candidate genes in wild tomatoes. *Mol Ecol* 2010, 19:4144–4154.
2. Feldman CR, Brodie ED, Brodie ED, Pfrender ME: The evolutionary origins of beneficial alleles during the repeated adaptation of garter snakes to deadly prey. *Proc Natl Acad Sci* 2009, 106:13415–13420.
3. Van't Hof AE, Edmonds N, Dalikova M, Marec F, Saccheri IJ: Industrial melanism in British peppered moths has a singular and recent mutational origin. *Science* 2011, 332:958–960.

4. Stapley J, Reger J, Feulner PG, Smadja C, Galindo J, Ekblom R, Bennison C, Ball AD, Beckerman AP, Slate J: **Adaptation genomics: the next generation.** *Trends Ecol Evol* 2010, **25**:705–712.
5. Colbourne JK, Pfrender ME, Gilbert D, Thomas WK, Tucker A, Oakley TH, Tokishita S, Aerts A, Arnold GJ, Basu MK, Bauer DJ, Cáceres CE, Carmel L, Casola C, Choi JH, Detter JC, Dong Q, Dusheyko S, Eads BD, Fröhlich T, Geiler-Samerotte KA, Gerlach D, Hatcher P, Jogdeo S, Krijgsveld J, Kriventseva EV, Kultz D, Laforsch C, Lindquist E, Lopez J, et al: **The ecoresponsive genome of *Daphnia pulex*.** *Science* 2011, **331**:555–561.
6. Jeyasingh PD, Ragavendran A, Paland S, Lopez JA, Sterner RW, Colbourne JK: **How do consumers deal with stoichiometric constraints? Lessons from functional genomics using *Daphnia pulex*.** *Mol Ecol* 2011, **20**:2341–2352.
7. Latta LC, Weider LJ, Colbourne JK, Pfrender ME: **The evolution of salinity tolerance in *Daphnia*: a functional genomics approach.** *Ecol Lett* 2012, **15**:794–802.
8. Sarnelle O, Wilson AE: **Local adaptation of *Daphnia pulicaria* to toxic cyanobacteria.** *Limnol Oceanogr* 2005, **50**:1565–1570.
9. Cousyn C, De Meester L, Colbourne JK, Brendonck L, Verschuren D, Volckaert F: **Rapid, local adaptation of zooplankton behavior to changes in predation pressure in the absence of neutral genetic changes.** *Proc Natl Acad Sci U S A* 2001, **98**:6256–6260.
10. Blom JF, Baumann H, Codd GA, Jüttner F: **Sensitivity and adaptation of aquatic organisms to oscillapeptin J and [D-Asp3(E)-Dhb7]microcystin-RR.** *Archiv fuer Hydrobiologie* 2006, **167**:547–559.
11. Miner BG, De Meester L, Pfrender ME, Lampert W, Hairston NG: **Linking genes to communities and ecosystems: *Daphnia* as an ecogenomic model.** *Proc Roy Soc Lond B Biol Sci* 2013, **279**:1873–1882.
12. Asselman J, De Coninck DIM, Glaholt S, Colbourne JK, Janssen CR, Shaw JR, De Schampheleire KAC: **Identification of pathways, gene networks, and paralogous gene families in *Daphnia pulex* responding to exposure to the toxic cyanobacterium *Microcystis aeruginosa*.** *Environ Sci Technol* 2012, **46**:8448–8457.
13. DeMott WR: **Foraging strategies and growth inhibition in five daphniids feeding on mixtures of a toxic cyanobacterium and a green alga.** *Freshwat Biol* 1999, **42**:263–274.
14. Schwarzenberger A, Küster CJ, Von Elert E: **Molecular mechanisms of tolerance to cyanobacterial protease inhibitors revealed by clonal differences in *Daphnia magna*** *Mol Ecol* 2012, **21**(19):4898–4911. doi: 10.1111/j.1365-294X.2012.05753.x
15. Lürling M: **Effects of microcystin-free and microcystin-containing strains of the cyanobacterium *Microcystis aeruginosa* on growth of the grazer *Daphnia magna*.** *Environ Toxicol* 2003, **18**:202–210.
16. Threlkeld ST: **Midsummer dynamics of 2 *Daphnia* species in Wintergreen Lake, Michigan.** *Ecology* 1979, **60**:165–179.
17. Ghadouani A, Pinel-Alloul B, Prepas EE: **Effects of experimentally induced cyanobacterial blooms on crustacean zooplankton communities.** *Freshwat Biol* 2003, **48**:363–381.
18. Hansson LA, Gustafsson S, Rengefors K, Bomark L: **Cyanoabacterial chemical warfare affects zooplankton community composition.** *Freshwat Biol* 2007, **52**:1290–1301.
19. Sarnelle O: **Initial conditions mediate the interaction between *Daphnia* and bloom-forming cyanobacteria.** *Limnol Oceanogr* 2007, **52**:2120–2127.
20. Hairston NG, Holtmeier CL, Lampert W, Weider LJ, Post DM, Fischer JM, Cáceres CE, Fox JA, Gaedke U: **Natural selection for grazer resistance to toxic cyanobacteria: evolution of phenotypic plasticity?** *Evolution* 2001, **55**:2203–2214.
21. Gustafsson S, Hansson LA: **Development of tolerance against toxic cyanobacteria in *Daphnia*.** *Aquat Ecol* 2004, **38**:37–44.
22. Smith VH, Schindler DE: **Eutrophication science: where do we go from here?** *Trends Ecol Evol* 2009, **24**:201–207.
23. Agrawal MK, Bagchi D, Bagchi SN: **Acute inhibition of protease and suppression of growth in zooplankton, *Moina macrocopa*, by *Microcystis* blooms collected in Central India.** *Hydrobiologia* 2001, **464**:37–44.
24. Schwarzenberger A, D'Hondt S, Vyverman W, Von Elert E (Eds): **Seasonal Succession of Cyanobacterial Protease Inhibitors and *Daphnia magna* Genotypes in A Eutrophic Swedish Lake;** 2013. in press.
25. Von Elert E, Zitt A, Schwarzenberger A: **Inducible tolerance in *Daphnia magna* to dietary protease inhibitors.** *J Exp Biol* 2012, **215**:2051–2059.
26. DeMott WR, Dhawale S: **Inhibition of in-vitro protein phosphatase-activity in three zooplankton species by microcystin-Ir, a toxin from cyanobacteria.** *Arch Hydrobiol* 1995, **134**:417–424.
27. Pflugmacher S, Wiegand C, Oberemm A, Beattie KA, Krause E, Codd GA, Steinberg CEW: **Identification of an enzymatically formed glutathione conjugate of the cyanobacterial hepatotoxin microcystin-LR: the first step of detoxification.** *Biochim Biophys Acta* 1998, **1425**:527–533.
28. Amado LL, Monserrat JM: **Oxidative stress generation by microcystins in aquatic animals: why and how.** *Environ Int* 2010, **36**:226–235.
29. Gustafsson S: **Zooplankton Response to Cyanotoxins,** Ph.D. thesis. Lund University; 2007.
30. Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiyutin B, Koonin EV, Krylov DM, Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS, Smirnov S, Sverdlov AV, Vasudevan S, Wolf YI, Yin JJ, Natale DA: **The COG database: an updated version includes eukaryotes.** *BMC Bioinformatics* 2003, **4**. doi:10.1186/1471-2105-4-41.
31. Sturm A, Cunningham P, Dean M: **The ABC transporter gene family of *Daphnia pulex*.** *BMC Genomics* 2009, **10**:170.
32. Sturm A, George SS, Dean M, Cunningham P, Treuner-Freeman A: **ABC transporters in the *Daphnia pulex* genome: implications for ecotoxicology and drug resistance in crustacean parasites.** *Comp Biochem Physiol Mol Integr Physiol* 2009, **153A**:S109.
33. Takacova M, Imrichova D, Cernicka J, Gbelska Y, Subik J: **KNQ1, a *Kluyveromyces lactis* gene encoding a drug efflux permease.** *Curr Genet* 2004, **45**:1–8.
34. Mouches C, Pasteur N, Berge JB, Hyrien O, Raymond M, Desaintvincent BR, Desilvestri M, Georgiou GP: **Amplification of an esterase gene is responsible for insecticide resistance in a California *Culex* mosquito.** *Science* 1986, **233**:778–780.
35. Scoville AG, Pfrender ME: **Phenotypic plasticity facilitates recurrent rapid adaptation to introduced predators.** *Proc Natl Acad Sci U S A* 2010, **107**:4260–4263.
36. Schwarzenberger A, Von Elert E: **Cyanobacterial protease inhibitors lead to maternal transfer of increased protease gene expression in *Daphnia*.** *Oecologia* 2013, **172**:11–20.
37. Brown E, Pilkington J, Nussey D, Watt K, Hayward A, Tucker R, Graham A, Paterson S, Beraldi D, Pemberton J, Slate J: **Detecting genes for variation in parasite burden and immunological traits in a wild population: testing the candidate gene approach.** *Mol Ecol* 2013, **22**:757–773.
38. Von Elert E, Jüttner F: **Phosphorus limitation not light controls the exudation of allelopathic compounds by *Trichormus doliolum*.** *Limnol Oceanogr* 1997, **42**:1796–1802.
39. Dittmann E, Neilan BA, Erhard M, Von Doehren H, Börner T: **Insertional mutagenesis of a peptide synthetase gene that is responsible for hepatotoxin production in the cyanobacterium *Microcystis aeruginosa* PCC 7806.** *Mol Microbiol* 1997, **26**:779–787.
40. Agrawal MK, Zitt A, Bagchi D, Weckesser J, Bagchi SN, Von Elert E: **Characterization of proteases in guts of *Daphnia magna* and their inhibition by *Microcystis aeruginosa* PCC 7806.** *Environ Toxicol* 2005, **20**:314–322.
41. Martin C, Oberer L, Ino T, Koenig WA, Busch M, Weckesser J: **Cyanopeptolins, new depsipeptides from the cyanobacterium *Microcystis* sp. PCC 7806.** *J Antibiot* 1993, **46**:1550–1556.
42. Wacker A, Von Elert E: **Polyunsaturated fatty acids: evidence for non-substitutable biochemical resources in *Daphnia galeata*.** *Ecology* 2001, **82**:2507–2520.
43. Threlkeld ST (E): **Estimating cladoceran birth rates: the importance of egg mortality and the egg age distribution.** *Limnol Oceanogr* 1979, **24**:601–612.
44. Colbourne JK: **wFleaBase: the *Daphnia* genome database.** *BMC Bioinformatics* 2005, **6**:45.
45. Koonin EV, Fedorova ND, Jackson JD, Jacobs AR, Krylov DM, Makarova KS, Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS, Rogozin IB, Smirnov S, Sorokin AV, Sverdlov AV, Vasudevan S, Wolf YI JJ, Natale DA: **A comprehensive evolutionary classification of proteins encoded in complete eukaryotic genomes.** *Genome Biol* 2004, **5**:R7.
46. Kanehisa M, Araki M, Goto S, Hattori M, Hirakawa M, Itoh M, Katayama T, Kawashima S, Okuda S, Tokimatsu T, Yamanishi Y: **KEGG for linking genomes to life and the environment.** *Nucl Acids Res* 2008, **36**:D480–D484.
47. Martin-Creuzburg D, Von Elert E, Hoffmann KH: **Nutritional constraints at the cyanobacteria-*Daphnia magna* interface: the role of sterols.** *Limnol Oceanogr* 2008, **53**:456–468.
48. Schwarzenberger A, Courts C, Von Elert E: **Target gene approaches: gene expression in *Daphnia magna* exposed to predator-borne kairomones or to microcystin-producing and microcystin-free *Microcystis aeruginosa*.** *BMC Genomics* 2009, **10**:527.

49. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT: The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 2009, 55:611–622.
50. Heckmann LH, Connon R, Hutchinson TH, Maund SJ, Sibly RM, Callaghan A: Expression of target and reference genes in *Daphnia magna* exposed to ibuprofen. *BMC Genomics* 2006, 7:175–182.

doi:10.1186/1471-2164-15-776

Cite this article as: Schwarzenberger et al.: Deciphering the genetic basis of microcystin tolerance. *BMC Genomics* 2014 15:776.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

