

Effects of Reiki Versus Physiotherapy on Relieving Lower Back Pain and Improving Activities Daily Living of Patients With Intervertebral Disc Hernia

Journal of Evidence-Based Integrative Medicine

Volume 23: 1-5

© The Author(s) 2018

Reprints and permission:

sagepub.com/journalsPermissions.nav

DOI: 10.1177/2515690X18762745

journals.sagepub.com/home/cam



Farnaz Jahantigh, BSN¹, Abdolghani Abdollahimohammad, PhD²,
Mohammadreza Firouzkouhi, PhD², and Vahid Ebrahiminejad, MD³

Abstract

Patients with intervertebral disc herniation (IVDH) seek complementary and conventional medical therapies to manage related problems. This study aimed to determine the effectiveness of Reiki compared with physiotherapy to relieve the lower back pain intensity and to improve the activities of daily living (ADL) in the IDVH patients. In this clinical trial study, 60 patients with IVDH were randomly assigned to one of the Reiki, physiotherapy, and drug therapy groups. The severity of pain and the ADL were measured using visual analog scale (VAS) pain and ADL-Instrumental ADL questionnaire before and after the intervention. A significant difference was found in pain intensity and ADL improvement between Reiki and the drug therapy. However, there was no significant difference between Reiki and physiotherapy groups in managing pain and improving ADL. Reiki and physiotherapy are effective methods in managing pain and improving ADL in patients with IVDH; however, Reiki is more cost-effective and faster treatment method than physiotherapy.

Keywords

energy therapy, intervertebral disc hernia, activities of daily living, lower back pain, physical therapy

Received December 20, 2017. Accepted for publication February 6, 2018.

An intervertebral disc herniation (IVDH) is one of the causes of low back pain.^{1,2} A chronic lumbar pain may last for at least 12 weeks.³ This annoying pain leads to physical and psychological disturbances as well as financial problems, which have negative effects on the family and society.⁴ Back pain refrains people from doing their daily activities due to fear of pain recurrence, and this may increase their disabilities.⁵

Different methods such as drug therapy, physiotherapy, surgery, and complementary and alternative procedures are used to relieve back pain. Pharmaceutical treatments are prescribed for pain management, although some drugs may produce both dependence and tolerance.⁶

Physiotherapy is known as an effective method in improving the quality of life and reducing the lower back pain using heat therapy and low-frequency vibrations at the lumbar region, ultrasound, abdominal and lower back exercises based on patients' status.^{3,7}

Surgical treatment is recommended when conventional or preservative therapies are ineffective and the root of the nerve is under pressure.⁸ Common surgical procedures for IVDH include laminectomy, discectomy, microdiscectomy, artificial disc surgery, and spinal fusion.⁹

The use of complementary therapies as nonsurgical procedures for the treatment of chronic pain is increasing.¹⁰ Reiki is one of the therapies, which was approved by the National Center for Complementary and Alternative Medicine for pain relieving. Reiki is categorized as a biofield treatment.¹¹ Reiki can balance and alignment the energy chakras and auras; hence, promoting health in the individuals. One of the nursing theorists, Martha E. Rogers, described the theory of energy fields, and she argued that there are energy fields in human beings and their environments and when the energy flow in humans is

¹ Student Research Committee, Zabol University of Medical Sciences, Zabol, Iran

² Faculty of Nursing and Midwifery, Zabol University of Medical Sciences, Zabol, Iran

³ Khatam Al-Anbia Hospital, Zahedan University of Medical Sciences, Zahedan, Iran

Corresponding Author:

Abdolghani Abdollahimohammad, PhD, Faculty of Nursing and Midwifery, Zabol University of Medical Sciences, Zabol, Iran.

Email: abdolqani@gmail.com



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

disrupted, energy therapy is used to rebuild these disrupted energy fields.¹² Energy therapists use direct touch or distance healing to realignment the energy fields of individuals to promote physical, emotional, mental, and spiritual recovery.¹³ Reiki energy therapy has been applied for treating many physical, emotional, and psychological disorders such as blood pressure, pain, headache, mood disorders, anxiety, osteoarthritis, wound healing, and sleep disturbances.¹²

As the patients with IVDH experience severe pain, which affects their activities of daily living (ADL) and there is also a paucity of research on the effect of distance Reiki on pain management, this study aimed to determine the effectiveness of distance Reiki versus physiotherapy on lower back pain and the ADL of patients with IVDH.

Methods

This clinical trial study was conducted to compare the effects of Reiki energy therapy with physiotherapy on the managing of low back pain and the improving daily activities in patients with IVDH in Physiotherapy Clinic of Khatam Al-Anbia Hospital, Zahedan, Iran.

The inclusion criteria were willingness to participate in the study, being literate, and having a herniated disc between the lumbar vertebra confirmed by an orthopedic surgeon using computed tomography or magnetic resonance imaging. Exclusion criteria were having other spinal diseases (lumbar stenosis, relocation of lumbosacral spine stenosis), length discrepancy in the lower extremities, acute lumbosacral muscle spasm, kidney disorders, pelvic diseases, drug addiction, pregnancy, and having experience of energy therapy.

Data were collected using a demographic questionnaire, visual analog scale (VAS) pain, and ADL-Instrumental ADL (ADL-IADL) questionnaire to assess the degree of independence and dependence on ADL. Demographic questionnaire includes age, sex, marital status, educational level, occupation, duration of illness, number of hospital admissions due to low back pain, history of other diseases, except for IVDH, and the drug addiction. The VAS was used to assess the pain before and after the intervention, and scoring was 0 (no pain) to 10 (the highest pain intensity). The ADL-IADL questionnaire has 25 items, and is rated using a 4-point scale, from 1 (no), 2 (with help), 3 (only and hardly), and to 4 (lonely).¹⁴⁻¹⁷

Sampling was carried out from July to October 2017. Of 63 patients, 3 were excluded due to other spinal diseases (a patient) and unwilling to fill in the check-up questionnaires (2 patients). The patients were randomly assigned to 3 groups, including Reiki (20 patients), physiotherapy (20 patients), or drug therapy (20 patients). The patients were matched based on gender and age (Figure 1).

The Reiki group received three 15-minute distance energy-healing sessions on consecutive days by a master of Reiki. The physiotherapy group underwent 7 to 10 sessions of physiotherapy for 60 to 90 minutes using heat therapy, transcutaneous electrical nerve stimulation, pelvic traction, and physical exercises under supervision a physiotherapist. Indomethacin capsule 75 mg and methocarbamol tablet 500 mg every 8 hours daily for a week was prescribed for the drug group as well as Reiki and physiotherapy groups.

The questionnaires were completed before and one week after the intervention. The data were analyzed by SPSS version 23 using chi-square, independent *t* test, analysis of variance (ANOVA), and repeated-measures ANOVA.

Ethical approval was obtained from the Zabol University of Medical Sciences, Zabol, Iran. Written consent was obtained through a cover letter before asking participants to fill in the questionnaires.

Results

The demographic features (Table 1) show that the majority of participants (60%) were female, married (>90%), and had a higher education level ($\geq 45\%$). The average age of patients in the Reiki (45.10 years), physiotherapy (42.45 years), and drug therapy (48.50 years) groups were not significantly different among the study groups ($P = .246$). There was no significant difference in demographic data among the study groups (all P s > .05).

Table 2 shows the comparison of the mean scores of pain intensity and ADL in the study groups before and after interventions using ANOVA and repeated-measures ANOVA tests. Pain intensity and ADL were not significantly different among the study groups before the interventions ($P > .05$). However, Pain was significantly relieved in physiotherapy and Reiki groups. Cohen's *f* showed a large effect size for Reiki versus physiotherapy as well as drug therapy for managing pain. Besides, there was no significant difference in pain intensity after treatment using physiotherapy and Reiki. However, the severity of pain was significantly different between the drug therapy and Reiki ($P = .002$).

There was no significant difference in the ADL between the Reiki and the physiotherapy groups posttreatment, but a significant difference was found in the ADL improvement between the Reiki and the drug therapy groups ($P = .011$). Cohen's *f* revealed a large effect size for Reiki versus drug therapy and moderate ones for Reiki versus physiotherapy as well as physiotherapy versus drug therapy (Table 2).

Discussion

The aim of this study was to determine the effect of Reiki energy therapy compared with physiotherapy on relieving lumbar pain and on the ADL of patients with IVDH and improvement of their lives. The results showed that Reiki reduces the severity of back pain and improve the level of activities among patients with IVDH. The results showed that there was no significant difference in the improvement of daily activities in Reiki and physiotherapy groups, but in this regard, the Reiki group had a significant difference with drug therapy, so that Reiki was more effective in improving the activities compared with drug therapy. Pain relief in Reiki group was more than that of the physiotherapy and drug therapy groups, which indicates that this method is more effective in controlling pain and improving the daily activities in the patients with IVDH.

Back pain is associated with physical, psychological, emotional, social changes, and even inappropriate exercise.^{3,5,18} Pain as an emotional state could be regulated by the artificial stimulation of autonomic nervous system. Vagus nerve plays an important role in physical and mental well-being based on polyvagal theory.¹⁹ Therefore, stimulation of vagus reduces the




Figure 1. Samples allocation.

perception of pain and improves mood and quality of life in the patients with chronic health problems. The effectiveness of Reiki in managing pain could be explained by polyvagal theory. On the other hand, according to the Ki theory of energy therapy, Reiki opens and balances the centers and the path of energy flow in individuals and improves physical, psychological, and emotional problems.²⁰ Studies show that Reiki reduces pain²¹ and improves the quality of life.^{11,13,22-24}

Although studies are varying in the effectiveness of distance Reiki, the present study showed a positive effect of distance

Reiki on pain and ADL in patients with IVHD, which is consistent with the previous studies.^{13,21,23,24}

Although the present study compared 3 different interventions and showed a significant improvement in the pain relief and ADL in patients with IVDH, it had some limitations, which were beyond the control of the researchers. The pain perception may be affected by ambient factors such as environmental noise.²¹ Despite the recommendation of lying or sitting down in a comfortable position and quiet place during distance Reiki, some patients had not found such a place in their house. Some

Table 1. Frequency, Mean, and Standard Deviation of Demographic Data.

Demographics	Reiki	Physiotherapy	Drug Therapy	P
Age, years, mean (SD)	45.10 (13.07)	42.45 (9.64)	48.50 (10.95)	.426
Gender, n (%)				1.00
Female	12 (60)	12 (60)	12 (60)	
Male	8 (40)	8 (40)	8 (40)	
Marital status, n (%)				.766
Single	2 (10)	0 (0)	1 (5)	
Married	18 (90)	20 (100)	20 (100)	
Education level, n (%)				.484
Less than diploma	9 (45)	7 (35)	4 (20)	
Diploma	2 (10)	3 (15)	2 (10)	
Higher education	9 (45)	10 (50)	14 (70)	
Job, n (%)				.926
Housewife	8 (40)	9 (45)	9 (45)	
Employee	7 (35)	8 (40)	8 (40)	
Free job	5 (25)	3 (15)	3 (15)	

Table 2. Mean, Standard Deviation, Marginal Mean, 95% Confidence Interval, and Effect Size of the Study Groups Before and After Interventions.

Groups	Pain			Activities of Daily Living		
	Before, Mean (SD)	After, Mean (SD)	Marginal Means (95% CI)	Before, Mean (SD)	After, Mean (SD)	Marginal Means (95% CI)
^a Reiki	8.20 (1.36)	3.80 (1.47)	6.00 (5.35, 6.65)	62.20 (14.24)	77.60 (9.96)	69.90 (65.16, 74.64)
^b Physiotherapy	7.75 (1.48)	5.60 (1.85)	6.67 (6.02, 7.33)	55.10 (13.14)	70.45 (11.37)	62.77 (58.03, 67.52)
^c Drug	8.40 (1.53)	7.00 (1.68)	7.70 (7.05, 8.35)	58.50 (12.45)	61.50 (8.75)	59.67 (54.93, 64.42)
<i>F</i> (<i>df</i> = 2, 57)	1.04	18.33	6.94	1.67	13.67	4.90
<i>P</i>	.361	<.001	<.002	.076	<.001	<.011
		<i>c</i> > <i>b</i> > <i>a</i>	<i>c</i> > <i>a</i>		<i>c</i> < <i>a</i> , <i>b</i>	<i>c</i> < <i>a</i>
	Effect size (<i>f</i>):			Effect size (<i>f</i>):		
	Reiki vs physiotherapy = 0.44			Reiki vs physiotherapy = 0.29		
	Reiki vs drug = 0.78			Reiki vs drug = 0.65		
	Physiotherapy vs Drug = 0.34			Physiotherapy vs drug = 0.36		

Abbreviations: SD, standard deviation; CI, confidence interval; *df*, degrees of freedom.

patients refused to perform daily activities regularly due to fear of pain recurrence in the drug therapy group. The other limitation was no blindness in the involved groups.

Conclusion

Reiki as one of the complementary methods can be used by nurses for managing pain and improving ADL. Reiki is a cost-effective, safe, and without known side effects can be used with other therapeutics. In future studies, blindness of study populations and adding a sham group as the control group is recommended.

Acknowledgments

The authors appreciate and thank the staff at the Physiotherapy Center of Khatam al-Anbia Hospital of Zahedan, Iran and the patients who participated in the study.

Author Contributions

FJ and AA wrote the preliminary draft and contributed toward data gathering and first idea of starting this project. AA also performed the Reiki therapy. MF rewrote the draft and contributed toward writing the final version of the article. VE contributed toward data gathering, guidance, and correction of the article.


Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Abdolghani Abdollahimohammad, PhD  <http://orcid.org/0000-0002-7929-5539>

Mohammadreza Firouzkouhi, PhD  <http://orcid.org/0000-0002-5122-195X>

Ethical Approval

Ethical approval (ZBMU.1.REC.1396.36) was obtained from the Ethical Committee of Zabol University of Medical Sciences, Zabol, Iran. Written consent was obtained through a cover letter before asking participants to fill in the questionnaires.

References

- Adams MA, Dolan P. Lumbar intervertebral disk injury, herniation and degeneration. In: Pinheiro-Franco JL, Vaccaro AR, Benzel EC, Mayer HM, eds. *Advanced Concepts in Lumbar Degenerative Disk Disease*. Berlin, Germany: Springer; 2016: 23-39.
- Hinkle JL, Cheever KH. *Brunner and Suddarth's Textbook of Medical-Surgical Nursing*. 13th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2014.
- Chan CW, Mok NW, Yeung EW. Aerobic exercise training in addition to conventional physiotherapy for chronic low back pain: a randomized controlled trial. *Arch Phys Med Rehabil*. 2011;92: 1681-1685.
- Gordon R, Bloxham S. A systematic review of the effects of exercise and physical activity on non-specific chronic low back pain. *Healthcare (Basel)*. 2016;4:22.
- Bunzli S, Gillham D, Esterman A. Physiotherapy-provided operant conditioning in the management of low back pain disability: a systematic review. *Physiother Res Int*. 2011;16:4-19.
- Ross GB, Mavor M, Brown SH, Graham RB. The effects of experimentally induced low back pain on spine rotational stiffness and local dynamic stability. *Ann Biomedical Eng*. 2015;43:2120-2130.
- Magalhães MO, Comachio J, Ferreira PH, Pappas E, Marques AP. Effectiveness of graded activity versus physiotherapy in patients with chronic nonspecific low back pain: midterm follow up results of a randomized controlled trial. *Braz J Phys Ther*. 2018;22:82-91.
- Deyo RA, Mirza SK. Clinical practice. Herniated lumbar intervertebral disk. *N Engl J Med*. 2016;374:1763-1772.
- Yang AJ, Coronado RA, Hoffecker L, et al. Conservative care in lumbar spine surgery trials: a descriptive literature review. *Arch Phys Med Rehabil*. 2017;98:165-172.
- Murthy V, Sibbritt DW, Adams J. An integrative review of complementary and alternative medicine use for back pain: a focus on prevalence, reasons for use, influential factors, self-perceived effectiveness, and communication. *Spine J*. 2015;15:1870-1883.
- Thrane S, Cohen SM. Effect of Reiki therapy on pain and anxiety in adults: an in-depth literature review of randomized trials with effect size calculations. *Pain Manag Nurs*. 2014;15:897-908.
- Aust MP. *Complementary and Alternative Therapies in Nursing*. 7th ed. New York, NY: Springer; 2014.
- Fazzino DL, Griffin MTQ, McNulty SR, Fitzpatrick JJ. Energy healing and pain: a review of the literature. *Holist Nurs Pract*. 2010;24:79-88.
- Haghani H, Nikpour S, Sola AH. Examining health promotion behaviors and quality of life in the elderly. *J Ardabil Univ Med Sci*. 1387;8:29-36.
- Johnson N, Barion A, Rademaker A, Rehkemper G, Weintraub S. The activities of daily living questionnaire: a validation study in patients with dementia. *Alzheimer Dis Assoc Disord*. 2004;18: 223-230.
- Ogon M, Krismer M, Söllner W, Kantner-Rumplmair W, Lampe A. Chronic low back pain measurement with visual analogue scales in different settings. *Pain*. 1996;64:425-428.
- Price DD, McGrath PA, Rafii A, Buckingham B. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. *Pain*. 1983;17:45-56.
- McGirt MJ, Bydon M, Archer KR, et al. An analysis from the quality outcomes database, part 1. Disability, quality of life, and pain outcomes following lumbar spine surgery: predicting likely individual patient outcomes for shared decision-making. *J Neurosurg Spine*. 2017;27:357-369.
- McManus DE. Reiki is better than placebo and has broad potential as a complementary health therapy. *J Evid Based Complementary Altern Med*. 2017;22:1051-1057.
- Midilli TS, Eser I. Effects of reiki on post-cesarean delivery pain, anxiety, and hemodynamic parameters: a randomized, controlled clinical trial. *Pain Manag Nurs*. 2015;16:388-399.
- Shaybak E, Abdollahimohammad A, Rahnama M, Masinaeizhad N, Azadi-Ahmadabadi C, Firouzkohi M. The effect of reiki energy healing on CABG postoperative chest pain caused by coughing and deep breathing. *Indian J Public Health Res Dev*. 2017;8:305.
- Alarcão Z, Fonseca JRS. The effect of Reiki therapy on quality of life of patients with blood cancer: results from a randomized controlled trial. *Eur J Integrat Med*. 2016;8:239-249.
- Demir M, Can G, Kelam A, Aydinler A. Effects of distant reiki on pain, anxiety and fatigue in oncology patients in turkey: a pilot study. *Asian Pac J Cancer Prev*. 2015;16:4859-4862.
- Baldwin AL, Fullmer K, Schwartz GE. Comparison of physical therapy with energy healing for improving range of motion in subjects with restricted shoulder mobility. *Evid Based Complement Alternat Med*. 2013;2013:329731.