

The effect of ascorbic acid and H₂O₂ treatment on the stability of anthocyanin pigments in berries

Elham NIKKHAH^{1,*}, Masoud KHAIAMY¹, Reza HEIDARY¹, Ali Sorayaie AZAR²

¹Department of Biology, Faculty of Science, Urmia University, IRAN

²Islamic Azad University, Maragheh Branch, IRAN

Received: 13.05.2008

Abstract: Anthocyanins are natural pigments widely distributed in nature. Anthocyanin pigment molecules are a subclass of flavonoids. They are responsible for the red, purple, and blue colors observed in many flowers, fruits, and vegetables. Fruits and berries are the main sources of anthocyanins in nature. Anthocyanins are thought to contribute to the nutritive value of fruits and berries due to their antioxidative, anti-carcinogenic, anti-inflammatory, and anti-angiogenic properties. Anthocyanins can also improve the nutritional value of processed foods by preventing the oxidation of lipids and proteins. As such, identification of agents that can affect the stability of anthocyanins and the protective effect of anthocyanins is very important. In the present study anthocyanin pigment was extracted from 3 different berries (*Morus nigra* L., *Morus alba* var. *nigra*, and *Fragaria* L.). After soaking and wetting in ethanol (1% acidified), the extracted anthocyanin pigments were exposed to 3 different concentrations of ascorbic acid (AA) (10%, 25%, and 50%) and H₂O₂ (9.31, 18.61, and 27.92 mmol/L). Six groups of anthocyanin solutions were refrigerated and kept in darkness for 63 days, and every 3 weeks anthocyanin absorbance was recorded at 526 nm. AA absorbance decreased relative to the blank in all the treated samples. These results indicate the destructive effect of AA on anthocyanins. In the samples treated with H₂O₂ anthocyanin degradation increased and the intensity of color decreased as the concentration of H₂O₂ increased.

Key words: Anthocyanin, ascorbic acid, H₂O₂, degradation, berries

Dutlarda anthosiyenin pigmentlerinin stabilitesi üzerine askorbik asit ve H₂O₂ davranışının etkisi

Özet: Anthosiyeninler, doğada geniş çapta dağılmış doğal pigmentlerdir. Anthosiyenin renk molekülleri flavonoidlerin bir alt sınıfıdır. Bunlar birçok çilek, meyve ve sebzelerde kırmızı, mor ve mavi renklerden sorumludur. Meyve ve çilekler doğada anthosiyeninlerin ana kaynağıdır. Anthosiyeninlerin anti-oksidatif, anti-kanserojen, anti-inflamatuvar ve anti-anjiyogenetik bakımından meyve ve çileklerin sağlığa katkıda bulunduğu düşünülmektedir. Anthosiyeninler aynı zamanda gıda ürünlerinde lipit ve proteinlerin oksidasyonunu engelleyerek işlenmiş yiyeceklerin besin değerini artırabilmektedir. Böylece bazı etkenlerin tanımı, bunların anthosiyenin stabilitesi ve anthosiyenin kararlılığında etkili olması bakımından çok önemlidir. Bu çalışmada anthosiyenin pigment üç farklı çilekten (*Morus nigra* L., *Morus alba* var. *nigra* ve *Fragaria* L.) ekstre edildi. Etanolde (% 1 asidite) emdirilerek ve daha sonra ısıtılarak ekstre edilen anthosiyen pigmentler askorbik asit (% 10, % 25 ve % 50) ve H₂O₂ (9,31, 18,61 ve 27,92 mmol/L) nin üç farklı konsantrasyonuna maruz bırakıldı. Anthosiyenin altı çözeltisi karanlıkta ve 63 gün ve her hafta soğutucuda tutuldu, anthosiyenin miktarı 526 nm kaydedildi. Tamamıyla, AA durumunda tüm örneklerde absorbans miktarı blankla bağlantılı olarak azaldı. Bu sonuçlarla anthosiyenine askorbik asidin yıkıcı etkisi gösterildi. H₂O₂ durumunda, H₂O₂ 'nun yüksek konsantrasyonunda, anthosiyenin miktarı azalması yüksektir ve renk yoğunluğu düşüktür.

Anahtar sözcükler: Anthosiyenin, askorbik acid, H₂O₂, indirgeme, dut

Introduction

Anthocyanins belong to flavonoid groups and are responsible for the attractive colors—ranging from red to blue—of flowers and fruits (1). Interest in the field of anthocyanin chemistry has been generated by restricted and limited use of synthetic dyes as food ingredients. Because of low toxicity, anthocyanins have great potential as food coloring and have replaced synthetic red dyes. Recently, anthocyanins have been reported to have pharmacological effects, such as lowering the atherogenic index (2), and lowering triglyceride and free fatty acid levels (3). Moreover, Kamei et al. reported that anthocyanins are more effective in inhibiting the growth of tumor cells than other flavonoids (4).

Nonetheless, many commercial limitations exist for the use of anthocyanin extracts in food products, including low stability, which is influenced by pH, temperature, oxygen, light, polymeric forms, and concentration, and the presence of phenolic compounds and some chemical structures (5). Some anthocyanins are more stable than others, depending on their molecular structure. One example is malvidin, a major anthocyanin in grapes that is more stable than other anthocyanins due to dimethyl oxylation of the molecule (6). Hydroxylation of organic acids, in most cases, makes anthocyanins more stable molecules (7,8). Anthocyanins have a C₆C₃C₆-skeleton typical of flavonoids. They are glycosylated polyhydroxy and polymethoxy derivatives of the 2-phenylbenzopyrylium cation, i.e. the flavylium cation. The main constituent of anthocyanins is aglycone, the flavylium cation, which contains conjugated double bonds responsible for absorption of light at around 500 nm, causing the pigments to appear red to the human eye. Aglycones are referred to as anthocyanidins, which are usually penta- (3,5,7,3',4') or hexa-substituted (3,5,7,3',4',5').

Currently, 22 different anthocyanidins are known, but only 6 are commonly used in foods. The most important anthocyanidins are pelargonidin, cyanidin, peonidin, delphinidin, malvidin, and petunidin. These aglycones differ in the number of hydroxyl and

methoxyl groups in the B-ring of the flavylium cation (5). One of the general methods for combating oxidation and increasing the nutritional value of food products is the use of fruit juices with ascorbic acid (AA), but studies have shown that the presence of AA has a negative effect on anthocyanin stability, leading to the mutual degradation of these compounds (9-11). Additionally, hydrogen peroxide (H₂O₂) has been used in foods and food packaging materials for various purposes in many European countries for over 30 years. H₂O₂ is the most commonly used packaging sterilant in aseptic processing systems (12). As berries contain large amounts of anthocyanins and transformation of these pigments into other forms by variety of agents, such as enzymes, oxidation, light, temperature, and some chemicals, during storage causes their color to change from red to brown, which has a negative impact on the appearance of products and diminishes their usefulness, the aim of the present study was to investigate the effects of AA and H₂O₂ on the color appearance and color stability of fruit juice.

Materials and methods

Sample preparation

Samples of berries were obtained locally. These samples included *Morus nigra*, *Morus alba* var. *nigra*, and *Fragaria* L. The berries were washed with distilled water and kept frozen at -18 °C until use. The experiments were performed in 2007 at the biochemistry lab of Urmia University, Iran.

Extraction of anthocyanins

Extraction was carried out according to Chiriboga and Francis (13). Briefly, after taking the samples out of the freezer, they were left at room temperature for 30 min to defrost. Then 1000 g of each sample was placed in a mixer and after adding ethanol solvent was mixed for 10 min. Next, the products were filtered in a Büchner funnel vacuum and Whatman filter paper (grade 1); the remains of each mixture left on the filter paper was again washed with the above-mentioned solvent and filtered. The quantity of solvent used was sufficient to completely remove all color from the

berries, leaving a clear liquid. Then the filtered product was placed in a balloon container within a vacuum evaporator at 35 °C to separate the ethanol-acid solvent. The balloon container was separated from the vacuum evaporator and distilled water was added to dissolve the concentrated extract that formed at the bottom of the balloon container. The product was then transferred to a 1000-mL container, brought to the volume of 1000 mL using distilled water, and was centrifuged at 8000 rpm. The supernatant was separated and kept for further analysis.

Treatment with ascorbic acid (AA)

To examine the effect of AA, 3 concentrations of AA (10%, 25%, and 50%) were selected. Anthocyanin extracts (90 mL) were poured into 3 groups of test tubes, each with 3 replicates. At first, the pH of the anthocyanin extracts was regulated with a pH meter (pH 2). Subsequently, different concentrations of AA were added to the anthocyanin extracts. The 3 groups of anthocyanin extracts were refrigerated and kept in darkness for 63 days, and every 3 weeks the anthocyanin absorbance at 526 nm was recorded. The selected doses and wavelength were selected according to Duangmal et al. (14).

Treatment with hydrogen peroxide (H₂O₂)

To examine the effect of H₂O₂, 3 concentrations (9.31, 18.61, and 27.92 µmol/L) were selected. Anthocyanin extracts (90 mL) were poured into 3 groups of test tubes, each with 3 replicates. At first, the pH of the anthocyanin extracts was regulated with a pH meter (pH 2). Subsequently, different concentrations of H₂O₂ were added to the anthocyanin extracts. The 3 groups of anthocyanin extracts were refrigerated and kept in darkness for 63 days, and every 3 weeks anthocyanin absorbance at 520 nm was recorded. The selected doses and wavelength were selected according to Özkan et al. (15).

Statistical analysis

Statistical analysis of the data was performed with ANOVA using Microsoft SAS. All experiments were repeated 3 times.

Results and discussion

The effect of ascorbic acid (AA)

The results of statistical analysis show that during the 63-day experimental period the greatest value of absorbance was at zero time, whereas the lowest absorbance was linked to secondary time in *Morus nigra*, and third time in *Morus alba* var. *nigra* and *Fragaria* L. (Figure 1). These results highlight the destructive effect of time on the stability of anthocyanins. In the case of concentration, there was a significant difference between the primary concentration and other concentrations. In total, the greatest value of absorbance occurred with zero concentration (blank) and the lowest absorbance occurred with the primary concentration. Nonetheless, as the AA concentration increased absorbance gradually increased relative to the primary concentration. In all the samples absorbance decreased relative to the blank; therefore, results of this study suggest that AA had a destructive effect on the anthocyanins.

Other studies have shown that AA plays various roles in color stability (16,17). Poei-Langston and Wrolstad observed that addition of AA to a model system of anthocyanins resulted in the loss of pigment stability (11). Skrede et al. also reported that AA caused a decrease in pigment stability (18). Moreover, Marti et al. reported that decomposition of anthocyanin accelerated in the presence of AA (11,19). Duangmal et al. reported that fortification with vitamin C (AA) reduced the half-life values of pigments. They have observed that vitamin C promoted anthocyanin degradation (14). Duangmal et al. reported marked destruction of anthocyanins and vitamin C in Mao juice during storage. Statistical analysis showed that mean anthocyanin absorbance in the presence of various concentrations of AA significantly differed (14). The presence of AA was observed to have a negative impact on anthocyanin stability, leading to the mutual degradation of these compounds (9,10,20).

The results of the present study are in accord with all of the above-mentioned studies. These results




Figure 1. Changes in the anthocyanin content of extracts treated with AA during storage at 0, 21, 42, and 63 days. According to the 3 graphs, anthocyanin absorbance decreased as the concentration of AA increased, relative to the blank (the mean \pm SE of 3 measurements).

might have been due to AA's enhancing polymer pigment formation and bleaching anthocyanin pigments. Direct condensation between anthocyanins and AA has been suggested as a mechanism for anthocyanin degradation (11). The mechanism proposed by Jurd (21) for the degradation of anthocyanins in the presence of AA, which was subsequently supported by Pœi-Langston and Wrolstad (11), is the direct condensation of AA on the carbon 4 of the anthocyanin molecule, causing the loss of both. On the other hand, according to Iacobucci and Sweeny, the loss of anthocyanin color in the presence of AA occurs due to oxidative cleavage of the pyrylium ring by a free radical mechanism in which AA acts as a molecular oxygen activator and

produces free radicals (22). One of these free radicals is H₂O₂. Formation of H₂O₂ due to AA oxidation affects anthocyanin stability (16,17) and leads to a decrease in red color (9). As AA and its derivatives are used in many foods (including fruit juices) to improve their nutritional quality and to prevent enzymatic browning reactions (23), and because of its potent antioxidant capacity by acting as a singlet oxygen quencher (24), use of this substance in some cases is inevitable. Nonetheless, there are some methods for decreasing AA's negative effects on anthocyanin stability. Shrikhande and Francis reported that the presence of flavonol exerts a protective effect with respect to the degradation of anthocyanins in the presence of AA—probably by competing with

anthocyanins in preference for condensation reactions (25)—and also observed that the stability of acylated anthocyanins increases in the presence of AA (26). It seems that AA protects anthocyanins from enzymatic degradation (17). Sapers and Simons (27), and Sapers et al. reported that the stability of acylated anthocyanins increases in the presence of AA and that AA protects anthocyanins from enzymatic destruction (17).

The effect of H_2O_2

In the present study higher concentrations of H_2O_2 increased anthocyanin degradation and decreased the intensity of color. The results of statistical analysis show that during the 63-day study period, absorbance in the 3 groups of anthocyanins (each with a different

concentration of H_2O_2) was significantly different, as in the above-mentioned reports. Based on our statistical analysis, there were significant differences between the means of various times, from 0 to 3. With time, absorbance and anthocyanin content gradually decreased. This result was observed in 3 samples (Figure 2). Moreover, there were significant differences between various concentration means and H_2O_2 concentrations. Anthocyanin degradation increased and, as a result, absorbance decreased. In all samples the highest level of absorbance was at zero concentration and the lowest was at the 3rd concentration (27.92 mmol/L).

One of the chemical agents that affect the stability of anthocyanins is H_2O_2 . The deleterious effect of




Figure 2. Changes in the anthocyanin content of extracts treated with H_2O_2 during storage at 0, 21, 42, and 63 days. Based on the 3 graphs, anthocyanin absorbance decreased as the concentration of H_2O_2 increased, relative to blank (mean \pm SE of 3 measurements).

H₂O₂ on anthocyanin stability in fruit juices is well known. Degradation of anthocyanins in the presence of H₂O₂ has been demonstrated in strawberry (29), sour cherry juice (30), and in orange, grape, and pomegranate juice (31). Different sensitivities to various concentrations of H₂O₂ have been reported (27). Sapers and Simmons observed rapid decolorization of strawberry, raspberry, and cherry anthocyanins in the presence of H₂O₂. They used H₂O₂ as a surface sterilizer in sweet cherries, raspberries, and strawberries, and observed rapid bleaching of anthocyanins. Statistical analysis shows that the concentrations of anthocyanins in the presence of various H₂O₂ concentrations were significantly different, and in the presence of high concentrations of H₂O₂ destruction of anthocyanins was rapid, as was observed in the present study (27).

The susceptibility of anthocyanins to H₂O₂ has been known for a long time. Sondheimer and Kertesz were among the first to investigate the kinetics of anthocyanin degradation by H₂O₂ in both strawberry juice and in a pure solution of the major strawberry anthocyanin (pelargonidin-3-glucoside) (29). According to their research, oxidative degradation of anthocyanins occurs in 2 steps: an initial reversible reaction with the formation of anthocyanin-H₂O₂ adduct, followed by a slower irreversible one. It was shown that the decomposition and dissociation products of H₂O₂ are responsible for the oxidation and subsequent degradation of phenolic compounds (27,28). In fact, De et al. reported that the °OH radical is the main reactive species to cleave the benzene ring in phenolic compounds, and degrades the substrate into CO₂ and H₂O (32). Von Elbe and Schwartz reported that quinones, formed by the oxidation of phenols, also have deleterious effects on anthocyanins

(33). Thus, 2 factors can primarily affect the degradation of anthocyanins by H₂O₂ in fruit juices, which generally contain copious amounts of phenolic compounds: (a) the amount of free radicals and the HOO anion formed by the decomposition and dissociation of H₂O₂, respectively; and (b) the amount of quinones formed by the H₂O₂-catalyzed oxidation of phenolic compounds (33).

Conclusion

Attractive color is one of the most important sensory characteristics of fruit and berry products; however, the red color of berry products is unstable and susceptible to degradation. Maintaining a strong and stable color in berry wines and juices is problematic during processing and storage. In the present study we observed that there was marked destruction of anthocyanins in berry juices containing H₂O₂ during storage. Destruction of anthocyanins in berry juices containing AA was also observed, but with less intensity. As such, aseptic systems should be frequently controlled to ensure the effective removal of residual H₂O₂ from food contact surfaces and fortification of aseptically packed anthocyanin-rich fruit juices with AA should be avoided or carried out very carefully.

Corresponding author:

Elham NIKKHAH

Department of Biology,

Faculty of Science,

Urmia University,

IRAN

E-mail: tu8084@yahoo.com

References

1. Strack D, Wray V. The anthocyanins. In: Harborne JB. Ed. The Flavonoids: Advances in Research since 1986. Chapman and Hall, London; 1993: pp. 1-19.
2. Igarashi K, Abe S, Sato J. Effects of Atsumi-Kabu (red turnip, *Brassicacampestris* L.) Anthocyanin on serum cholesterol levels in cholesterol-fed rats. *Agric Biol Chem* 54: 171-175, 1990.
3. Igarashi K, Inagaki K. Effects of the major anthocyanin of wild grape (*Vitis coignetiae*) on serum lipid levels in rats. *Agric Biol Chem* 55: 285-287, 1991.
4. Kamei H, Kojima T, Hasegawa M et al. Suppression of tumor cell growth by anthocyanins in vitro. *Cancer Investig* 13: 590-594, 1995.

5. Rein M. Copigmentation reactions and color stability of berry anthocyanins, PhD, Academic Dissertation, University of Helsinki, Department of Applied Chemistry and Microbiology, Food Chemistry Division, 2005.
6. Bridle P, Timberlake CF. Anthocyanins as natural food colors-selected aspects. *Food Chem* 58: 103-109, 1997.
7. Bassa IA, Francis FG. Stability of anthocyanins from sweet potatoes in a model beverage. A research note. *J Food Science* 52: 1753-1754, 1987.
8. Francis F. Food colorants: Anthocyanins. *Critical Reviews in Food Science and Nutrition* 28: 273-314, 1989.
9. Brenes CH, Del Pozo-Insfran D, Talcott S. Stability of copigmented Anthocyanins and ascorbic acid in a grape juice model system. *Journal of Agricultural and Food Chemistry* 53: 49-56, 2005.
10. Garzon GA, Wrolstad ER. Comparison of the stability of pelargonidin based anthocyanins in strawberry juice and concentrate. *Journal of Food Science* 67 (4): 1288-1299, 2002.
11. Poesi-Langston MS, Wrolstad RE. Color degradation in an ascorbic acid anthocyanin flavanol model system. *J Food Sci* 46: 1218-1236, 1981.
12. Özkan M. Degradation of anthocyanins in sour cherry and pomegranate juices by hydrogen peroxide in the presence of added ascorbic acid. *Food Chemistry* 78: 499-504, 2002.
13. Chiriboga C, Francis FJ. An anthocyanin recovery system from cranberry pomace. *J Am Soc Hort Sci* 9: 223. 1970.
14. Duangmal K, Wongsiri S, Sueprasarn S. Color appearance of fruit juice affected by vitamin C. Color and Paints, Interim Meeting of the International Color Association, Proceedings: 121-124, 2004.
15. Özkan M, Yemenicioglu A, Asef N et al. Degradation kinetics of anthocyanins from sour cherry, pomegranate and strawberry juices by hydrogen peroxide. *Journal of Food Science* 67(2): 525-529, 2002.
16. Markakis P. Stability of anthocyanins in foods. In: Markakis P. ed. *Anthocyanins as food colors*. Academic Press, New York; 1982: pp. 163-180.
17. Talcott ST, Brines CH, Piers DM et al. Photochemical stability and color retention of co pigmented and processed Muscatine grape juice. *J Agric Food Chem*. 51: 957-963, 2003.
18. Skrede G, Wrolstad RE, Lea P et al. Color stability of strawberry and black currant syrups. *J Food Sci* 57: 172-177, 1992.
19. Marti N, Perez-Vicente A, Garcia-Viguera C. Influence of storage temperature and ascorbic acid addition on pomegranate juice. *J Sci Food Agric* 82: 217-221, 2002.
20. Rodriguez-Saona LE, Giusti MM, Wrolstad RE. Color and pigment stability of red radish and red-fleshed potato anthocyanins in juice model systems. *Journal of Food Science* 64: 451-456, 1999.
21. Jurd L. Some advances in the chemistry of anthocyanins-type plant pigments. In: Chichester CO. Ed. *The chemistry of plant pigments*. New York: Academic Press; 1972: pp. 123-142.
22. Iacobucci GA, Sweeny JG. The chemistry of anthocyanins and related flavylium salts. *Tetrahedron* 39: 3005-3038, 1983.
23. Freedman L, Francis FJ. Effect of ascorbic acid on color of jellies. *Journal of Food Science* 49(4): 1212-1213, 1984.
24. Elliott JG. Application of antioxidant vitamins in foods and beverages. *Food Technology* 53(2): 46-48, 1999.
25. Shrikhande AJ, Francis FJ. Effect of flavonols on ascorbic acid and anthocyanin stability in model systems. *J Food Sci* 39: 904-906, 1974.
26. Del Pozo-Insfran D, Brenes CH, Talcott ST. Phytochemical composition and pigment stability of acai (*Euterpe oleracea* Mart.). *J Agric Food Chem* 52: 1539-1545, 2004.
27. Sapers GM, Simmons G. Hydrogen peroxide disinfection of minimally processed fruits and vegetables. *Food Technology* 52: 48-52, 1998.
28. Sapers GM, Miller RL, Choi SW et al. Structure and composition of mushrooms as affected by hydrogen peroxide wash. *J Food Sci* 64: 889-892, 1999.
29. Sondheimer E, Kertesz ZI. The kinetics of the oxidation of strawberry anthocyanin by hydrogen peroxide. *Food Research* 17: 288-298, 1952.
30. Özkan M, Yemenicioglu A, Citak B et al. Effect of hydrogen peroxide on sour cherry anthocyanins. *J Food Qual* 23: 421-428, 2000.
31. Özkan M, Kirca A, Cemeroglu B. Effects of hydrogen peroxide on the stability of ascorbic acid during storage in various fruit juices. *Food Chemistry* 88: 591-597, 2004.
32. De Ancos B, Cano MP, Hernandez A, Monreal M. Effects of microwave heating on pigment composition and color of fruit purees. *J Sci Food Agric* 79: 663-670, 1999.
33. Von Elbe JH, Schwartz SJ. Colorants. In: Fennema OR. Ed. *Food chemistry*. 3rd Ed. New York, Marcel Dekker; 1996: pp. 651-722.