

## The Effects of Oral Administration of an Aqueous Extract of *Ficus bengalensis* Stem Bark on Some Hematological and Biochemical Parameters in Rats with Streptozotocin-Induced Diabetes

Mahalingam GAYATHRI, Krishnan KANNABIRAN

Biomolecules and Genetics Division, School of Biotechnology, Chemical, and Biomedical Engineering  
VIT University, Vellore-632 014, Tamil Nadu - INDIA

Received: 13.05.2008

**Abstract:** The effects of sub-chronic administration of an aqueous extract of *Ficus bengalensis* stem bark on hematological and biochemical parameters in rats with streptozotocin (STZ)-induced diabetes were evaluated. The possible protective effect of the *F. bengalensis* (500 mg/kg of body weight per day) extract on STZ-induced changes was evaluated over 12 weeks. The parameters evaluated were plasma proteins, including total protein albumin and globulin, non-protein nitrogenous substances, including urea, uric acid, and creatinine, and hematological indices, including total hemoglobin content, erythrocytes, leucocytes, and platelets counts. The aqueous extract had a significant ( $P < 0.05$ ) protective effect against STZ-induced changes in terms of the markers of protein metabolism and hematological indices. Furthermore, increases in levels of non-protein nitrogenous substances were normalized in response to the administration of the extract. Thus, the aqueous extract of *F. bengalensis* had a significant protective effect on STZ-induced changes to biochemical and hematological parameters in the diabetic rats.

**Key Words:** *Ficus bengalensis*, streptozotocin, non-protein nitrogenous substances, hematological indices

### Oral Yolla Uygulanan *Ficus bengalensis* Gövde Kabuğu Sıvı Ekstresinin Streptozotosin ile İndüklenmiş Diyabetik Ratların Bazı Hematolojik ve Biyokimyasal Parametreleri Üzerine Etkisi

**Özet:** Bu çalışmada *Ficus bengalensis* gövde kabuğu sıvı ekstresinin sub kronik uygulamasının streptozotosin (STZ) ile indüklenmiş diyabetik ratların hematolojik ve biyokimyasal parametrelerine etkisi değerlendirilmiştir. STZ ile indüklenmişdeki değişimlere *F. bengalensis* (500 mg/kg vücut ağırlığı/gün) ekstraktının olası koruyucu etkisi 12 hafta için test edilmiştir. Diyabetik ratlarda parametre olarak total protein, albumin, globulini kapsayan plazma proteinleri, üre, ürik asit, kreatinini kapsayan protein olmayan azotlu bileşikler ve toplam hemoglobin miktarı, eritrosit, lökosit, trombosit miktarını kapsayan hematolojik veriler değerlendirilmiştir. Protein metabolizması ve hematolojik veriler ışığında STZ ile indüklenmiş değişimlere karşı sıvı ekstrakt önemli ( $P < 0.05$ ) koruyucu etki göstermiştir. Ayrıca protein olmayan azotlu maddelerin artan düzeyleri ekstrakt uygulaması ile normal düzeylere indirilmiştir. Bu nedenle *F. bengalensis*'in sıvı ekstresi, ratlarda STZ ile indüklenmiş değişimlerde biyokimyasal ve hematolojik parametreler üzerine üstün koruyucu etkiye sahiptir.

**Anahtar Sözcükler:** *Ficus bengalensis*, streptozotocin, protein olmayan azotlu bileşikler, hematolojik veriler

### Introduction

Streptozotocin (STZ) is often used to induce diabetes mellitus in animals due to its toxic effects on pancreatic beta cells (1). STZ is known for its acute cytotoxic effect on cells and molecules, which results in the generation of reactive oxygen species that cause oxidative damage (2) by

depleting the antioxidant defense system in experimental animals (3). Furthermore, STZ is a selective beta-cell genotoxicant; upon administration of a single dose STZ rapidly produces acute diabetes by generating DNA adducts, including N3-methyladenine and O(6)-methylguanine adducts, leading to subsequent beta-cell

death by necrosis (4). It was previously reported that high doses of STZ produce marked steatosis in the liver and tubular injury in the kidneys, as well as increased blood urea nitrogen and creatinine levels in experimental animals (5). STZ produces dose-dependent mortality in experimental animals and was reported to cause 15%-20% mortality at a dose of 65 mg/kg of body weight (6). *Ficus bengalensis* (Banyan tree) is a large tree with many aerial roots that grows wild in the lower region of the Himalayas and throughout India. Different parts of the tree are reported to possess medicinal properties. The leaves are good for ulcers, aerial roots are useful in the treatment of gonorrhea, and seeds and fruit are known for their cooling effect and are often used as a tonic (7). A water extract of the bark of this plant was shown to have a hypoglycemic effect (8), as well as hypocholesterolemic and hypolipidemic effects (9,10).

The objective of the present study was to investigate the possible protective effect of an aqueous extract *F. bengalensis* on STZ-induced changes in levels of total protein, albumin, and non-protein nitrogenous substances, as well as hematological indices in diabetic rats.

## Materials and Methods

### Animals

Male albino rats (Wistar strain, 150-200 g) were purchased from Tamil Nadu Veterinary and Animal Sciences University, Madhavaram, Chennai. The rats were housed under standard husbandry conditions ( $30 \pm 2$  °C, 60%-70% relative humidity, and 12:12 h day/night cycle), and were allowed standard pellet rat feed and water ad libitum. Animal experiments were designed and conducted in accordance with the guidelines of the Institutional Animal Ethical Committee (IAEC, VIT University).

### Preparation of Plant Extract

The bark of *F. bengalensis* was collected from the Morappur Forest, Dharmapuri District, Tamil Nadu, during April 2006. A voucher specimen was prepared and submitted to the Forest Department. The bark was washed with distilled water, shade dried, powdered, and stored in an air-tight container until further use.

The aqueous extract was prepared as described by Vats et al. (11). *F. bengalensis* stem bark (100 g) was powdered in an electric grinder. The powder was then soaked in 500 ml of sterile water, stirred intermittently,

and left overnight. The macerated pulp was then filtered through a coarse sieve and the filtrate was dried at reduced temperature. This dry mass served as the aqueous extract of *F. bengalensis* used for experimentation. To increase the shelf life and uniformity of the extract, it was completely lyophilized by a continuous freeze drying operation for 54 h (Christ freeze dryer, Germany) and the yield was calculated as 3.7% (w/v).

### Induction of Diabetes Mellitus

A single dose of a freshly prepared solution (35 mg/kg of bodyweight in 0.1 M cold citrate buffer, pH 4.5) of STZ (Sigma, USA) was injected (intraperitoneally) into the rats. After 72 h, blood was collected from the tail vein under ether anesthesia. Blood glucose levels were determined using a Microlab 2000 autoanalyzer (Hamilton, UK). The animals were considered to be diabetic if blood glucose values were > 250 mg/dl and only those animals were used for the study. STZ-induced diabetes developed and stabilized over a period of 7 days (12). Untreated control rats were given 0.1 M citrate buffer (pH 4.5) only.

### Experimental Design

Animals were divided into 4 groups, each consisting of 6 animals. Group 1 served as a control, group 2 included rats with STZ-induced diabetes, group 3 served as a positive control and received the standard hypoglycemic drug, tolbutamide (100 mg/kg of body weight), and group 4 included rats with STZ-induced diabetes that were treated with the aqueous extract (500 mg/kg of body weight per day) for 12 weeks via oral intubation. Rats were sacrificed at the end of 12 weeks and blood samples were collected to estimate plasma proteins, non-protein nitrogenous substances, and hematological indices.

### Biochemical and hematological Parameters

Total hemoglobin was estimated by the cyanmethemoglobin method (13). Plasma total protein was estimated by the method of Lowry et al. (14) and albumin by the method of Peters et al. (15). Plasma urea and uric acid were estimated by the method of Tomas (16), and creatinine was estimated by the method of Fossati et al. (17). Hematological indices, including erythrocytes count, leucocytes count, and platelets count, were also determined using standard methods (18).

### Acute Toxicity

The *F. bengalensis* aqueous extract was tested for acute toxicity by administering oral doses ranging from 100 mg

to 1 g/kg of body weight per day and from 2 to 5 g/kg of body weight per day for 7 weeks to 2 groups of rats (n = 6) prior to the present investigation. The rats were monitored for any lethal effects due to administration of the drug.

### Statistical Analysis

Statistical analysis was performed using SPSS v.9.05. Data were analyzed by one-way analysis of variance (ANOVA), followed by Duncan's multiple range test. All the results were expressed as mean  $\pm$  SD for the 6 rats in each group. A P value < 0.05 was considered significant.

### Results and Discussion

The extract (administered at 20-50 times the effective dose) was observed to be non-toxic to the rats. Experimental animals appeared normal throughout the 7-week study period; there were no physical or behavioral changes observed, and there were no deaths in any of the experimental groups.

The levels of total protein and albumin in plasma decreased significantly ( $P < 0.05$ ) in the rats with STZ-induced diabetes, in comparison to the untreated control rats (Table 1). The levels of total protein and albumin returned to normal after the administration of the extract (500 mg/kg of body weight per day) for 12 weeks; however, a mild globulin level increase ( $P < 0.05$ ) in the diabetic rats was normalized by the administration of the plant extract. Blood levels of urea, uric acid, and creatinine increased significantly ( $P < 0.05$ ) in the rats with STZ-induced diabetes, in comparison to the untreated control

rats. Administration of the plant extract (500 mg/kg of body weight per day) for 12 weeks normalized levels of non-protein nitrogenous substances. The observed effect of the *F. bengalensis* aqueous extract was equivalent to the effect of tolbutamide.

Decreases in erythrocyte, leucocyte, and platelet counts were observed, and hemoglobin levels returned to normal following administration of the extract (500 mg/kg of body weight per day) for 12 weeks (Table 2). The decrease in hemoglobin level in the diabetic rats may have been due to a reduction in protein synthesis in all tissues, with decreased production of ATP and absolute or relative insulin deficiency (19).

The observed increase in plasma levels of urea, uric acid, and creatinine may have been due to STZ-induced metabolic disturbances, as well as renal dysfunction. Levels of non-protein nitrogenous substances are always used as significant markers for the assessment of renal dysfunction (20). Increased protein glycation in STZ-induced diabetes was reported to be associated with increased muscle wasting and, thereby, increased release of purines. The elevated levels of purine in diabetes are reported to be the main source of uric acid, in addition to xanthine oxidase activity (21). Our data revealed that the *F. bengalensis* aqueous extract significantly reduced the levels of serum urea, uric acid, and creatinine in rats with STZ-induced diabetes.

The observed changes in blood cell counts may have been due to the necrotic effect of STZ, which could also have been the cause of the drop in total hemoglobin. Some of the constituents present in the extract might reduce

Table 1. The effect of *F. bengalensis* extract on plasma proteins, urea, uric acid, and creatinine in rats with STZ-induced diabetes.

Groups	Total Protein (g/dl)	Albumin (g/dl)	Urea (m mol/l)	Uric acid (m mol/l)	Creatinine ( $\mu$ mol/l)
Normal	6.87 $\pm$ 0.81	4.60 $\pm$ 2.9	8.31 $\pm$ 0.27	65.43 $\pm$ 0.1	64.3 $\pm$ 2.3
Diabetic control <sup>#</sup>	4.25 $\pm$ 0.85*	3.90 $\pm$ 2.8*	9.41 $\pm$ 0.26*	142.8 $\pm$ 0.3*	74.8 $\pm$ 2.4*
Diabetic + Tolbutamide (100 g/kg per day)	6.10 $\pm$ 0.53	4.73 $\pm$ 1.9	8.75 $\pm$ 0.25 <sup>c</sup>	112.9 $\pm$ 0.3	63.2 $\pm$ 2.1
Diabetic + <i>F. bengalensis</i> (500 mg/kg per day)	6.45 $\pm$ 0.51 <sup>a</sup>	4.42 $\pm$ 2.1 <sup>a</sup>	8.25 $\pm$ 0.28 <sup>a</sup>	65.24 $\pm$ 0.1 <sup>a</sup>	64.0 $\pm$ 2.3 <sup>a</sup>

Each value is the mean  $\pm$  SD for 6 rats in each group.

<sup>#</sup>STZ: Streptozotocin (35 mg/kg body weight in 0.1 M cold citrate buffer, pH 4.5).

\*Values are statistically significant when compared to controls at  $F > 0.05$  (ANOVA) and  $P < 0.05$  (DMRT)

<sup>a</sup>Values are statistically significant when compared to diabetic controls at  $F > 0.05$  (ANOVA) and  $P < 0.05$  (DMRT).

Table 2. The effect of the *F. bengalensis* aqueous extract on hematological indices in rats with STZ-induced diabetes.

Groups	Hb (g/dl)	RBC X10 <sup>12</sup> /l	WBC X10 <sup>12</sup> /l	Platelets X10 <sup>9</sup> /l
Normal	15.50 ± 0.5	5.10 ± 0.7	9.90 ± 0.7	163 ± 1.8
Diabetic control <sup>#</sup>	12.95 ± 1.0*	3.95 ± 0.6*	8.10 ± 0.5*	151 ± 1.3*
Diabetic + Tolbutamide (100 mg/kg per day)	14.10 ± 1.1	5.20 ± 0.7	8.62 ± 0.6	160 ± 1.2
Diabetic + <i>F. bengalensis</i> (500 mg/kg per day)	15.95 ± 0.9 <sup>a</sup>	5.39 ± 0.6 <sup>a</sup>	8.85 ± 0.5 <sup>a</sup>	162 ± 2.0 <sup>a</sup>

Each value is the mean ± SD for 6 rats in each group.

STZ: Streptozotocin (35 mg/kg body weight in 0.1 M cold citrate buffer, pH 4.5).

\*Values are statistically significant when compared to controls at  $F > 0.05$  (ANOVA) and  $P < 0.05$  (DMRT).

<sup>a</sup>Values are statistically significant when compared to diabetic controls at  $F > 0.05$  (ANOVA) and  $P < 0.05$  (DMRT).

STZ-induced changes in total protein, albumin, non-protein nitrogenous substances, and hematological indices. The antioxidant activity of polyphenolic flavonoids, leucopelargonidin, and leucocynidin isolated from *F. bengalensis* has been previously reported (22). Isolation of 3 ketones (20-tetratriacontene-2-one, 6-heptatriacontene-10-one, and pentatriacontan-5-one) and 2 other compounds (beta-sitosterol-alpha-D-glucose and meso inositol) from the stem bark of *F. bengalensis* have been reported (23). Thus, the ameliorative potential of the aqueous extract may be due to the synergistic effect of several of the phytochemicals present in the extract.

## Conclusion

Findings from the present study show that the aqueous extract of *F. bengalensis* stem bark has some useful biological properties, as indicated by the significant changes observed in the biochemical parameters of rat plasma. Further studies are required to assess in greater detail the

biochemical/pharmacological properties of purified *F. bengalensis* extract(s) on blood sugar levels in vivo.

## Acknowledgements

The authors wish to thank the management of VIT University for providing the necessary facilities for the completion of this study. We are also grateful to the Principal, Voorhees College, Vellore, and Dr. David Reader, Dept. of Zoology, for permitting us to utilize their animal house facility to conduct this study.

### Corresponding Author:

Krishnan KANNABIRAN

Biomolecules and Genetics Division,

School of Biotechnology,

Chemical and Biomedical Engineering, VIT University,

Vellore-632 014, Tamil Nadu - INDIA

E-mail: kkb@vit.ac.in

## References

- Kim MJ, Ryu GR, Chung JS, et al. Protective effects of epicatechin against the toxic effects of STZ on rat pancreatic islets: in vivo and in vitro. *Pancreas* 26(3): 292-299, 2003.
- Szkudelski T. The mechanism of alloxan and streptozotocin action in  $\beta$ -cells of the rat pancreas. *Physiol Res* 50(6): 537-546, 2001.
- Yue KK, Chung WS, Leung AW et al. Redox changes precede the occurrence of oxidative stress in eyes and aorta, but not in kidneys of diabetic rats. *Life Sci* 73(20): 2557-2570, 2003.
- Burns N, Gold B. The effect of 3-methyladenine DNA glycosylase-mediated DNA repair on the reduction of toxicity and diabetes by the beta-cell toxicant streptozotocin. *Toxicol Sci* 95: 391-400, 2007.
- Koulmanda M, Qipo A, Chebrolu S et al. The effect of low versus high dose of Streptozotocin in cynomolgus monkeys (*Macaca fascicularis*). *Am J Transplant* 3: 267- 272, 2003.

6. Kordowiak AM, Goc A, Drozdowska E et al. Sodium orthovanadate exerts influence on liver Golgi complexes from control and streptozotocin-diabetic rats. *J Inorg Biochem*, 99(5): 1083-1089, 2005.
7. Satyavati GV, Raina MK, Sharma M. Medicinal plants of India, Indian Council of Medical Research, Vol. I, New Delhi, 1976.
8. Shukla R, Prabhu KM, Murthy PS. Hypoglycaemic effect of the water extract of *Ficus bengalensis* in alloxan recovered, mildly diabetic and severely diabetic rabbits. *Int J Diab Dev Ctries* 14: 78-81, 1994.
9. Shukla R, Anand K, Prabhu KM et al. Hypocholesterolemic effect of water extract of the bark of Banyan tree, *Ficus bengalensis*. *Indian J Clin Biochem* 10: 14-18, 1995.
10. Shukla R, Anand K, Prabhu KM et al. Hypolipidaemic effect of water extract of the bark of *Ficus bengalensis* in alloxan induced diabetes mellitus in rabbits. *Indian J Clin Biochem* 10: 119-121, 1995.
11. Vats V, Grover GK, Rathi SS. Evaluation of anti-hyperglycemic effect of *Trigonella foenum-graecum* Linn, *Ocimum sanctum* Linn and *Pterocarpus marsipium* Linn in normal and alloxanized diabetic rats. *J Ethnopharmacol* 79: 95-100, 2002.
12. Sarkar S, Pranava M, Marita RA. Demonstration of the hypoglycemic action of *Momordica charantia* in a validated animal model of diabetes. *Pharmacol Res* 33: 1-4, 1996.
13. Drabkin DL, Austin JM. Spectrophotometric studies, spectrophotometric constants for common hemoglobin derivatives in human, dog and rabbit blood. *J Biol Chem* 98: 719-733, 1932.
14. Lowery OH, Rosenbrough NJ, Farr AL et al. Protein determination using folin-Ciocalteu reagent. *J Biol Chem* 193: 265-276, 1951.
15. Peter T, Biamonte GT, Doumas BT. Protein (total protein) in serum, urine and cerebrospinal fluid; albumin in serum. In: Faulkner WR, Meites D (Eds.). 9<sup>th</sup> ed. *Selected Methods of Clinical Chemistry*. American Association for Clinical Chemistry, Washington, D.C.; 1982, pp 125-132.
16. Tomas L, Clinical Laboratory Diagnostics, Verlagsgesellschaft, 1<sup>st</sup> ed, Frankfurt, Germany; 1998.
17. Fossati P, Principe L, Berti G. Use of 3,5-dichloro-2-hydroxybenzenesulfonic acid-aminophenazone chromogenic system in direct enzymatic assay of uric acid in serum and urine. *Clin Chem* 26: 227-231, 1980.
18. Rao GM, Morghom LO. Erythrocyte count and hemoglobin levels in diabetic women. *Eur J Appl Physiol* 52: 272-275, 1984.
19. Chatterjee MN, Shinda R. Metabolism of carbohydrate. In: *Textbook of Medical Biochemistry*, Jaypee Brothers Medical Publishers Pvt. Ltd.; 1994, pp. 421-422.
20. Almadal TP, Vilstrup H. Strict insulin treatment normalizes the organic nitrogen contents and the capacity of urea-nitrogen synthesis in experimental diabetes in rats. *Diabetologica* 31:114-118, 1988.
21. Anwar MM, Meki AMR. Oxidative stress in streptozotocin-induced diabetic rats: effects of garlic oil and melatonin. *Comparative Biochem Physiol* 135: 539-547, 2003.
22. Mathew BC, Devi KS, Augusti KT. Antioxidant effect of two flavonoids from the bark of *Ficus bengalensis* Linn in hypolipidemic rats. *Indian J Exp Biol* 36: 902-906, 1998.
23. Subramanian PM, Misra CS. Chemical constituents of *Ficus bengalensis* (part II). *Polish J Pharmacol* 30: 559-562, 1978.