

Research Article

Activity of Ca^{2+} -activated Cl^- channels contributes to regulating receptor- and store-operated Ca^{2+} entry in human pulmonary artery smooth muscle cells

Aya Yamamura, Hisao Yamamura, Amy Zeifman, and Jason X.-J. Yuan

Department of Medicine, Section of Pulmonary, Critical Care, Sleep and Allergy, Institute for Personalized Respiratory Medicine, Center for Cardiovascular Research, and Department of Pharmacology, University of Illinois at Chicago, Chicago, Illinois, USA

ABSTRACT

Intracellular Ca^{2+} plays a fundamental role in regulating cell functions in pulmonary arterial smooth muscle cells (PASMCs). A rise in cytosolic Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{cyt}}$) triggers pulmonary vasoconstriction and stimulates PASMC proliferation. $[\text{Ca}^{2+}]_{\text{cyt}}$ is increased mainly by Ca^{2+} release from intracellular stores and Ca^{2+} influx through plasmalemmal Ca^{2+} -permeable channels. Given the high concentration of intracellular Cl^- in PASMCs, Ca^{2+} -activated Cl^- (Cl_{Ca}) channels play an important role in regulating membrane potential and cell excitability of PASMCs. In this study, we examined whether activity of Cl_{Ca} channels was involved in regulating $[\text{Ca}^{2+}]_{\text{cyt}}$ in human PASMCs via regulating receptor- (ROCE) and store- (SOCE) operated Ca^{2+} entry. The data demonstrated that an angiotensin II (100 nM)-mediated increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ via ROCE was markedly attenuated by the Cl_{Ca} channel inhibitors, niflumic acid (100 μM), flufenamic acid (100 μM), and 4,4'-diisothiocyanatostilbene-2,2'-disulfonic acid (100 μM). The inhibition of Cl_{Ca} channels by niflumic acid and flufenamic acid significantly reduced both transient and plateau phases of SOCE that was induced by passive depletion of Ca^{2+} from the sarcoplasmic reticulum by 10 μM cyclopiazonic acid. In addition, ROCE and SOCE were abolished by SKF-96365 (50 μM) and 2-aminoethyl diphenylborinate (100 μM), and were slightly decreased in the presence of diltiazem (10 μM). The electrophysiological and immunocytochemical data indicate that Cl_{Ca} currents were present and TMEM16A was functionally expressed in human PASMCs. The results from this study suggest that the function of Cl_{Ca} channels, potentially formed by TMEM16A proteins, contributes to regulating $[\text{Ca}^{2+}]_{\text{cyt}}$ by affecting ROCE and SOCE in human PASMCs.

Key Words: angiotensin II, Ca^{2+} signaling, Ca^{2+} -activated Cl^- current, niflumic acid, TMEM16A

INTRODUCTION

In pulmonary artery smooth muscle cells (PASMCs), cytosolic Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{cyt}}$) is mainly regulated by a balance of Ca^{2+} release from intracellular stores and Ca^{2+} influx through plasmalemmal Ca^{2+} -permeable channels, as well as Ca^{2+} sequestration into intracellular stores by the Ca^{2+} - Mg^{2+} ATPase on the sarcoplasmic/endoplasmic reticulum membrane (SERCA) and Ca^{2+} extrusion via the Ca^{2+} - Mg^{2+} ATPase and $\text{Na}^+/\text{Ca}^{2+}$ exchanger

on the plasma membrane.^[1,2] PASMCs functionally express various Ca^{2+} -permeable channels including (a) voltage-dependent Ca^{2+} channels (VDCCs) that are activated by membrane depolarization,^[3] and (b) receptor-operated Ca^{2+} (ROC) channels that are stimulated and activated by vasoconstrictors, such as endothelin-1,^[4] serotonin,^[5] phenylephrine,^[6] and histamine,^[7] and by growth factors, including epidermal growth factor^[8] and platelet-derived growth factor.^[9] The activation of ROC channels by interaction between ligands and membrane receptors

Address correspondence to:

Prof. Jason X.-J. Yuan
Department of Medicine
University of Illinois at Chicago
COMRB Rm. 3131 (MC 719)
909 South Wolcott Avenue
Chicago IL 60612 USA
Phone: (312) 355-5911
Fax: (312) 996-1793
Email: jxyuan@uic.edu

Access this article online

Quick Response Code:



Website: www.pulmonarycirculation.org

DOI: 10.4103/2045-8932.83447

How to cite this article: Yamamura A, Yamamura H, Zeifman A, Yuan JX. Activity of Ca^{2+} -activated Cl^- channels contributes to regulating receptor- and store-operated Ca^{2+} entry in human pulmonary artery smooth muscle cells. *Pulm Circ* 2011;1:269-79.

results in receptor-operated Ca^{2+} entry (ROCE) that greatly contributes to increases in $[\text{Ca}^{2+}]_{\text{cyt}}$ in PSMCs exposed to vasoconstrictors and growth factors.^[1,10,11] PSMCs also possess (c) store-operated Ca^{2+} (SOC) channels that are opened by the depletion of Ca^{2+} from the sarcoplasmic reticulum (SR), which leads to capacitative Ca^{2+} entry, or store-operated Ca^{2+} entry (SOCE). SOCE is an important mechanism involved in maintaining a sustained elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ and refilling Ca^{2+} into the depleted SR.^[1,10-12] We showed previously that increased Ca^{2+} influx through SOC or SOCE contributes to stimulating PSMC proliferation; inhibition of SOCE significantly attenuated growth factor-mediated PSMC proliferation. These results suggest that SOCE plays a significant role in regulating proliferation in vascular smooth muscle cells.^[9,13,14]

It has been well demonstrated that the activity of Ca^{2+} -activated Cl^- (Cl_{Ca}) channels play an important role in regulating contraction, migration, and apoptosis in many cell types.^[15,16] In vascular smooth muscle cells, Cl_{Ca} channels are activated by a rise in $[\text{Ca}^{2+}]_{\text{cyt}}$ following agonist-induced Ca^{2+} release from the SR through inositol-1,4,5-trisphosphate receptors (IP_3Rs). In addition, the activation of Cl_{Ca} channels is evoked by spontaneous Ca^{2+} release through ryanodine receptors in the SR and is responsible for eliciting spontaneous transient inward currents in several types of vascular smooth muscle cells. The intracellular Cl^- concentration in vascular smooth muscle cells (including PSMCs) is estimated to be 30 to 60 mM,^[15-17] so the reversal potential for Cl^- is supposed to be much less negative (ranging from -20 to -30 mV) than that for K^+ (approximately -80 mV). Therefore, an increase in Cl^- conductance in PSMCs under these conditions would generate inward currents (due to Cl^- efflux) and cause membrane depolarization which subsequently induces Ca^{2+} influx by opening VDCCs and ultimately results in vasoconstriction. The molecular composition of Cl_{Ca} channels in vascular smooth muscle cells (including PSMCs), however, is not fully identified. Recently, a transmembrane protein encoded by TMEM16A gene has been demonstrated to form Cl_{Ca} channels in vascular smooth muscle cells.^[18-20]

In this study, we examined whether Cl_{Ca} channel activity was involved in the regulation of $[\text{Ca}^{2+}]_{\text{cyt}}$ via ROCE and SOCE in human PSMCs using digital imaging fluorescence microscopy. We also examined the functional expression of Cl_{Ca} channels (TMEM16A) in human PSMCs using electrophysiological and immunocytochemical approaches.

MATERIALS AND METHODS

Cell culture

Human PSMCs (passage 5 to 10) from normal subjects

were purchased from Lonza (Walkersville, MD). Cells were cultured in Medium 199 (Invitrogen-GIBCO, Grand Island, NY) supplemented with 10% fetal bovine serum (Invitrogen-GIBCO), 100 U/ml penicillin plus 100 $\mu\text{g}/\text{ml}$ streptomycin (Invitrogen-GIBCO), 50 $\mu\text{g}/\text{ml}$ D-valine (Sigma-Aldrich, St. Louis, MO), and 20 $\mu\text{g}/\text{ml}$ endothelial cell growth supplement (BD Biosciences, Franklin Lakes, NJ) at 37°C. All cells were incubated in a humidified 5% CO_2 atmosphere at 37°C. After reaching confluence, the cells were sub-cultured by trypsinization with 0.05% trypsin-EDTA (Invitrogen-GIBCO), plated onto 25-mm cover slips (Fisher Scientific, Pittsburgh, PA), and incubated at 37°C for 1-3 days before electrophysiological and fluorescence microscopy experiments.

$[\text{Ca}^{2+}]_{\text{cyt}}$ measurement

Human PSMCs cultured on 25-mm cover slips were placed in a recording chamber on the stage of an invert fluorescent microscope (Eclipse Ti-E; Nikon, Tokyo, Japan) equipped with an objective lens (S Plan Fluor 20 \times /0.45 ELWD; Nikon) and an EM-CCD camera (Evolve; Photometrics, Tucson, AZ). $[\text{Ca}^{2+}]_{\text{cyt}}$ was monitored using a membrane-permeable Ca^{2+} -sensitive fluorescent indicator, fura-2 acetoxymethyl ester (fura-2/AM; Invitrogen-Molecular Probes, Eugene, OR) and imaged with NIS Elements 3.2 software (Nikon). Cells were loaded by incubation in HEPES-buffered solution containing 4 μM fura-2/AM for 60 min. at room temperature (25°C). The loaded cells were then washed with HEPES-buffered solution for 10 min. to remove excess extracellular indicator and allow sufficient time for intracellular esterase to cleave acetoxymethyl ester from fura-2. Cells were then excited at 340-nm and 380-nm wavelengths (D340 \times v2 and D380 \times v2 filters, respectively; Chroma Technology, Bellows Falls, VT) by a xenon arc lamp (Lambda LS; Sutter Instrument, Novato, CA) and an optical filter changer (Lambda 10-B; Sutter Instrument). Emission of Fura-2 was collected through a dichroic mirror (400DCLP; Chroma Technology) and a wide band emission filter (D510/80m; Chroma Technology). $[\text{Ca}^{2+}]_{\text{cyt}}$ within a region of interest (5 \times 5 μm) that was placed at the peripheral region of each cell was measured as the ratio of fluorescence intensities (F_{340}/F_{380}) every 2 sec. The HEPES-buffered solution had an ionic composition of 137 mM NaCl, 5.9 mM KCl, 2.2 mM CaCl_2 , 1.2 mM MgCl_2 , 14 mM glucose, and 10 mM HEPES. The pH was adjusted to 7.4 with 10 N NaOH. The external Ca^{2+} -free solution was prepared by removing extracellular CaCl_2 and adding 1 mM EGTA (to chelate the residual Ca^{2+} in the bath solution). The recording chamber was continuously perfused with HEPES-buffered solution at a flow rate of 2 ml/min. using a mini-pump (Model 3385; Control, Friendswood, TX). $[\text{Ca}^{2+}]_{\text{cyt}}$ measurements were carried out at 32°C using an automatic temperature controller (TC-344B, Warner Instruments, Holliston, MA).

Electrophysiological recording

The whole-cell Cl_{Ca} current in a single PSMC was recorded using the patch-clamp technique with an Axopatch-1D amplifier (Molecular Devices-Axon, Foster City, CA), an analog-digital converter (Digidata 1200; Molecular Devices-Axon), and pCLAMP 8 software (Molecular Devices-Axon). The extracellular (bath) solution had an ionic composition of 137 mM NaCl, 10 mM tetraethylammonium (TEA) chloride, 5 mM 4-aminopyridine (4-AP), 2.2 mM CaCl_2 , 1.2 mM MgCl_2 , 14 mM glucose, and 10 mM HEPES. The pH was adjusted to 7.4 with 10 N NaOH. The low Cl^- concentration solution was prepared by substituting 117 mM NaCl of the extracellular solution with the equal molar of sodium gluconate. The pipette (intracellular) solution contained 120 mM CsCl, 20 mM TEA chloride, 4.3 mM CaCl_2 , 2.8 mM MgCl_2 , 2 mM Na_2ATP , 10 mM HEPES, and 5 mM EGTA. The pCa was fixed to 6.0, which was estimated by the Maxchelator program (<http://www.stanford.edu/~cpatton/maxc.html>). The pH was adjusted to 7.2 with 1 N CsOH. The recording chamber was continuously superfused with extracellular solution at a flow rate of 2 ml/min. using a perfusion system (VC-6; Warner Instrument, Hamden, CT). Electrophysiological recordings were carried out at room temperature (25°C).

Immunocytochemical staining

Cultured cells on 35-mm culture dishes with 14-mm glass bottom (MatTek, Ashland, MA) were fixed with 4% paraformaldehyde in Dulbecco's phosphate buffered saline (DPBS; Invitrogen-GIBCO) for 10 min. at room temperature (25°C). Excessive paraformaldehyde was removed thoroughly with DPBS. The cells were then treated with DPBS containing 0.2% Triton X-100, 1% normal goat serum (Dako Denmark, Glostrup, Denmark), and TMEM16A antibody (pre-diluted, ab53213, Abcam, Cambridge, MA; or 1:100 dilution, ab53212, Abcam) for 12 hr. at 4°C. After washing repeatedly in DPBS, the cells were covered with DPBS containing 0.2% Triton X-100, 1% normal goat serum, and Alexa Fluor 488-labeled secondary antibody (1:100 dilution; Invitrogen-Molecular Probes) for 1 hr. at room temperature and then rinsed with DPBS. Then cells were mounted in VECTASHIELD hard-set mounting medium with 4',6-diamidino-2-phenylindole (DAPI, 1.5 µg/ml) (Vector Laboratories, Burlingame, CA) and placed on the stage of an invert fluorescent microscope (Eclipse Ti-E; Nikon) equipped with an objective lens (Plan Apo 60×/1.40 oil immersion; Nikon), a CCD camera (CoolSNAP ES²; Photometrics), and NIS Elements 3.2 software (Nikon). Immunocytochemical images were obtained using the specific filter sets for DAPI (Ex340-380/DM400/Em435-485; Chroma Technology) and Alexa Fluor 488 (Ex460-500/DM505/Em510-560; Chroma Technology).

Drugs

Pharmacological reagents were obtained from Sigma-Aldrich. All hydrophobic compounds were dissolved in dimethyl sulfoxide (DMSO) at the concentration of 10 or 100 mM as a stock solution. It was confirmed that up to 0.1 % of DMSO did not affect these responses.

Statistical analysis

Pooled data are shown as the mean±SE. The statistical significance between two groups was determined by Student's *t*-test. The statistical significance among groups was determined by Scheffé's test after one-way analysis of variance. Significant difference is expressed in the figures as **P*<0.05 or ***P*<0.01.

RESULTS

Inhibition of agonist-induced Ca^{2+} influx or ROCE by Cl_{Ca} channel blockers in human PSMCs

The increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ evoked by agonist stimulation was imaged in human PSMCs loaded with 4 µM fura-2/AM and quantitated in arbitrary units (au) by the change in F_{340}/F_{380} ratio. Short-term application (2 min.) of 100 nM angiotensin II induced a transient increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ (by 0.49 ± 0.01 au, *n*=150) (Figs. 1 and 2). The angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increase was attenuated by 100 µM niflumic acid, a fenamate compound that is most frequently used as a blocker of Cl_{Ca} channels (from 0.51 ± 0.03 to 0.13 ± 0.02 au, *n*=29, *P*<0.01) (Fig. 1a and d). The inhibitory effect of niflumic acid on the angiotensin II-mediated increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ was reversible upon washout (0.42 ± 0.04 au, *n*=29). Pretreatment with 100 µM flufenamic acid, another fenamate compound that blocks Cl_{Ca} channels, markedly reduced the angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increase (from 0.51 ± 0.02 to 0.13 ± 0.03 au, *n*=33, *P*<0.01) (Fig. 1 b and e). A different type of Cl^- channel blocker, 4,4'-diisothiocyanatostilbene-2,2'-disulfonic acid (DIDS)—one of the stilbene derivatives that is structurally unrelated to fenamates—also caused a significant inhibition of the angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increase (from 0.43 ± 0.04 to 0.06 ± 0.01 au, *n*=18, *P*<0.01) (Fig. 1c and f). These data indicated that the function of Cl_{Ca} channels is involved in regulating ROCE in human PSMCs; inhibition of Cl_{Ca} channels significantly and reversibly attenuates the agonist-mediated Ca^{2+} entry.

Effects of Ca^{2+} channel blockers on ROCE in human PSMCs

To elucidate the Ca^{2+} signal pathway for angiotensin II-induced ROCE, effects of the inhibitors for several different types of Ca^{2+} channels were examined in human PSMCs. The angiotensin II-mediated increase in




Figure 1: Attenuation of ROCE by Cl_{Ca} channel blockers in human PSMCs. Angiotensin II (AngII, 100 nM) was used to induce ROCE in human PSMCs. (a-c) Representative traces showing angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increases in human PSMCs before, during, and after application of 100 μM niflumic acid (NFA; a), flufenamic acid (FFA; b), and DIDS (c). Blockage of Cl_{Ca} channels reduces angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation in human PSMCs. (d-f) Summarized data showing the reversible inhibitory effects of niflumic acid (d), flufenamic acid (e), and DIDS (f) on angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ rises in human PSMCs. Statistical significance (versus control) is indicated as ** $P < 0.01$.

$[\text{Ca}^{2+}]_{\text{cyt}}$ was reduced by treatment with 50 μM SKF-96365, an inhibitor of non-selective cation channels (from 0.59 ± 0.05 to 0.28 ± 0.04 au, $n=13$, $P < 0.01$) (Fig. 2 a and d). The application of 100 μM 2-aminoethoxydiphenylborate (2-APB), which blocks IP_3 Rs and also non-selective cation channels, abolished the angiotensin II-induced increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ (from 0.46 ± 0.03 to 0.08 ± 0.01 au, $n=22$, $P < 0.01$) (Fig. 2 b and e). Blockage of VDCC with 10 μM diltiazem, however, had a trend to inhibit the angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increase (from 0.45 ± 0.02 to 0.36 ± 0.03 au, $n=35$, $P > 0.05$ by Scheffé's test, but $P < 0.01$ by Student's *t*-test) (Fig. 2 c and f). These pharmacological data indicate that the angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increase was mainly caused by Ca^{2+} release from the SR through IP_3 R followed by Ca^{2+} influx via non-selective cation channels in human PSMCs. Ca^{2+} influx through the diltiazem-sensitive L-type VDCCs slightly contributes to the angiotensin II-induced rise of $[\text{Ca}^{2+}]_{\text{cyt}}$ in human PSMCs.

Inhibitory effect of Cl_{Ca} channel blockers on SOCE in human PSMCs

In the next set of experiments, we examined the effect of Cl_{Ca} channel blockers on SOCE in human PSMCs (Fig. 3). SOCE was induced by passive depletion of Ca^{2+} from the SR with 10 μM cyclopiazonic acid (CPA), a blocker of SERCA. In the absence of extracellular Ca^{2+} , application of CPA induced a transient increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ that was due predominantly to Ca^{2+} leakage from the SR to the cytosol. Restoration of extracellular Ca^{2+} after approximately 10 min. treatment with CPA caused another increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ that was apparently due to Ca^{2+} influx through store-operated cation (or Ca^{2+}) channels or SOCE.

As shown in Figure 3, there were no significant differences in the resting $[\text{Ca}^{2+}]_{\text{cyt}}$ (0.59 ± 0.02 versus 0.62 ± 0.02 au, $n=55$, $P=0.25$) and the amplitude of the increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ due to CPA-induced Ca^{2+} leakage from the SR to




Figure 2: Effects of Ca^{2+} channel blockers on ROCE in human PSMCs. Angiotensin II (AngII, 100 nM) was used to induce ROCE in human PSMCs. (a-c) Representative traces showing angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ rises in human PSMCs before, during, and after application of 50 μM SKF-96365 (SKF, an inhibitor of nonselective cation channels; a), 100 μM 2-APB (which blocks IP_3Rs and also non-selective cation channels; b), and 10 μM diltiazem (a VDCC blocker; c). (d-f) Summarized data showing effects of SKF-96365 (d), 2-APB (e), and diltiazem (f) on angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increase in human PSMCs. Statistical significance (versus control) is indicated as **P<0.01.

the cytosol when cells were treated with 100 μM niflumic acid or vehicle (0.1% DMSO) (Fig. 3 a and b). The transient and plateau phases of CPA-induced increases in $[\text{Ca}^{2+}]_{\text{cyt}}$ due to SOCE, as well as the amplitude and “rise-speed” of SOCE were all significantly decreased by 100 μM niflumic acid. The rise-speed of SOCE in the presence of niflumic acid (0.18 ± 0.01 ratio/s, $n=55$) was significantly slower than that in the absence (0.36 ± 0.04 ratio/s, $P<0.01$) (Fig. 3c). In addition, application of niflumic acid significantly reduced the amplitude of the transient (from 0.42 ± 0.02 to 0.31 ± 0.02 au, $n=55$; $P<0.01$) and plateau (from 0.15 ± 0.01 to 0.10 ± 0.01 au, $n=55$; $P=0.02$) phases of SOCE (Fig. 3a and d). Treatment of the cells with 100 μM flufenamic acid also significantly decreased the amplitude of the transient and plateau phases of CPA-induced SOCE in human PSMCs ($n=34$). These data clearly suggest that the activity of Cl_{Ca} channels is involved in regulating SOCE in human PSMCs.

Effects of Ca^{2+} channel blockers on SOCE in human PSMCs

To functionally define the Ca^{2+} channels responsible for CPA-mediated SOCE, we examined the effects of different Ca^{2+} channel blockers on SOCE in human PSMCs. The transient component of CPA-induced SOCE was significantly reduced by 50 μM SKF-96365 (0.21 ± 0.01 au, $n=51$, versus vehicle control, 0.37 ± 0.01 au, $n=58$; $P<0.01$) (Fig. 4 a, b and e) and 100 μM 2-APB (0.10 ± 0.01 au, $n=46$, $P<0.01$) (Fig. 4 c and e). Application of 10 μM diltiazem slightly (but significantly) affected the transient component of SOCE (from 0.37 ± 0.01 to 0.32 ± 0.01 au, $n=42$, $P<0.01$), but had no effect on the plateau phase of SOCE (from 0.16 ± 0.01 au, $n=58$, to 0.14 ± 0.01 au, $n=42$, $P=0.22$) (Fig. 4 a, d and e). Similar to their effects on the transient phase of SOCE, 50 μM SKF-96365 (from 0.16 ± 0.01 , $n=58$, to 0.06 ± 0.01 au, $n=51$; $P<0.01$) (Fig. 4a, b and e) or 100 μM 2-APB (to 0.05 ± 0.01 au, $n=46$, $P<0.01$)




Figure 3: Attenuation of SOCE by Cl_{Ca} channel blocker in human PSMCs. SOCE was induced by passive depletion of Ca^{2+} from the SR with 10 μM CPA in human PSMCs. (a) Typical trace of CPA-induced Ca^{2+} release and SOCE in the absence (vehicle, 0.1% DMSO) and presence of 100 μM niflumic acid (NFA) in a human PSMC. Blockage of Cl_{Ca} channels caused a reduction of SOCE in human PSMCs. (b-d) Summarized data showing effects of niflumic acid on the resting $[\text{Ca}^{2+}]_{\text{cyt}}$ (b), the rise speed of SOCE (c), the transient and plateau amplitudes of CPA-induced SOCE (d) in human PSMCs. Statistical significance versus vehicle control is indicated as * $P < 0.05$ or ** $P < 0.01$.

(Fig. 4c and e) also significantly reduced the plateau phase of CPA-induced SOCE. These pharmacological data indicate that the Ca^{2+} signaling pathway for CPA-induced SOCE was mainly dependent on Ca^{2+} influx through non-selective cation channels in human PSMCs. The activity of VDCCs might be, in part, involved in the Ca^{2+} influx pathway in human PSMCs.

Whole-cell Cl_{Ca} currents in human PSMCs

Electrophysiological and pharmacological properties of Cl_{Ca} currents in human PSMCs were analyzed by

whole-cell patch-clamp configuration using a pipette (intracellular) solution containing 120 mM Cs^+ , 20 mM TEA ($\text{pCa}=6.0$) and a bath (extracellular) solution containing 10 mM TEA and 5 mM 4-AP. The mean cell-capacitance was 9.0 ± 1.2 pF ($n=14$). Depolarizing pulses (500 ms) were applied from a holding potential of -60 mV to a series of test potentials ranging from -80 to +100 mV by 20-mV increments every 15 sec. Outward currents were elicited by depolarization from the holding potential to the positive potentials above 0 mV, and the averaged current density at +100 mV was 86 ± 7 pA/pF ($n=14$) (Fig. 5a and b). The reversal potential of the whole-cell currents was -3.4 ± 1.3 mV ($n=14$) (Fig. 5b), which is close to the theoretical (or calculated) equilibrium potential of Cl^- (+0.1 mV). In addition, the reversal potential was shifted positively (to the right) by approximately 30 mV (positive shift by 36 mV in theory) by changing the extracellular Cl^- concentration from 153.8 to 36.8 mM (data not shown). Importantly, we were able to detect inward "tail" currents when cells were repolarized to the holding potential (Fig. 5a), which is an important characteristic of Cl_{Ca} currents. Furthermore, we analyzed the tail currents and the current-voltage relationship using the pulse protocol as follows: depolarizing pre-pulses were applied from a holding potential of -60 to +100 mV for 100 ms and subsequently test pulses were applied between -40 to +40 mV by 10-mV increments for 500 ms every 15 sec. (Fig. 5c). The current-voltage relationship revealed that the reversal potential of the tail currents (1.9 ± 0.9 mV, $n=4$; Fig. 5d) was also very close to the theoretical (or calculated) equilibrium potential of Cl^- .

Application of 100 μM niflumic acid significantly attenuated both the outward Cl_{Ca} current and the inward tail current in human PSMCs (Fig. 6). Niflumic acid decreased the outward currents elicited by depolarization from a holding potential of -60 to +60 mV (for 500 ms every 15 s) (from 37.4 ± 5.3 to 16.5 ± 1.2 pA/pF, $n=3$; $P=0.046$). The inward tail currents were also inhibited by the pretreatment with niflumic acid (-5.6 ± 1.3 pA/pF, $n=3$, versus control of -22.2 ± 4.7 pA/pF, $P=0.040$). These electrophysiological data indicate that Cl_{Ca} channels sensitive to niflumic acid are functionally expressed in human PSMCs.

Expression of TMEM16A in human PSMCs

The molecular basis of Cl_{Ca} channels in human PSMCs was identified by an immunocytochemical approach using two specific primary antibodies of TMEM16A (ab53213 and ab53212 from Abcam), a potential protein candidate for Cl_{Ca} channels. Immunocytochemical experiments (Fig. 7) revealed that specific fluorescent signals of TMEM16A protein were localized in the cell membrane. Qualitatively, the same images were obtained from 3




Figure 4: Effects of Ca^{2+} channel blockers on SOCE in human PSMCs. SOCE was induced by passive depletion of Ca^{2+} from the SR with 10 μM CPA in human PSMCs. (a-d) Typical traces showing CPA-induced Ca^{2+} release and SOCE in the absence (vehicle, 0.1% DMSO; a) and presence of 50 μM SKF-96365 (SKF, a blocker for non-selective cation channels; b), 100 μM 2-APB (a blocker of IP_3Rs and non-selective cation channels; c), and 10 μM diltiazem (an inhibitor of VDCCs; d) in human PSMCs. (e) Summarized data showing effects of Ca^{2+} channel blockers on the transient and plateau amplitudes of CPA-induced SOCE in human PSMCs. The number of cells examined is given in parentheses. Statistical significance versus vehicle control is indicated as **P<0.01.

separate sets of experiments. This result indicates that the activity of Cl_{Ca} channels in human PSMCs is potentially due to channels formed by TMEM16A.

DISCUSSION

In vascular smooth muscle cells, Cl_{Ca} channels are present for diverse physiological and pathological functions. In this study, we showed that the blockage of Cl_{Ca} channels using pharmacological tools markedly attenuated both ROCE and SOCE in human PSMCs. Our electrophysiological and immunocytochemical data also indicated that the activity of Cl_{Ca} channels functionally expressed in human PSMCs was due potentially to channels formed by TMEM16A proteins.

Intracellular free Ca^{2+} plays an important role in the regulation of contraction, proliferation, and migration of PSMCs. An increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ in PSMCs is a major trigger for pulmonary vasoconstriction and an important stimulus for PSMC proliferation that leads to pulmonary vascular remodeling under pathological conditions. Elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ in PSMCs results from Ca^{2+} release from intracellular stores, such as the SR, and Ca^{2+} influx through plasmalemmal Ca^{2+} channels, such as ROC channels, SOC channels, and VDCCs.^[1,2]

Angiotensin II is a vasoconstrictor that is commonly used for eliciting agonist-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ rises in PSMCs and other vascular smooth muscle cells.^[21,22] In this study, we used angiotensin II to induce ROCE because, at the concentration of 100 nM, it caused an increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ in a large number of human PSMCs (>70%). The angiotensin II-mediated increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ via ROCE was markedly reduced by two different types of Cl_{Ca} channel inhibitors, fenamates (niflumic acid and flufenamic acid) and one of the stilbene derivatives (DIDS). Both niflumic acid and flufenamic acid are well known fenamates that are most frequently used as Cl_{Ca} channel blockers in electrophysiological and pharmacological studies. However, these compounds have been reported to also act on other types of ion channels such as non-selective cation channels,^[23] large-conductance Ca^{2+} -activated K^+ channels,^[24] and transient receptor potential canonical subfamily (TRPC) channels.^[25] Therefore, to confirm whether or not the inhibitory effects of niflumic acid and flufenamic acid on the angiotensin II-evoked increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ in human PSMCs were mediated by the blockage of Cl_{Ca} channels, we analyzed the effects of another type of Cl_{Ca} channel blocker, DIDS, a stilbene derivative that is structurally unrelated to fenamates, on the angiotensin II-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increase. Similar to niflumic acid and flufenamic acid, DIDS also significantly suppressed the angiotensin II-




Figure 5: Whole-cell Cl_{Ca} currents in human PSMCs. Cl_{Ca} currents were measured using a pipette solution containing 120 mM Cs^+ , 20 mM TEA and pCa 6.0, and a bath solution containing 10 mM TEA and 5 mM 4-AP in human PSMCs. (a) Representative outward currents (black arrowhead), elicited by depolarization from a holding potential of -60 mV to a series of test potentials (-80 to +100 mV) for 500 ms every 15 sec. and inward tail currents (gray arrowhead), induced by repolarization to -60 mV in a human PSMC. (b) I-V relationship at peak amplitude during depolarization (black arrowhead in "a"). The current reverses at about 0 mV, the theoretical equilibrium potential of Cl^- . (c) Representative tail currents (gray arrowhead) at a series of test potentials (-40 to +40 mV) for 500 ms after depolarization from a holding potential of -60 to +100 mV for 100 ms every 15 sec. in a human PSMC. (d) I-V relationship of tail currents (gray arrowhead in "c"). The reversal potential is close to 0 mV.

induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increase, although it might have interfered with the fura-2 fluorescence due to its faint yellow colored solution at a concentration of 100 μM . These results, by using two different types of Cl_{Ca} channel blockers, strongly suggest that the function of Cl_{Ca} channels is involved in regulating ROCE in human PSMCs.

SOCE is essential for maintaining a high level of $[\text{Ca}^{2+}]_{\text{cyt}}$ and for refilling intracellular Ca^{2+} stores (i.e., SR) in smooth muscle cells.^[1,10-12] High levels of $[\text{Ca}^{2+}]_{\text{cyt}}$ and sufficient levels of Ca^{2+} in the SR are required for proliferation of vascular smooth muscle cells.^[1,10] SOCE is enhanced while SOC channels are upregulated during PSMC proliferation to increase Ca^{2+} influx and provide sufficient Ca^{2+} for activation of the intracellular mechanisms responsible for cell proliferation and growth.^[9,13,14] In




Figure 6: Inhibition of Cl_{Ca} currents by niflumic acid in human PSMCs. Effect of niflumic acid on whole-cell Cl_{Ca} currents was examined in human PSMCs. (a) Representative outward currents (black arrowhead), elicited by depolarization from a holding potential of -60 to +60 mV, and inward tail currents (gray arrowhead), induced by repolarization to -60 mV in the absence (black line) and presence (gray line) of 100 μM niflumic acid (NFA) in a human PSMC. (b) Summarized data showing the effect of niflumic acid on outward (black arrowhead in "a") and tail (gray arrowhead in "a") currents in human PSMCs. The number of cells examined is given in parentheses. Statistical significance versus control is indicated as *P < 0.05.




Figure 7: Expression of TMEM16A in human PSMCs. Immunocytochemical analysis of TMEM16A, a potential candidate for the Cl_{Ca} channel subunit, was performed in human PSMCs using two specific primary antibodies (ab53213 and ab53212 from Abcam). Alexa Fluor 488 and DAPI were used as a secondary antibody and a nuclear marker, respectively. Specific signals of TMEM16A protein were detected on the plasma membranes of human PSMCs. Similar immunocytochemical images were obtained from 3 sets of independent experiments.

the present study, we demonstrated that the blockage of Cl_{Ca} channels by niflumic acid and flufenamic acid reduced both the transient and plateau components of SOCE as well as the rise speed of SOCE in human PSMCs. These data indicate that the function of Cl_{Ca} channels is also involved in regulating SOCE in human PSMCs (in addition to the effect on ROCE). Suppression of SOCE by blockage of Cl_{Ca} channels is thought to cause reduced PSMC proliferation, which may be a novel strategy for

preventing the abnormal proliferation under pathological conditions. TRPC channels have been demonstrated to be involved in agonist- or growth factor-mediated Ca^{2+} entry in PSMCs,^[1,10,26] while functional coupling of stromal interaction molecule (STIM) proteins (STIM1 and STIM2) with TRPC and/or Orai channels have recently been suggested as a novel candidate for SOC channel subunits in PSMCs.^[27-29] TRPC channel genes are thought to encode pore-forming subunits that compose ROC^[30-32] and SOC^[33-35] channels in many cell types of vascular smooth muscles including PSMCs.^[9,13,14,36-38] Ca^{2+} entry via ROC and SOC channels is modulated by second messengers, phosphorylation of signal transduction proteins, and transcription factors.^[2,10,39] The protein expression levels of TRPC, STIM and Orai are changed under pathological conditions such as in pulmonary arterial hypertension.^[1,10,39]

Smooth muscle cells contain a high concentration of Cl^- in the intracellular space, which is considerably different from other cell types, such as neurons, cardiomyocytes, and skeletal muscle myocytes.^[17] Therefore, increases in Cl^- conductance across the plasma membrane (e.g., as a result of activation of Cl_{Ca} channels when $[\text{Ca}^{2+}]_{\text{cyt}}$ is increased) lead to Cl^- efflux and inward currents, which consequently causes membrane depolarization, enhanced Ca^{2+} influx through VDCCs, increased $[\text{Ca}^{2+}]_{\text{cyt}}$, and vasoconstriction.^[15,16] In the present study, electrophysiological data indicated that Cl_{Ca} channels were functionally expressed in human PSMCs and the Cl^- currents through Cl_{Ca} channels were sensitive to niflumic acid. The electrophysiological properties (e.g., the time-dependent outward current during membrane depolarization, the inward tail current during repolarization, the outward rectification, and the shift of reversal potential based on the change in extracellular Cl^- concentration) and pharmacological properties (e.g., the dependency on intracellular Ca^{2+} concentration and the sensitivity to niflumic acid) of whole-cell Cl_{Ca} currents obtained from human PSMCs were consistent with the same properties reported previously in PSMCs from rabbits^[40] and rats.^[41,42] The $[\text{Ca}^{2+}]_{\text{cyt}}$ increase mediated by ROCE and SOCE also activates the Cl_{Ca} channels, resulting in membrane depolarization followed by additional Ca^{2+} influx through VDCCs. Slight decreases in ROCE and SOCE by diltiazem, a VDCC blocker, suggested that VDCCs only partly contributed to the regulation of ROCE and SOCE, although the Ca^{2+} influx pathway was mainly due to non-selective Ca^{2+} channels sensitive to SKF-96365 and 2-APB in human PSMCs.

Cl_{Ca} channels play important roles in diverse functions in vascular smooth muscle cells. In spite of its physiological and pathological significances, the molecular architecture of Cl_{Ca} channels in vascular smooth muscle cells has not been clearly demonstrated. More recently, the TMEM16

family, consisting of 10 genes in mammals, has been found as a novel candidate for Cl_{Ca} channel subunits.^[18-20,43] Heterologous expression of TMEM16A has been shown to generate Cl^- currents sensitive to intracellular Ca^{2+} and with the degree of outward rectification, ion selectivity, and pharmacological profile^[18-20] similar to the activity of native Cl_{Ca} channels observed in many tissues containing interstitial cells of Cajal in gastrointestinal muscles,^[44-46] airway epithelial cells,^[47,48] as well as vascular smooth muscle cells.^[42,49] The distribution pattern of TMEM16A in interstitial cells of Cajal in gastrointestinal muscles,^[44-46,50] airway epithelial cells,^[47,48] and vascular smooth muscle cells^[49] implies the functional expression of Cl_{Ca} conductance. In this study, TMEM16A protein was localized in the plasma membrane of human PSMCs, indicating that the activity of the Cl_{Ca} channel in human PSMCs was, at least in part, due to channels formed by TMEM16A. It has been reported that the TMEM16A gene also has some splice variants^[18,42,49,51] and TMEM16B, a closely related analogue, also can generate Cl^- currents activated by Ca^{2+} .^[19,52-54] It is unclear whether TMEM16B is another subunit that forms Cl_{Ca} channels in human PSMCs.

Pulmonary arterial hypertension is a fatal and progressive disease characterized pathologically by severe pulmonary vascular remodeling. A central aspect of pulmonary vascular remodeling is adventitial, medial, and intimal hypertrophy caused by excessive proliferation of fibroblasts and myofibroblasts in the adventitia, PSMCs in the media and endothelial cells in the intima. The concentric pulmonary vascular wall remodeling or thickened arterial and arteriole wall, narrows the intra-arterial lumen, increases pulmonary vascular resistance and ultimately causes pulmonary hypertension.^[2,39] Since SOC and ROC channels are upregulated in PSMC isolated from patients with idiopathic pulmonary arterial hypertension (IPAH) and from animals with hypoxia-mediated pulmonary hypertension, Ca^{2+} entry through these upregulated cation channels may play an important pathogenic role in the initiation and progression of pulmonary vascular remodeling under the pathological conditions.^[1,10,14,26,28,37] It remains unclear, however, whether the activity of Cl_{Ca} channels is also involved in the sustained pulmonary vasoconstriction and excessive pulmonary vascular remodeling in patients with IPAH and animals with hypoxia-induced pulmonary hypertension. Based on the observations from this study, the attenuation of SOCE and ROCE by Cl_{Ca} channel blockers (e.g., niflumic acid, flufenamic acid, and DIDS) may serve as a potential therapeutic approach for pulmonary vascular disease. Although it is suggested that Cl^- channels are involved in SOCE and proliferation in PSMCs,^[55,56] further experiments are necessary to elucidate the mechanism underlying the regulation of ROCE and SOCE by Cl_{Ca} channels in human PSMCs.

REFERENCES

- Firth AL, Remillard CV, Yuan JX. TRP channels in hypertension. *Biochim Biophys Acta* 2007;1772:895-906.
- Morrell NW, Adnot S, Archer SL, Dupuis J, Jones PL, MacLean MR, et al. Cellular and molecular basis of pulmonary arterial hypertension. *J Am Coll Cardiol* 2009;54:S20-31.
- Firth AL, Remillard CV, Platoshyn O, Fantozzi I, Ko EA, Yuan JX. Functional ion channels in human pulmonary artery smooth muscle cells: Voltage-dependent cation channels. *Pulm Circ* 2011;1:48-71.
- Shimoda LA, Sylvester JT, Sham JS. Inhibition of voltage-gated K^+ current in rat intrapulmonary arterial myocytes by endothelin-1. *Am J Physiol* 1998;274:L842-53.
- Pitt BR, Weng W, Steve AR, Blakely RD, Reynolds I, Davies P. Serotonin increases DNA synthesis in rat proximal and distal pulmonary vascular smooth muscle cells in culture. *Am J Physiol* 1994;266:L178-86.
- Doi S, Damron DS, Horibe M, Murray PA. Capacitative Ca^{2+} entry and tyrosine kinase activation in canine pulmonary arterial smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 2000;278:L118-30.
- Mauban JR, Wilkinson K, Schach C, Yuan JX. Histamine-mediated increases in cytosolic $[Ca^{2+}]$ involve different mechanisms in human pulmonary artery smooth muscle and endothelial cells. *Am J Physiol Cell Physiol* 2006;290:C325-36.
- Ohlstein EH, Douglas SA, Sung CP, Yue TL, Loudon C, Arleth A, et al. Carvedilol, a cardiovascular drug, prevents vascular smooth muscle cell proliferation, migration, and neointimal formation following vascular injury. *Proc Natl Acad Sci USA* 1993;90:6189-93.
- Yu Y, Sweeney M, Zhang S, Platoshyn O, Landsberg J, Rothman A, et al. PDGF stimulates pulmonary vascular smooth muscle cell proliferation by upregulating TRPC6 expression. *Am J Physiol Cell Physiol* 2003;284:C316-30.
- Yang XR, Lin MJ, Sham JS. Physiological functions of transient receptor potential channels in pulmonary arterial smooth muscle cells. *Adv Exp Med Biol* 2010;661:109-22.
- Wang Y, Deng X, Hewavitharana T, Soboloff J, Gill DL. Stim, ORAI and TRPC channels in the control of calcium entry signals in smooth muscle. *Clin Exp Pharmacol Physiol* 2008;35:1127-33.
- Albert AP, Saleh SN, Peppiatt-Wildman CM, Large WA. Multiple activation mechanisms of store-operated TRPC channels in smooth muscle cells. *J Physiol* 2007;583:25-36.
- Sweeney M, Yu Y, Platoshyn O, Zhang S, McDaniel SS, Yuan JX. Inhibition of endogenous TRP1 decreases capacitative Ca^{2+} entry and attenuates pulmonary artery smooth muscle cell proliferation. *Am J Physiol Lung Cell Mol Physiol* 2002;283:L144-55.
- Golovina VA, Platoshyn O, Bailey CL, Wang J, Limsuwan A, Sweeney M, et al. Upregulated TRP and enhanced capacitative Ca^{2+} entry in human pulmonary artery myocytes during proliferation. *Am J Physiol Heart Circ Physiol* 2001;280:H746-55.
- Large WA, Wang Q. Characteristics and physiological role of the Ca^{2+} -activated Cl^- conductance in smooth muscle. *Am J Physiol* 1996;271:C435-54.
- Leblanc N, Ledoux J, Saleh S, Sanguinetti A, Angermann J, O'Driscoll K, et al. Regulation of calcium-activated chloride channels in smooth muscle cells: a complex picture is emerging. *Can J Physiol Pharmacol* 2005;83:541-56.
- Chipperfield AR, Harper AA. Chloride in smooth muscle. *Prog Biophys Mol Biol* 2000;74:175-221.
- Caputo A, Caci E, Ferrera L, Pedemonte N, Barsanti C, Sondo E, et al. TMEM16A, a membrane protein associated with calcium-dependent chloride channel activity. *Science* 2008;322:590-4.
- Schroeder BC, Cheng T, Jan YN, Jan LY. Expression cloning of TMEM16A as a calcium-activated chloride channel subunit. *Cell* 2008;134:1019-29.
- Yang YD, Cho H, Koo JY, Tak MH, Cho Y, Shim WS, et al. TMEM16A confers receptor-activated calcium-dependent chloride conductance. *Nature* 2008;455:1210-5.
- Cornfield DN, Stevens T, McMurtry IF, Abman SH, Rodman DM. Acute hypoxia increases cytosolic calcium in fetal pulmonary artery smooth muscle cells. *Am J Physiol* 1993;265:L53-6.
- Bonnet S, Belus A, Hyvelin JM, Roux E, Marthan R, Savineau JP. Effect of chronic hypoxia on agonist-induced tone and calcium signaling in rat pulmonary artery. *Am J Physiol Lung Cell Mol Physiol* 2001;281:L193-201.
- Gögelein H, Dahlem D, Englert HC, Lang HJ. Flufenamic acid, mefenamic acid and niflumic acid inhibit single nonselective cation channels in the rat exocrine pancreas. *FEBS Lett* 1990;268:79-82.
- Ottolia M, Toro L. Potentiation of large conductance K_{Ca} channels by niflumic, flufenamic, and mefenamic acids. *Biophys J* 1994;67:2272-9.
- Foster RR, Zadeh MA, Welsh GI, Satchell SC, Ye Y, Mathieson PW, et al. Flufenamic acid is a tool for investigating TRPC6-mediated calcium signalling in human conditionally immortalised podocytes and HEK293 cells. *Cell Calcium* 2009;45:384-90.
- Fuchs B, Dietrich A, Gudermand T, Kalwa H, Grimminger F, Weissmann N. The role of classical transient receptor potential channels in the regulation of hypoxic pulmonary vasoconstriction. *Adv Exp Med Biol* 2010;661:187-200.
- Lu W, Wang J, Shimoda LA, Sylvester JT. Differences in STIM1 and TRPC expression in proximal and distal pulmonary arterial smooth muscle are associated with differences in Ca^{2+} responses to hypoxia. *Am J Physiol Lung Cell Mol Physiol* 2008;295:L104-13.
- Song MY, Makino A, Yuan JX. STIM2 contributes to enhanced store-operated Ca^{2+} entry in pulmonary artery smooth muscle cells from patients with idiopathic pulmonary arterial hypertension. *Pulm Circ* 2011;1:84-94.
- Ng LC, McCormack MD, Airey JA, Singer CA, Keller PS, Shen XM, et al. TRPC1 and STIM1 mediate capacitative Ca^{2+} entry in mouse pulmonary arterial smooth muscle cells. *J Physiol* 2009;587:2429-42.
- Jung S, Strotmann R, Schultz G, Plant TD. TRPC6 is a candidate channel involved in receptor-stimulated cation currents in A7r5 smooth muscle cells. *Am J Physiol Cell Physiol* 2002;282:C347-59.
- Tai K, Hamaide MC, Debaix H, Gailly P, Wibo M, Morel N. Agonist-evoked calcium entry in vascular smooth muscle cells requires IP_3 receptor-mediated activation of TRPC1. *Eur J Pharmacol* 2008;583:135-47.
- Maruyama Y, Nakanishi Y, Walsh EJ, Wilson DP, Welsh DG, Cole WC. Heteromultimeric TRPC6-TRPC7 channels contribute to arginine vasopressin-induced cation current of A7r5 vascular smooth muscle cells. *Circ Res* 2006;98:1520-7.
- Peng G, Lu W, Li X, Chen Y, Zhong N, Ran P, et al. Expression of store-operated Ca^{2+} entry and transient receptor potential canonical and vanilloid-related proteins in rat distal pulmonary venous smooth muscle. *Am J Physiol Lung Cell Mol Physiol* 2010;299:L621-30.
- Xu SZ, Beech DJ. TrpC1 is a membrane-spanning subunit of store-operated Ca^{2+} channels in native vascular smooth muscle cells. *Circ Res* 2001;88:84-7.
- Brueggemann LI, Markun DR, Henderson KK, Cribbs LL, Byron KL. Pharmacological and electrophysiological characterization of store-operated currents and capacitative Ca^{2+} entry in vascular smooth muscle cells. *J Pharmacol Exp Ther* 2006;317:488-99.
- McDaniel SS, Platoshyn O, Wang J, Yu Y, Sweeney M, Krick S, et al. Capacitative Ca^{2+} entry in agonist-induced pulmonary vasoconstriction. *Am J Physiol Lung Cell Mol Physiol* 2001;280:L870-80.
- Lin MJ, Leung GP, Zhang WM, Yang XR, Yip KP, Tse CM, et al. Chronic hypoxia-induced upregulation of store-operated and receptor-operated Ca^{2+} channels in pulmonary arterial smooth muscle cells: A novel mechanism of hypoxic pulmonary hypertension. *Circ Res* 2004;95:496-505.
- Ng LC, Gurney AM. Store-operated channels mediate Ca^{2+} influx and contraction in rat pulmonary artery. *Circ Res* 2001;89:923-9.
- Hassoun PM, Mouthon L, Barberá JA, Eddahibi S, Flores SC, Grimminger F, et al. Inflammation, growth factors, and pulmonary vascular remodeling. *J Am Coll Cardiol* 2009;54:S10-9.
- Greenwood IA, Ledoux J, Leblanc N. Differential regulation of Ca^{2+} -activated Cl^- currents in rabbit arterial and portal vein smooth muscle cells by Ca^{2+} -calmodulin-dependent kinase. *J Physiol* 2001;534:395-408.
- Yuan XJ. Role of calcium-activated chloride current in regulating pulmonary vasomotor tone. *Am J Physiol* 1997;272:L959-68.
- Manoury B, Tamuleviciute A, Tammaro P. TMEM16A/anocatin 1 protein mediates calcium-activated chloride currents in pulmonary arterial smooth muscle cells. *J Physiol* 2010;588:2305-14.
- Hartzell HC, Yu K, Xiao Q, Chien LT, Qu Z. Anocatin/TMEM16 family members are Ca^{2+} -activated Cl^- channels. *J Physiol* 2009;587:2127-39.
- Hwang SJ, Blair PJ, Britton FC, O'Driscoll KE, Hennig G, Bayguinov YR, et al. Expression of anocatin 1/TMEM16A by interstitial cells of Cajal is fundamental for slow wave activity in gastrointestinal muscles. *J Physiol* 2009;587:4887-904.
- Huang F, Rock JR, Harfe BD, Cheng T, Huang X, Jan YN, et al. Studies on expression and function of the TMEM16A calcium-activated chloride channel. *Proc Natl Acad Sci USA* 2009;106:21413-8.
- Zhu MH, Kim TW, Ro S, Yan W, Ward SM, Koh SD, et al. A Ca^{2+} -activated Cl^- conductance in interstitial cells of Cajal linked to slow wave currents and pacemaker activity. *J Physiol* 2009;587:4905-18.
- Rock JR, O'Neal WK, Gabriel SE, Randell SH, Harfe BD, Boucher RC, et al. Transmembrane protein 16A (TMEM16A) is a Ca^{2+} -regulated Cl^- secretory channel in mouse airways. *J Biol Chem* 2009;284:14875-80.

48. Ousingsawat J, Martins JR, Schreiber R, Rock JR, Harfe BD, Kunzelmann K. Loss of TMEM16A causes a defect in epithelial Ca^{2+} -dependent chloride transport. *J Biol Chem* 2009;284:28698-703.
49. Davis AJ, Forrest AS, Jepps TA, Valencik ML, Wiwchar M, Singer CA, et al. Expression profile and protein translation of TMEM16A in murine smooth muscle. *Am J Physiol Cell Physiol* 2010;299:C948-59.
50. Gomez-Pinilla PJ, Gibbons SJ, Bardsley MR, Lorincz A, Pozo MJ, Pasricha PJ, et al. Ano1 is a selective marker of interstitial cells of Cajal in the human and mouse gastrointestinal tract. *Am J Physiol Gastrointest Liver Physiol* 2009;296:G1370-81.
51. Ferrera L, Caputo A, Ubbly I, Bussani E, Zegarra-Moran O, Ravazzolo R, et al. Regulation of TMEM16A chloride channel properties by alternative splicing. *J Biol Chem* 2009;284:33360-8.
52. Stöhr H, Heisig JB, Benz PM, Schöberl S, Milenkovic VM, Strauss O, et al. TMEM16B, a novel protein with calcium-dependent chloride channel activity, associates with a presynaptic protein complex in photoreceptor terminals. *J Neurosci* 2009;29:6809-18.
53. Pifferi S, Dibattista M, Menini A. TMEM16B induces chloride currents activated by calcium in mammalian cells. *Pflügers Arch* 2009;458:1023-38.
54. Stephan AB, Shum EY, Hirsh S, Cygnar KD, Reisert J, Zhao H. ANO2 is the cilia calcium-activated chloride channel that may mediate olfactory amplification. *Proc Natl Acad Sci USA* 2009;106:11776-81.
55. Forrest AS, Angermann JE, Raghunathan R, Lachendro C, Greenwood IA, Leblanc N. Intricate interaction between store-operated calcium entry and calcium-activated chloride channels in pulmonary artery smooth muscle cells. *Adv Exp Med Biol* 2010;661:31-55.
56. Liang W, Ray JB, He JZ, Backx PH, Ward ME. Regulation of proliferation and membrane potential by chloride currents in rat pulmonary artery smooth muscle cells. *Hypertension* 2009;54:286-93.

Source of Support: This work was supported, in part, by grants from the National Heart, Lung, and Blood Institute of the National Institutes of Health (HL066012 and HL098053 to JX-JY), **Conflict of Interest:** None declared.

New features on the Journal's website


Optimized content for mobile and hand-held devices

HTML pages have been optimized of mobile and other hand-held devices (such as iPad, Kindle, iPod) for faster browsing speed. Click on **[Mobile Full text]** from Table of Contents page. This is simple HTML version for faster download on mobiles (if viewed on desktop, it will be automatically redirected to full HTML version)

E-Pub for hand-held devices

EPUB is an open e-book standard recommended by The International Digital Publishing Forum which is designed for reflowable content i.e. the text display can be optimized for a particular display device. Click on **[EPub]** from Table of Contents page. There are various e-Pub readers such as for Windows: Digital Editions, OS X: Calibre/Bookworm, iPhone/iPod Touch/iPad: Stanza, and Linux: Calibre/Bookworm.

E-Book for desktop

One can also see the entire issue as printed here in a 'flip book' version on desktops. Links are available from Current Issue as well as Archives pages. Click on  View as eBook