Check for updates

DE GRUYTER OPEN

NAD

Research report

Anabolic-androgenic steroid use in the Nordic countries: A meta-analysis and meta-regression analysis

DOMINIC SAGOE & TORBJØRN TORSHEIM & HELGE MOLDE & CECILIE SCHOU ANDREASSEN & STÅLE PALLESEN

ABSTRACT

AIMS – To investigate the lifetime prevalence and moderators of non-medical AAS use in the five Nordic countries. METHODS – We conducted a meta-analysis and meta-regression using studies gathered from searches in PsycINFO, PubMed, ISI Web of Science, Google Scholar, and reference checks. Included were 32 studies that provided original data on 48 lifetime prevalence rates based on a total of 233,475 inhabitants of the Nordic countries. RESULTS – The overall lifetime prevalence obtained was 2.1% [95% confidence interval (CI): 1.3-3.4, $I^2 = 99.5$, $P < 0.001$]. The prevalence for males, 2.9% (95% CI: 1.7-4.8, $I^2 = 99.2$, $P < 0.001$), was significantly higher ($Q_{bet} = 40.5$, $P < 0.001$) than the rate for females, 0.2% (95% CI: 0.1-0.4, $I^2 = 90.5$, $P < 0.001$). Sweden has the highest prevalence of AAS use: 4.4%, followed by Norway: 2.4%, Finland: 0.8%, Iceland: 0.7%, and Denmark: 0.5%. Country, sample type, and male sample percentage significantly predicted AAS use prevalence in a meta-regression analysis. No indication of publication bias was found. CONCLUSION – Though subject to some limitations, our findings suggest that non-medical AAS use should be regarded as a serious public health problem in the Nordic countries.

KEYWORDS – anabolic steroids, Nordic countries, Scandinavia, prevalence, meta-analysis, meta-regression

Submitted 19.03.2014 Final version accepted 21.05.2014

Introduction

Anabolic–androgenic steroids (AAS) are a group of hormones including testosterone and its synthetic derivatives. These hormones are used clinically to treat conditions such as reproductive system dysfunction, breast cancer, and anemia. Increasingly, some healthy individuals have been using AAS for non-medical purposes (Boyadjiev, Georgieva, Massaldjieva, & Gueorguiev, 2000; Sagoe, Molde, Andreassen, Torsheim, & Pallesen, 2014a). Non-medical AAS use was mainly restricted to elite athletes and bodybuilders in the 1960s and 1970s as a means to enhance strength and athletic performance (Yesalis & Bahrke, 1995). Non-medical AAS use has however spread into the general population in the last few decades (Kanayama, Hudson, & Pope, 2008) mainly driven by a desire for boosting physical strength and improving appearance (Kanayama, Hudson, & Pope, 2010; Parkinson & Evans, 2006; Tanner, Miller, & Alongi, 1995). Indeed, results from a recent global epidemiological investigation indicates that recreational sportspeople constitute the largest group of AAS users (Sagoe et al., 2014a).

There is extensive evidence connecting

Unauthenticated
Download Date | 12/16/16 1:04 PM

long-term AAS use to criminality (Lood, Eklund, Garle, & Ahlner, 2012; Lundholm, Haggård, Möller, Hallqvist, & Thiblin, 2013) and several debilitating physical and psychological disorders as well as mortality (Bahrke & Yesalis, 2004; Dodge & Hoagland, 2011; Hoffman & Ratamess, 2006; Kanayama et al., 2008; Pallesen, Jøsendal, Johnesen, Larsen, & Molde, 2006; Pope, & Kanayama, 2012; Pope, Kanayama, & Hudson, 2012; Urhausen et al., 2004). Despite such evidence of the harmful consequences of non-medical use, AAS seem to be among the least studied of the world's major abused drugs (Pope et al., 2013). Hence, Degenhardt and Hall (2012) did not investigate the prevalence of AAS use in their study of the global prevalence of illicit drug use and dependence because of the scarcity of information regarding the AAS use epidemiology compared to drugs such as cocaine and cannabis.

Kanayama, Hudson, and Pope (2012) suggest that the prevalence of AAS use is higher in the Nordic countries compared to most other parts of the world. Several epidemiological investigations of AAS use have been conducted in the Nordic countries. However, to our knowledge, there has never been a pan-Nordic meta-analytic investigation of the prevalence of AAS use. Thus, we conducted a meta-analysis on the lifetime prevalence of non-medical AAS use in the Nordic countries. We calculated overall prevalence estimates across the Nordic countries, gender, publication year, sample type, and sampling method. We further investigated the predictive effect of the above study variables on the overall lifetime prevalence rate using a meta-regression analysis.

Methods

Literature search strategy and inclusion criteria


We conducted a systematic and comprehensive literature search in PsycINFO, PubMed, ISI Web of Science, and Google Scholar for articles published between 1970 and July 2013. The following keywords: 'anabol*', 'steroid*', 'doping' were each used in combination with 'preval*', 'epidem*', and 'incidence' for the search. Studies were included if they satisfied the following criteria: (a) they were published between 1970 and July 2013 (b) they presented original data on the lifetime prevalence rate of AAS use, and (c) they were published in English or a Nordic language. From an initial pool of 16,626 hits, 311 full-text papers were retrieved for further evaluation. After screening the 311 full-text papers for eligibility, 162 met the inclusion criteria.

In addition, we checked the references of identified studies in search of potential unidentified studies conducted for any Nordic country or countries. We also searched online databases and websites for data on lifetime prevalence rates of AAS use in general population or household surveys, school surveys, government reports, and regional reports for any Nordic country or countries. Twenty-five (25) new articles were identified through this grey literature search. Thus, we screened a total of 187 studies for eligibility. After screening the 187 articles, 32 articles presented original data on lifetime prevalence of AAS use for the Nordic countries (Denmark, Finland, Iceland, Norway, and Sweden) and were consequently included in the present study.

In the search for grey literature, we ad-

Unauthenticated
Download Date | 12/16/16 1:04 PM

Figure 1. Flow diagram of systematic literature search.



hered to Calabria et al.'s (2009) strategy that if data from a representative national study existed for a country, data from a study with similar a methodology and target age group was not included in order to prevent duplicates. Thus, for adolescents, we relied on the European School Survey Project on Alcohol and Other Drugs (ESPAD). ESPAD is a survey of European adolescents (of about 35 countries) conducted every fourth year since 1995 (Hibell et al., 2000, 2004, 2007, 2009, 2012). The literature search was in line with the guidelines of Preferred Reporting Items for Systematic Reviews and Meta-Analyses (Moher, Liberati, Tetzlaff, & Altman, 2009) and the Meta-analysis of Observational Studies in Epidemiology (MOOSE) group (Stroup et al., 2000). Figure 1 presents the literature search and selection process.

Description of studies

Five articles (Hibell et al., 2000, 2004, 2009, 2012; Nilsson, Baigi, Marklund, & Fridlund, 2001a) out of the 32 articles identified presented prevalence rates of AAS use for multiple studies totaling up to 16 separate studies. Thus, a total of 48 separate studies were identified which provided original data on lifetime prevalence rates of AAS use for the Nordic countries. A total of 233,475 inhabitants [61,329 females, 85,313 males – some studies do not present a sample breakdown in terms of gender] of

Unauthenticated
Download Date | 12/16/16 1:04 PM

the Nordic countries participated in these studies. The year of publication of the studies ranged from 1974 (Haug & Ingvaldsen, 1974; Solberg, 1974) to 2013 (Nøkleby, 2013; Singhammer, 2013; Lindqvist et al., 2013). Most studies were conducted in Sweden (n = 20), followed by Norway (n = 13), Finland (n = 7), Iceland (n = 5), and Denmark (n = 3). The study characteristics are presented in Table 1.

Data extraction

Studies were scrutinized and selected based on their titles, abstracts, and subject matter by the first author. We developed a standardized data extraction form. Using this form, the first author and another reviewer independently extracted data from the identified studies. The following data were extracted from the included studies: author name and publication year, country, and region of research, type of sample (prisoners and arrestees, recreational sportspeople, drug users, athletes, non-athletes, and high school), assessment method (questionnaires or interview), sampling method (random or non-random), sample size (total, male, and female), age of participants (range, mean, and standard deviation), response rate, and lifetime prevalence rate of AAS use reported (male, female, and overall). The inter-reviewer reliability for the reviewers for 162 studies identified in the first part of the search was found to be Kappa = 0.854 ($P < 0.001$) indicating an almost perfect agreement between the two reviewers (Viera & Garrett, 2005). Consensus was reached on discrepant extractions between the two reviewers through further review and discussion of the articles. A final table of all studies is presented in Table 1.

Publication bias

We assessed publication bias visually by funnel plot and statistically by the trim and fill procedure (Duval & Tweedie, 2000) in Comprehensive Meta-Analysis version 3.0 (Biostat Inc., 2014). The point estimate and 95 percent confidence interval (95% CI) for the combined studies under the random-effects model was 2.1% (95% CI: 1.3-3.4). These values were unchanged when the trim and fill function was applied indicating the absence of publication bias. The absence of publication bias was further confirmed as inspection of the funnel plot showed a symmetrical distribution of studies in terms of prevalences.

Statistical analysis

We conducted a meta-analysis and meta-regression analysis to estimate the lifetime prevalence, as well as predictors of lifetime AAS use prevalence, in the Nordic countries. The meta-analysis and meta-regression analysis were conducted using Comprehensive Meta-Analysis, version 3.0 (Biostat Inc., 2014). In calculating overall prevalence figures, relevant study characteristics (see Table 1) were coded for each study in Comprehensive Meta-Analysis, version 3.0. Thus, we were able to calculate pooled prevalence figures for relevant study characteristics based on the coded data for each study.

The calculation of prevalence rates and 95% CI was done using a random-effects model because it is most useful when the studies included in the meta-analysis may not be representative of the entirety of studies on the topic. Thus, results generated from the use of the random-effects model have more external validity than results generated from the use of the fixed-

Unauthenticated
Download Date | 12/16/16 1:04 PM

Table 1. Characteristics of studies on the lifetime prevalence of AAS use in the Nordic countries.

First author, year	Country	Sample type	Assessment method	Sampling method	*N*	Sample size (male)	Sample size (female)	Age range (y)	Age mean	Age SD	Prevalence (male) %	Prevalence (female) %	Prevalence (overall) %	Response rate %
Gårevik 2010	Sweden	Arrested drug users	I	NR	56	-	-	-	30†	7.4†	-	-	81	-
Hakansson 2012	Sweden	General	Q	R	21211	14160	7051	15-64	29	-	1.7	0.3	1.7	32
Haug 1974	Norway	Weightlifters	Q	NR	15	15	-	-	-	-	87	-	87	38
Hibell 1997	Sweden	High school	Q	R	3472	1746	1725	15-16	-	-	2	0	1	86
Hibell 2000	Denmark	High school	Q	R	1790	875	915	15-16	-	-	1	0	1	92
Hibell 2000	Finland	High school	Q	R	3286	1646	1640	15-16	-	-	1	0	1	90
Hibell 2000	Iceland	High school	Q	R	3524	1758	1766	15-16	-	-	1	0	1	89
Hibell 2000	Norway	High school	Q	R	3918	1980	1811	15-16	-	-	2	0	1	90
Hibell 2000	Sweden	High school	Q	R	3455	1715	1730	15-16	-	-	2	0	1	87
Hibell 2004	Denmark	High school	Q	R	2978	1504	1474	15-16	-	-	2	1	1	89
Hibell 2004	Finland	High school	Q	R	3543	1739	1804	15-16	-	-	1	0	1	91
Hibell 2004	Iceland	High school	Q	R	3348	1728	1604	15-16	-	-	1	1	1	81
Hibell 2004	Norway	High school	Q	R	3833	1945	1888	15-16	-	-	2	0	1	87
Hibell 2004	Sweden	High school	Q	R	3232	1592	1640	15-16	-	-	1	0	1	87
Hibell 2007	Sweden	High school	Q	R	4245	1960	2285	15-16	-	-	1	0	1	84
Hibell 2009	Finland	High school	Q	R	4988	2297	2691	15-16	-	-	1	0	1	91
Hibell 2009	Iceland	High school	Q	R	3510	1797	1713	15-16	-	-	1	0	1	81
Hibell 2009	Norway	High school	Q	R	3482	1778	1704	15-16	-	-	2	0	1	89
Hibell 2009	Sweden	High school	Q	R	3179	1550	1629	15-16	-	-	2	0	1	84
Hibell 2012	Denmark	High school	Q	R	2181	979	1202	15-16	-	-	0	1	1	89
Hibell 2012	Finland	High school	Q	R	3744	1815	1929	15-16	-	-	1	0	1	90
Hibell 2012	Iceland	High school	Q	R	3333	1717	1616	15-16	-	-	1	0	1	81
Hibell 2012	Norway	High school	Q	R	2938	1498	1440	15-16	-	-	1	0	1	88
Hibell 2012	Sweden	High school	Q	R	2569	1311	1258	15-16	-	-	1	0	1	85
Kindlundh 1998	Sweden	Adolescents	Q	R	2742	1353	1364	16-19	-	-	1.7	0.1	0.9	80.8
Klötz 2010	Sweden	Prisoners	I	NR	59	59	-	21-52†	30.1†	6.5†	55.9	-	55.9	50
Korte 1998	Finland	Prisoners	Q	R	354	354	-	18-76	32	10	9.6	-	9.6	82.5
Leifman 2011	Sweden	Gym users	Q	R	1752	1183	563	16 – 49*	33	-	3.9	0.2	2.05	91.1
Lindqvist 2013	Sweden	Former athletes	Q	NR	683	683	-	39-82	57	-	21	-	21	68.6
Lindström 1990	Sweden	Bodybuilders	Q	NR	171	138	33	15+	25	-	38.4	9.09	32.7	90
Ljungqvist 1975	Sweden	Athletes	Q	NR	99	-	-	-	-	-	-	-	31	69
Lundholm 2010	Sweden	Prisoners	I	NR	3597	3201	396	-	-	-	28.1	5	26	-
Mattila 2009	Finland	Adolescents	Q	R	22519	-	-	12-18	-	-	0.5	0.2	0.3	74
Mattila 2010	Finland	Military conscripts	Q	NR	10396	10396	-	18-29	-	-	0.9	-	0.9	96
Nilsson 1995	Sweden	High school	Q	R	1383	688	695	14-19	-	-	5.8	1.0	3.4	96
Nilsson 2001a	Sweden	Adolescents	Q	R	345	345	-	16-17	-	-	8.7	-	8.7	96
Nilsson 2001a	Sweden	Adolescents	Q	R	451	451	-	16-17	-		5.76	-	5.76	-
Nilsson 2001b	Sweden	Adolescents	Q	R	5827	-	-	16-17	-	-	2.9	0	1.4	95
Nilsson 2005	Sweden	Adolescents	Q	R	4049	4049	-	14-18	-	-	1.2	-	1.2	92.7
Nøkleby 2013	Norway	Drug users	Q	NR	109	79	30	17-50	-	-	40.5	20	35	85
Pallesen 2006	Norway	Adolescents	Q	R	1351	703	642	-	17.5	2.2	3.6	0.6	2.1	69.8
Sagoe 2014b	Norway	Adolescents	Q	R	2055	963	1088	17	17	0	0.52	0.09	0.30	70.4
Singhammer 2013	Denmark	Recreational sportspeople	Q	R	5010	2006	3004	15-60	-	-	-	-	1.8	-
SNIPH 2009	Sweden	General	Q	R	58000	-	-	15-64	-	-	1	0	1	-
Solberg 1974	Norway	Weightlifters	Q	NR	25	25	-	-	-	-	24	-	24	68
Thorlindsson 2010	Iceland	High school	Q	R	10918	5195	5585	15-24	17.7	1.84	1.6	0.2	0.9	-
Wichstrøm 2001	Norway	Adolescents	Q	R	8508	3931	4577	14-25	17.33	2.18	1.2	0.6	0.8	78
Wichstrøm 2006	Norway	High school	Q	R	2924	1277	1647	15-24	22.1	1.9	-	-	1.9	68

I: Interview; Q: Questionnaire; NR: Non-random sampling; R: Random sampling; SNIPH: Swedish National Institute of Public Health: †: AAS users; *: Majority of participants

Unauthenticated Download Date | 12/16/16 1:04 PM

Table 2. Prevalence rates, confidence intervals, and heterogeneity statistics.

	N	p%	95% CI	*Q*	df(Q)	I^2
Overall	48	2.1	1.3–3.4	8724.4*	47	99.5
Gender						
Male	41	2.9	1.7–4.8	5107.5*	40	99.2
Female	32	0.2	0.1–0.4	326.8*	31	90.5
Country						
Sweden	20	4.4	2.0–9.4	6209.4*	19	99.7
Norway	13	2.4	1.2–4.7	519.6*	12	97.7
Finland	7	0.8	0.3–1.8	284.5*	6	97.9
Iceland	5	0.7	0.5-0.9	13.3**	4	69.9
Denmark	3	0.5	0.4–0.6	0.1ns	2	0
Sample type						
Drug users	2	59.2	16.5-91.4	26.9*	1	96.3
Athletes	5	32.3	22.0–44.7	28.8*	4	86.1
Prisoners and arrestees	3	26.2	11.5–49.3	67.5*	2	97.0
Recreational sportspeople	1	2.1	1.5–2.8	0ns	0	0
Non-athletes	3	1.2	0.8–1.7	71.7*	2	97.2
High school	34	0.9	0.7–1.1	611.8*	33	94.6
Sampling method						
Non-random	12	18.7	6.0–45.5	5857.6*	11	99.8
Random	36	1.0	0.8–1.3	707.0*	35	95.0
Publication year						
1970–1989	3	44.8	18.5–74.4	12.5**	2	84.1
1990–1999	7	3.8	1.3–11.1	800.4*	6	99.3
2000–2013	38	1.4	0.8–2.6	7195.7*	37	99.5

* $p < 0.001$, ** $p < 0.01$, *ns* = not significant.
N: number of studies included in the analysis. p%: prevalence (%). 95% CI: 95% confidence interval. *Q*: heterogeneity statistic. df(*Q*): *Q*'s degrees of freedom. I^2: heterogeneity index.

effect model (Borenstein, Hedges, Higgins, & Rothstein, 2009).The *Q* statistic and the I^2 index were used to assess the heterogeneity of the prevalences.

Furthermore, in order to identify moderator variables that could explain the variance in the overall prevalence rate, a meta-regression analysis was performed under a random-effects model. The moderator variables included in the meta-regression analysis were: country (Sweden, Norway, Finland, Iceland, and Denmark), sample type (drug users, athletes, prisoners and arrestees, recreational sportspeople, non-athletes, and high school), sampling method (non-random and random), publication year (1970–1989, 1990–1999, and 2000–2013), and the percentage of males in the sample [lower than or equal to fifty percent (≤ 50%), greater than seventy-five percent (> 75%), greater than fifty percent but lower than or equal to seventy-five percent (> 50% to ≤ 75%), and percentage not provided]. The category with the highest number of studies was used as a contrast for each moderator variable.

Results

Prevalence rates and heterogeneity testing

Table 2 presents the total number of studies, the prevalence rates and confidence limits as well as the heterogeneity statistics (*Q* and I^2) for the overall population of the five Nordic countries, males, females, countries, sample type, sampling method, and publication year.

From Table 2, the overall prevalence rate obtained from 48 studies was 2.1% (95% CI : 1.3-3.4%, $I^2 = 99.5$, $P < 0.001$). In addition, the prevalence rate for males, 2.3%, was significantly higher ($Q_{bet} = 40.5$, *df* =

Unauthenticated
Download Date | 12/16/16 1:04 PM

Table 3. Meta-regression analysis of the predictors of AAS use prevalence.

	B	SE	95% CI		*Z*	*P*
Country						
Sweden†						
Norway	-.14	.29	-.70	.43	-.48	.633
Finland	-1.05	.29	-1.62	-.50	-3.68	.000
Iceland	-.48	.36	-1.18	.23	-1.33	.184
Denmark	-.75	.43	-1.60	.09	-1.75	.080
Sample type						
High school†						
Drug users	4.93	.64	3.68	6.17	7.76	.000
Athletes	2.34	.69	.99	3.68	3.41	.000
Prisoners and arrestees	2.65	.53	1.60	3.69	4.97	.000
Recreational sportspeople	.61	.67	-.70	1.93	.92	.360
Non-athletes	-.06	.47	-.97	.85	-.13	.899
Sampling method						
Non-random†						
Random	-.19	.48	-1.13	.75	-.40	.691
Publication year						
2000–2013†						
1970–1989	1.29	.67	-.02	2.61	1.93	.054
1990–1999	.28	.29	-.28	.84	.99	.325
Percentage of males in sample						
≤ 50%†						
> 75%	1.29	.36	.59	1.99	3.60	.000
> 50% to 75%	.13	.25	-.36	.62	.53	.59
% not provided	.39	.44	-.47	1.25	.89	.372

R^2 = 89.0%.
† = Reference category.

1, $P < 0.001$) than the prevalence rate for females, 0.2%.

When prevalence was investigated by country, Sweden had the highest overall prevalence rate of AAS use: 4.4%, followed by Norway: 2.4%, Finland: 0.8%, Iceland: 0.7%, and Denmark: 0.5%. In addition, apart from Denmark, the heterogeneity statistic (*Q*) for the overall prevalence rates reached statistical significance ($P < 0.001$).

With reference to sample type, drug users had the highest overall prevalence rate: 59.2%, followed by athletes: 32.3%, prisoners and arrestees: 26.2%, and recreational sportspeople 2.1%. Moreover, prevalence rate for non-athletes was 1.2% while high school students had the lowest prevalence rate: 0.9%. With the exception of recreational sportspeople, the heterogeneity statistic (*Q)* for the overall prevalence rates for all sample types reached statistical significance ($P < 0.001$).

In addition, studies that used non-random sampling methods had a higher overall prevalence rate, 18.7%, than studies that used random sampling methods, 1.0%.

Furthermore, publication years 1970 to 1989 had the highest overall prevalence rate: 44.8%, followed by 1990 to 1999: 3.8%, and 2000 to 2013: 1.4%. The heterogeneity statistic for the prevalence rates, *Q*, also reached statistical significance ($P < 0.001$) for all publication years.

Meta-regression analysis

We performed a meta-regression analysis to evaluate the effect of country, sample type, sampling method, publication year, and the percentage of males in the sample on the overall prevalence of AAS use.

Unauthenticated
Download Date | 12/16/16 1:04 PM

Of these variables, country, sample type (athletes, drug users, and prisoners and arrestees), and percentage of males in the sample [greater than seventy-five percent (> 75%)] significantly predicted AAS use prevalence. Together, they accounted for 89.0% of the variance in the overall AAS use prevalence rate. The results are presented in Table 3.

Discussion

This paper presents, to our knowledge, the first-ever meta-analysis and meta-regression analysis of the lifetime prevalence of AAS use specifically for the Nordic countries. The lifetime prevalence rate across all studies was 2.1%. Moreover, the overall lifetime prevalence rate for males, 2.3%, was significantly higher than the overall lifetime prevalence rate for females, 0.2% confirming the preponderance of available evidence (Andrade et al., 2012; Johnson, Jay, Shoup, & Rickert, 1989; Kindlundh, Isacson, Berglund, & Nyberg, 1998; Sagoe et al., 2014a).

We found further support for this result in the finding that percentage of males in samples significantly predicted prevalence in the meta-regression analysis consistent with results from a worldwide prevalence study (Sagoe et al., 2014a). In addition, we found that Sweden has the highest prevalence rate of AAS use: 4.4%, followed by Norway: 2.4%, Finland: 0.8%, Iceland: 0.7%, and Denmark: 0.5%. In corroboration of this finding, country was a significant predictor of prevalence in the meta-regression analysis. Moreover, the prevalence for Finland was significantly lower than prevalence for Sweden in the meta-regression analysis.

Our finding that outside the arena of competitive athletics, prevalence of AAS use is highest among drug users and prisoners and arrestees is further consistent with a recent study that found a very high incidence rate of AAS and polydrug use in a laboratory testing of the urine samples of Swedish prisoners (Lood et al., 2012). Similarly, consistent with Striegel et al. (2006) who found that athletic involvement has a significant positive correlation with AAS use, we found that athletes and recreational sportspeople have higher prevalence of AAS use than non-athletes. This result corroborates evidence suggesting that the odds of AAS use increases by about 91% with participation in at least one sport (Dodge & Jaccard, 2006; Lorang et al., 2011). Moreover, consistent with results from a global investigation of AAS use prevalence (Sagoe et al., 2014a), sample type significantly predicted prevalence in the meta-regression analysis.

The finding that studies using non-random sampling methods report a higher prevalence rate than studies based on random sampling methods seems to be related to the fact that the predominance of non-randomly selected samples comprised athletes, prisoners, arrestees, and drug users among whom AAS use prevalence is relatively higher compared to high school students and non-athletes, as previously found (Baker, Thomas, Davies, & Graham, 2008; Bojsen-Møller & Christiansen, 2010; Grace, Baker, & Davies, 2001; Lood et al., 2012). However, sampling method was insignificant in the meta-regression analysis indicating that other moderators better accounted for the heterogeneity in prevalence.

Unauthenticated
Download Date | 12/16/16 1:04 PM

Study strengths and limitations

This study, to our knowledge, is the first to have systematically examined the lifetime prevalence rate of non-medical AAS use specifically in the Nordic countries by a quantitative meta-analytic approach. Thus, the prevalence estimates in the present study constitute the best currently available basis for policymaking and planning in the Nordic countries. The systematic nature of this research, the large number of included studies and participants, and the analysis of the data using meta-analysis and meta-regression analysis are also notable assets.

The present meta-analysis, however, has some limitations worth noting in the interpretation of findings. First, the prevalence rates reported in the studies included in our meta-analysis may have exaggerated our final prevalence estimates. Kanayama et al. (2007) suggest that prevalence rates of AAS use are sometimes exaggerated because some respondents answer that they have used AAS when in fact they have used some non-AAS supplement they believed was an AAS. This problem has exacerbated with the proliferation of supplements since the 1990s as it has become more difficult to determine whether a person is using AAS or some non-AAS substance. It is important to note, however, that this problem with false-positive responses may be minimal in the Nordic countries where most questionnaires are administered in Nordic languages rather than English. This is because questionnaires administered in non-English languages may be better at differentiating substances containing AAS from non-AAS substances. In addition, the overall prevalence figures reported in the present study may have been influenced by the inclusion of studies on some groups/populations noted for AAS use such as athletes, offenders, and drug users (Sagoe et al., 2014a). Still, we break down the estimates for these different groups/populations in order to present nuanced information about the prevalence estimates.

Furthermore, the present study investigated the lifetime prevalence of AAS use which, expectedly, should be higher than current prevalence because lifetime prevalence estimates due to their retrospective nature and wider period of coverage are more vulnerable to recall bias compared to current prevalence which cover a shorter period of use (Gmel & Daeppen, 2007). Moreover, in contrast to current prevalences, lifetime prevalence estimates cannot be validated against objective measures such as urine testing (Pagonis et al., 2006).

Although subject to the limitations noted above, our prevalence estimates suggest that non-medical AAS use should be considered a major public health problem in the Nordic countries and must require the attention of policy makers and researchers. In this regard, efforts need to be made in all the Nordic countries not only to deal with this problem, but also to monitor trends in the incidence and prevalence of AAS use. This research provides a strong foundation that can be built upon with the emergence of more evidence on AAS use in the Nordic countries.

Unauthenticated
Download Date | 12/16/16 1:04 PM

Declaration of interest None.

Dominic Sagoe, PhD Cand.
Department of Psychosocial Science
University of Bergen, Norway
E-mail: Dominic.Sagoe@psysp.uib.no

Torbjørn Torsheim, PhD
Department of Psychosocial Science
University of Bergen, Norway
E-mail: Torbjoern.Torsheim@psysp.uib.no

Helge Molde, PhD
Department of Clinical Psychology
University of Bergen, Norway
E-mail: Helge.Molde@psysp.uib.no

Cecilie Schou Andreassen, PhD
Department of Psychosocial Science
University of Bergen, Norway;
The Competence Center,
Bergen Clinics Foundation, Norway
E-mail: Cecilie.Andreassen@psych.uib.no

Ståle Pallesen, PhD
Department of Psychosocial Science
University of Bergen, Norway
E-mail: Staale.Pallesen@psysp.uib.no

REFERENCES

Andrade, A. G., Duarte, P. C. A. V., Barroso, L. P., Nishimura, R., Alberghini, D. G., & Oliveira, L. G. (2012). Use of alcohol and other drugs among Brazilian college students: Effects of gender and age. *Revista Brasileira de Psiquiatria, 34*(3), 294–305.

Bahrke, M. S., & Yesalis, C.E. (2004). Abuse of anabolic androgenic steroids and related substances in sport and exercise. *Current Opinion in Pharmacology, 4*(6), 614–620.

Baker, J. S., Thomas, N. E., Davies, B., & Graham, M. R. (2008). Anabolic androgenic steroid (AAS) abuse: Not only an elite performance issue? *Open Sports Medicine Journal, 2*, 38–39.

Biostat Inc. (2014). *Comprehensive Meta-Analysis Version 3.0*. Englewood, NJ.: Biostat Inc.

Bojsen-Møller, J., & Christiansen, A. V. (2010). Use of performance-and image-enhancing substances among recreational athletes: A quantitative analysis of inquiries submitted to the Danish anti-doping authorities. *Scandinavian Journal of Medicine and Science in Sports, 20*(6), 861–867.

Borenstein, M., Hedges, L., Higgins, J., & Rothstein, H. (2009*). Introduction to Meta-Analysis*. Chichester, West Sussex: Wiley-Blackwell.

Boyadjiev, N., Georgieva, K., Massaldjieva, R., & Gueorguiev, S. (2000). Reversible hypogonadism and azoospermia as a result of anabolic-androgenic steroid use in a bodybuilder with personality disorder. A case report. *Journal of Sports Medicine and Physical Fitness, 40*(3), 271–274.

Calabria, B., Degenhardt, L., Nelson, P., Bucello, C., Lynskey, M., Hall, W., Roberts, A., & McLaren, J. (2009). *Methodology used in a systematic review of evidence on the prevalence of cannabis use and dependence.* Global Burden of Disease Mental Disorders and Illicit Drug Use Expert group. Illicit drugs discussion paper No. 14. Sydney: National Drug and Alcohol Research Centre, University of New South Wales.

Degenhardt, L., & Hall, W. (2012). The extent of illicit drug use, dependence, and their contribution to global burden of disease.

Unauthenticated
Download Date | 12/16/16 1:04 PM

Lancet, 379(9810), 55–70.
Dodge, T., & Hoagland, M. F. (2011). The use of anabolic androgenic steroids and polypharmacy: a review of the literature. *Drug and Alcohol Dependence, 114*(2-3), 100–109.
Dodge, T. L., & Jaccard, J. J. (2006). The effect of high school sports participation on the use of performance-enhancing substances in young adulthood. *Journal of Adolescent Health, 39*(3), 367–373.
Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot–based method of testing and adjusting for publication bias in meta-analysis. *Biometrics, 56*(2), 455–463.
Gårevik, N., & Rane, A. (2010). Dual use of anabolic-androgenic steroids and narcotics in Sweden. *Drug and Alcohol Dependence, 109*(1), 144–146.
Gmel, G., & Daeppen, J. B. (2007). Recall bias for seven-day recall measurement of alcohol consumption among emergency department patients: Implications for case-crossover designs. *Journal of Studies on Alcohol and Drugs, 68*, 303–310.
Grace, F., Baker, J., & Davies, B. (2001). Anabolic androgenic steroid use in recreational gym users: a regional sample of the Mid-Glamorgan area. *Journal of Substance Use, 6*(3), 189–195.
Hakansson, A., Mickelsson, K., Wallin, C., & Berglund, M. (2012). Anabolic androgenic steroids in the general population: user characteristics and associations with substance use. *European Addiction Research, 18*(2), 83–90.
Haug, J., & Ingvaldsen, R. (1974). *Noen synspunkter pa bruken av hormonpreparater blant aktive idrettsutøvere* [Some views on the use of hormone drugs among active athletes]. Oslo: Institute of Psychology, University of Oslo.
Hibell, B., Anderson, B., Ahlström, S., Balakieva, O., Bjarnason, T., Kokkevi, A., & Morgan, M. (2000). *The 1999 ESPAD report. Alcohol and other drug use among students in 30 European countries.* Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN).
Hibell, B., Anderson, B., Bjarnason, T., Ahlström, S., Balakieva, O., Kokkevi, A., & Morgan, M. (2004). *The ESPAD Report 2003. Alcohol and other drug use among students in 35 European countries*. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN).
Hibell, B., Andersson, B., Bjarnason, T., Kokkevi, A., Morgan, M., & Narusk, A. (1997). *Alcohol and other drug use among students in 26 European countries*. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN) and the Pompidou Group at the Council of Europe.
Hibell, B., Guttormsson, U., Ahlström, S., Balakireva, O., Bjarnason, T., Kokkevi, A., & Kraus, L. (2012). *The 2011 ESPAD report: substance use among students in 36 European countries*. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN).
Hibell, B., Guttormsson, U., Ahlström, S., Balakireva, O., Bjarnason, T., Kokkevi, A., & Kraus, L. (2007). *The 2007 ESPAD report. Substance use among students in 35 European countries*. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN), The European Monitoring Centre for Drugs and Drug Addiction (EMCDDA).
Hibell, B., Guttormsson, U., Ahlström, S., Balakireva, O., Bjarnason, T., Kokkevi, A., & Kraus, L. (2009). *The 2007 ESPAD report. Substance use among students in 35 European countries*. The Swedish Council for Information on Alcohol and Other Drugs, Stockholm, Sweden.
Hoffman, J. R., & Ratamess, N. A. (2006). Medical issues associated with anabolic steroid use: are they exaggerated. *Journal of Sports Science and Medicine, 5*(2), 182–193.
Johnson, M. D., Jay, M. S., Shoup, B., & Rickert, V. I. (1989). Anabolic steroid use by male adolescents. *Pediatrics, 83*(6), 921–924.
Kanayama, G., Boynes, M., Hudson, J. I., Field, A. E., & Pope Jr., H. G., (2007). Anabolic steroid abuse among teenage girls: An illusory problem? *Drug and Alcohol Dependence, 88*, 156–162.

Unauthenticated
Download Date | 12/16/16 1:04 PM

Kanayama, G., Hudson, J.I., & Pope Jr., H.G. (2008). Long-term psychiatric and medical consequences of anabolic–androgenic steroid abuse: A looming public health concern? *Drug and Alcohol Dependence, 98*(1), 1–12.

Kanayama, G., Hudson, J.I., & Pope Jr., H.G. (2010). Illicit anabolic–androgenic steroid use. *Hormones and Behavior, 58*(1), 111–121.

Kanayama, G., Hudson, J.I., & Pope Jr., H.G. (2012). Culture, psychosomatics and substance abuse: The example of body image drugs. *Psychotherapy and Psychosomatics, 81*(2), 73–78.

Kindlundh, A.M., Isacson, D.G., Berglund, L., & Nyberg, F. (1998). Doping among high school students in Uppsala, Sweden: A presentation of the attitudes, distribution, side effects, and extent of use. *Scandinavian Journal of Public Health, 26*(1), 71–74.

Klötz, F., Petersson, A., Hoffman, O., & Thiblin, I. (2010). The significance of anabolic androgenic steroids in a Swedish prison population. *Comprehensive Psychiatry, 51*(3), 312–318.

Korte, T., Pykäläinen, J., & Seppälä, T. (1998). Drug abuse of Finnish male prisoners in 1995. *Forensic Science International, 97*(2-3), 171–183.

Leifman, H., Rehnman, C., Sjöblom, E., & Holgersson, S. (2011). Anabolic androgenic steroids—use and correlates among gym users—an assessment study using questionnaires and observations at gyms in the Stockholm region. *International Journal of Environmental Research and Public Health 8*(7), 2656–2674.

Lindqvist, A.S., Moberg, T., Eriksson, B.O., Ehrnborg, C., Rosén, T., & Fahlke, C. (2013). A retrospective 30-year follow-up study of former Swedish-elite male athletes in power sports with a past anabolic androgenic steroids use: A focus on mental health. *British Journal of Sports Medicine, 47*, 965–969.

Lindström, M., Nilsson, A., Katzman, P., Janzon, L., & Dymling, J.F. (1990). Use of anabolic-androgenic steroids among body builders—frequency and attitudes. *Journal of Internal Medicine, 227*(6), 407–411.

Ljungqvist, A. (1975). The use of anabolic steroids in top Swedish athletes. *British Journal of Sports Medicine, 9*(2), 82.

Lood, Y., Eklund, A., Garle, M., & Ahlner, J. (2012). Anabolic androgenic steroids in police cases in Sweden 1999–2009. *Forensic Science International, 219*(1-3), 199–204.

Lorang, M., Callahan, B., Cummins, K.M., Achar, S., & Brown, S.A. (2011). Anabolic androgenic steroid use in teens: Prevalence, demographics, and perception of effects. *Journal of Child and Adolescent Substance Abuse, 20*(4), 358–369.

Lundholm, L., Haggård, U., Möller, J., Hallqvist, J., & Thiblin, I. (2013). The triggering effect of alcohol and illicit drugs on violent crime in a remand prison population: A case-crossover study. *Drug and Alcohol Dependence, 129*(1-2), 110–115.

Lundholm, L., Käll, K., Wallin, S., & Thiblin, I. (2010). Use of anabolic androgenic steroids in substance abusers arrested for crime. *Drug and Alcohol Dependence, 111*(3), 222–226.

Mattila, V., Parkkari, J., Laakso, L., Pihlajamäki, H., & Rimpelä, A. (2009). Use of dietary supplements and anabolic-androgenic steroids among Finnish adolescents in 1991–2005. *European Journal of Public Health, 20*(3), 306–311.

Mattila, V., Rimpelä, A., Jormanainen, V., Sahi, T., & Pihlajamäki, H. (2010). Anabolic-androgenic steroid use among young Finnish males. *Scandinavian Journal of Medicine and Science in Sport, 20*(2), 330–335.

Moher, D., Liberati, A., Tetzlaff, J., & Altman, D.G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA Statement. *PLoS Medicine, 6*(6), e1000097.

Nilsson, S. (1995). Androgenic anabolic steroid use among male adolescents in Falkenberg. *European Journal of Clinical Pharmacology, 48*, 9–11.

Nilsson, S., Baigi, A., Marklund, B., & Fridlund, B. (2001a). Trends in the misuse of androgenic anabolic steroids among boys

Unauthenticated
Download Date | 12/16/16 1:04 PM

16-17 years old in a primary health care area in Sweden. *Scandinavian Journal of Primary Health Care, 19*(3), 181–182.

Nilsson, S., Baigi, A., Marklund, B., & Fridlund, B. (2001b). The prevalence of the use of androgenic anabolic steroids by adolescents in a county of Sweden. *European Journal of Public Health, 11*(2), 195–197.

Nilsson, S., Spak, F., Marklund, B., Baigi, A., & Allebeck, P. (2005). Attitudes and behaviors with regards to androgenic anabolic steroids among male adolescents in a county of Sweden. *Substance Use and Misuse, 40*(1), 1–12.

Nøkleby, H. (2013). Use of doping agents and symptoms of eating disorders among male and female patients in drug addiction treatment. *Nordic Studies on Alcohol and Drugs, 30*(4), 331–346.

Pagonis, T.A., Angelopoulos, N.V., Koukoulis, G.N., & Hadjichristodoulou, C.S. (2006). Psychiatric side effects induced by supraphysiological doses of combinations of anabolic steroids correlate to the severity of abuse. *European Psychiatry, 21*, 551–562.

Pallesen, S., Jøsendal, O., Johnsen, B.-H., Larsen, S., & Molde, H. (2006). Anabolic steroid use in high school students. *Substance Use and Misuse, 41*(13), 1705–1717.

Parkinson, A.B., & Evans, N.A. (2006). Anabolic androgenic steroids: a survey of 500 users. *Medicine and Science in Sports and Exercise, 38*(4), 644–651.

Pope Jr., H.G., & Kanayama, G. (2012). Anabolic–androgenic steroids. In J. Verster, K. Brady, M. Galanter, & P. Conrod (Eds.), *Drug abuse and addiction in medical illness: Causes, consequences and treatment* (pp. 251–264). New York: Springer.

Pope Jr., H.G., Kanayama, G., & Hudson, J.I. (2012). Risk factors for illicit anabolic-androgenic steroid use in male weightlifters: A cross-sectional cohort study. *Biological Psychiatry, 71*(3), 254–261.

Pope Jr., H.G., Kanayama, G., Athey, A., Ryan, E., Hudson, J.I., & Baggish, A. (2013). The lifetime prevalence of anabolic-androgenic steroid use and dependence in Americans: Current best estimates. *American Journal of Addictions*, doi: 10.1111/j.1521-0391.2013.12118.x [Epub ahead of print]

Sagoe, D., Molde, H., Andreassen, C.S., Torsheim, T., & Pallesen, S. (2014a). The global epidemiology of anabolic-androgenic steroid use: A meta-analysis and meta-regression analysis. *Annals of Epidemiology, 24*(5), 383–398.

Sagoe, D., Andreassen, C.S., Molde, H., Torsheim, T., & Pallesen, S. (2014b). Prevalence and correlates of anabolic-androgenic steroid use in a nationally representative sample of 17 year-old Norwegian adolescents. *Substance Use and Misuse*, doi: 10.3109/10826084.2014.958859 [Epub ahead of print].

Singhammer, J. (2013). Attitudes toward anabolic-androgenic steroids among non-competing athletes in various types of sports–a cross-sectional study. *Sport Science Review, 22*(1), 109–128.

Solberg, S. (1974). Unpublished data. In: Solberg, S., 1982. Anabolic steroids and Norwegian weightlifters. *British Journal of Sports Medicine, 16*(3), 169–171.

Striegel, H., Simon, P., Frisch, S., Roecker, K., Dietz, K., Dickhuth, H.-H., et al. (2006). Anabolic ergogenic substance users in fitness-sports: A distinct group supported by the health care system. *Drug and Alcohol Dependence, 81*(1), 11–19.

Stroup, D.F., Berlin, J.A., Morton, S.C., Olkin, I., Williamson, G.D., Rennie, D., et al. (2000). *Meta-analysis of observational studies in epidemiology: A proposal for reporting. Meta-analysis of observational studies in epidemiology (MOOSE) group. JAMA 283*, 2008–2012.

Swedish National Institute of Public Health, & Lund University. (2009). *Sub study in the Swedish National Institute of Public Health's prevalence project with the aim of surveying narcotics use in Sweden.* Unpublished study. Stockholm: Swedish National Institute of Public Health.

Swedish National Institute of Public Health. (1991). *Lagen om förbud mot vissa dopningsmedel*. [Act prohibiting certain doping substances.] Stockholm:

Unauthenticated
Download Date | 12/16/16 1:04 PM

Government of Sweden.
Tanner, S.M., Miller, D.W., & Alongi, C. (1995). Anabolic steroid use by adolescents: Prevalence, motives, and knowledge of risks. *Clinical Journal of Sport Medicine, 5*(2), 108–115.
Thorlindsson, T., & Halldorsson, V. (2010). Sport, and use of anabolic androgenic steroids among Icelandic high school students: a critical test of three perspectives. *Substance Abuse Treatment, Prevention, and Policy, 5*(1), 1–11.
Urhausen, A., Albers, T., & Kindermann, W. (2004). Are the cardiac effects of anabolic steroid abuse in strength athletes reversible? *Heart, 90*(5), 496–501.
Viera, A.J., & Garrett, J.M. (2005). Understanding interobserver agreement: the kappa statistic. *Family Medicine, 37*(5), 360–363.
Wichstrøm, L. (2006). Predictors of future anabolic androgenic steroid use. *Medicine and Science in Sports and Exercise, 38(*9), 1578–1583.
Wichstrøm, L., & Pedersen, W. (2001). Use of anabolic-androgenic steroids in adolescence: Winning, looking good or being bad? *Journal of Studies on Alcohol, 62*(1), 5–13.
Yesalis, C.E., & Bahrke, M.S. (2008). Anabolic-androgenic steroids. *Sports Medicine, 19*(5), 326–340.



Unauthenticated
Download Date | 12/16/16 1:04 PM