

RESEARCH

Open Access



Operator iteration on the Young inequality

Xianhe Zhao*, Le Li and Hongliang Zuo

*Correspondence:
zhaoxianhe989@163.com
College of Mathematics and
Information Science, Henan Normal
University, Xinxiang, Henan 453007,
China

Abstract

In this paper, we employ iteration on operator version of the famous Young inequality and obtain more arithmetic-geometric mean inequalities and the reverse versions for positive operators. Concretely, we obtain refined Young inequalities with the Kantorovich constant, the reverse ratio type and difference type inequalities for arithmetic-geometric operator mean under different conditions.

MSC: 47A30; 47A63

Keywords: operator iteration; Young inequality; arithmetic-geometric means; Kantorovich constant

1 Introduction

Throughout this paper, A, B are both positive operators on a Hilbert space H , and $\mathcal{B}_h(H)$ is the semi-space of all bounded linear self-adjoint operators on H . In addition, notation $\mathcal{B}^+(H)$ is written as the set of all positive operators in $\mathcal{B}_h(H)$. Besides, we may assume that A and B are invertible without loss of generality,

$$A\nabla_\mu B = (1 - \mu)A + \mu B \quad \text{and} \quad A\sharp_\mu B = A^{1/2}(A^{-1/2}BA^{-1/2})^\mu A^{1/2}, \quad \text{where } 0 \leq \mu \leq 1.$$

When $\mu = 1/2$ we write $A\nabla B$ and $A\sharp B$ for brevity; see Kubo and Ando [1]. The Kantorovich constant is defined by $K(t, 2) = \frac{(t+1)^2}{4t}$ for $t > 0$, while the Specht ratio [2] is denoted by

$$S(t) = \frac{t^{\frac{1}{t-1}}}{e \log t^{\frac{1}{t-1}}} \quad \text{for } t > 0, t \neq 1, \quad \text{and} \quad S(1) = \lim_{t \rightarrow 1} S(t) = 1,$$

and has the following properties:

- (i) $S(t) = S(\frac{1}{t}) \geq 1$ for $t > 0$.
- (ii) $S(t)$ is a monotone increasing function on $(1, +\infty)$.
- (iii) $S(t)$ is a monotone decreasing function on $(0, 1)$.

We start from an improvement of the famous Young inequality as follows.

Theorem ZS ([3]) For $a, b > 0$, we have

$$a\nabla_\mu b \geq K(h, 2)^r a^{1-\mu} b^\mu$$

for all $\mu \in [0, 1]$, where $r = \min\{\mu, 1 - \mu\}$ and $h = \frac{b}{a}$. It admits an operator extension

$$A \nabla_{\mu} B \geq K(h, 2)^r A \sharp_{\mu} B$$

for positive operators A and B on a Hilbert space.

Next, we show the reverse arithmetic-geometric mean inequality with the Specht ratio for two positive operators.

Theorem T ([4]) For invertible operators A and B with $0 < aI_H \leq A, B \leq bI_H$, we have

- (i) $A \nabla_{\mu} B \leq S(h) A \sharp_{\mu} B$,
- (ii) $A \nabla_{\mu} B - A \sharp_{\mu} B \leq L(1, h) \log S(h) A$,

where $L(1, h)$ is defined by $L(a, b) = \frac{a-b}{\log a - \log b}$ ($a \neq b$); $L(a, a) = a$, $h = \frac{b}{a}$.

These inequalities have recently been improved by Furuichi [5] as follows.

Theorem F ([5]) If $0 < aI_H \leq A, B \leq bI_H$, then

- (i) $A \nabla_{\mu} B - 2r(A \nabla B - A \sharp B) \leq S(\sqrt{h}) A \sharp_{\mu} B$,
- (ii) $A \nabla_{\mu} B - A \sharp_{\mu} B - 2r(A \nabla B - A \sharp B) \leq L(\sqrt{h}, 1) \log S(\sqrt{h}) bI_H$,

where $r = \min\{\mu, 1 - \mu\}$, $L(a, b) = \frac{a-b}{\log a - \log b}$, $h = \frac{b}{a}$.

Afterwards, Krnić *et al.* [6] introduced Jensen’s operator and established some bounds for the spectra of Jensen’s operator. The obtained results were then applied to operator means. In such a way, they get refinements and converses of numerous mean inequalities for Hilbert space operators. See [1, 2, 4, 6–10] for more related developments.

See also [11] for another improvement of the reverse weighted arithmetic-geometric operator mean inequalities. Their proof is independent of [5].

Theorem ZF ([11]) If $0 < aA \leq B \leq bA$ with $0 < a < 1 < b$, then

- (i) $A \nabla_{\mu} B - 2r(A \nabla B - A \sharp B) \leq \max\{S(\sqrt{a}), S(\sqrt{b})\} A \sharp_{\mu} B$,
- (ii) $A \nabla_{\mu} B - A \sharp_{\mu} B - 2r(A \nabla B - A \sharp B)$
 $\leq \max\left\{L\left(\frac{1}{\sqrt{a}}, 1\right) \log S(\sqrt{a}), L\left(\frac{1}{\sqrt{b}}, 1\right) \log S(\sqrt{b})\right\} bA$,

where $r = \min\{\mu, 1 - \mu\}$, $L(a, b) = \frac{a-b}{\log a - \log b}$.

This paper aims to provide a method to obtain more arithmetic-geometric mean inequalities and the reverse version for positive operators. In Section 2, we introduce the main lemmas. In Section 3, utilizing the refined Young inequality and iteration method, we establish some weighted arithmetic-geometric mean inequality for two positive operators. We also obtain reverse ratio type and difference type inequalities for positive operators by means of iteration under different conditions in Section 4 and Section 5, respectively.

2 Main lemmas

First of all, the main lemmas including operator iteration can be stated as follows.

Lemma 2.1 ([6]) *If $A, B \in \mathcal{B}^+(H)$, $0 \leq \mu, \nu \leq 1$, then*

$$2 \min\{\mu, 1 - \mu\}(A \nabla B - A \sharp B) \leq A \nabla_{\mu} B - A \sharp_{\mu} B \leq 2 \max\{\mu, 1 - \mu\}(A \nabla B - A \sharp B). \tag{2.1}$$

Lemma 2.2 *If A and B are positive operators on a Hilbert space, $0 \leq \mu, \nu \leq 1$, then*

$$A \nabla_{\mu}(A \sharp_{\nu} B) = A \nabla_{\mu\nu} B - \mu(A \nabla_{\nu} B - A \sharp_{\nu} B).$$

Proof

$$\begin{aligned} A \nabla_{\mu}(A \sharp_{\nu} B) &= (1 - \mu)A + \mu A \sharp_{\nu} B \\ &= A - \mu A + \mu \nu A - \mu \nu A + \mu \nu B - \mu \nu B + \mu A \sharp_{\nu} B \\ &= \mu \nu B + (1 - \mu \nu)A - \mu [(1 - \nu)A + \nu B - A \sharp_{\nu} B] \\ &= A \nabla_{\mu\nu} B - \mu(A \nabla_{\nu} B - A \sharp_{\nu} B). \end{aligned} \quad \square$$

3 Further refinement of the Young inequalities

In this section, we use the scalar ratio type arithmetic-geometric mean inequality to get a series of operator versions.

Lemma 3.1 ([3]) *If $a, b > 0$ and $\mu \in [0, 1]$, then*

$$a \nabla_{\mu} b \geq K(h, 2)^r a^{1-\mu} b^{\mu},$$

where $r = \min\{\mu, 1 - \mu\}$ and $h = \frac{b}{a}$.

Using the method of the proof of Theorem 7 in [3] we can obtain the following theorem.

Theorem 3.2 *If $A, B \geq 0$, $1 < h \leq A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \leq h'$ or $0 < h' \leq A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \leq h < 1$, then*

$$A \nabla_{\mu} B \geq K(h, 2)^r A \sharp_{\mu} B \tag{3.1}$$

for all $\mu \in [0, 1]$, where $r = \min\{\mu, 1 - \mu\}$.

Proof For the case of $1 < h \leq A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \leq h'$, by Lemma 3.1 we have

$$(1 - \mu) + \mu x \geq K(x, 2)^r x^{\mu}$$

for any $x > 0$. And hence

$$(1 - \mu)I + \mu X \geq \min_{1 < h \leq x \leq h'} K(x, 2)^r X^{\mu}$$

for a positive operator X such that $1 < hI \leq X \leq h'I$.

Substituting $A^{-\frac{1}{2}}BA^{-\frac{1}{2}}$ for X in the above inequality we have

$$(1 - \mu)I + \mu A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \geq \min_{1 < h \leq x \leq h'} K(x, 2)^r (A^{-\frac{1}{2}}BA^{-\frac{1}{2}})^\mu.$$

It is easy to check that the function $K(x, 2)$ is increasing for $x > 1$, then

$$(1 - \mu)I + \mu A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \geq K(h, 2)^r (A^{-\frac{1}{2}}BA^{-\frac{1}{2}})^\mu. \tag{3.2}$$

Multiplying both sides by $A^{\frac{1}{2}}$ to inequality (3.2), we obtain the required inequality.

For the case of $0 < h' \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq h < 1$, since the function $K(x, 2)$ is decreasing for $0 < x < 1$ and $K(\frac{1}{h}, 2) = K(h, 2)$, similarly we obtain inequality (3.1). \square

Theorem 3.3 For two operators $A, B \geq 0$ and $1 < h \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq h'$ or $0 < h' \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq h < 1$, we have

$$A \nabla_\mu B - 2r(A \nabla B - A \sharp B) \geq K(\sqrt{h}, 2)^R A \sharp_\mu B \tag{3.3}$$

for all $\mu \in [0, 1]$, where $r = \min\{\mu, 1 - \mu\}$ and $R = \min\{2r, 1 - 2r\}$.

Proof If $0 \leq \mu \leq \frac{1}{2}$, then $0 \leq 2\mu \leq 1$.

Here $1 < h \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq h'$ ensures that $1 < \sqrt{h} \leq A^{-\frac{1}{2}}(A \sharp B)A^{-\frac{1}{2}} \leq \sqrt{h'}$, and $0 < h' \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq h < 1$ ensures that $0 < \sqrt{h'} \leq A^{-\frac{1}{2}}(A \sharp B)A^{-\frac{1}{2}} \leq \sqrt{h} < 1$, respectively. Substituting B by $A \sharp B$ and μ by 2μ in (3.1), it follows that

$$A \nabla_{2\mu} (A \sharp B) \geq K(\sqrt{h}, 2)^{\min\{2\mu, 1-2\mu\}} A \sharp_{2\mu} (A \sharp B).$$

By Lemma 2.1 and $A \sharp_{2\mu} (A \sharp B) = A \sharp_\mu B$, we have

$$A \nabla_{2\mu} (A \sharp B) = A \nabla_\mu B - 2\mu(A \nabla B - A \sharp B),$$

and then

$$A \nabla_\mu B - 2\mu(A \nabla B - A \sharp B) \geq K(\sqrt{h}, 2)^{\min\{2\mu, 1-2\mu\}} A \sharp_\mu B.$$

If $\frac{1}{2} \leq \mu \leq 1$, then $0 \leq 1 - \mu \leq \frac{1}{2}$. By the above inequality we have

$$B \nabla_{1-\mu} A - 2(1 - \mu)(B \nabla A - B \sharp A) \geq K(\sqrt{h}, 2)^{\min\{2(1-\mu), 1-2(1-\mu)\}} B \sharp_{1-\mu} A.$$

Therefore, for $0 \leq \mu \leq 1$, we have

$$A \nabla_\mu B - 2 \min\{\mu, 1 - \mu\}(A \nabla B - A \sharp B) \geq K(\sqrt{h}, 2)^{\min\{2r, 1-2r\}} A \sharp_\mu B.$$

This completes the proof. \square

Now, applying the same iteration method as in Theorem 3.2 to inequality (3.3), we obtain the following.

Corollary 3.4 *Assume the conditions as in Theorem 3.3. Then*

$$A \nabla_{\mu} B - (2r - R)(A \nabla B - A \sharp B) - 2R(A \nabla_{\frac{2[2\mu]+1}{4}} B - A \sharp_{\frac{2[2\mu]+1}{4}} B) \geq K(\sqrt{h}, 2)^{R'} A \sharp_{\mu} B$$

for all $\mu \in [0, 1]$, where $r = \min\{\mu, 1 - \mu\}$, $R = \min\{2r, 1 - 2r\}$, $R' = \min\{2R, 1 - 2R\}$, $h = \frac{M}{m}$, and $[x]$ is the greatest integer less than or equal to x .

Corollary 3.5 ([12]) *Suppose that two operators A, B and positive real numbers m, m', M, M' satisfy either $0 < m'I \leq A \leq mI < MI \leq B \leq M'I$ or $0 < m'I \leq B \leq mI < MI \leq A \leq M'I$.*

(I) *If $0 \leq \mu \leq \frac{1}{2}$, then*

$$A \nabla_{\mu} B - (2r - R)(A \nabla B - A \sharp B) - 2R(A \nabla_{\frac{1}{4}} B - A \sharp_{\frac{1}{4}} B) \geq K(\sqrt{h}, 2)^{R'} A \sharp_{\mu} B.$$

(II) *If $\frac{1}{2} \leq \mu \leq 1$, then*

$$A \nabla_{\mu} B - (2r - R)(A \nabla B - A \sharp B) - 2R(A \nabla_{\frac{3}{4}} B - A \sharp_{\frac{3}{4}} B) \geq K(\sqrt{h}, 2)^{R'} A \sharp_{\mu} B,$$

where $r = \min\{\mu, 1 - \mu\}$, $R = \min\{2r, 1 - 2r\}$, $R' = \min\{2R, 1 - 2R\}$, $h = \frac{M}{m}$, and $h' = \frac{M'}{m'}$.

Proof In the case of (i), $1 < h = \frac{M}{m} \leq A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \leq \frac{M'}{m'} = h'$; In the case of (ii), $0 < \frac{1}{h'} = \frac{m'}{M'} \leq A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \leq \frac{m}{M} = \frac{1}{h} < 1$. Then Corollary 3.4 leads to the required inequality. This completes the proof. □

4 Reverse ratio type arithmetic-geometric mean inequalities

In the following, we show a refinement of reverse arithmetic-geometric mean inequality by applying the main lemmas in Section 2.

Theorem 4.1 *If $0 < aI \leq A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \leq bI$ with $a < 1 < b$, and $0 \leq \mu \leq 1$, then*

$$A \nabla_{\mu} B - (2r - R)(A \nabla B - A \sharp B) - 2R(A \nabla_{\frac{2[2\mu]+1}{4}} B - A \sharp_{\frac{2[2\mu]+1}{4}} B) \leq \max\{S(\sqrt[4]{a}), S(\sqrt[4]{b})\} A \sharp_{\mu} B,$$

where $r = \min\{\mu, 1 - \mu\}$, $R = \min\{2r, 1 - 2r\}$.

Proof If $0 \leq \mu \leq \frac{1}{2}$, then $0 \leq 2\mu \leq 1$. Since $0 < aI \leq A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \leq bI$ ensures that $\sqrt{a}A \leq A \sharp B \leq \sqrt{b}A$, by substituting B by $A \sharp B$ and μ by 2μ in (i) of Theorem ZF, it follows that

$$A \nabla_{2\mu}(A \sharp B) - 2 \min\{2\mu, 1 - 2\mu\} [A \nabla(A \sharp B) - A \sharp(A \sharp B)] \leq \max\{S(\sqrt[4]{a}), S(\sqrt[4]{b})\} A \sharp_{2\mu}(A \sharp B).$$

Lemma 2.2 leads to the following equalities:

$$A \nabla_{2\mu}(A \sharp B) = A \nabla_{\mu} B - 2\mu(A \nabla B - A \sharp B),$$

$$A \nabla(A \sharp B) - A \sharp(A \sharp B) = A \nabla_{\frac{1}{4}} B - A \sharp_{\frac{1}{4}} B - \frac{1}{2}(A \nabla B - A \sharp B).$$

Then it follows by Lemma 2.1 that

$$\begin{aligned}
 & A\nabla_{\mu}B - 2\mu(A\nabla B - A\sharp B) - 2\min\{2\mu, 1 - 2\mu\} \left[A\nabla_{\frac{1}{4}}B - A\sharp_{\frac{1}{4}}B - \frac{1}{2}(A\nabla B - A\sharp B) \right] \\
 & \leq \max\{S(\sqrt[4]{a}), S(\sqrt[4]{b})\}A\sharp_{\mu}B.
 \end{aligned} \tag{4.1}$$

If $\frac{1}{2} \leq \mu \leq 1$, then $0 \leq 1 - \mu \leq \frac{1}{2}$. The hypothesis $0 < aI \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq bI$ admits $\frac{1}{\sqrt{b}}B \leq B\sharp A \leq \frac{1}{\sqrt{a}}B$. Then by the inequality (4.1) we have

$$\begin{aligned}
 & B\nabla_{1-\mu}A - 2(1 - \mu)(B\nabla A - B\sharp A) - 2R \left[B\nabla_{\frac{1}{4}}A - B\sharp_{\frac{1}{4}}A - \frac{1}{2}(B\nabla A - B\sharp A) \right] \\
 & \leq \max\left\{S\left(\frac{1}{\sqrt[4]{b}}\right), S\left(\frac{1}{\sqrt[4]{a}}\right)\right\}B\sharp_{1-\mu}A.
 \end{aligned}$$

Notice that $S(\frac{1}{\sqrt[4]{a}}) = S(\sqrt[4]{a})$ and $S(\frac{1}{\sqrt[4]{b}}) = S(\sqrt[4]{b})$, so it follows that

$$\begin{aligned}
 & A\nabla_{\mu}B - 2(1 - \mu)(A\nabla B - A\sharp B) - 2R \left[A\nabla_{\frac{3}{4}}B - A\sharp_{\frac{3}{4}}B - \frac{1}{2}(A\nabla B - A\sharp B) \right] \\
 & \leq \max\{S(\sqrt[4]{a}), S(\sqrt[4]{b})\}A\sharp_{\mu}B.
 \end{aligned}$$

This completes the proof. □

Since $0 < aI_H \leq A, B \leq bI_H$ with $a < b$ admits that $\frac{1}{\sqrt{h}} \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq \sqrt{h}$ where $\frac{1}{\sqrt{h}} < 1 < \sqrt{h}$, we obtain the counterpart of Theorem 4.1.

Corollary 4.2 *If $0 < aI \leq A, B \leq bI$ with $a < b$ and $0 \leq \mu \leq 1$, then*

$$A\nabla_{\mu}B - (2r - R)(A\nabla B - A\sharp B) - 2R(A\nabla_{\frac{2[2\mu]+1}{4}}B - A\sharp_{\frac{2[2\mu]+1}{4}}B) \leq S(\sqrt[4]{h})A\sharp_{\mu}B,$$

where $r = \min\{\mu, 1 - \mu\}$, $R = \min\{2r, 1 - 2r\}$, and $h = \frac{b}{a}$.

5 Reverse difference type arithmetic-geometric mean inequalities

In the following theorem we show the corresponding difference type analogs of Theorem 4.1.

Theorem 5.1 *If $0 \leq \mu \leq 1$ and $0 < aI \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq bI$ with $a < 1 < b$, then*

$$\begin{aligned}
 & A\nabla_{\mu}B - A\sharp_{\mu}B - (2r - R)(A\nabla B - A\sharp B) - 2R(A\nabla_{\frac{2[2\mu]+1}{4}}B - A\sharp_{\frac{2[2\mu]+1}{4}}B) \\
 & \leq \max\{L(\sqrt[4]{a}, 1) \log S(\sqrt[4]{a}), L(\sqrt[4]{b}, 1) \log S(\sqrt[4]{b})\} \frac{b}{\sqrt{a}}A,
 \end{aligned}$$

where $r = \min\{\mu, 1 - \mu\}$, $R = \min\{2r, 1 - 2r\}$.

Proof (I) If $0 \leq \mu \leq \frac{1}{2}$, then $0 \leq 2\mu \leq 1$. Since $0 < aI \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq bI$ ensures that $\sqrt{a}A \leq A\sharp B \leq \sqrt{b}A$, substitute B by $A\sharp B$ and μ by 2μ in (ii) of Theorem ZF, so we obtain

$$\begin{aligned}
 & A\nabla_{2\mu}(A\sharp B) - A\sharp_{2\mu}(A\sharp B) - 2\min\{2\mu, 1 - 2\mu\}[A\nabla(A\sharp B) - A\sharp(A\sharp B)] \\
 & \leq \max\{L(\sqrt[4]{1/a}, 1) \log S(\sqrt[4]{a}), L(\sqrt[4]{1/b}, 1) \log S(\sqrt[4]{b})\} \sqrt{b}A.
 \end{aligned}$$

As showed in the proof of Theorem 4.1, the following inequality holds:

$$\begin{aligned}
 & A\nabla_{\mu}B - A\sharp_{\mu}B - (2r - R)(A\nabla B - A\sharp B) - 2R(A\nabla_{\frac{1}{4}}B - A\sharp_{\frac{1}{4}}B) \\
 & \leq \max\{L(\sqrt[4]{1/a}, 1) \log S(\sqrt[4]{a}), L(\sqrt[4]{1/b}, 1) \log S(\sqrt[4]{b})\} \sqrt{b}A \\
 & \leq \max\{L(\sqrt[4]{a}, 1) \log S(\sqrt[4]{a}), L(\sqrt[4]{b}, 1) \log S(\sqrt[4]{b})\} b/\sqrt{a}A, \tag{5.1}
 \end{aligned}$$

since $a < 1 < b$ and $L(\frac{1}{\sqrt[4]{a}}, 1) = \frac{1}{\sqrt[4]{a}}L(\sqrt[4]{a}, 1)$.

(II) If $\frac{1}{2} \leq \mu \leq 1$, then $0 \leq 1 - \mu \leq \frac{1}{2}$. The hypothesis $0 < aI \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq bI$ ensures $\frac{1}{\sqrt{b}}B \leq B\sharp A \leq \frac{1}{\sqrt{a}}B$. Then by the inequality (5.1) and $S(\frac{1}{\sqrt[4]{a}}) = S(\sqrt[4]{a})$, we have

$$\begin{aligned}
 & A\nabla_{\mu}B - A\sharp_{\mu}B - (2r - R)(A\nabla B - A\sharp B) - 2R(A\nabla_{\frac{3}{4}}B - A\sharp_{\frac{3}{4}}B) \\
 & = B\nabla_{1-\mu}A - B\sharp_{1-\mu}A - (2r - R)(B\nabla A - B\sharp A) - 2R(B\nabla_{\frac{1}{4}}A - B\sharp_{\frac{1}{4}}A) \\
 & \leq \max\{L(\sqrt[4]{b}, 1) \log S(\sqrt[4]{1/b}), L(\sqrt[4]{a}, 1) \log S(\sqrt[4]{1/a})\} \sqrt{1/a}B \\
 & \leq \max\{L(\sqrt[4]{b}, 1) \log S(\sqrt[4]{b}), L(\sqrt[4]{a}, 1) \log S(\sqrt[4]{a})\} b/\sqrt{a}A.
 \end{aligned}$$

The proof is done. □

Similarly, we replace the hypothesis $0 < aI \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq bI$, where $a < 1 < b$ by $0 < aI \leq A, B \leq bI$ with $a < b$, and we obtain the following.

Corollary 5.2 *If $0 < aI \leq A, B \leq bI$ with $a < b$, and $0 \leq \mu \leq 1$, then*

$$\begin{aligned}
 & A\nabla_{\mu}B - A\sharp_{\mu}B - (2r - R)(A\nabla B - A\sharp B) - 2R(A\nabla_{\frac{2[2\mu]+1}{4}}B - A\sharp_{\frac{2[2\mu]+1}{4}}B) \\
 & \leq b\sqrt{h}L(\sqrt[4]{h}, 1) \log S(\sqrt[4]{h})I,
 \end{aligned}$$

where $r = \min\{\mu, 1 - \mu\}$, $R = \min\{2r, 1 - 2r\}$, and $h = \frac{b}{a}$.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

Acknowledgements

We wish to thank the referee for careful reading and valuable comments for the original draft. This research was supported by the National Natural Science Foundation of China (11501176) and the Science and Technology Planning Project of the Education Department of Henan Province (14B110009).

Received: 7 September 2016 Accepted: 15 November 2016 Published online: 24 November 2016

References

1. Kubo, F, Ando, T: Means of positive operators. *Math. Ann.* **246**(3), 205-224 (1980)
2. Tominaga, M: Specht's ratio in the Young inequality. *Sci. Math. Jpn.* **55**(3), 583-588 (2002)
3. Zuo, HL, Shi, GH, Fujii, M: Refined Young inequalities with Kantorovich constant. *J. Math. Inequal.* **5**(4), 551-556 (2011)
4. Pečarić, J, Beesack, PR: On Jensen's inequality for convex functions II. *J. Math. Anal. Appl.* **118**, 125-144 (1986)
5. Furuichi, S: On refined Young inequalities and reverse inequalities. *J. Math. Inequal.* **5**(1), 21-31 (2011)

6. Krnić, M, Lovričević, N, Pečarić, J: Jensen's operator and applications to mean inequalities for operators in Hilbert space. *Bull. Malays. Math. Soc.* **35**(1), 1-14 (2012)
7. Hirzallah, O, Kittaneh, F, Krnić, M, Lovričević, N, Pečarić, J: Refinements and reverses of means inequalities for Hilbert space operators. *Banach J. Math. Anal.* **7**(2), 15-19 (2013)
8. Klarić, M, Pečarić, J, Perić, J: On the converse Jensen inequality. *Appl. Math. Comput.* **218**, 6566-6575 (2012)
9. Krnić, M, Lovričević, N, Pečarić, J: Improved arithmetic-geometric and Heinz means inequalities for Hilbert space operators. *Publ. Math. (Debr.)* **80**, 3-4 (2012)
10. Specht, W: Zer Theorie der elementaren Mittel. *Math. Z.* **74**, 91-98 (1960)
11. Zuo, HL, Fujii, M, Fujii, JI, Seo, Y: Upper bound for spectra of Jensen operator and its application to reverse arithmetic-geometric means. *Math. Inequal. Appl.* **17**(2), 641-648 (2014)
12. Liao, WS, Wu, JL: Improved Young and Heinz inequalities with the Kantorovich constant. *J. Math. Inequal.* **10**(2), 559-570 (2016)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
