

## Opinions

# Competitive Exclusion between Piroplasmosis and Anaplasmosis Agents within Cattle

Loubna Dib, Idir Bitam, Maja Tahri, Mourad Bensouilah, Thierry De Meeüs\*

There are about 869 species of ticks described so far [1], most of which are responsible for the transmission of a huge diversity of microorganisms belonging to almost all the trees of life (viruses, bacteria, protozoa, fungi, and nematodes) [2–4]. Despite the strong opportunity for interaction between these pathogenic species, little is known about the competitive interactions between tick-borne diseases within the vector and the vertebrate host (see [3] for review).

Severity of symptoms is often found associated with co-infection by different pathogens (e.g., [3,5–7]). Moreover, dual infections may affect the therapeutic strategies [5]. Exploring the incidence of co-infections and testing for possible interactions (positive or negative) thus represent important objectives.

*Theileria annulata*, *Babesia bovis*, and *Anaplasma marginale* are among the most economically important haemoparasitic tick-borne diseases of ruminants worldwide [8] and represent a serious economic challenge, particularly in developing countries [9–11]. In this paper we analyse the co-occurrence of these pathogens within individual bovine hosts in northeastern Algeria (North Africa) from one original dataset and one published dataset from the same region [12] that we have reanalysed, and test for the existence of positive or negative associations between these pathogens. We then discuss the implications of our findings in terms of therapeutic strategies and further studies.

All samples were collected around Boutheldja, a small town located on the northeast border of Algeria (close to Tunisia) and east of Annaba in Algeria (GPS coordinates 36° 45' 7.0 N; 8° 10' 0 E) corresponding to an area of 113.53 km<sup>2</sup> (Figure S1). The sampling period extended from July to December 2004. Ill cattle were diagnosed by private veterinarians, who were contacted directly by their owners. The main criteria used for diagnosis were the presence of the ticks, a hyperthermic condition, presence of icterus, ganglionic hypertrophy, feebleness, lack of appetite, anorexia, anaemia, and dehydration. In case of positive diagnostic a blood smear was carried out, as described in Table 1. Parasites were then identified according to morphology, localisation in the erythrocytes, and using the key proposed by Morel in 1981 [13]. We also reanalysed the data from a published descriptive work [12] undertaken in 2002 in the same part of Algeria on 54 cattle with an identical approach as described above.

Four species were identified in the samples: *T. annulata*, *B. bovis*, *A. marginale*, and *Theileria orientalis*. Note that *Anaplasma* are rickettsial bacteriae while the others are apicomplexan sporozoan (piroplasm), and that transovarial transmission is only known for *Babesia* sp. and not for the three other species.

In the area investigated, *Theileria* sp. are transmitted by different *Hyalomma* ticks, *B. bovis* is transmitted by the tick *Boophilus annulatus*, and *A. marginale* by *B. annulatus* and *Rhipicephalus bursa* [4]. All are intracellular parasites [14].

In 2004, the most frequently encountered pathogenic agent was *T. annulata* (47.6% of infected cattle), followed by *A. marginale* (40.5%), *B. bovis* (33.3%), and *T. orientalis* (2.4%). In 2002, as presented by [12], some differences were observed with 74%, 24%, 8%, and 16% for *T. annulata*, *A. marginale*, *B. bovis*, and *T. orientalis*, respectively. The comparison of microbe prevalences between 2002 and 2004 samples was undertaken with an approximation of the Fisher exact test with 1,000,000 randomisations using the module Struc of Genepop version 3.1.c [15]. Randomisation tests gave significant differences ( $p$ -values  $\leq 0.036$ ) for all species but *A. marginale* ( $p$ -value = 0.117). In 2004 (original sample), 12 hosts appeared negative, 15 were infected by *T. annulata* only, ten by *B. bovis* only, seven by *A. marginale*, and four were infected both by *B. bovis* and *A. marginale*, five by *T. annulata* and *A. marginale*, and one by *T. orientalis* and *A. marginale*. In 2002 [12], four hosts appeared negative, 29 were infected by *T. annulata* only, three by *B. bovis* only, four by *A. marginale*, three by *T. orientalis*, and one was infected both by *B. bovis* and *A. marginale*, five by *T. annulata* and *A. marginale*, two by *T. orientalis* and *A. marginale*, and three by *T. annulata* and *T. orientalis*. Presence or absence of a particular micropathogen was treated as a phenotypic character. Thus there were four such phenotypic characters each taking the value 1 (absence) or 2 (presence). To study the association between these four characters a correlation coefficient, initially designed for genetic data, was adapted to our data. This coefficient of correlation is noted  $R_{ij}$ , where  $i$  and  $j$  stands for the pair of

**Editor:** Marianne Manchester, The Scripps Research Institute, United States of America

**Citation:** Dib L, Bitam I, Tahri M, Bensouilah M, De Meeüs T (2008) Competitive exclusion between piroplasmosis and anaplasmosis agents within cattle. PLoS Pathog 4(1): e7. doi:10.1371/journal.ppat.0040007

**Copyright:** © 2008 Dib et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Loubna Dib is with the Institut Vétérinaire, Centre Universitaire El Tarf, Route de Matroha, El Tarf, Algeria. Idir Bitam is with the Laboratoire d'Entomologie Médicale, Annexe Sidi Fredj, Institut Pasteur d'Alger, Algeria. Maja Tahri and Mourad Bensouilah are with the Laboratoire d'Écobiologie des Milieux Marin et Littoraux, Université Badji Mokhtar, Annaba, Algeria. Thierry De Meeüs is with the Génétique et Evolution des Maladies Infectieuses (GEMI), UMR 2724 IRD/CNRS, Centre IRD, Montpellier, France.

\* To whom correspondence should be addressed. E-mail: demeeus@mpl.ird.fr

**Table 1.** Within-Cattle Occurrence Correlation between Pathogens

		<i>Babesia bovis</i>	<i>Anaplasma marginale</i>	<i>Theileria orientalis</i>
<b>Present study (2004)</b>	<i>Theileria annulata</i>	−0.691 (0.0001)	−0.151 (0.064)	−0.023 (1)
	<i>Babesia bovis</i>	—	−0.081 (0.33)	−0.113 (1)
	<i>Anaplasma marginale</i>	—	—	0.194 (0.404)
<b>2002 sample</b>	<i>Theileria annulata</i>	−0.508 (0.003)	−0.423 (0.007)	−0.371 (0.02)
	<i>Babesia bovis</i>	—	0.01 (1)	−0.131 (0.601)
	<i>Anaplasma marginale</i>	—	—	0.007 (1)
<b>Combination</b>	<i>Theileria annulata</i>	−0.599 (0.0001)	−0.287 (0.004)	−0.197 (0.099)
	<i>Babesia bovis</i>	—	−0.035 (0.696)	−0.122 (0.907)
	<i>Anaplasma marginale</i>	—	—	0.101 (0.771)

Correlation between the different pairs of species was measured by  $R_{ij}$  and their significance is presented within parentheses. Values are given for the 2004 sample (present study), the 2002 sample published in [12], and for the combination where  $R_{ij}$  is averaged over both samples and  $p$ -values obtained by the Fisher procedure. Negative values are indicative of avoidance between pathogens (see text for more detailed explanations). Sampling methods: Blood sampling was undertaken from a peripheral ear vein after alcohol disinfection of the skin. The blood was collected on a de-greased strip of glass on which it was spread into a thin layer with another strip. The strip was labelled and its smear quickly dried and stored into an appropriate box that was sent to the laboratory. In the laboratory, the preparation was fixed with methanol and then coloured with a Giemsa solution (1/10 diluted with distilled water) during 25–30 min. Smears were then washed with tap water and left to dry at room temperature, after which they were examined under a microscope ( $\times 100$  magnification). doi:10.1371/journal.ppat.0040007.t001

loci (or as here, phenotypic characters), the association of which is under study. It was computed by Genetix 4.02 [16]. This parameter is described in a series of articles cited in the Genetix help menu (i.e., [17–19]). If  $n_{11}$  is the number of cattle without pathogens  $i$  or  $j$ ,  $n_{12}$  the number of cattle infected by  $j$  but not by  $i$ ,  $n_{21}$  those infected by  $i$  but not  $j$ ,  $n_{22}$  those infected by both species, and  $n$  the total sample size, then for a given pathogen pair  $ij$ :

$$R_{ij} = \frac{\frac{n}{n-1} \left[ \frac{n_{11}}{n} - \frac{(n_{11}+n_{12})(n_{11}+n_{21})}{n^2} \right]}{\sqrt{\frac{(n_{11}+n_{12})(n_{22}+n_{21})(n_{11}+n_{21})(n_{22}+n_{12})}{n^4}}}$$

The significance of  $R_{ij}$  was tested by a permutation procedure. Phenotypes (1 or 2) of the two characters studied (e.g., *T. annulata* and *B. bovis*) are associated at random a number of times (10,000) and a statistic  $R$  is recalculated on the randomised data set. The exact  $p$ -value is estimated as the proportion of statistics from randomised data sets that are larger or equal to the one observed in the real data set. This test was undertaken under Genetix 4.02, which uses Weir's  $R$  [20] as the statistic. However, given how we coded data, this statistic simply corresponds to  $R_{ij}$  absolute value (bi-lateral testing). The 12 ill cows that were negative according to the cytological tests in 2004 were not taken into account for this study, considering that microbes might have been missed (false negative). The four negative cattle from [12] were also ignored. However, undertaking the same tests and including these data as true negatives (no microbe) did not change the results described below. To unite both data sets (2004 and 2002), we simply averaged  $R_{ij}$  (unweighted mean) and combined  $p$ -values with the Fisher procedure [21] where the quantity  $-2 \sum_{i=1}^n \log_e(p_i)$  can be compared to a chi-square distribution with  $2n$  degrees of freedom, with  $n$  the number of tests (here  $n = 2$ ) and  $p_i$  the  $p$ -value of the  $i$ th test. The results of the correlation study between pairs of pathogenic agents provided mainly negative correlations (five out of six for combined data, Table 1). For the original sample (2004), only one of the correlations, between *T. annulata* and *B. bovis*, was highly significant. In fact, no cattle individual was found co-infected by both pathogens despite an expected co-

occurrence of  $0.476 \times 0.333 \times 42 = 7$  co-infected individuals. The second interesting result, though marginally non-significant, was observed between *T. annulata* and *A. marginale*, for which only five individuals were found co-infected for an expected frequency of height. All other pairs were not significant, but we can notice that with only one case reported for *T. orientalis*, the rarity of this species in 2004 makes any conclusion impossible for this pathogen. Nevertheless, the analysis combining 2002 [12] and 2004 samples confirms all the tendencies observed (Table 1). An absolute exclusion exists between *T. annulata* and *B. bovis* (no co-infection observed), a strong avoidance characterises the relationship between *T. annulata* and *A. marginale*, and a moderate one (if any) between this species and *T. orientalis*.

Because  $R_{ij}$  is oriented and because we coded absence as 1 and presence as 2 for each phenotype, a positive value would bear witness to a positive association between the two pathogenic agents within the same individual hosts, while a negative value gives evidence for avoidance between the pairs of microbes studied. In the first case, different interpretations can be formulated as cooperation between pathogens (immunocompromising by one agent, opening the gate to infection by the other) or identical ecological needs (e.g., same vector). Immunosuppressive effects are what seem to be most of the time observed for various pathogenic agents, in particular in tick-borne diseases, as babesioses with other parasites in mice, or anaplasmosis with louping ill virus in sheep and goats (reviewed in [22]). In the case of a negative association, a hypothesis for competitive exclusion or at least a negative interference (cross-immunisation) can be advanced. This is the case for *T. annulata* and *B. bovis*, which appear to strictly avoid co-infection. This is also the case for *T. annulata* and *A. marginale*, though to a lower extent and probably also for *T. annulata* and *T. orientalis*, though the rarity of the latter species (low statistical power) forbids a definitive conclusion. This may seem to contradict the conclusions found in other studies on similar systems [23–25], though prevalence differences, lack of detailed data, and absence of specific analysis may be the main causes of this apparent discrepancy. Significant negative correlations always involve *T. annulata* against the other pathogen species. In the area

studied, *Theileria* sp. and the other pathogens (*B. bovis*, *A. marginale*) are transmitted by different tick species [4]. Thus, competition must occur within the vertebrate host. Within-host competition between microbes (mostly intra-specific) was reviewed in Read and Taylor [26], with the best documented examples apparently found in *Plasmodium* sp. Competition was poorly studied in tick-borne diseases and only suspected once (to our knowledge) between *Anaplasma phagocytophilum*, *Ehrlichia muris*, and *Babesia microti* in unfed adults of *Ixodes persulcatus* in northwestern Russia [27]. This might confirm a generalised weak compatibility between these three kinds of intracellular pathogens within the vector [27] as well as within the vertebrate host (present study). Weak compatibility between these pathogens was never evidenced in vertebrate hosts for which specific studies are scarce or difficult to interpret in that perspective. Immunological surveys are not conclusive enough, as they may better reflect the history of sequential infections, not necessarily all successful, experienced by one host. Many studies report the co-occurrence of tick-borne micro-pathogens (e.g., [3,5,12,23,24] and references therein), and specific analyses seemed to conclude with positive associations or even suggested cooperation in multi-infected mice: between *Borrelia burgdorferi* (Lyme disease agent) and *B. microti* (transmission to tick is enhanced) [6] or between *B. burgdorferi* and *A. phagocytophilum* (*Borrelia* number is increased in co-infected mice) [28] (see also [29] for review and references therein). Synchronous infections with *Babesia divergens* and *A. phagocytophila* seem very common, but their interaction remains poorly understood and a suppression of *Babesia* by *Anaplasma* even seems possible (reviewed in [25]). As suggested above, such exclusion may be mediated by host immune system (concomitant immunity) or by direct interference (one pathogen tends to eliminate the other). Within the vector, most studies only report the co-occurrence of different microbes [10,29–37]. Associations between different pathogenic species were rarely tested, and only positive associations [6,28,38,39] or no effect [40] between borrelia spirochaetes and co-infecting microbes were detected.

Exclusion between pathogens may appear beneficial to the host, as severity of symptoms is often found associated with co-infection by different pathogens (e.g., [3,5–7]). Nevertheless, depending on the mechanisms involved and which, if any, of the pathogens is competitively dominant, this phenomenon may severely affect the outcome of prophylactic or vaccination campaigns. If one pathogenic species is less sensitive to treatment but at a competitive disadvantage against the most sensitive one, this may lead to the opening of the widest gate to the most pathogenic microbe. It is thus essential to better understand the origin and mechanisms of such competitive interactions, not only for babesioses, theilerioses, and anaplasmoses that represent a real threat to livestock on a global scale, particularly in developing countries where they constrain economic improvement [8–10], but also for other vector-borne or other diseases affecting animals or humans. This parameter is thus worth investigating further, because it may open new perspectives in the design of therapeutic strategies of economically and medically important pathogenic agents, particularly in tick-borne and vector-borne diseases, where it has attracted little attention so far.

## Supporting Information

**Figure S1.** A Typical Landscape of El-Tarf Wilaya around Boutheldja, on the Northeast Border of Algeria (Close to Tunisia), East of Annaba in Algeria

Several local breed cattle, as those studied in the present study, can be seen. All cattle belong to the so-called Atlas brown breed. The Cheurfa and Guelmoise sub-breeds harbor various clear coat colors, almost white for some. The Sétifiennne displays a uniform black coat and the Chelefiennne a fawn coat. Photo credit: Loubna Dib.

Found at doi:10.1371/journal.ppat.0040007.sg001 (2.8 MB TIF).

## Acknowledgments

We thank Franck Prugnolle for very useful and stimulating discussions and one anonymous referee for helpful suggestions.

**Author contributions.** MAB conceived and designed the experiments. LD, IB, and MT performed the experiments. IB and TDM analyzed the data. IB contributed reagents/materials/analysis tools. TDM wrote the paper.

**Funding.** Funding of this work was supported by the Centre Universitaire El Tarf, the CNRS, and the IRD. Thierry De Meeûs is supported by the CNRS and IRD.

**Competing interests.** The authors have declared that no competing interests exist.

## References

1. Camicas JL, Hervy JP, Adam F, Morel PC (1998) The ticks of the world (Acarida, Ixodida). Paris: ORSTOM. 233 p.
2. Aeschlimann A, Burgdorfer W, Matile H, Peter O, Wyler R (1979) Aspects nouveaux du rôle de vecteur joué par *Ixodes ricinus* L. en Suisse. Acta Trop 36: 181–191.
3. Belongia EA (2002) Epidemiology and impact of coinfections acquired from *Ixodes* ticks. Vector Borne Zoonot Dis 2: 265–273.
4. Jongejan F, Uilenberg G (2004) The global importance of ticks. Parasitology 129: S3–S14.
5. Nadelman RB, Horowitz HW, Hsieh TC, Wu JM, Agüero-Rosenfeld ME, et al. (1997) Simultaneous human granulocytic ehrlichiosis and Lyme borreliosis. N Engl J Med 337: 27–30.
6. Thompson C, Spielman A, Krause PJ (2001) Coinfecting deer-associated zoonoses: Lyme disease, babesiosis, and ehrlichiosis. Clin Infect Dis 33: 676–685.
7. Le Hesran JY, Akiana J, El Ndiaye HM, Dia M, Senghor P, et al. (2004) Severe malaria attack is associated with high prevalence of *Ascaris lumbricoides* infection among children in rural Senegal. Trans R Soc Trop Med Hyg 98: 397–399.
8. Uilenberg G (1995) International collaborative research: significance of tick-borne hemoparasitic diseases to world animal health. Vet Parasitol 57: 19–41.
9. D'oliveira C, Vanderweide M, Habela MA, Jacquet P, Jongejan F (1995) Detection of *Theileria annulata* in blood-samples of carrier cattle by PCR. J Clin Microbiol 33: 2665–2669.
10. Cringoli G, Otranto D, Testini G, Buono V, Di Giulio G, et al. (2002) Epidemiology of bovine tick-borne diseases in southern Italy. Vet Res 33: 421–428.
11. Barros SL, Madrugá CR, Araújo FR, Menk CF, de Almeida MAO, et al. (2005) Serological survey of *Babesia bovis*, *Babesia bigemina*, and *Anaplasma marginale* antibodies in cattle from the semi-arid region of the state of Bahia, Brazil, by enzyme-linked immunosorbent assays. Mem Inst Oswaldo Cruz 100: 613–617.
12. Ziam H, Benaouf H (2004) Prevalence of blood parasites in cattle from wilayates of Annaba and El Tarf east Algeria. Arch Inst Pasteur Tunis 81: 27–30.
13. Morel PC (1981) Maladies à tiques du bétail en Afrique. In: Troncy PM, Itard J, Morel PC, editors. Précis de parasitologie vétérinaire tropicale. Maisons-Alfort Ministère de la Coopération et du Développement. pp. 473–473.
14. Uilenberg G (1976) Tick-borne livestock diseases and their vectors. 2. Epizootiology of tick-borne diseases. World Animal Review 17: 8–15.
15. Raymond M, Rousset F (1995) Genepop (version 1.2): population genetics software for exact tests and ecumenicism. J Hered 86: 248–249.
16. Belkhir k, Borsa P, Chikhi L, Raufaste N, Bonhomme F (2004) GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations. Montpellier: Laboratoire Génome, Populations, Interactions CNRS UMR 5000, Université de Montpellier II.
17. Cockerham CC, Weir BS (1979) Digenic descent measures for finite populations. Genet Res 30: 121–147.
18. Black IVWC, Krafur ES (1985) A FORTRAN program for the calculation and analysis of two-locus linkage disequilibrium coefficients. Theor Appl Genet 70: 491–496.

19. Garnier-Géré P, Dillmann C (1992) A computer program for testing pairwise linkage disequilibria in subdivided populations. *J Hered* 83: 239.
20. Weir BS (1979) Inferences about linkage disequilibrium. *Biometrics* 35: 235–254.
21. Fisher RA (1970) *Statistical methods for research workers*, 14th edition. Edinburgh: Oliver and Boyd. 362 p.
22. Persing DH (1997) The cold zone: a curious convergence of tick-transmitted diseases. *Clin Infect Dis* 25: S35–S42.
23. Georges K, Loria GR, Riili S, Greco A, Caracappa S, et al. (2001) Detection of haemoparasites in cattle by reverse line blot hybridisation with a note on the distribution of ticks in Sicily. *Vet Parasitol* 99: 273–286.
24. Almeria S, Castella J, Ferrer D, Gutierrez JF, Estrada-Pena A, et al. (2002) Reverse line blot hybridization used to identify hemoprotozoa in Minorcan cattle. *Ann N Y Acad Sci* 969: 78–82.
25. Zintl A, Mulcahy G, Skerrett HE, Taylor SM, Gray JS (2003) *Babesia divergens*, a bovine blood parasite of veterinary and zoonotic importance. *Clin Microbiol Rev* 16: 622–636.
26. Read AF, Taylor LH (2001) The ecology of genetically diverse infections. *Science* 292: 1099–1102.
27. Alekseev AN, Dubinina HV, Jushkova OV (2004) First report on the coexistence and compatibility of seven tick-borne pathogens in unfed adult *Ixodes persulcatus* Schulze (Acarina: Ixodidae). *Int J Med Microbiol* 293: 104–108.
28. Holden K, Hodzic E, Feng S, Freet KJ, Lefebvre RB, et al. (2005) Coinfection with *Anaplasma phagocytophilum* alters *Borrelia burgdorferi* population distribution in C3H/HeN mice. *Infect Immun* 73: 3440–3444.
29. Korenberg EI (2004) Problems in the study and prophylaxis of mixed infections transmitted by ixodid ticks. *Int J Med Microbiol* 293: 80–85.
30. Schwartz I, Fish D, Daniels TJ (1997) Prevalence of the rickettsial agent of human granulocytic ehrlichiosis in ticks from a hyperendemic focus of Lyme disease. *N Engl J Med* 337: 49–50.
31. Leutenegger CM, Pusterla N, Mislin CN, Weber R, Lutz H (1999) Molecular evidence of coinfection of ticks with *Borrelia burgdorferi* sensu lato and the human granulocytic ehrlichiosis agent in Switzerland. *J Clin Microbiol* 37: 3390–3391.
32. Levin ML, des Vignes F, Fish D (1999) Disparity in the natural cycles of *Borrelia burgdorferi* and the agent of human granulocytic ehrlichiosis. *Emerg Infect Dis* 5: 204–208.
33. Schouls LM, Van de Pol I, Rijpkema SGT, Schot CS (1999) Detection and identification of *Ehrlichia*, *Borrelia burgdorferi* sensu lato, and *Bartonella* species in Dutch *Ixodes ricinus* ticks. *J Clin Microbiol* 37: 2215–2222.
34. Adelson ME, Rao RVS, Tilton RC, Cabets K, Eskow E, et al. (2004) Prevalence of *Borrelia burgdorferi*, *Bartonella* spp., *Babesia microti*, and *Anaplasma phagocytophilum* in *Ixodes scapularis* ticks collected in northern New Jersey. *J Clin Microbiol* 42: 2799–2801.
35. Halos L, Jamal T, Maillard R, Beugnet F, Le Menach A, et al. (2005) Evidence of *Bartonella* sp in questing adult and nymphal *Ixodes ricinus* ticks from France and co-infection with *Borrelia burgdorferi* sensu lato and *Babesia* sp. *Vet Res* 36: 79–87.
36. Halos L, Vourc'h G, Cotte V, Gasqui P, Barnouin J, et al. (2006) Prevalence of *Anaplasma phagocytophilum*, *Rickettsia* sp and *Borrelia burgdorferi* sensu lato DNA in questing *Ixodes ricinus* ticks from France. *Ann N Y Acad Sci* 1078: 316–319.
37. Molad T, Mazuz ML, Fleiderovitz L, Fish L, Savitsky I, et al. (2006) Molecular and serological detection of *A. centrale*- and *A. marginale*-infected cattle grazing within an endemic area. *Vet Microbiol* 113: 55–62.
38. Jenkins A, Kristiansen BE, Allum AG, Aakre RK, Strand L, et al. (2001) *Borrelia burgdorferi* sensu lato and *Ehrlichia* spp. in *Ixodes* ticks from southern Norway. *J Clin Microbiol* 39: 3666–3671.
39. De Meets T, Lorimier Y, Renaud F (2004) Lyme borreliosis agents and the genetics and sex of their vector, *Ixodes ricinus*. *Microbes Infect* 6: 299–304.
40. Levin ML, Fish D (2000) Acquisition of coinfection and simultaneous transmission of *Borrelia burgdorferi* and *Ehrlichia phagocytophilum* by *Ixodes scapularis* ticks. *Infect Immun* 68: 2183–2186.

The Creative Commons  
Attribution License



allows anyone to download,  
reuse, reprint, distribute, or  
copy articles in PLoS journals, so long  
as the original author and source are  
credited. View the license at  
[creativecommons.org/licenses/by/2.0](http://creativecommons.org/licenses/by/2.0).