

Original Article

Is a Successful High-K Fitness Strategy Associated with Better Mental Health?

Cezar Giosan, Department of Psychiatry, Weill Medical College of Cornell University, New York, USA.
Email: cezar@giosan.com (corresponding author)

Katarzyna Wyka, Department of Psychiatry, Weill Medical College of Cornell University, New York, USA.

Abstract: This study examined the associations between a high-K fitness strategy and mental health. These associations were tested on a sample of 1400 disaster workers who had exposure to a singular traumatic event and who underwent psychological evaluations. The results showed that high-K was an important negative predictor of psychopathology, accounting for significant variance in PTSD, general psychopathology, functional disability, anger, and sleep disturbances. Implications of the results are discussed.

Keywords: Fitness, adaptedness, high-K, r/K, life history theory, psychopathology, PTSD, anger, sleep disturbance, disability.

Introduction

Clinical psychology is one area where evolutionary psychology is having a major impact (see Nesse, 2005, for a review). The persistent, cross-cultural, relatively high incidence of mental disorder in the general population suggests that it might have evolutionary significance. The attempts to explain mental disorder from an evolutionary perspective centered on the possible functions of distinct mental conditions (e.g., Crow, 1991, 1997, 1995; Fiske and Haslam, 1997; Mealey, 2000) or on the definitions or taxonomy of mental disorder (Wakefield, 1997; Wakefield and First, 2003). To this end, disorder is seen either as evolutionary dysfunction—failure of natural design, or as naturally selected conditions having present or past fitness advantages (Nesse, 2005).

For instance, very much unlike the prevalent view that it is a brain disorder (Andreasen, 1984; Valenstein, 1998; Wolpert, 1999), evolutionary psychology attempts to explain depression through the functions it may serve, with some authors arguing that low mood is associated with less likelihood of engaging in risk-taking behaviors (Leith and Baumeister, 1996) and others arguing that low levels of depression seem to be “normal” or functional states (McGuire, Troisi, and Raleigh, 1997; Nesse, 1998; Nesse and Williams, 1994).

Likewise, phobias are explained as “prepared” fears gone awry. Prepared fears are fears of things that were a threat to survival during the environment of evolutionary adaptedness. Most phobias are exaggerations of such fears (e.g., spiders, heights, storms, snakes, strangers, blood, unfamiliar places) (Marks, 1987; Marks and Nesse, 1994) and few develop phobias of modern threats to survival (e.g., cars, cigarettes, or alcohol).

Mental illness has negative effects on fitness. Severe depression is associated with reduced fitness (McGuire et al., 1997), as is schizophrenia (Avila, Thaker, and Adami, 2001; Haukka, Suvisaari, and Lönqvist, 2003). Women with psychotic disorders have been shown to have lower fertility than matched normal comparison subjects (Howard, Kumar, Lesse, and Thornicroft, 2002). Likewise, people suffering from non-affective psychoses have been shown to have significantly fewer children than their unaffected siblings, with the effect more pronounced for male patients versus well brothers (McGrath et al., 1998).

Mental illness of the parents seems not only to affect their resultant fitness translated in sheer number of children but it also compromises offspring quality. Children of parents with severe mental illness are themselves at increased risk of childhood psychiatric disorders (Cowling, Luk, Mileskin, and Birlson, 2004) and the risk of offspring with impaired intellectual handicap is significantly increased for schizophrenia and affective psychoses mothers compared with controls (Jablensky et al., 1998).

Purpose of the study

The purpose of this study was to examine the associations between a high-K reproductive strategy and mental illness. According to Life History Theory (e.g., Bogaert and Rushton, 1989; Figueredo et al., 2006; Mac Arthur and Wilson, 1967; Pianka, 1970; Wilson, 1975), for any organism, its available resources are finite, which translates into trade-offs in their allocation for solving various fitness-relevant tasks. For instance, the resources allocated to grow a bigger body cannot be used to pursue mates, and vice versa. The extremes of these fundamental dimensions of reproduction are traditionally termed the r/K theory, with r representing the maximum egg output and no investment and K representing the opposite. Rushton (1985, 1990) extrapolated the r/K continuum of life history strategies to human differences from Wilson’s (1975) analysis of cross-species differences in reproductive strategies. In humans, since they appear to be highly K-selected, these dimensions are often referred to as “differential K.”

A high-K strategy in humans is theorized as a specific and cohesive fitness strategy, in which the individual invests in somatic and parental effort to produce a fewer number of “fitter” and more competitive offspring. Measures of high-K strategy have been shown to correlate with offspring quality and with resultant fitness (Giosan, 2006). Traits associated with a high-K strategy are long-term thinking and planning, commitment to long-term relationships, extensive parental investment, existence of social support structures, adherence to social rules (e.g., altruism and cooperation) and careful consideration of risks (Figueredo et al., 2006; Giosan, 2006).

Because a high-K strategy is associated with greater somatic effort as opposed to reproductive effort, “slow” strategists should manifest the benefits of this in terms of better mental and physical health. Likewise, low-K (“fast”) strategists should exhibit increased psychopathology and increased physical illness, because these individuals do not allocate sufficient somatic effort to prevent such problems. Indeed, in one of the first studies on this

topic, high-K strategy was shown to be an important negative predictor of depressive symptomatology, accounting for 20% in the variance in the Beck Depression Inventory scores, after controlling for risk factors in depression such as demographics, prior trauma, prior psychopathology, or recent negative life events (Giosan, 2007).

Expanding on these findings, the principal aim of this study was to explore the associations between a high-K fitness strategy and a larger array of mental conditions, including PTSD and related psychopathology, sleep disturbances, anger, and functional disability. The overarching hypothesis tested in this study was that a successful high-K strategy is a negative predictor of mental illness.

Materials and Methods

Participants were a sample of $n = 1400$ disaster workers who were deployed to the World Trade Center in the immediate aftermath of the 9/11 attack. They were administered a full battery of self-reports and a subsample of 436 (31%) respondents were also administered a structured clinical interview conducted by doctoral level clinicians.

Measures

Clinician-Administered Measures

- 1) *Clinician-Administered PTSD Scale*. The Clinician-Administered PTSD scale (CAPS; Blake et al., 1995) is a structured interview for posttraumatic stress disorder (PTSD) that yields both a dichotomous (present/absent) diagnosis of PTSD and a continuous measure of its severity. The CAPS assesses the frequency and intensity of each PTSD symptom on separate 5-point rating scales (0 - 4). Frequency and intensity may be summed for each item to yield a nine-point (0 - 8) severity rating, and these ratings may be summed across items to yield a severity score for the full PTSD syndrome.
- 2) *The Structured Clinical Interview for the DSM-IV* (SCID; First, Spitzer, Williams, and Gibbon, 1997) is a semi-structured clinical interview designed to determine DSM-IV diagnoses, and its psychometric properties have been well-established (First et al., 1997). The SCID and selected modules for the DSM-IV, which included past major depression, past panic disorder, and past generalized anxiety disorder, were administered and collapsed into one variable (past psychopathology present or absent).
- 3) *The WTC Exposure Questionnaire* was developed by a panel of trauma experts to assess occupational exposure through performance of disaster work as well as personal exposure to the WTC attacks (e.g., loss of loved ones). The development of the exposure index in this study began with variables that were shown in the disaster literature to predict PTSD and revised with questions that emerged as unique to working at the WTC site. For the purposes of this study, a total exposure score was calculated by summing up the items that the participants reported as having experienced.

Self-reports

- 4) *High-K Strategy Scale* (HKSS; Giosan, 2006). HKSS is a 26-item scale tapping into various indicators of high-K strategy. These indicators include upward mobility (high-K strategists seek opportunities to gain resources for greater investment in their offspring), health and attractiveness (high-K strategists invest in somatic, rather than reproductive, effort, therefore they should enjoy better health and be, consequently, more attractive), social capital, or risk consideration. As predicted, this construct correlated significantly with perceived offspring quality (through the fact that high-K strategists invest their time and resources in parental effort). It also correlated negatively with number of previous marriages (because high-K strategists are committed to long-term relationships). Also, it correlated with objective measures of health, educational level and social support (because high-K strategists invest in somatic, educational opportunities and social capital for gaining better access to resources for their offspring). As expected, this construct showed a weak but significant correlation with actual number of children (the net result of a successful high-K strategy is fitness, but characterized through smaller number of children). Lastly, since a high-K strategy is theorized as a cohesive reproductive strategy, the indicators that make it up were predicted to have high internal consistency. Indeed, HKSS's Cronbach's α was .92 (Giosan, 2006). The items of this scale are summed up to yield a total score.
- 5) *The PTSD Checklist* (PCL; Weathers, Herman, Huska, and Keane, 1993) is a DSM-correspondent, 17-item self-report measure of PTSD. Using a five-point rating scale (1 = "not at all," 5 = "extremely"), respondents indicate how much they were bothered in the past month by each of the DSM-IV PTSD symptoms. The PCL is psychometrically sound and has been used in nearly two hundred studies across a variety of trauma populations (Weathers et al., 1993). The PCL yields both a continuous severity score and a PTSD diagnosis.
- 6) *Brief Symptom Inventory*. The Brief Symptom Inventory (BSI; Derogatis, 1993) is a 53-item abbreviated version of the Symptom Checklist 90-Revised (Derogatis and Spencer, 1982) that assesses a broad range of symptoms of psychopathology. The BSI yields scores for nine symptom dimensions as well as global indices, based on the calculation of mean item responses. In the present study we used the Global Severity Index (GSI) from the BSI, where higher scores indicate more severe overall psychopathology.
- 7) *The State Trait Anger Expression Inventory-2* (STAXI-2). The STAXI-2, a revised 57-item version of the STAXI (Spielberger, 1988), provides concise measures of the experience, expression, and control of anger. For the purposes of this study, the 15-item STAXI State Anger (S-Ang) scale was used to assess total state anger. The Cronbach's α for this scale in this study was 0.93.
- 8) *Sheehan Disability Scale*. The Sheehan Disability Scale (SDS; Sheehan, 1983) is a 10-point visual analogue scale that assesses disability across three domains: work, social life, and family life. The three items may be summed into a single dimensional measure of global functional impairment that ranges from 0 (unimpaired) to 30 (highly impaired), which was used in this study. This scale

has been widely used in psychopharmacology randomized controlled trials and has strong internal consistency (Cronbach's $\alpha = 0.89$ for the three-item scale).

- 9) *The Sleep Index* is a modified version of the Pittsburgh Sleep Quality Index (PSQI) (Buysse, Reynolds, Monk, Berman, and Kupfer, 1989), and was used to assess sleep disturbances and their frequency. PSQI has been widely used to measure sleep difficulty in different populations (Chesson et al., 2000) and has good reliability and validity (Buysse et al., 1989; Doi et al., 2000). The Cronbach's α of the sleep index in this study was acceptable (Cronbach's $\alpha = 0.77$).

Statistical approach

Examining the independent effects of high-K on measures of psychopathology was the main goal of the study. To this end, correlations between high-K, demographics (age, race, education, marital status), past psychopathology (dichotomized: "yes" for presence of past psychopathology, "no" for absence), prior trauma, self-reported (PCL) and clinician-administered (CAPS) PTSD severity score, general psychopathology (GSI), anger (STAXI-2), disability (Sheehan), sleep disturbance (total Sleep Index score) and Exposure to 9/11 total score, were first performed.

Simultaneous regressions of the outcomes examined (PCL, CAPS, GSI, STAXI-2, Sheehan and total Sleep Index score) on the variables that were significantly correlated with them were then performed to identify the variables that account for significant variance (tolerances greater than .8) in these outcomes. To examine the unique contribution of high-K, variables that showed tolerances greater than .8 were then controlled for in regressions of the outcomes on high-K.

Results

Demographics

Table 1 depicts the demographic characteristics of the sample ($n = 1400$). Overall, the sample consisted of primarily middle-aged, white, married men with the majority having at least a high-school education.

High-K and psychopathology

The correlation table for all the measures is presented in Table 2. Significant correlations between high-K and age, marital status, prior trauma, Clinician-Administered PTSD Scale total severity score, PTSD checklist total severity score, Global Severity Index, State-Trait Anger Expression Inventory score and Sheehan Disability Scale score, were found (all $ps \leq .01$).

Table 1. Sociodemographic characteristics

Characteristics of sample (<i>n</i> = 1400)		%
Age, <i>M</i> (<i>SD</i>)	45.88 (8.86)	
Gender		
	Male	97.6
	Female	2.4
Race		
	Caucasian	66.4
	African American	18.4
	Hispanic	13.6
	Asian	.9
	Other	.7
Education		
	Some or no high-school	2.0
	High-School graduate	48.7
	Some college or training	34.9
	College graduate	10.7
	More than college	3.7
Marital Status		
	Married	72.7
	Separated or divorced	11.7
	Single	13.7
	Cohabiting	1.6
	Widowed	.3

Regressions of PTSD Checklist (PCL) scores on high-K, after controlling for marital status, exposure and past psychopathology—variables that had tolerances greater than .8 in simultaneous regressions of PCL on variables that correlated significantly with it—showed that high-K accounts for an important 10% of the variance in the PCL (negative relationship) (Table 3).

Regressions of Clinician-Administered PTSD Scale (CAPS) severity scores on high-K, after controlling for past psychology, prior trauma, and exposure to 9/11 score, showed that high-K accounts for 5% in the variance (negative relationship) (Table 3).

Regressions of Sheehan total disability score on high-K, after controlling for age, race, marital status and prior trauma, showed that high-K accounts for 7% of the variance (negative relationship) (Table 4).

Table 2. Correlation matrix

	1	2	3	4	5	6	7	8	9	10	11	12	13
1 Age													
2 Race	-.09**												
3 Education	-.11**	0.05											
4 Marital	.20**	-.14**	.01										
5 Trauma	.10**	0.01	.03	-.01									
6 Pastpsych	-.13**	-0.09	.03	-.08	.21**								
7 CAPS	.01	-0.02	-.06	-.08	.21**	.39**							
8 PCL	-.03	-0.02	.01	-.06*	.21**	.30**	.75**						
9 Sleep	-.01	-0.04	.01	-.02	.20**	.31**	.57**	.67**					
10 Sheehan	-.12**	0.03	-.01	.10**	.20**	.28**	.53**	.65**	.56**				
11 Staxi	-.07*	0.07	-.03	.06**	.11**	.16**	.36**	.56**	.38**	.59**			
12 GSI	.03	0.02	-.01	.07**	.21**	.30**	.67**	.85**	.67**	.77**	.65**		
13 Exposure	-.10*	-.13**	.04	.01	.25**	.19**	.22**	.27**	.16**	.05	.04	.14**	
14 High-K	-.07**	-.05	.05	.23**	-.08**	-.08	-.26**	-.30**	-.28**	-.27**	-.25**	-.35**	.01

** $p < .01$

* $p < .05$

Legend: Pastpsych = prior psychopathology

CAPS = Clinician-Administered PTSD Scale total severity score

PCL = The PTSD Checklist severity score

Sleep = Sleep Disturbances score

Sheehan = Sheehan Disability Scale total score

Staxi = The State Trait Anger Expression Inventory score

GSI = Global Severity Index, where higher scores indicate more severe psychopathology.

Exposure = The WTC Exposure score

Table 3. Regressions of PCL and CAPS on high-K

	PCL					CAPS			
	R^2	F	Beta	t		R^2	F	Beta	t
Step 1	.17	20.14			Step 1	.19	30.80		
-Marital status			.02	.40	- Past psych			.33	7.30**
-Prior trauma			.14	3.01**	- Prior trauma			.10	2.15*
- Past psych			.22	4.96**	- Exposure			.14	3.08**
- Exposure			.19	4.32**					
Step 2					Step 2				
- High-K	.27	28.81	-.33	-7.28**	- High-K	.24	31.6	-.23	-5.26**

** $p < .01$

* $p < .05$

Regressions of Sleep disturbance on high-K, after controlling for total exposure score, prior trauma and past psychology showed that high-K accounts for 7% of the variance (negative relationship) in the sleep index (Table 4).

Table 4. Regressions of Sleep disturbance and Sheehan on high-K

SLEEP DISTURBANCE					SHEEHAN				
	R^2	F	Beta	t		R^2	F	Beta	t
Step 1 - Prior trauma - Past psych - Exposure	.16	18.40	.21 .24 .11	3.79** 4.46 2.01*	Step 1 -Age -Race -Marital status -Trauma	.07	15.90	-.16 .01 -.02 .17	-4.83** 2.5 -4.5 6.14
Step 2 - High-K	.23	20.99	-.26	-4.93**	Step 2 - High-K	.14	27.97	-.28	-8.43**

** $p < .01$

* $p < .05$

Regressions of the Global Severity Index (GSI) on high-K, after controlling for total exposure score, marital status, past psychopathology and prior trauma, showed that high-K is a negative predictor of general psychopathology, accounting for an important 12% in the variance (Table 5).

Regressions of State Trait Anger Expression Inventory (STAXI-2) scores on high-K, after controlling for age, race, marital status and prior trauma, showed that high-K is an independent negative predictor of anger, accounting for 6% in the variance (Table 5).

Table 5. Regressions of STAXI-2 and GSI on high-K

STAXI-2					GSI				
	R^2	F	Beta	t		R^2	F	Beta	t
Step 1 - Age - Race - Marital Status - Prior trauma	.03	8.06	-.10 .06 .01 .11	-3.93** 2.01* .17 3.65**	Step 1 - Marital status - Past psych - Prior trauma - Exposure	.13	15.00	.16 .24 .13 .07	.75 5.25** 2.92** 1.51
Step 2 - High-K	.09	21.06	-.26	-8.29**	Step 2 - High-K	.25	26.86	-.42	-8.06**

** $p < .01$

* $p < .05$

Discussion

The present study examined the associations between a high-K fitness strategy and psychopathology, assessed both through clinician-administered measures and self-reports. It was found that people who had higher scores on HKSS were also those who reported lower levels of psychopathology, and that high-K emerged as a significant predictor of psychopathology. This pattern was not singular, but was found across a range of psychopathologies, including PTSD, sleep disturbances, functional disability, anger, or overall psychopathology.

These findings, along with previous findings that found high-K to be strongly inversely related to depressive symptomatology (Giosan, 2007), suggest that high-K appears to be an important buffer of psychopathology. In our study, high-K accounted for

variance ranging from 5% to 12% in the different psychopathologies assessed, the greatest variance being found in the General Symptom Inventory total score.

The strong negative association between high-K and PTSD seems to speak against an evolutionary view that argues for the adaptive value of PTSD symptoms (e.g., Nesse, 2005), at least for this particular reproductive strategy. Our results show that lower scores on high-K, which are associated with reduced fitness (Giosan, 2006), are also associated with increased severity in PTSD symptoms. Since the findings documented in this study show that this reproductive strategy correlates negatively with the severity of PTSD, this result may offer indirect support to the view that holds that PTSD is the expression of an overlearned survival response in vulnerable individuals (Silove, 1998), which, when it develops, may have significant negative effects on those individuals' fitness.

Despite these important findings, several caveats are in order: First, as in any correlational study, the direction of causality cannot be established with certainty. One can argue that people score highly on the high-K precisely because they enjoyed better mental health in the first place and their resources could thus be channeled to the indicators that make up the high-K factor. This is a legitimate claim and testing the direction of causality is not a small feat. Further longitudinal studies examining the associations between changes in high-K and changes in psychopathology are needed to address such questions. Second, the logistics of the data collection biased the sample toward white males, which may raise questions about the generalizability of these results to other populations. Further studies on different populations need to be done to address this aspect. Third, one might argue that the relationships found in this study might be the result of a broad association between general mate quality/value and psychopathology. The design of the study did not have mate quality in the list of the variables examined therefore it could not address this question. Further research aimed at clarifying these complex associations should be done.

Despite these limitations, the present study, to our knowledge the first to link high-K and psychopathology, offers evidence that this specific life history strategy is strongly negatively linked to, and a negative predictor of, a broad range of psychopathologies.

Directions for further research

The ramifications of the findings in the present study are multiple. One avenue of research is to examine the effects of interventions on high-K on psychopathology. Indeed, since many of the indicators making up the high-K independent criterion of fitness are modifiable (e.g., appearance, social capital, personal safety, status) the question of whether active interventions on such indicators would translate into decreased psychopathology gains practical and clinical significance. A positive answer to this empirical question would likely aid therapists in their quests of finding more effective interventions.

Also, the high-K measure used in this study is understood as a unidimensional construct tapping into “slow” life history strategy, with high scores representing successful resolution of the challenges associated with a high-K strategy, and low scores representing a faulty high-K fitness strategy and not necessarily “fast” life history strategy. It would be important and complementary to this work to examine whether faulty fitness strategies in general—being they faulty high-K or faulty low-K—are associated with mental illness.

Received 4 August 2008; Revision submitted 22 September 2008; Accepted 17 November 2008

References

- Andreasen, N.C. (1984). *The broken brain: The biological revolution in psychiatry*. New York: Harper and Row.
- Avila, M., Thaker, G., and Adami, H. (2001). Genetic epidemiology and schizophrenia: A study of reproductive fitness. *Schizophrenia Research*, 47, 233-241.
- Blake, D.D., Weathers, F.W., Nagy, L.M., Kaloupek, D.G., Gusman, F.D., Charney, D.S., and Keane, T.M. (1995). The development of a clinician-administered PTSD scale. *Journal of Traumatic Stress*, 8, 75-90.
- Bogaert, A.F., and Rushton, J.P. (1989). Sexuality, delinquency and r/K reproductive strategies: Data from a Canadian university sample. *Personality and Individual Differences*, 10, 1071-1077.
- Buyse, D.J., Reynolds, C.F., Monk, T.H., Berman, S.R., and Kupfer, D. J. (1989). The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Research*, 28, 193-213.
- Chesson, A., Hartse, K., Anderson, W.M., Davila, D., Johnson, S., Littner, M., et al. (2000). Practice parameters for the evaluation of chronic insomnia: an American Academy of Sleep Medicine Report. *Standards of Practice Committee of the American Academy of Sleep Medicine*, 23, 247-41.
- Cowling, V., Luk, E.S.L., Mileskin, C., and Birleson, P. (2004). Children of adults with severe mental illness: Mental health, help seeking and service use. *Psychiatric Bulletin*, 28, 43-46.
- Crow, T.J. (1991). Origins of psychosis and the "Descent of Man". *British Journal of Psychiatry*, 159, 76-82.
- Crow, T.J. (1995). A Darwinian approach to the origins of psychosis. *British Journal of Psychiatry*, 167, 12-25.
- Crow, T.J. (1997). Is schizophrenia the price that *Homo sapiens* pays for language? *Schizophrenia Research*, 28, 127-141.
- Derogatis, L.R. (1993). *Brief Symptom Inventory: Administration Scoring and Procedures Manual*, 3rd ed. Minneapolis, MN: National Computer Systems, Minneapolis, MN.
- Derogatis, L.R., and Spencer, M.S. (1982). *The Brief Symptom Inventory (BSI): Administration, scoring, and procedures manual*. Baltimore, MD: Johns Hopkins University School of Medicine, Clinical Psychometrics Research Unit.
- Doi, Y., Minowa, M., Uchiyama, M., Okawa, M., Kim, K., Shibui, K., et al. (2000). Psychometric assessment of subjective sleep quality using the Japanese version of the Pittsburgh Sleep Quality Index (PSQI-J) in psychiatric disordered and control subjects. *Psychiatry Research*, 97, 165-172.
- Figueredo, J.A., Vasquez, G., Brumbach, B.H., Schneider, S.M.R., Sefcek, J.A., Tal, I.R., Hill, D., Wenner, C.J., Jacobs, W.J. (2006). Consilience and life history theory: From genes to brain to reproductive strategy. *Developmental Review*, 26, 243-275.
- First M.B., Spitzer R.L., Williams J.B.W., Gibbon M. (1997). *Structured Clinical Interview for DSM-IV (SCID)*. Washington, DC: American Psychiatric Association.

- Fiske, A.P., and Haslam, N. (1997). Is obsessive-compulsive disorder a pathology of the human disposition to perform socially meaningful rituals? Evidence of similar content. *The Journal of Mental Disease*, 185, 211-222.
- Giosan, C. (2006). High-K strategy scale: A measure of the high-K independent criterion of fitness. *Evolutionary Psychology*, 4, 394-405.
- Giosan, C. (2007). High-K reproductive strategy as a negative predictor of depressive symptomatology. Paper presented at the Northeast Evolutionary Psychology Society conference, SUNY at New Paltz, April 2007.
- Haukka, J., Suvisaari, J., Lönnqvist, J. (2003). Fertility of patients with schizophrenia, their siblings, and the general population: A cohort study from 1950 to 1959 in Finland. *The American Journal of Psychiatry*, 160, 460-463.
- Howard, L.M., Kumar, C., Lesse, M., and Thornicroft, G. (2002). The general fertility rate in women with psychotic disorders. *Psychiatry*, 159, 991-997.
- Jablensky, A., Zubrick, S., Morgan, V., Bower, C., Yellachich, L., Nguyen, H., Pinder, T., Montgomery, P., Croft, M., and Castle, D. (1998). Reproductive pathology in women with schizophrenia and affective psychoses. *Schizophrenia Research*, 29, 20-21.
- Leith, K.P., and Baumeister, R. (1996). Why do moods increase self-defeating behavior? Emotion, risk-taking and self-regulation. *Journal of Personality and Social Psychology*, 71, 1250-1267.
- Mac Arthur R.H., and Wilson, E.O. (1967). *The theory of island biogeography*. Princeton, NJ: Princeton University Press.
- Marks, I.M. (1987). *Fears, phobias, and rituals*. New York: Oxford University Press.
- Marks, I.M., and Nesse, R.M. (1994). Fear and Fitness: An evolutionary analysis of anxiety disorders. *Ethology and Sociobiology*, 15, 247-261.
- McGrath, J., Hearle, J., Lenner, L., Plant, K., Drummond, A., and Barkla, J. (1998). Fertility of patients with psychoses versus their unaffected siblings. *Schizophrenia Research*, 29, 20-21.
- McGuire, M.T., Troisi, A., and Raleigh, M.M. (1997). Depression in evolutionary context. In S. Baron-Cohen, (Ed.). *The maladapted mind: Classic readings in evolutionary psychopathology* (pp. 255-282). London: Erlbaum/Taylor and Francis.
- Mealey, L. (2000). Anorexia: A “losing” strategy? *Human Nature*, 11, 105-116.
- Nesse, R. (1998). Emotional disorders in evolutionary perspective. *British Journal of Medical Psychology*, 71, 397-415.
- Nesse, R.M. and Williams, G.C. (1994). *Why we get sick: The new science of Darwinian medicine*. New York: Times Books.
- Nesse, R.M. (2005). Evolutionary psychology and mental health. In D.M. Buss (Ed.). *The handbook of evolutionary psychology* (pp. 903-927). New Jersey: Wiley.
- Pianka, E.R. (1970). On r- and K-selection. *American Naturalist*, 104, 592-596.
- Rushton, J.P. (1985). Differential K Theory: The sociobiology of individual and group differences. *Personality and Individual Differences*, 6, 441-452.
- Rushton, J.P. (1990). Sir Francis Galton, epigenetic rules, genetic similarity theory, and human life history analysis. *Journal of Personality*, 58, 117-140.
- Sheehan, D.V. (1983). *The anxiety disease*. New York: Scribner's.
- Silove, D. (1998). Is posttraumatic stress disorder an overlearned survival response? An evolutionary-learning hypothesis. *Psychiatry*, 61, 181- 190.

- Spielberger, C.D. (1988). Anger expression scale. In M. Hersen and A.S. Bellack (Eds.), *Dictionary of behavioral assessment techniques*. (pp. 446-448). New York: Pergamon Press.
- Valenstein, E.S. (1998). *Blaming the brain: The truth about drugs and mental health*. New York: Free Press.
- Wakefield, J.C. and First, M. (2003). Clarifying the distinction between disorder and non-disorder: Confronting the overdiagnosis ("false positives") problem in DSM-V. In K.A. Phillips, M.B. First, and H.A. Pincus (Eds.), *Advancing DSM: Dilemmas in psychiatric diagnosis* (pp. 23-56). Washington, D.C.: American Psychiatric Press.
- Wakefield, J.C. (1997). Diagnosing DSM: Pt. 1. DSM and the concept of mental disorder. *Behavior Research and Therapy*, 35, 633-650.
- Weathers F.W., Litz B.T., Herman D.S., Huska J.A., Keane T.M. (1993). *The PTSD Checklist (PCL): Reliability, validity, and diagnostic utility*. Paper presented at the Annual Convention of the International Society for Traumatic Stress Studies, San Antonio, TX, October 1993.
- Wilson, E.O. (1975). *Sociobiology: The new synthesis*. Cambridge, MA: Harvard University Press.
- Wolpert, L. (1999). *Malignant sadness: The anatomy of depression*. New York: Free Press.