

**SOME ADDITIONAL NOTES ON THE GENUS *QUEDIUS*
STEPHENS, 1829 (COLEOPTERA: STAPHYLINIDAE:
STAPHYLININAE) FAUNA OF TURKEY**

İnanç Özgen*, Eduard A. Khachikov and Semih Örgel*****

* Baskil Vocational School, Firat University, Elazığ, TURKEY.

** Rostov Branch of Russian Entomological Society, 344030 Rostov-on-Don, RUSSIA.

*** Department of Biology, Faculty of Science, University of Ege, 35100 Bornova, İzmir, TURKEY.

[**Özgen, İ., Khachikov, E. A. & Örgel, S.** 2016. Some additional notes on the genus *Quedius* Stephens, 1829 (Coleoptera: Staphylinidae: Staphylininae) fauna of Turkey. *Munis Entomology & Zoology*, 11 (2): 619-621]

ABSTRACT: In this study, 10 species of the genus *Quedius* Stephens were recorded from different regions of Turkey. Additional notes on most of them new to certain Turkish regions and provinces are given. *Quedius sofiri* Khachikov, 2005 is recorded for the first time from Turkey.

KEY WORDS: Coleoptera, Staphylinidae, Staphylininae, *Quedius*, fauna, new record, Turkey

The beetle family Staphylinidae is the most diverse group of the order Coleoptera, comprising more than 56.000 valid species belonging to 33 subfamilies in the world (Newton, 2007; Grebennikov & Newton, 2009). According to recent contributions, about 1900 species of the family Staphylinidae are known from Turkey (Anlaş, 2007, 2009; Schülke & Smetana, 2015).

78 species of the genus *Quedius* Stephens are currently distributed in Turkey (Anlaş, 2009; Anlaş & Rose, 2009; Abacigil et al., 2013; Firat & Sert, 2016), 20 of them occur only in Turkey and represent more than 25% of the Turkish *Quedius* fauna. Nevertheless, the *Quedius* fauna of Turkey and its distribution are still poorly investigated and many species are known only from a small number of localities.

The aim of this study is to enhance scientific knowledge on the distribution of Turkish *Quedius*.

MATERIALS AND METHODS

The present paper is based on material collected between the dates of 2007-2010 in the different parts of Turkey. The reference specimens in this study are deposited in the private collection of the first author. Classification and nomenclature of the genus *Quedius* suggested by Herman (2001) and Schülke & Smetana (2015) has been followed in this study.

RESULTS

***Quedius cinctus* Paykull, 1790**

Material examined: Adiyaman: 1 ex., 05.IV.2008, Gölbaşı, Karakuyu village, 1 km W, 1210 m, 37°41'52"N, 37°38'14"E, leg. Yağmur. Kastamonu: 2 exs., 06.IX.2010, Ilgaz Dağı, 1858 m, 41°04'41"N, 33°43'56"E, leg. Kunt.

Distribution in Turkey: Ankara, Antalya, Bolu, Eskişehir, İstanbul, İzmir, Karaman, Kırşehir, Mersin, Nevşehir, Rize, Siirt (Anlaş, 2009; Anlaş & Rose, 2009; Özgen & Anlaş, 2010; Assing, 2013; Çiftçi & Hasbeli, 2016; Firat & Sert, 2016).

Quedius cruentus (Olivier, 1795)

Material examined: *Bursa*: 2 exs., 28.VIII.2007, Uludağ, Aynalı lake, leg. Koç.

Distribution in Turkey: Balıkesir and Manisa (Anlaş, 2009; Abacigil et al., 2013).

Quedius lateralis (Gravenhorst, 1802)

Material examined: *Uşak*: 2 exs., Eşme, Kısık 2 km NE, Gediz river bank, 470 m., 38°38'06"N, 28°57'19"E, 23.IV.2010, leg. Anlaş.

Distribution in Turkey: Balıkesir, Eskişehir, İsparta, İstanbul, İzmir (Anlaş, 2009; Assing, 2013; Abacigil et al., 2013, Fırat & Sert, 2016).

Quedius levicollis Brulle, 1832

Material examined: *Elaçığ*: 2 exs., 17.V.2011, Harput, leg. Özgen, *Izmir*: 10 exs., 10.XII.2008, Bayındır, Yakapınar, leg. Anlaş; 6 exs., 29.V.2010, Bozdağ Kayak Merkezi, 2000 m, leg. Anlaş; 2 exs., 28.V.2010, Buca, Kaynaklar, leg. Anlaş; 3 exs., 03.IV.2010, Menemen, Emiralem, pitfall traps, leg. Yağmur; 2 exs., 11.XII.2010, Buca, Kaynaklar, pitfall traps, leg. Yağmur. *Kütahya*: 6 exs., 24.IV.2010, Şaphane, İlicasu 1 km N, 723 m, 38°56'58"N, 29°17'29"E, leg. Anlaş. *Malatya*: 2 exs., 14.IX.2007, Arapgir 3 km NE, 912 m, 39°04'13"N, 38°30'34"E, leg. Anlaş & Yağmur. *Manisa*: 2 exs., 30.X.2010, Turgutlu, Çikrikçi-Baktrılı road, leg. Anlaş; 1 ex., 30.X.2010, Gölstmarmara, Beyler 1 km NE, 120 m, 38°42'10"N, 27°58'56"E, leg. Anlaş.

Distribution in Turkey: Adana, Ankara, Balıkesir, Çanakkale, Erzurum, Eskişehir, İzmir, Manisa (Anlaş, 2009; Anlaş & Rose, 2009; Kesdek et al., 2009; Abacigil et al., 2013; Surgut & Varlı, 2012; Çiftçi & Hasbenli, 2016; Fırat & Sert, 2016).

Quedius ochripennis Menetries, 1802

Material examined: *Izmir*: 1 ex., 29.V.2010 Bozdağ Kayak Merkezi, ca. 2000 m, leg. Anlaş.

Distribution in Turkey: Adana, Ankara, Çankırı, Eskişehir, Hatay, İsparta, İstanbul, Kayseri, Kırşehir, Mardin, Nevşehir, Niğde, Yozgat (Anlaş, 2009; Japoshvili & Anlaş, 2011; Çiftçi & Hasbenli, 2016; Fırat & Sert, 2016).

Remarks: First record for the Aegean Region of Turkey.

Quedius nitipennis (Stephens, 1833)

Material examined: *Adiyaman*: 1 ex., 02.VII.2007, Gölbasi, Akçalar road inside 1 km, leg. Yağmur; 1 ex., 05.IV.2008, Gölbasi, Karakuyu village, 1 km W, 1210 m, 37°41'52"N, 37°38'14"E, leg. Yağmur.

Distribution in Turkey: Adana, Ardahan, Artvin, Erzurum, Kayseri, Kocaeli, Mersin, Niğde (Anlaş, 2009; Kesdek et al., 2009; Assing, 2013).

Quedius semiobscurus Marsham, 1802

Material examined: *Izmir*: 7 exs., 04.V.2010, 03.IV.2010 and 11.XII.2010, Buca, Kaynaklar, pitfall traps, leg. Yağmur; Menemen, Emiralem, pitfall traps, leg. Yağmur.

Distribution in Turkey: Balıkesir and Bursa (Anlaş, 2009; Abacigil et al., 2013).

Remarks: First record for the Aegean Region of Turkey.

Quedius sofiri Khachikov, 2005

Material examined: *Kastamonu*: 2 exs., 06.IX.2010, Ilgaz Dağı, 1858 m, 41°04'41"N, 33°43'56"E, leg. Kunt.

Remarks: This species is known only from its type locality in Russian South European Territory (Khachikov, 2005). Thus, this species is recorded for the first time from Turkey.

Quedius suramensis Eppelsheim, 1880

Material examined: *Kastamonu*: 1 ex., 06.IX.2010 Ilgaz Dağı, 1858 m, 41°04'41"N, 33°43'56"E, leg. Kunt.

Distribution in Turkey: Artvin, Bolu, Bursa, Gümüşhane, Kastamonu, Ordu, Rize, Sakarya, Sinop, Trabzon (Coiffait, 1977, 1978; Anlaş, 2009).

***Quedius vicinus* Menetries, 1832**

Material examined: *Şanlıurfa*: 2 exs, 11.XI.2010, Birecik, Kelaynak Valley, leg. Anlaş & Yağmur.

Distribution in Turkey: Bolu, Giresun, Kastamonu, Kırıkkale, Ordu, Rize, Sakarya, Trabzon (Korge, 1964, Bordoni, 1976, Anlaş, 2009; Firat & Sert, 2016).

Remarks: First record for the southeastern Anatolia.

ACKNOWLEDGEMENTS

Authors would like to thank our colleagues for making their staphylinid by-catches to us.

LITERATURE CITED

- Abacıgil, T. Ö., Varlı, S. & Tezcan, S. 2013. Faunistic studies on Staphylininae (Coleoptera: Staphylinidae) in Kazdağıları (Balıkesir province) in Turkey. *Munis Entomology & Zoology*, 8 (1): 415-433.
- Anlaş, S. 2007. The present situation Staphylinidae (Coleoptera) fauna of Turkey. *Linzer biologische Beiträge*, 39 (2): 5-9.
- Anlaş, S. 2009. Distributional checklist of the Staphylinidae (Coleoptera) of Turkey, with new and additional records. *Linzer biologische Beiträge*, 41: 215-342.
- Anlaş, S. & Rose, A. 2009. Some additional notes about Staphylininae (Coleoptera: Staphylinidae) fauna of Turkey. *Munis Entomology & Zoology*, 4 (2): 327-333.
- Bordoni, A. 1976. Quarto contributo alla conoscenza degli Staphylinidae della Turchia (Coleoptera). *Fragmента Entomologica*, 12: 221-239.
- Çiftçi, D. & Hasbeli, A. 2016. Faunistic Studies on Steninae, Paederinae and Staphylinae (Coleoptera: Staphylinidae) from Sündiken Mountains (Turkey). *Entomofauna*, 37 (6): 101-136.
- Coiffait, H. 1977. Note sur quelques *Quedius* et *Heterothops* nouveaux ou mal connus (Col. Staphylinidae). *Nouvelle Revue d'Entomologie*, 7 (2): 133-143.
- Firat, S. & Sert, O. 2016. Faunistic and zoogeographical composition and preliminary evaluations of some ecological features of the subfamily Staphylininae (Coleoptera: Staphylinidae) of the Central Anatolian Region of Turkey. *Turkish Journal of Zoology*, 40 (1): 164-185.
- Grebennikov, V. V. & Newton, A. F. 2009. Good-bye Scydmaenidae, or why the ant-like stone beetles should become megadiverse Staphylinidae sensu latissimo (Coleoptera). *European Journal of Entomology*, 106 (2): 275-301.
- Japoshvili, G. & Anlaş, S. 2011. Notes on the Family Staphylinidae (Coleoptera) Collected by Pitfall Traps in Gölcük Natural Park, Isparta Province of Turkey. *J. Entomol. Res. Soc.*, 13: 41-48.
- Kesdek, M., Yıldırım, E. & Anlaş, S. 2009. Contribution to the knowledge of Staphylinidae (Coleoptera) fauna of Turkey. *Munis Entomology & Zoology*, 4 (2): 355-364.
- Khachikov, E. A. 2005. Novye zhukhi-stafilinidy (Coleoptera, Staphylinidae) s yuga Rossii. New species of rovebeetles (Coleoptera, Staphylinidae) from the Southern Russia. *Caucasian Entomological Bulletin*, 1 (2): 119-122.
- Korge, H. 1964. Carabiden- und Staphylinidenfunde in den Pontischen Gebirgen Kleinasiens und in Mazedonien (Coleoptera). *Reichenbachia*, 4 (14): 105-126.
- Newton, A. F. 2007. Documenting biodiversity: how well are we doing in Staphyliniformia (Coleoptera) Entomological Society of America poster presentation. 2007; Do471. Available (ESA members only) at <http://esa.confex.com/esa/2007/techprogram/paper_32168.htm>.
- Özgen, İ. & Anlaş, S. 2010. A cow dung investigation on Staphylinidae (Coleoptera) with a new record from Turkey. *Munis Entomology Zoology*, 5 (2): 642-645.
- Schülke, M. & Smetana, A. 2015. Staphylinidae, pp. 304-1134. – In: Löbl I. & Löbl D. (eds), Catalogue of Palaearctic Coleoptera. Volume 2. Hydrophiloidae – Staphylinoidae. Revised and updated edition. – Leiden: Brill: xxvi + 1702 pp.
- Surgut, H. & Varlı, S. V. 2012. An evaluation on Coleoptera (Insecta) species collected by pitfall traps in Karabiga (Çanakkale province) of Turkey. *Munis Entomology & Zoology*, 7 (1): 449-461.