

**SOME ADDITIONAL NOTES ON THE GENUS
PHILONTHUS STEPHENS (COLEOPTERA:
STAPHYLINIDAE: STAPHYLININAE) IN TURKEY**

İnanç Özgen*, Eduard A. Khachikov,
Semih Örgel*** and Çağatay Altın******

* Baskil Vocational School, Fırat University, Elazığ, TURKEY. E-mail: inancozgen@gmail.com

** Rostov Branch of Russian Entomological Society, 59 Alexandrovsky spusk str., 344030 Rostov-on-Don, RUSSIA.

*** Department of Biology, Faculty of Science, University of Ege, 35100 Bornova, Izmir, TURKEY.

**** Alaşehir Vocational School, Celal Bayar University, 45600, Alaşehir, Manisa, TURKEY.

[Özgen, İ., Khachikov, E. A., Örgel, S. & Altın, Ç. 2016. Some additional notes on the genus *Philonthus* Stephens (Coleoptera: Staphylinidae: Staphylininae) in Turkey. *Munis Entomology & Zoology*, 11 (1): 263-267]

ABSTRACT: In this study, 24 species of the genus *Philonthus* were recorded from different regions of Turkey. Additional notes on most of them new to certain Turkish regions and provinces are given. Among them *P. concinnus* (Gravenhorst, 1802), *P. nitidicollis* (Lacordaire, 1835) and *P. rufimanus* (Heer, 1839) are found the most common and abundant species.

KEY WORDS: Coleoptera, Staphylinidae, Staphylininae, *Philonthus*, fauna, Turkey

The Staphylinidae is a widespread and rather large family of the order Coleoptera, comprising about 60.000 species belonging to 33 subfamilies in all zoogeographical regions of the world (Newton, 2007). 1600 species of Staphylinidae are listed in Turkey by Anlaş (2009), 346 of them are belong to the subfamily Staphylininae.

The genus *Philonthus* is one of the largest genus within Staphylininae, at the present state of knowledge comprising some 1200 species in the world. (Schillhammer, 1998). In Turkey, 65 *Philonthus* species are known (Anlaş, 2009, updated).

The *Philonthus* fauna of Turkey are still poorly investigated. The aim of this study is to enhance scientific knowledge on the distribution of Turkish *Philonthus*.

MATERIAL AND METHODS

The present paper is based primarily on material collected in different parts of Turkey by using different collection methods in 2006-2011. Material is deposited in the private collection of the first author. Classification and nomenclature of the *Philonthus* suggested by Herman (2001) and Smetana (2004) has been followed in this study.

RESULTS

Philonthus alberti (Schillhamer, 2000)

Material examined: **Afyonkarahisar:** 1 ex., Şuhut, Dadak 2 km N, 1320 m, 38°36'18"N, 30°26'59"E, 11.VIII.2010, leg. Anlaş. **Erzincan:** 2 exs., Refahiye, Şahverdi- Aydıncık, 1753 m, 39°50'58"N, 38°48'57"E, 17.V.2011, leg. Anlaş, Khachikov & Özgen; 2 exs., Üzümlü, Küçük Sarıkaya creek bank, 1713 m, 39°14'20"N, 39°50'02"E, 18.V.2011, leg. Anlaş, Khachikov & Özgen. **Kırklareli:** 1 ex., Demirköy, Rhodendron forest, 25.V.2010, leg. Kunt. **Distribution in Turkey:** Artvin, Bolu (Anlaş, 2009).

Remarks: This species is recorded for the first time from Aegean Region.

***Philonthus atratus* (Gravenhorst, 1802)**

Material examined: **Diyarbakır:** 1 ex., Silvan, Boyunlu, 1084 m, 38°13'45"N, 40°58'50"E, 15.IV.2010, leg. Özgen; 2 exs., Çermik, Çakmak, Çamaltı, 29.IX.2010, leg. Özgen. **Muş:** 2 exs., Buğlan Geçidi, 30.V.2011, leg. Khachikov & Kasatkin.

Distribution in Turkey: Ankara, Batman, Çankırı, Diyarbakır, Erzurum, Konya, Malatya, Mersin, Niğde (Anlaş, 2009; Anlaş & Rose, 2009; Kesdek et al., 2009; Özgen et al., 2010).

***Philonthus carbonarius* (Gravenhorst, 1802)**

Material examined: **Bursa:** 1 ex., Uludağ, Kilimli Lake, 2279 m, 40°04'57"N, 23°13'31"E, 28.VII.2008, leg. Koç. **Kahramanmaraş:** 2 exs., Başkonuş Yaylası, 41°59'58"N, 37°28'61"E, 1291 m, 21.VI.2007, leg. Yağmur.

Distribution in Turkey: Antalya, Ardahan, Bingöl, Elazığ, Erzurum, Kars, Manisa (Anlaş, 2009; Anlaş & Rose, 2009a; Kesdek et al., 2009; Özgen & Anlaş, 2010).

***Philonthus cochleatus* (Scheerpeltz, 1937)**

Material examined: **Muş:** 3 exs., Varto, 31.V.2011, leg. Khachikov & Kasatkin.

Distribution in Turkey: Adana (Smetana, 1954; Anlaş, 2009).

Remarks: This species is recorded for the first time from Eastern Anatolia.

***Philonthus cognatus* (Stephens, 1832)**

Material examined: **Afyonkarahisar:** 3 exs., Şuhut, Dadak 2 km N, 1320 m, 38°36'18"N, 30°26'59"E, 11.VIII.2010, leg. Anlaş. **Gaziantep:** 2 exs., Şahinbey, Sarısalkım 1 km S, 12.XI.2006, leg. Yağmur. **Izmir:** 4 exs., Bozdağ Zirve, 2500 m, 28.III.2007, leg. Anlaş. **Manisa:** 1 ex., Turgutlu, Ovacık Yaylası, 14.VII.2006, leg. Anlaş.

Distribution in Turkey: Ardahan, Artvin, Erzurum, Kars, Mersin, Trabzon, (Anlaş, 2009; Kesdek et al., 2009).

Remarks: This species is recorded for the first time from Western Anatolia.

***Philonthus concinnus* (Gravenhorst, 1802)**

Material examined: **Afyonkarahisar:** 3 exs., Sinanpaşa 15 km SW, Uluköy 2 km N, 1600 m, 38°42'48"N, 30°04'40"E, 23.IV.2010, leg. Anlaş. **Bursa:** 4 exs., Uludağ, Kovukdere Lake (Saklıgöl) 2206 m, 40°05'12"N 29°12'22"E 29.VII.2008, leg. Koç; 1 ex., Uludağ, Aynalı Gölü, 2310 m, 40°04'13"N, 29°14'01"E, 30.VIII.2008, leg. Koç; 2 exs., Uludağ, Kilimli Lake, 2279 m, 40°04'57"N, 23°13'31"E, 28.VII.2008, leg. Koç. **Diyarbakır:** 1 ex., Silvan, Boyunlu, 1084 m, 38°13'45"N, 40°58'50"E, 15.IV.2010, leg. Özgen; 2 exs., Silvan, Boyunlu 4 km S, 21.V.2010, leg. Özgen; 2 exs., Ergani pass, 26.V.2010, leg. Özgen; 2 exs., Eğil road 3 km inside, 04.VI.2010, leg. Özgen; 5 exs., Eğil, Kalkan, 21.V.2010, leg. Özgen & Yağmur; 3 exs., Çermik, Artuklu, 775 m., 38°08'26"N, 39°31'33"E, 01.VI.2010, leg. Özgen. **Elazığ:** 1 ex., Doğukent, 12.VI.2010, leg. Özgen; 1 ex., Gümüşkavak, 13.VI.2010, leg. Özgen. **Erzincan:** 6 exs., 17.V.2011, Refahiye, Şahverdi-Aydıncık, 1753 m, 39°50'58"N, 38°48'57"E, leg. Anlaş, Khachikov & Özgen. **Gaziantep:** 3 exs., Şahinbey, Sofalıcı village, Sof Mts., 37°08'42"N, 37°07'44"E, 23.III.2008, leg. Yağmur. **Gümüşhane:** 4 exs., 16.V.2011, Kelkit, Çimenli, 1689 m, 39°58'06"N, 39°22'48"E, leg. Anlaş, Khachikov & Özgen. **Izmir:** 6 exs., Ödemiş, Bozdağlar, road to ski resort, ca. 1600 m, 38°21'N, 28°06'E, 21.V.2006, leg. Anlaş. **Malatya:** 1 ex., Hekimhan 2 km NW, İpekyolu beldesi, Girmana mevki, 25.VII.2007, leg. Yağmur. **Manisa:** 3 exs., Spil National Park, ca. 1200 m, 38°33'43"N, 27°23'25"E, 30.IX.2006, leg. Anlaş; 2 exs., Spil National Park, ca. 1000 m, 38°33'N, 27°23'E, 27.IX.2008, leg. Anlaş. **Mersin:** 4 exs., Çamlıyayla, Korucak 1 km E, 715 m, 37°08'56"N, 34°42'51"E, 22.VII.2010, leg. Anlaş. **Muş:** 2 exs., Buğlan Geçidi, 30.V.2011, leg. Khachikov & Kasatkin; 2 exs., Varto, 31.V.2011, leg. Khachikov & Kasatkin. **Siirt:** 5 exs., Baykan 4 km NE, ca. 770 m, 38°11'42"N, 41°49'03"E, 17.XI.2010, leg. Anlaş. **Tunceli:** 2 exs., Pülümür 3 km SE, 19.V.2011, leg. Anlaş, Khachikov & Özgen. **Uşak:** 4 exs., Eşme, Kısıcık 2 km NE, Gediz river bank, 470 m., 38°38'06"N, 28°57'19"E, 29.V.2010, leg. Anlaş.

Distribution in Turkey: Adana, Ankara, Antalya, Ardahan, Bingöl, Bolu, Diyarbakır, Erzincan, Erzurum, Iğdır, Kayseri, Konya, Manisa, Mardin, Mersin, Tunceli (Anlaş, 2009; Anlaş & Rose, 2009; Kesdek et al., 2009; Özdemir & Sert, 2009; Özgen & Anlaş, 2010; Özgen et al., 2010).

***Philonthus corruscus* (Gravenhorst, 1802)**

Material examined: **Gümüşhane:** 2 exs., Kelkit, Çimenli, 1689 m, 39°58'06"N, 39°22'48"E, 16.V.2011, Anlaş, Khachikov & Özgen. **Manisa:** 1 ex., Otoman, 330 m, 38°44'47"N, 27°10'34"E, 04.X.2008, leg. Anlaş. **Muş:** 1 ex., Buğlan Geçidi, 30.V.2011, leg. Khachikov & Kasatkin.

Distribution in Turkey: Adana, Ankara, Izmir, Mardin, Mersin? (Anlaş, 2009; Özdemir & Sert, 2009; Özgen & Anlaş, 2010).

***Philonthus cruentatus* (Gmelin, 1790)**

Material examined: **Gümüşhane:** 2 exs., Kelkit, Yukarı Özlüce 4 km S, 1615 m, 39°58'27"N, 39°29'42"E, 16.V.2011, Anlaş, Khachikov & Özgen.

Distribution in Turkey: Antalya, Denizli, İstanbul (Anlaş, 2009; Anlaş & Rose, 2009).

Remarks: This species is recorded for the first time from Eastern Anatolia.

***Philonthus debilis* (Gravenhorst, 1802)**

Material examined: **Afyonkarahisar:** 3 exs., Şuhut, Dadak 2 km N, 1320 m, 38°36'18"N, 30°26'59"E, 11.VIII.2010, leg. Anlaş. **Erzincan:** 1 ex., Refahiye, Şahverdi-Aydıncık, 1753 m, 39°50'58"N, 38°48'57"E, 17.V.2011, Anlaş, Khachikov & Özgen.

Distribution in Turkey: Adana, Ankara, Denizli, Mersin (Anlaş, 2009).

Remarks: This species is recorded for the first time from Eastern Anatolia.

***Philonthus ebeninus* (Gravenhorst, 1802)**

Material examined: **Elazığ:** 2 exs., Doğukent, 12.VI.2010, leg. Özgen. **Muş:** 2 exs., Buğlan Geçidi, 30.V.2011, leg. Khachikov & Kasatkin; 1 ex., Varto, 31.V.2011, leg. Khachikov & Kasatkin.

Distribution in Turkey: Adana, Antalya, Bursa, İzmir, Mersin (Anlaş, 2009; Anlaş & Rose, 2009).

Remarks: This species is recorded for the first time from Eastern Anatolia.

***Philonthus fumarius* (Gravenhorst, 1806)**

Material examined: **Muş:** 1 ex., Buğlan Geçidi, 30.V.2011, leg. Khachikov & Kasatkin; 1 ex., Varto, 31.V.2011, leg. Khachikov & Kasatkin.

Distribution in Turkey: İstanbul (Apfelbeck, 1901; Anlaş, 2009).

Remarks: This species is recorded for the first time from Eastern Anatolia.

***Philonthus intermedius* (Lacordaire, 1835)**

Material examined: **Adıyaman:** 4 exs., Gölbaşı, Akçalar road inside 1 km, 02.VII.2007, leg. Yağmur; 3 exs., Gölbaşı, Karakuyu village, 1 km W, 1210 m, 37°41'52"N, 37°38'14"E, 05.IV.2008, leg. Yağmur. **Erzincan:** 1 ex., Üzümlü, Küçük Sarıkaya creek bank, 1713 m, 39°14'20"N, 39°50'02"E, 18.V.2011, Anlaş, Khachikov & Özgen. **Gümüşhane:** 2 exs., Kelkit, Yukarı Özlüce 4 km S, 1615 m, 39°58'27"N, 39°29'42"E, 16.V.2011, Anlaş, Khachikov & Özgen. **İzmir:** 6 exs., Ödemiş, Bozdağlar, road to ski resort, ca. 1600 m, 38°21"N, 28°06"E, 21.V.2006, leg. Anlaş. **Muş:** 1 ex., Buğlan Geçidi, 30.V.2011, leg. Khachikov & Kasatkin.

Distribution in Turkey: Ankara, Antalya, Denizli, İzmir, Kahramanmaraş, Mardin, Mersin (Anlaş, 2009; Anlaş & Rose, 2009; Özgen & Anlaş, 2010).

***Philonthus micans* (Gravenhorst, 1802)**

Material examined: **Kırklareli:** 2 exs., Demirköy, Hamam Gölü, 01.IX.2009, leg. Kunt.

Distribution in Turkey: Mersin [Tarsus (=Tarsous), Caramania] (Peyron, 1858; Fauvel, 1874; Herman, 2001; Anlaş, 2009).

Remarks: This species is recorded for the second time from Turkey.

***Philonthus nitidicollis* (Lacordaire, 1835)**

Material examined: **Bursa:** 2 exs., Uludağ, Aynalı Gölü, 28.VIII.2007, leg. Koç.

Denizli: 3 exs., Babadağ, 1450 m 37°47'16"N, 28°47'45"E, 12.IV.2009, leg. Yağmur.

Dişarbakır: 1 ex., Dicle university campus, 05.VI.2010, leg. Özgen. **Gaziantep:** 3 exs., Şehitkamil, Sofalıcı 5 km SW, 37°07'44"N, 37°05'13"E, 23.III.2008, leg. Yağmur. **İzmir:** 7 exs., Bozdağ Zirve, 2500 m, 28.III.2007, leg. Anlaş; 5 exs., Bozdağ Kayak Merkezi, ca. 2000 m, 29.V.2010 and 28.VII.2009, leg. Anlaş; 5 exs., Karaburun, Hades cave, 06.04.2009, leg. Yağmur & Durmuş. **Manisa:** 4 exs., Spil National Park, ca. 1200 m, 38°33'43"N, 27°23'25"E, 30.IX.2006, leg. Anlaş. **Mardin:** 2 exs., Mazıdağı, Gürgöze, 950 m., 37°29'16"N, 40°31'06"E, 31.V.2010, leg. Özgen. **Siirt:** 8 exs., Baykan 4 km NE, ca. 770 m, 38°11'42"N, 41°49'03"E, 21.V.2010 and 17.XI.2010, leg. Anlaş & Yağmur. **Şırnak:** 2 exs., Dicle river 4 km W, Yalın-tepe village, 12.V.2007, leg. Yağmur.

Distribution in Turkey: Adana, Ankara, Antalya, Bingöl, Gaziantep, Isparta, İzmir, Konya, Siirt (Anlaş, 2009; Anlaş & Rose, 2009; Özgen & Anlaş, 2009; Japoshvili & Anlaş, 2011).

***Philonthus parvicornis* (Gravenhorst, 1802)**

Material examined: **Muş:** 2 exs., Varto, 31.V.2011, leg. Khachikov & Kasatkin.

Distribution in Turkey: Isparta, Muğla (Anlaş, 2009; Assing, 2013).

Remarks: This species is recorded for the first time from Eastern Anatolia.

***Philonthus quisquiliarius* (Gyllenhal, 1810)**

Material examined: **Dişarbakır:** 2 exs., Eğil, Kalkan 1 km SW, 38°08'37"N, 40°03'37"E, 13.IV.2008, leg. Yağmur; 1 ex., Eğil road 3 km inside, 04.VI.2010, leg. Özgen; 2 exs., Eğil, Kalkan, 21.V.2010, leg. Özgen & Yağmur.

Distribution in Turkey: Adana, Ankara, Izmir, Mersin (Anlaş, 2009; Özgen et al., 2010).
Remarks: This species is recorded for the first time from Eastern Anatolia.

***Philonthus rotundicollis* Menetries, 1832**

Material examined: **Erzincan:** 2 exs., Üzümlü, Küçük Sarıkaya creek bank, 1713 m, 39°14'20"N, 39°50'02"E, 18.V.2011, Anlaş, Khachikov & Özgen.

Distribution in Turkey: Gaziantep (Anlaş, 2007, 2009).

***Philonthus rubripennis* (Stephens, 1832)**

Material examined: **Mardin:** 2 exs., 31.V.2010, Mazıdağı, Gürgöze, 950 m, 37°29'08"N, 40°31'38"E, leg. Özgen. **Tunceli:** 1 ex., 13.IX.2007, Ovacık, Munzur Gözeleri, leg. Anlaş. 2 exs., Pülümür 3 km SE, 19.V.2011, Anlaş, Khachikov & Özgen. **Uşak:** 3 exs., Eşme, Kısık 2 km NE, Gediz river bank, 470 m, 38°38'06"N, 28°57'19"E, 23.IV.2010, leg. Anlaş.

Distribution in Turkey: Erzurum, Mersin (Anlaş, 2009; Kesdek et al., 2009).

***Philonthus rufimanus* (Heer, 1839)**

Material examined: **Bursa:** 3 exs., Uludağ, Aynalı Gölü, 28.VIII.2007, leg. Koç. **Elazığ:** 2 exs., Doğukent, 1080 m, 38°40'50"N, 39°15'42"E, 27.V.2010 and 01.VIII.2010, leg. Özgen; 1 ex., Gümüşkavak, 02.VIII.2010, leg. Özgen; 2 exs., Keban, Ulupınar 2 km NW, 1009 m, 38°43'58"N, 38°49'46"E, 20.V.2011, Anlaş, Khachikov & Özgen. **Gümüşhane:** 1 ex., Manastır 3 km N, Cehennem valley, 1867 m, 40°31'56"N, 39°36'16"E, 15.V.2011, Anlaş, Khachikov & Özgen. **Kahramanmaraş:** 2 exs., Andırın, Boztopraklı, ca. 700 m, ca. 37°31'N, 36°22'E, 23.VI.2007, leg. Yağmur. **Kilis:** 3 exs., Musabeyli, Akbayır, 08.VII.2006, leg. Anlaş. **Kütahya:** 4 exs. Simav 10 km NW, near Simav Lake, 720 m, 39°10'32"N, 28°55'21"E, 24.IV.2010, leg. Anlaş; 2 exs., Simav 5 km W, Yeşilköy 1 km N, 24.IV.2010, leg. Anlaş. **Malatya:** 2 exs., Arapgir 3 km NE, 912 m, 39°04'13"N, 38°30'34"E, 14.IX.2007, leg. Anlaş. **Manisa:** 3 exs., Soma, Yağcılı, 256 m, 39°20'07"N, 27°40'34"E, 08.IV.2007, leg. Anlaş; 1 ex., Selendi, Eskin, 29.VII.2007, leg. Anlaş; 4 exs., Demirci çayı, Bozköy 2 km E, 342 m, 38°52'55"N, 28°31'47"E, 24.IV.2010, leg. Anlaş. **Mardin:** 2 exs., Mazıdağı, Gürgöze, 950 m, 37°29'08"N, 40°31'38"E, 31.V.2010, leg. Özgen. **Muş:** 1 ex., Buğlan, 30.V.2011, leg. Khachikov & Kasatkin; 2 exs., Varto, 31.V.2011, leg. Khachikov & Kasatkin. **Siirt:** 7 exs., Baykan 4 km NE, ca. 770 m, 38°11'42"N, 41°49'03"E, 21.V.2010, leg. Anlaş. **Tunceli:** 1 ex., Ovacık, Munzur Gözeleri, 13.IX.2007, leg. Anlaş; 3 exs., Ovacık, Ağaçpınar 6 km E, Munzur river banks, 1197 m, 39°21'28"N, 39°15'51"E, 13.IX.2007, leg. Anlaş; 4 exs., Tunceli 5 km N, Anafatma, Munzur river bank, 920 m ca. 39°07'N, 39°30'E, 13.IX.2007, leg. Anlaş & Yağmur; 6 exs., Çemişgezek 1,5 km NW, Ormanyolu creek, 948 m, 39°04'06"N, 38°54'18"E, 14.IX.2007, leg. Anlaş; 3 exs., Çemişgezek, Payamdüzü, 955 m, 38°59'57"N, 39°02'28"E, 14.IX.2007, leg. Anlaş & Yağmur; 1 ex., 19.V.2011, Pülümür 3 km SE; 19.V.2011, leg. Anlaş; 2 exs., Uzuntarla 2 km E, leg. Anlaş. **Uşak:** 8 exs., Eşme, Kısık 2 km NE, Gediz river bank, 470 m, 38°38'06"N, 28°57'19"E, 23.IV.2010 and 29.V.2010, leg. Anlaş.

Distribution in Turkey: Aydın, Bayburt, Izmir, Kilis, Mersin? Manisa, Tunceli (Anlaş, 2009; Anlaş & Rose, 2009).

***Philonthus succicola* (Thomson, 1860)**

Material examined: **Kırklareli:** 1 ex., Demirköy, Dupnisa-Sarpdere road, 01.X.2009, leg. Kunt.

Distribution in Turkey: Istanbul (Apfelbeck, 1901; Anlaş, 2009).

***Philonthus svanetiensis* (Coiffait, 1974)**

Material examined: **Erzincan:** 1 ex., Üzümlü, Küçük Sarıkaya creek bank, 1713 m, 39°14'20"N, 39°50'02"E, 18.V.2011, Anlaş, Khachikov & Özgen.

Distribution in Turkey: Artvin, Erzurum, Rize (Assing, 2007; Anlaş, 2009).

***Philonthus tenuicornis* Mulsant & Rey, 1853**

Material examined: **Kırklareli:** 1 ex., İğneada-Demirköy, Siglioba, 03.X.2009, leg. Kunt. **Manisa:** 1 ex., Otoman, 330 m, 38°44'47"N, 27°10'34"E, 04.X.2008, leg. Anlaş.

Distribution in Turkey: Not cited (Smetana, 2004; Anlaş, 2009).

Remarks: This species is reported for the first time as exact locality records.

***Philonthus umbratilis* (Gravenhorst, 1802)**

Material examined: **Tunceli:** 1 ex., Tunceli 5 km N, Anafatma, Munzur river bank, 920 m ca. 39°07'N, 39°30'E, 13.IX.2007, leg. Anlaş & Yağmur.

Distribution in Turkey: Not cited (Smetana, 2004; Anlaş, 2009).

Remarks: This species is reported for the first time as exact locality records.

***Philonthus varians* (Paykull, 1789)**

Material examined: **Gümüşhane:** 1 ex., Torul 10 km SW, 1314 m, 40°30'34"N, 39°17'16"E, 14.V.2011, Anlaş, Khachikov & Özgen.

Distribution in Turkey: Mersin, Sinop (Peyron, 1858; Anlaş, 2009; Assing, 2010).

ACKNOWLEDGEMENTS

Authors would like to thank our colleagues for making their staphylinid by-catches to us.

LITERATURE CITED

- Anlaş, S.** 2007. The present situation of the Staphylinidae fauna of Turkey (Coleoptera). Linzer biologische Beiträge, 39 (1): 5-9.
- Anlaş, S.** 2009. Distributional checklist of the Staphylinidae (Coleoptera) of Turkey, with new and additional records. Linzer biologische Beiträge, 41: 215-342.
- Anlaş, S. & Rose, A.** 2009. Some additional notes about Staphylininae (Coleoptera: Staphylinidae) fauna of Turkey. Munis Entomology & Zoology, 4 (2): 327-333.
- Apfelbeck, V.** 1901. Bericht über eine Entomologische Forschungsreise nach der Türkei und Griechenland im Jahre 1900. Wiss Mitt Bosnisch-Herzegowin Landesmus, 8: 447-469 (in German).
- Assing, V.** 2007. On the Xantholinini of Turkey and adjacent regions (Coleoptera: Staphylinidae, Staphylininae). Zootaxa, 1474: 1-54.
- Assing, V.** 2010. On the Staphylinidae of Turkey. VII. Five new species and additional records (Coleoptera: Staphylinidae). Koleopterologische Rundschau, 80: 71-102.
- Assing, V.** 2013. On the Staphylinidae (Coleoptera) of Turkey IX. Five new species, a new synonymy, and additional records. Stuttgarter Beiträge zur Naturkunde A, Neue Serie, 6: 103-125.
- Fauvel, A.** 1874. Faune Gallo-Rhénane ou descriptions des insectes qui habitent la France, la Belgique, la Hollande, le Luxembourg, les provinces Rhénanes et la Valais avec tableaux synoptiques et planches gravées. Bull. Soc. Linn. Normandie, 8: 167-340 (in French).
- Herman, L. H.** 2001. Catalog of the Staphylinidae (Insecta: Coleoptera). 1758 to the end of the second millennium. V. Staphylinine Group (Part 2). Staphylininae: Diochini, Maerothiini, Othiini, Platyprosopini, Staphylinini (Amblyopinina, Anisolinina, Hyptomina, Philonthina). Bulletin of the American Museum of Natural History, 265: 2441-3020.
- Japoshvili, G. & Anlaş, S.** 2011. Notes on the Family Staphylinidae (Coleoptera) Collected by Pitfall Traps in Gölcük Natural Park, Isparta Province of Turkey. J. Entomol. Res. Soc., 13: 41-48.
- Özdemir, S. & Sert, O.** 2009. Determination of Coleoptera fauna on carcasses in Ankara province, Turkey. Forensic Science International, 183: 24-32.
- Özgen, İ. & Anlaş, S.** 2010. A cow dung investigation on Staphylinidae (Coleoptera) with a new record from Turkey. Munis Entomology Zoology, 5 (2): 642-645.
- Özgen, İ., Anlaş, S. & Eren, S.** 2010. Contribution to the knowledge of Staphylinidae (Coleoptera) Fauna of Cotton and Pistachio Fields in Southeastern Anatolia. Anadolu Doğa Bilimleri Dergisi, 1 (1): 20-26.
- Kesdek, M., Yıldırım, E. & Anlaş, S.** 2009. Contribution to the knowledge of Staphylinidae (Coleoptera) fauna of Turkey. Munis Entomology & Zoology, 4 (2): 355-364.
- Newton, A. F.** 2007. Documenting biodiversity: how well are we doing in Staphyliniformia (Coleoptera) Entomological Society of America poster presentation. 2007; Do471. Available (ESA members only) at http://esa.confex.com/esa/2007/techprogram/paper_32168.htm.
- Peyron, E.** 1858. Catalogue des Coléoptères des environs de Tarsous (Caramanie), avec la description des espèces nouvelles. Ann. Soc. Entomol. Fr., 3: 353-434 (in French).
- Schilhammer, H.** 1998. Revision of the East Palaearctic and Oriental species of *Philonthus* Stephens - Part 1. The *cyanipennis* group (Coleoptera: Staphylinidae, Staphylininae). Koleopterologische Rundschau, 68: 101-118.
- Smetana, A.** 1954. V ý sledky zoologické expedice Národního musea v Praze do Turecka. 17. Coleoptera VI. Staphylinidae (genera *Philonthus* Curt., *Gabrieus* Steph.). Acta Entomologica Musei Nationalis Pragae, 29 (439): 177-180.
- Smetana, A.** 2004. Family Staphylinidae (except subfamilies Pselaphinae and Scaphidiinae). In: Löbl I. & A. Smetana (eds), Catalogue of Palaearctic Coleoptera. Volume 2. Hydrophiloidea, Histeroidea, Staphyliniidea. Apollo Books, Stenstrup: 237-698.