

The Effects of Negative Reinforcement on Increasing Patient Adherence to Appointments at King Abdullah University Hospital in Jordan

INQUIRY: The Journal of Health Care
Organization, Provision, and Financing
Volume 53: 1–9
© The Author(s) 2016
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/0046958016660411
inq.sagepub.com

Mohammad Alyahya, MSc, PhD¹, Heba H. Hijazi, MSc, PhD¹,
and Farid T. Nusairat, MSc, PhD¹

Abstract

Appointment nonadherence is a health behavior that represents a burden to health care systems. On March 1, 2015, a new negative reinforcement intervention involving “service fees” for a visit without appointment was implemented at King Abdullah University Hospital in Jordan. To evaluate the effect of this intervention in improving patient adherence to medical appointment, a retrospective preintervention and postintervention analysis was used, including all patients ($n = 65\,535$) who had scheduled appointments at 39 outpatient clinics. A repeated-measures analysis of variance was first performed. Then, a multivariate linear regression model was used to identify factors that might predict individuals who are likely to attend or miss their appointments and those who have a greater tendency to visit the hospital with or without appointments. Although the average percentage of appointments attended was more than missed preintervention and postintervention, the decrease in percentage of missed appointments was more pronounced postintervention. Also, the average percentage of visits without appointments was less than visits with appointments in both times, but the decrease in the percentage of visits without appointments was more prominent after. The regression analysis revealed that younger, married and male patients were more likely to miss their appointment before and after the intervention. Also, younger patients had a tendency to attend without appointments. Conversely, patients with the lower copayment rate had a tendency to adhere to appointment times. In conclusion, negative reinforcement interventions could improve patient appointment adherence rates. Accordingly, interventions designed that consider evidence and are theory-based are needed to change patient behavior.

Keywords

appointment adherence, no-show, negative reinforcement, patient behavior, hospital

Introduction

Over the last years, there has been revived interest in adherence to medical appointments; this was strongly influenced by the high prevalence of no-show behavior and its subsequent impact.^{1,2} No-show prevalence rates vary across health care settings and populations. In the structured review conducted by Turkcan et al,¹ the mean no-show prevalence across the 62 studies included was 23.8%, with the highest rate occurring in North America (27.1%) and the lowest (14.9%) in Europe, whereas the rate was 24.3% for studies undertaken in Asia.¹

No-show behavior affects all parties involved in the delivery of health care, including patients, families, providers, and insurance agencies. For patients, missed appointments result in poor treatment adherence, disruption of multidisciplinary health care, and difficulty in patient-provider relationship.³ They are also associated with increased risk of

rehospitalization,^{4,5} poor control of chronic conditions,^{3,5} and can affect the health of other patients who lose the opportunity to receive timely care.⁶ For health care providers, missed appointments reduce the chance of junior physicians to learn from different cases,⁷ and could lead to staff underutilization, lower productivity, and increased costs.^{2,8–11}

Several studies were conducted to identify the risk factors associated with no-show behavior and characteristics of those

¹Jordan University of Science and Technology, Irbid, Jordan

Received 28 April 2016; revised June 26 2016; revised manuscript accepted 26 June 2016

Corresponding Author:

Mohammad Alyahya, Department of Health Management and Policy, Faculty of Medicine, Jordan University of Science and Technology, P.O. Box 3030, Irbid 22110, Jordan.
Email: Msalyahya@just.edu.jo



Creative Commons Non Commercial CC-BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 3.0 License (<http://www.creativecommons.org/licenses/by-nc/3.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

patients who are likely to miss their appointments. Many of these studies reached the same conclusion that being young, from minority racial or ethnic groups, or single, all significantly increased the likelihood of missing appointments.¹²⁻¹⁹ Other possible predictors were insurance status, gender, lower income, and day and time of appointment.^{18,19} A wide range of reasons why patients do not adhere to their medical appointments have also been extensively studied.

The most commonly self-reported reason for failure to attend was that patients “forgot” their scheduled medical appointments. This explains why reminding interventions (ie, phone calls, mails, and SMS) have received a substantial amount of attention, though reminders do not necessarily change patient behavior. Other identified reasons included lack of transportation, competing priorities with work time, family problems, difficulties with appointment scheduling systems, and patient’s health status such as feeling better or feeling too ill.²⁰⁻²²

To improve appointment adherence, several interventions have been utilized. Common interventions include appointment reminders, cancellation policy, patient education, and using a financial incentive.^{6,7,23-26} Despite the availability of these interventions, no-show behavior remains a burden to health care systems. Various policies have also attempted to curb the negative consequences of no-show behavior. Such policies include open-access or same day scheduling, follow-up after a no-show appointment, and overbooking appointments.²⁷⁻³⁰ Although these interventions and policies have been shown to reduce no-show rates, they are costly and labor-intensive, and their applicability is questionable.^{13,31}

Overbooking, for instance, increases patient wait time and patient dissatisfaction, and might lead to work overload at specific times.^{24,32} It has also been found that open-access scheduling does not always result in a significant reduction in no-show rates.³³⁻³⁵ Moreover, it is impractical to impose financial penalties on those patients who did not attend their scheduled appointment, as well as ethical and legal issue. Insurance agencies such as Medicaid in the United States do not permit health care providers to charge no-show patients.^{2,20}

New Intervention

Recently, on March 1, 2015, a new intervention was launched at King Abdullah University Hospital (KAUH), which is affiliated with the Jordan University of Science and Technology (JUST). Instead of using financial penalties, this intervention charges 15 Jordanian Dinars (US\$21) for a visit without an appointment. If a patient misses a scheduled appointment, he or she has to wait a long time to reschedule a new appointment, and may search for other ways to be seen by doctors. Many patients try to use their social relationships to be seen, which places a heavy burden on hospital resources. Unscheduled appointments are considered overbooking, and negatively affect those patients who already have a

scheduled appointment by increasing their wait time. Unscheduled patients also burden doctors by increasing the number of patients in the clinics, which causes doctors to have to rearrange patient time slots for those who have adhered to their appointments. The new policy applies to all outpatient services, irrespective of the nature of the visit. However, individual employees at KAUH and JUST, as well as their dependents, are exempt from the policy.

This intervention adopted by KAUH uses negative reinforcement to change patient behavior. In operant conditioning learning theory, negative reinforcement is defined as the removal, reduction, or escape of an unpleasant stimulus contingent upon a behavior, which results in an increased future rate of responding (behavior frequency).^{36,37} Negative reinforcement and punishment are often confused, because both involve the removal of an aversive stimulus. However, they differ from each other in how response is triggered and the effect on governing individual behavior.^{36,38} First, negative reinforcement entails avoiding a negative result when a desired behavior occurs, whereas punishment implies applying a negative consequence when undesired behavior is exhibited.^{36,39} Hence, by attending scheduled appointments, patients escape the negative consequences of a visit without an appointment. This in turn would decrease the appointment no-show rate. Second, negative reinforcement enhances the frequency of desired behavior, while whereas punishment reduces undesired target behavior.^{36,39,40} Instead of punishing those who do not attend their appointments, the intervention was introduced to increase the attendance rate.

Accordingly, the primary objective of this study is to identify whether the implementation of “service fees” for outpatient visits without a previously scheduled appointment as a negative reinforcer can improve patient attendance at scheduled appointments with the hospital. The study also aims to identify the main predictors of no-show behavior and visits without appointment.

Methods

Study Design and Data Collection

This is a preintervention and postintervention study⁴¹ of all patients who had scheduled appointments from September 1, 2014, till September 1, 2015—6 months before and 6 months after—at KAUH. KAUH is a multispecialty teaching hospital with 527 beds; it provides a wide range of health services and is considered as a referral hospital for 4 cities in the North of Jordan. Services provided by KAUH are in high demand, especially the outpatient clinics, which are renowned for their competent medical staff and availability of advanced technology.

The data of the study were extracted retrospectively from the database of the 39 outpatient specialist clinics at the hospital. For each patient, the following data were obtained:

Table 1. General Comparison Before and After the Intervention.

	Before		After	
	Mean of %	SD	Mean of %	SD
Attended appointments	54.87	41.79	58.72	39.36
Missed appointments	45.12	41.79	41.27	39.36
Visits with appointments	69.38	38.07	81.21	34.03
Visits without appointments	30.61	38.07	18.78	34.03

attended appointments 6 months before, missed appointments 6 months before, total appointments 6 months before, attended appointments 6 months after, missed appointments 6 months after, and total appointments 6 months after. Similarly, the data included numbers of total visits 6 months before, visits with appointments 6 months before, visits without appointments 6 months before, total visits 6 months after, visits with appointments 6 months after, and visits without appointments 6 months after. In addition, data on demographic variables and patient characteristics such as fee exemption status, patient age, gender, marital status, and copayment status were also collected.

Ethical approval. This study received ethical approval from the institutional review board (IRB) committee at KAUH and JUST (No. 20160098).

Data Analysis

Descriptive statistics are reported for appointments showed up and visits with appointments before and after the intervention. As the data before and after the intervention were observed on the same patients, a repeated-measures analysis of variance (ANOVA) was conducted to examine the effect of the intervention on appointment adherence and visits without appointments. The significance of collected factors and covariates on the probability of attending appointments and the probability of visits with appointments was tested. Factors considered included intervention, exemption status, gender, marital status, and copayment. Age of the patient was treated as a covariate.

To identify factors that may predict individuals who are likely to miss their appointments and those who have a greater tendency to visit the hospital without prescheduled appointments, a multivariate linear regression model was used to relate the probability of adherence to appointments and probability of visits with appointments to the significant factors found in the repeated-measures ANOVA. The analysis was performed using SPSS 20.0.

Results

A total of 65 535 patients who had scheduled appointments during the study period were identified. The mean age of

included patients was 36 (SD = 21) years of whom 37 487 were females (57.2%), 36 541 were married (55.8%), and 7431 of them had exemption (11.3%). Table 1 shows that the mean percentage of missed appointments decreased from 45.12% during the preintervention period to 41.27% during the postintervention period. This difference reflected a reduction of 8.53% and was statistically significant ($P < .001$). Interestingly, the mean percentage of visits without appointments reduced by 38.64% ($P < .001$), from 30.61% to 18.78% postintervention.

Effect of the Intervention on Appointment Adherence and Visits With Appointments

Based on the patients' characteristics, means and SDs for percentages of attended appointments and percentages of visits with appointments before and after the intervention are shown in Table 2. A repeated-measures ANOVA with intervention (before and after the intervention) as within-subjects factor and gender (male, female), marital status (single, married), exemption status (yes [with], no [without]), and copayment group (groups 1 through 4) as between-subjects along with patient age as a covariate was conducted. Two response variables were considered; the first was the probability of a show-up in an appointment calculated as the percentage of appointments the patient showed up of the total appointments (Table 3). The second response variable was the probability of a visit with appointment obtained as the percentage of visits with appointments of the total visits (Table 4).

For the probability of a show-up in an appointment, ANOVA results revealed that intervention ($P < .001$), age ($P = .018$), gender ($P < .001$), and copayment ($P < .001$) were the significant factors along with the 2-factor interactions: Intervention \times Age ($P < .001$), Intervention \times Copayment ($P = .002$), Gender \times Copayment ($P < .001$), and Marital status \times Copayment ($P = .004$). Exemption status was a marginally significant factor with a P value of .093.

For the probability of visits with appointment, the effect of intervention ($P < .001$), age ($P < .001$), gender ($P < .001$), marital status ($P < .001$), and copayment ($P < .001$) were all significant. The following 2-factor interactions were also significant: Intervention \times Gender ($P = .051$), Intervention \times Copayment ($P < .001$), Gender \times Copayment ($P = .056$),

Table 2. Percentages of Patients' Characteristics Who Attended Their Appointments and Had Visits With Appointments Before and After the Intervention.

Characteristic:	1				2			
	Before		After		Before		After	
	Mean of %	SD	Mean of %	SD	Mean of %	SD	Mean of %	SD
Gender								
Female	55.55	4.14	58.50	3.91	70.50	3.73	82.31	3.31
Male	53.95	4.22	56.61	3.97	67.84	3.89	79.66	3.52
Marital status								
Married	56.14	4.08	59.52	3.78	68.24	3.77	80.65	3.37
Single	53.10	4.30	54.73	4.03	71.12	3.85	82.22	3.44
Exemption								
Yes	53.24	4.16	57.93	3.93	66.75	3.90	76.59	3.74
No	55.08	4.18	56.07	3.94	69.72	3.79	81.82	3.35
Copayment								
Group 1 (copayment = 0%)	58.11	3.73	60.40	3.48	74.01	3.65	87.32	2.88
Group 2 (copayment = 6.5%)	55.29	4.19	58.61	3.99	58.01	3.67	78.86	3.34
Group 3 (copayment = 20%)	49.95	4.33	52.37	4.13	57.50	4.20	55.03	4.42
Group 4 (copayment = 100%)	53.48	4.46	50.46	4.13	75.16	3.76	80.46	3.49

Marital status \times Copayment ($P = .003$), and Exemption \times Copayment (marginally significant, $P = .062$).

Predictors of Appointment Adherence and Visits With/Without Appointments 6 Months Before and 6 Months After

To mathematically identify the size of the effect for the significant factors found in the ANOVA, multiple linear regression was utilized to build models that relate probability of show-up in appointments and probability of visits with appointments to their predictors. Table 5 shows the estimated regression parameters along with their 95% confidence intervals (CIs). Based on the parameter estimates, we found that female patients had a higher probability of adherence to appointments than males, with increased probability after the intervention. In contrast, married patients were more likely to miss their appointments than single patients. Married patients have shown a slight improvement in the adherence after the intervention. Among copayment groups, group 3 patients had the lowest probability of adherence compared with other groups (before and after). This means that patients within the lower range of copayment rate had a tendency to attend their appointments. However, all groups have shown a significant increased probability of adherence after the intervention. Patients with no exemption have shown a slightly reduced probability of adherence to appointments after the intervention compared with that before the intervention. The results also indicated that older patients had a tendency to attend appointments, particularly after the intervention.

In concordance with the above findings, female patients had a higher probability of visits with appointments than

male patients. Also, the probability of visits with appointments for married patients has improved after the intervention. Patients with the lower copayment rates were less likely to attend without previous appointment, whereas younger patients were much more likely to attend without appointments. Finally, a substantial improvement in the probability of visits with appointment has been achieved after the intervention for patients with no exemption.

Discussion

Appointment adherence is a health behavior that remains a challenge for health care systems because of its detrimental effects on patient treatment and outcomes.³⁻⁵ However, appointment adherence has received little attention as a behavioral issue, compared with, for example, treatment adherence, or immunization uptake.⁴²

This study examined whether an intervention implementing fees for unscheduled visits would result in improving appointment adherence. Although reducing visits without appointments has been an objective of the intervention, the ultimate aim is to increase the rate of appointment adherence. The study found that the mean percentage of visits without appointments decreased significantly from 30.61% to 18.78%, and that the mean percentage of the no-show rate dropped from 45.12% to 41.27% after the introduction of the intervention. In addition, our results demonstrated that both visits with appointments and attended appointments have improved after introducing the new intervention.

Although there are a number of other interventions that have been shown to decrease the rates of no-show behavior, these interventions may have limited impact and lead to

Table 3. Repeated-Measures ANOVA Within and Between Subjects for Shown Appointments.

Within-subjects contrasts					
Source	Sum of squares	df	Mean square	F	Significance
Intervention	78781.755	1	78781.755	62.936	.000
Intervention × Age	22576.220	1	22576.220	18.035	.000
Intervention × Gender	1060.922	1	1060.922	0.848	.357
Intervention × Marital Status	957.667	1	957.667	0.765	.382
Intervention × Exemption	340.389	1	340.389	0.272	.602
Intervention × Copayment	18933.597	3	6311.199	5.042	.002
Intervention × Gender × Marital Status	1381.632	1	1381.632	1.104	.293
Intervention × Gender × Exemption	113.157	1	113.157	0.090	.764
Intervention × Gender × Copayment	3999.901	3	1333.300	1.065	.362
Intervention × Marital Status × Exemption	1975.429	1	1975.429	1.578	.209
Intervention × Marital Status × Copayment	228.858	3	76.286	0.061	.980
Intervention × Exemption × Copayment	216.586	3	72.195	0.058	.982
Intervention × Gender × Marital Status × Exemption	2471.314	1	2471.314	1.974	.160
Intervention × Gender × Marital Status × Copayment	52.689	3	17.563	0.014	.998
Intervention × Gender × Exemption × Copayment	1937.919	3	645.973	0.516	.671
Intervention × Marital Status × Exemption × Copayment	1177.929	3	392.643	0.314	.816
Intervention × Gender × Marital Status × Exemption × Copayment	1333.630	3	444.543	0.355	.785
Tests of between-subjects effects					
Source	Sum of squares	df	Mean square	F	Significance
Intercept	13781447.711	1	13781447.711	8220.322	.000
Age	9392.568	1	9392.568	5.602	.018
Gender	20335.015	1	20335.015	12.129	.000
Marital status	366.166	1	366.166	0.218	.640
Exemption	4742.965	1	4742.965	2.829	.093
Copayment	95090.504	3	31696.835	18.906	.000
Gender × Marital Status	3904.447	1	3904.447	2.329	.127
Gender × Exemption	47.545	1	47.545	0.028	.866
Gender × Copayment	31706.037	3	10568.679	6.304	.000
Marital Status × Exemption	3758.568	1	3758.568	2.242	.134
Marital Status × Copayment	21923.179	3	7307.726	4.359	.004
Exemption × Copayment	6363.601	3	2121.200	1.265	.284
Gender × Marital Status × Exemption	1060.573	1	1060.573	0.633	.426
Gender × Marital Status × Copayment	6486.168	3	2162.056	1.290	.276
Gender × Exemption × Copayment	7477.574	3	2492.525	1.487	.216
Marital Status × Exemption × Copayment	2838.844	3	946.281	0.564	.638
Gender × Marital Status × Exemption × Copayment	3700.033	3	1233.344	0.736	.531

Note. ANOVA = analysis of variance.

dysfunctional consequences. For instance, overbooking and open-access policies might increase the risk of overloading the schedules of health care providers, which requires careful observation and management, and accurate prediction of demand.^{24,43} In addition, many previously discussed interventions have only been used in research studies, and have not necessarily been applied in health care facilities, or used to motivate individuals to change their behavior.^{24,26,44,45}

Nonetheless, the effectiveness of using financial incentives (rewards/penalties) in changing individual behavior has been theoretically substantiated and empirically supported in different health care settings.^{46,47} In a recent study, Kubanek et al³⁸ tried to address whether Thorndike's law of effect is symmetric or qualitatively distinct in regard to monetary reinforcement and punishment. They found that reinforcement and punishment play a distinct role in guiding individual

Table 4. Repeated-Measures ANOVA Within and Between Subjects for Visits With Appointments.

Tests of within-subjects contrasts					
Source	Sum of squares	df	Mean square	F	Significance
Intervention	140 297.388	1	140 297.388	171.362	.000
Intervention × Age	2676.046	1	2676.046	3.269	.071
Intervention × Gender	3110.976	1	3110.976	3.800	.051
Intervention × Marital Status	402.770	1	402.770	0.492	.483
Intervention × Exemption	593.961	1	593.961	0.725	.394
Intervention × Copayment	171 628.091	3	57 209.364	69.877	.000
Intervention × Gender × Marital Status	1085.078	1	1085.078	1.325	.250
Intervention × Gender × Exemption	1503.341	1	1503.341	1.836	.175
Intervention × Gender × Copayment	4546.986	3	1515.662	1.851	.135
Intervention × Marital Status × Exemption	1174.211	1	1174.211	1.434	.231
Intervention × Marital Status × Copayment	1555.539	3	518.513	0.633	.593
Intervention × Exemption × Copayment	2020.905	3	673.635	0.823	.481
Intervention × Gender × Marital Status × Exemption	204.896	1	204.896	0.250	.617
Intervention × Gender × Marital Status × Copayment	333.233	3	111.078	0.136	.939
Intervention × Gender × Exemption × Copayment	2240.589	3	746.863	0.912	.434
Intervention × Marital Status × Exemption × Copayment	1988.503	3	662.834	0.810	.488
Intervention × Gender × Marital Status × Exemption × Copayment	1507.626	3	502.542	0.614	.606
Tests of between-subjects effects					
Source	Sum of squares	df	Mean square	F	Significance
Intercept	21 752 809.608	1	21 752 809.608	14313.039	.000
Age	646 723.813	1	646 723.813	425.535	.000
Gender	23 997.730	1	23 997.730	15.790	.000
Marital status	21 529.054	1	21 529.054	14.166	.000
Exemption	82.637	1	82.637	0.054	.816
Copayment	1 723 224.589	3	574 408.196	377.952	.000
Gender × Marital Status	1525.399	1	1525.399	1.004	.316
Gender × Exemption	14.953	1	14.953	0.010	.921
Gender × Copayment	11 469.609	3	3823.203	2.516	.056
Marital Status × Exemption	6.630	1	6.630	0.004	.947
Marital Status × Copayment	21 191.658	3	7063.886	4.648	.003
Exemption × Copayment	11 127.199	3	3709.066	2.441	.062
Gender × Marital Status × Exemption	803.735	1	803.735	0.529	.467
Gender × Marital Status × Copayment	2263.996	3	754.665	0.497	.685
Gender × Exemption × Copayment	1148.201	3	382.734	0.252	.860
Marital Status × Exemption × Copayment	10343.410	3	3447.803	2.269	.078
Gender × Marital Status × Exemption × Copayment	8310.197	3	2770.066	1.823	.141

Note. ANOVA = analysis of variance.

behavior. Punitive policies might be effective in reducing the no-show rate but are difficult to implement and can affect access to health care service, particularly among those patients with limited income.¹⁷ Also, it might be unfair to charge those who had inevitable reason for not attending.^{2,13,48} In their qualitative study, Martin et al²⁰ argue that imposing financial penalties would have some effect in reducing missed

appointments but would not be significant; furthermore, it might be not acceptable to providers to fine nonattending patients. In addition, such policies might encourage patients to cancel or reschedule their appointments.⁴⁹

Instead of fining those patients who missed their scheduled appointment, KAUH introduced the “service fees” as a negative reinforce to those who want visits without appointments.

Table 5. Multiple Linear Regression Estimates for Predicting Probability of Shown Appointments and Probability of Visits With Appointments Before and After the Intervention.

Factor	Probability of show-up in appointments						Probability of visits with appointments					
	Before intervention			After intervention			Before intervention			After intervention		
	Coefficient	95% CI		Coefficient	95% CI		Coefficient	95% CI		Coefficient	95% CI	
		Lower	Upper		Lower	Upper		Lower	Upper		Lower	Upper
Intercept	60.230	56.669	63.792	44.681	41.081	48.281	81.285	73.927	88.643	88.906	82.411	95.400
Gender = female	4.397	0.639	8.155	4.980	1.182	8.779	6.449	2.173	10.725	1.018	-2.756	4.792
Marital = married	-3.634	-7.471	0.203	-2.519	-6.397	1.360	3.985	-0.330	8.300	7.905	4.096	11.714
Exemption = no	1.904	0.401	3.528	1.735	0.054	3.215	-6.080	-13.061	0.902	-0.733	-6.895	5.429
Copayment = group 1	1.155	-2.289	4.598	9.085	5.604	12.566	-2.811	-10.547	4.925	5.854	-0.974	12.682
Copayment = group 2	-1.129	-4.980	2.723	8.893	4.999	12.786	-14.673	-22.892	-6.455	2.348	-4.906	9.601
Copayment = group 3	-4.945	-9.256	-0.635	1.811	-2.546	6.169	-27.104	-35.472	-18.737	-33.469	-40.854	-26.084
Age	2.019	0.051	4.014	2.180	0.047	4.112	1.258	-0.291	1.924	2.300	0.329	3.870
Gender = Female × Copayment = group 1	-0.151	-4.473	4.171	-0.267	-4.636	4.102	-1.214	-5.966	3.537	5.245	1.051	9.439
Gender = Female × Copayment = group 2	-6.597	-11.612	-1.583	-2.439	-7.508	2.630	-4.012	-9.322	1.297	2.067	-2.619	6.754
Gender = Female × Copayment = group 3	-4.043	-8.006	-0.080	-4.923	-8.929	-0.917	-4.026	-8.514	.461	0.660	-3.301	4.621
Marital = Married × Copayment = group 1	5.183	0.835	9.531	1.882	-2.514	6.277	-1.130	-5.889	3.628	-8.832	-13.032	-4.632
Marital = Married × Copayment = group 2	8.650	3.723	13.577	7.106	2.126	12.087	2.577	-2.641	7.796	-0.594	-5.200	4.012
Marital = Married × Copayment = group 3	4.296	0.367	8.225	3.796	-0.176	7.768	-1.299	-5.740	3.141	-3.739	-7.659	0.180
Exemption = No × Copayment = group 1							4.339	-3.439	12.117	-0.447	-7.312	6.419
Exemption = No × Copayment = group 2							8.096	0.232	15.960	3.480	-3.461	10.421
Exemption = No × Copayment = group 3							7.543	0.208	14.878	1.659	-4.815	8.133

Note. CI = confidence interval.

Thus, if the patient attends his or her scheduled appointment, he or she will not need to pay for a visit without appointment. In addition, negative reinforcement enhances the frequency of desired behavior, whereas punishment reduces undesired target behavior.^{36,39,40} In this regard, the main reason for the intervention is increasing the rate of appointment attendance rather than punishing those who do not adhere to their scheduled appointments.

Identifying the main characteristics associated with appointment adherence can also inform the development of interventions to reduce the occurrence of no-show behavior.¹⁹ Our results confirm what has been previously found that younger patients are more likely to no-show than older patients.¹⁷⁻¹⁹ Older adults have more chronic conditions and long-term illnesses and missing scheduled appointments could adversely affect their health. They are more attentive to their health, and have enough free time compared with young adults.²⁶ Moreover, and in accordance with several previous studies,¹⁴⁻¹⁶ this study showed that male patients were more likely to no-show to their appointments before and after the intervention, though this has not been found

consistently.^{1,50,51} Some previous studies also indicated that being married is associated with a low no-show rate,^{1,17} while we found that single patients were more likely to attend their appointments. Singles usually have less family commitments and it is easier for them to manage their time. Interestingly, having full insurance coverage was found to be significantly associated with a tendency to attend appointments. A possible explanation is that most patients with 0% copayment had government insurance and were referred from other government hospitals, or they had full insurance coverage from the Royal Court or from other agencies. However, in both cases, full coverage is usually time limited. So, such patients try to enjoy the greatest possible benefits, especially as most of the patients are from lower income families.

Finally, there are other patient characteristics and predisposing factors that have been identified in previous studies as potential influences on patient adherence to prescheduled appointments including health status, income, and race and ethnicity.^{1,52} However, as most inhabitants in Jordan are Arabs, Jordanian ethnicity and culture are characterized by uniformity and there are no racial and ethnic differences in

Jordan. Other factors such as health status (chronic vs. acute) could also influence adherence to appointment. Chronic disease patients might be more apt to attend their appointments than patients with acute illnesses, because patients with chronic diseases usually have a good relationship with their caregivers, and they are more compliant with their treatment plan.^{1,17} In addition, the severity of health conditions might be associated with adherence to appointments; for instance, it has been shown that more ill patients are less likely to miss their appointments.^{1,17} Therefore, it is worthy to consider and examine the effect of these factors in future works to predict and improve appointment adherence.

Conclusion

This study demonstrates the ability of negative reinforcement to modify patient behavior and increase patient appointment adherence rates. Appointment adherence enhances the efficacy of health services. Besides using the scheduled appointment to provide adequate and timely patient care, health care settings maximize the utilization of existing resources. The study also highlights that developing an effective and socially appropriate intervention that could improve appointment attendance requires a clear explanation of the impact of such interventions from theoretical perspectives. Hence, health managers and policy makers need to consider evidence and theory-based interventions that go beyond merely the managerial practical issues. Further studies should focus on the empirical link between theory and practice in governing patient adherence behavior. Finally, we expect that visits without appointments and no-show rates could be further reduced than our reported results if the intervention is continued, as the current study examined impact only in the first 6 months of implementation.

Acknowledgments

We would like to thank Deanship of Research at Jordan University of Science and Technology for their keen help and support. Also, we are very grateful to Anas Matalkah, Manager of Information Systems Department at King Abdullah University Hospital (KAUH), for his technical support in data collection.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

1. Turkcan A, Nuti L, DeLaurentis P-C, et al. No-show modeling for adult ambulatory clinics. In: Denton B, ed. *Handbook*

of Healthcare Operations Management: Methods and Applications. New York, NY: Springer; 2013:251-288.

2. Guzek LM, Gentry SD, Golomb MR. The estimated cost of "no-shows" in an academic pediatric neurology clinic. *Pediatr Neurol*. 2015;52(2):198-201.
3. Jyun-You L, Chia-Fen M, Chao-Yu H. Medical appointment no-shows associated with poor glycaemic control among Taiwanese aborigines. *Aust J Rural Health*. 2012;20(6):312-317.
4. Nuti L, Turkcan A, Lawley MA, Zhang L, Sands L, McComb S. The impact of interventions on appointment and clinical outcomes for individuals with diabetes: a systematic review. *BMC Health Serv Res*. 2015;15(1):1-54.
5. Hwang AS, Atlas SJ, Cronin P, et al. Appointment "no-shows" are an independent predictor of subsequent quality of care and resource utilization outcomes. *J Gen Intern Med*. 2015;30(10):1426-1433. doi:1410.1007/s11606-11015-13252-11603.
6. Bech M. The economics of non-attendance and the expected effect of charging a fine on non-attendees. *Health Policy*. 2005;74(2):181-191.
7. Guse CE, Richardson L, Carle M, Schmidt K. The Effect of exit-interview patient education on no-show rates at a family practice residency clinic. *J Am Board Fam Pract*. 2003;16(5):399-404.
8. LaGanga LR, Lawrence SR. Clinic overbooking to improve patient access and increase provider productivity. *Decision Sciences*. 2007;38(2):251-276.
9. Berg B, Murr M, Chermak D, et al. Estimating the cost of no-shows and evaluating the effects of mitigation strategies. *Med Decis Making*. 2013;33(8):976-985.
10. Norris JB, Kumar C, Chand S, Moskowitz H, Shade SA, Willis DR. An empirical investigation into factors affecting patient cancellations and no-shows at outpatient clinics. *Decis Support Syst*. 2014;57:428-443.
11. Huang Y, Hanauer DA. Patient no-show predictive model development using multiple data sources for an effective overbooking approach. *Appl Clin Inform*. 2014;5(3):836-860.
12. Shimotsu S, Roehrl A, McCarty M, et al. Increased likelihood of missed appointments ("no shows") for racial/ethnic minorities in a safety net health system. *J Prim Care Community Health*. 2016;7(1):38-40. doi:10.1177/2150131915599980.
13. Corfield L, Schizas A, Williams A, Noorani A. Non-attendance at the colorectal clinic: a prospective audit. *Ann R Coll Surg Engl*. 2008;90(5):377-380.
14. Campbell K, Millard A, McCartney G, McCullough S. Who is least likely to attend? an analysis of outpatient appointment DNA data in NHS Dumfries & Galloway. In: NHS Health Scotland, ed. <http://www.scotpho.org.uk/downloads/scotphoreports/scotpho150319-DNA-analysis-NHS-Dumfries-and-Galloway.pdf>. Published March 2015. Accessed March 15, 2016.
15. Campbell K, Millard A, McCartney G, McCullough S. Who is least likely to attend? an analysis of outpatient appointment DNA data in NHS Greater Glasgow & Clyde. In: NHS Health Scotland, ed. <http://www.scotpho.org.uk/downloads/scotphoreports/scotpho150319-DNA-analysis-NHS-Greater-Glasgow-and-Clyde.pdf>. Published March 2015. Accessed March 15, 2016.

16. Lindauer SJ, Powell JA, Leypoldt BC, Tufekci E, Shroff B. Influence of patient financial account status on orthodontic appointment attendance. *Angle Orthod.* 2009;79(4):755-758.
17. Daggy J, Lawley M, Willis D, et al. Using no-show modeling to improve clinic performance. *Health Informatics J.* 2010;16(4):246-259.
18. Miller AJ, Chae E, Peterson E, Ko AB. Predictors of repeated "no-showing" to clinic appointments. *Am J Otolaryngol.* 2015;36(3):411-414. doi:10.1016/j.amjoto.2015.01.017.
19. Kaplan-Lewis E, Percac-Lima S. No-show to primary care appointments: why patients do not come. *J Prim Care Community Health.* 2013;4(4):251-255.
20. Martin C, Perfect T, Mantle G. Non-attendance in primary care: the views of patients and practices on its causes, impact and solutions. *Fam Pract.* 2005;22(6):638-643.
21. Samuels RC, Ward VL, Melvin P, et al. Missed appointments: factors contributing to high no-show rates in an urban pediatrics primary care clinic. *Clin Pediatr (Phila).* 2015;54(10):976-982.
22. Geiger SL. *Nonattendance Rates and Barriers to Health Care in Outpatient Clinic Settings* [Walden dissertations and doctoral studies]. Minneapolis, MN: College of Health Sciences, Walden University; 2015.
23. da Costa TM, Salomao PL, Martha AS, Pisa IT, Sigulem D. The impact of short message service text messages sent as appointment reminders to patients' cell phones at outpatient clinics in Sao Paulo, Brazil. *Int J Med Inform.* 2010;79(1):65-70. doi:10.1016/j.ijmedinf.2009.09.001.
24. Huang Y, Zuniga P. Effective cancellation policy to reduce the negative impact of patient no-show. *J Oper Res Soc.* 2014;65(5):605-615.
25. van Dieren Q, Rijckmans MJN, Mathijssen JJP, Lobbestael J, Arntz AR. Reducing no-show behavior at a community mental health center. *J Community Psychol.* 2013;41(7):844-850.
26. Parikh A, Gupta K, Wilson AC, Fields K, Cosgrove NM, Kostis JB. The effectiveness of outpatient appointment reminder systems in reducing no-show rates. *Am J Med.* 2010;123(6):542-548.
27. Zacharias C, Pinedo M. Appointment scheduling with no-shows and overbooking. *Prod Oper Manage.* 2014;23(5):788-801.
28. Molfenter T. Reducing appointment no-shows: going from theory to practice. *Subst Use Misuse.* 2013;48(9):743-749.
29. O'Connor ME, Matthews BS, Gao D. Effect of open access scheduling on missed appointments, immunizations, and continuity of care for infant well-child care visits. *Arch Pediatr Adolesc Med.* 2006;160(9):889-893.
30. Zeber J, Pearson D, Smith D. Analysis of health appointment no-shows. *Prim Health Care.* 2009;19(2):25-29.
31. Stubbs ND, Geraci SA, Stephenson PL, Jones DB, Sanders S. Methods to reduce outpatient non-attendance. *Am J Med Sci.* 2012;344(3):211-219. doi:10.1097/MAJ.0b013e31824997c6.
32. Tsai P-FJ, Teng G-Y. A stochastic appointment scheduling system on multiple resources with dynamic call-in sequence and patient no-shows for an outpatient clinic. *Eur J Oper Res.* 2014;239(2):427-436.
33. Samorani M, LaGanga LR. Outpatient appointment scheduling given individual day-dependent no-show predictions. *Eur J Oper Res.* 2015;240(1):245-257.
34. Liu N, Ziya S, Kulkarni VG. Dynamic scheduling of outpatient appointments under patient no-shows and cancellations. *M&SOM.* 2010;12(2):347-364.
35. Mehrotra A, Keehl-Markowitz L, Ayanian JZ. Implementation of open access scheduling in primary care: a cautionary tale. *Ann Intern Med.* 2008;148(12):915-922.
36. George JM, Jones GR. *Understanding and Managing Organizational Behavior.* 6th ed. Upper Saddle River, NJ: Prentice Hall/Pearson; 2012.
37. Iwata BA. Negative reinforcement in applied behavior analysis: an emerging technology. *J Appl Behav Anal.* 1987;20(4):361-378.
38. Kubanek J, Snyder LH, Abrams RA. Reward and punishment act as distinct factors in guiding behavior. *Cognition.* 2015;139:154-167.
39. Borkowski N. Process theories of motivation. In: Borkowski N, ed. *Organizational Behavior, Theory, and Design in Health Care.* 1st ed. London, England: Jones & Bartlett Learning; 2009, pp. 127-142.
40. Arvey RD, Ivancevich JM. Punishment in organizations: a review, propositions, and research suggestions. *Acad Manage Rev.* 1980;5(1):123-132.
41. Thiese MS. Observational and interventional study design types; an overview. *Biochem Med.* 2014;24(2):199-210.
42. Cameron E. *A Mixed Methods Investigation of Parental Factors in Non-attendance at General Paediatric Hospital Outpatient Appointments* [doctoral thesis]. Birmingham, UK: Aston University; 2015.
43. Vijayan M. *No Shows: Effectiveness of Termination Policy and Review of Best Practices* [master's thesis in public health]. Dayton, OH: Wright State University; 2014.
44. Arora S, Burner E, Terp S, et al. Improving attendance at post-emergency department follow-up via automated text message appointment reminders: a randomized controlled trial. *Acad Emerg Med.* 2015;22(1):31-37.
45. Taylor NF, Bottrell J, Lawler K, Benjamin D. Mobile telephone short message service reminders can reduce nonattendance in physical therapy outpatient clinics: a randomized controlled trial. *Arch Phys Med Rehabil.* 2012;93(1):21-26.
46. Giles EL, Robalino S, McColl E, Snichotta FF, Adams J. The effectiveness of financial incentives for health behaviour change: systematic review and meta-analysis. *PLoS ONE.* 2014;9(3):e90347.
47. Sutherland K, Christianson JB, Leatherman S. Impact of targeted financial incentives on personal health behavior: a review of the literature. *Med Care Res Rev.* 2008;65(6) (suppl):36S-78S.
48. Ratcliffe A, Gilland W, Maruchek A. Revenue management for outpatient appointments: joint capacity control and overbooking with class-dependent no-shows. *Flex Serv Manuf J.* 2012;24(4):516-548.
49. Chariatte V, Michaud P, Berchtold A, Akre C, Suris J. Missed appointments in an adolescent outpatient clinic: descriptive analyses of consultations over eight years. *Swiss Med Wkly.* 2007;137(47/48):677-681.
50. Giunta D, Briatore A, Baum A, Luna D, Waisman G, de Quiros FGB. Factors associated with nonattendance at clinical medicine scheduled outpatient appointments in a university general hospital. *Patient Prefer Adherence.* 2013;7:1163-1170.
51. Chang JT, Sewell JL, Day LW. Prevalence and predictors of patient no-shows to outpatient endoscopic procedures scheduled with anesthesia. *BMC Gastroenterol.* 2015;15(1):123.
52. Kheirkhah P, Feng Q, Travis LM, Tavakoli-Tabasi S, Sharafkhaneh A. Prevalence, predictors and economic consequences of no-shows. *BMC Health Serv Res.* 2016;16(1):13.