

Serum 25-hydroxyvitamin D and the risk of cardiovascular disease: dose-response meta-analysis of prospective studies^{1–3}

Runhua Zhang,^{4–7} Bohong Li,⁴ Xiang Gao,⁸ Rui Tian,⁴ Yuesong Pan,^{4–7} Yong Jiang,^{4–7} Hongqiu Gu,^{4–7} Yilong Wang,^{4–7} Yongjun Wang,^{4–7} and Gaifen Liu^{4–7*}

⁴Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, China; ⁵China National Clinical Research Center for Neurological Diseases, Beijing, China; ⁶Center of Stroke, Beijing Institute for Brain Disorders, Beijing, China; ⁷Beijing Key Laboratory of Translational Medicine for Cerebrovascular Disease, Beijing, China; and ⁸Department of Nutritional Sciences, Pennsylvania State University, State College, PA

ABSTRACT

Background: During the past decade, an increasing number of prospective studies have focused on the association between vitamin D and cardiovascular disease (CVD). However, the evidence on the relation between serum 25-hydroxyvitamin D [25(OH)D] and the risk of overt CVD is inconclusive.

Objective: We performed a dose-response meta-analysis to summarize and prospectively quantify the RR of low serum 25(OH)D concentration and total CVD (events and mortality).

Design: We identified relevant studies by searching PubMed and EMBASE up to December 2015 and by hand-searching reference lists. Prospective studies based on the general population and reported RRs and 95% CIs were included. A random-effects model was used to calculate the pooled RRs. Nonlinear association was assessed by using restricted cubic spline analyses.

Results: A total of 34 publications with 180,667 participants were eligible for the meta-analysis. We included 32 publications (27 independent studies) for total CVD events and 17 publications (17 independent studies) for CVD mortality. We observed an inverse association between serum 25(OH)D and total CVD events and CVD mortality, and the pooled RRs per 10-ng/mL increment were 0.90 (95% CI: 0.86, 0.94) for total CVD events and 0.88 (95% CI: 0.80, 0.96) for CVD mortality. A nonlinear association was detected for total CVD events (P -nonlinear < 0.001) and CVD mortality (P -nonlinear = 0.022).

Conclusion: Serum 25(OH)D concentration was inversely associated with total CVD events and CVD mortality from the observed studies. *Am J Clin Nutr* doi: 10.3945/ajcn.116.140392.

Keywords: 25-hydroxyvitamin D, cardiovascular disease, meta-analysis, dose-response, CVD mortality

INTRODUCTION

Historically, vitamin D is well known for its crucial role in skeletal disease, including bone metabolism (1, 2), fractures (3, 4), and falls (5). In recent decades, a large and rapidly expanding literature has focused on the associations between vitamin D and hypertension (6), diabetes (7, 8), metabolic syndrome (9), cardiovascular disease (CVD)⁹ (10, 11), cancer (12, 13), and mortality (14, 15). The Institute of Medicine released advice on Recommended Dietary Allowances for achieving a 25-hydroxyvitamin D

[25(OH)D] concentration ≥ 20 ng/mL, which was based on bone health (16). Unlike for skeletal disease, the evidence for vitamin D and the risk of extraskeletal chronic disease has been inconclusive. CVD is the leading cause of deaths and disability (17), and the prevalence of vitamin D insufficiency is high in the general population (18). If there is a cause-and-effect relation found between low vitamin D status and incident CVD, this would be of considerable benefit to public health (19). To date, there have been few clinical trials specifically designed to explore this relation. The VITAL (Vitamin D and Omega-3 Trial), which was designed to explore this relation, has not been completed and the results have not yet been reported (20). In fact, some (21, 22) but not all (23, 24) of the reviewed studies showed inverse associations between serum 25(OH)D concentration and the risk of CVD, and recent research showed a reverse J-shaped association (25). Some studies reported that high serum 25(OH)D was linked to an elevated risk of total mortality (26), and the association shown between high concentrations of 25(OH)D and CVD events was controversial (26, 27). Conflicting outcomes may be due in part to participant characteristics. In some studies, participants were enrolled from the hospital and were usually experiencing acute illness (28, 29). A

¹ This work was partially funded by the Beijing Municipal Science and Technology Commission (D131100002313002), the Beijing Municipal Commission of Health and Family Planning (2016-1-2041), the National Natural Science Foundation of China (81200914, 81471208), the Beijing high-level healthy human resource project (2014-3-033), and the Nova Program of the Beijing Science and Technology Commission (2008B30).

² The sponsors had no role in the study design; the collection, analysis, and interpretation of data; the writing of the report; or the decision to submit the manuscript for publication.

³ Supplemental Methods and Supplemental Figures 1–3 are available from the “Online Supporting Material” link in the online posting of the article and from the same link in the online table of contents at <http://ajcn.nutrition.org>.

*To whom correspondence should be addressed. E-mail: liugaifen1997@163.com.

⁹ Abbreviations used: CVD, cardiovascular disease; HF, heart failure; IHD, ischemic heart disease; MI, myocardial infarction; MMP, matrix metalloproteinase; 25(OH)D, 25-hydroxyvitamin D.

Received June 18, 2016. Accepted for publication January 26, 2017.

doi: 10.3945/ajcn.116.140392.

TABLE 1Characteristics of the included studies in the current dose-response meta-analysis¹

First author, year (ref)	Country	Endpoint	Age, y	Cases, <i>n</i>	Participants at risk, <i>n</i>	Follow-up, y	Design (study name)	Quality score
Giovannucci, 2008 (40)	USA	MI	63.8 (mean)	454	900	10	Nested case-control study (HPFS)	6
Wang, 2008 (10)	USA	CVD	59 (mean)	120	1739	5.4 (mean)	Cohort study (Framingham Offspring Study)	7
Kilkinen, 2009 (41)	Finland	CVD mortality Cerebrovascular deaths Cardiovascular deaths	49.4 (mean)	933 293 640	6219 6219 6219	27.1 (median)	Cohort study (Mini-Finland Health Survey)	7
Cawthon, 2010 (42)	USA	CVD mortality	≥65	110	1490	7.3 (mean)	Cohort study (MrOS)	6
Hutchinson, 2010 (43)	Norway	CVD mortality (nonsmokers) CVD mortality (smokers)	58.9 (mean)	325 188	4751 2410	11.7 (mean)	Cohort study (Tromsø study)	6
Michaëlsson, 2010 (26)	Sweden	CVD mortality	71 (mean)	196	1312	12.7 (median)	Cohort study (ULSAM)	6
Semba, 2010 (44)	Italy	CVD mortality	≥65	107	1006	6.5	Cohort study (InCHIANTI)	7
Eaton, 2011 (45)	USA	CVD mortality	20–79	79	2429	>10.5	Nested case-control (WHI)	6
Kestenbaum, 2011 (46)	USA	CVD mortality	≥65	389	2312	14 (median)	Cohort study (CHS)	8
Brøndum- Jacobsen, 2012 (21)	Denmark	Nonfatal IHD Nonfatal MI	57 (mean)	1578 802	10,170 10,170	21	Cohort study (Copenhagen City Heart Study)	8
Kritchevsky, 2012 (47)	USA	Fatal IHD/MI CVD mortality	74.7 (mean)	1522 228	10,170 2638	>8.5	Cohort study (Health ABC Study)	8
Lin, 2012 (48)	China	Cerebrovascular deaths Cardiovascular deaths	56.5 (mean)	279 200	1101 1101	>24	Cohort study (None)	6
Messenger, 2012 (49)	USA	CVD diseases	76.4 (mean)	140	813	4.4 (median)	Cohort study (MrOS)	6
Sun, 2012 (22)	USA	Stroke	60.8 (mean)	464	928	16	Nested case-control study (NHS)	7
Brøndum- Jacobsen, 2013 (50)	Denmark	Ischemic stroke	56 (mean)	1256	10,170	21 (median)	Cohort study (Copenhagen City Heart Study)	7
Deng, 2013 (51)	USA	CVD mortality	47 (mean)	1615	12,157	5	Cohort study (NHANES III cohort)	6
Kühn, 2013 (23)	Germany	MI Stroke Composite endpoint	50.6 (mean)	559 471 1030	2132 2132 2132	7.7 (mean)	Case-cohort study (EPIC-Germany)	7
Perna, 2013 (24)	Germany	Total CVD Total IHD Total stroke	50–74	1101 536 353	6709 6709 6709	6.5 (mean)	Cohort study (ESTHER study)	9
Robinson-Cohen, 2013 (52)	USA	IHD	62 (mean)	361	6436	8.5 (median)	Cohort study (MESA)	7
Rohrmann, 2013 (53)	Switzerland	CVD mortality	47.1 (mean)	122	3404	18 (mean)	Cohort study (Swiss National Cohort)	5
Schöttker, 2013 (54)	Germany	CVD mortality	62 (mean)	350	9254	9.5 (median)	Cohort study (ESTHER study)	8
Skaaby, 2013 (55)	Denmark	IHD Stroke	49.8 (mean)	478 316	9146 9146	10 (mean)	Cohort study (Monica10 and Inter99)	6
Bansal, 2014 (56)	USA	HF	62 (mean)	180	6459	8.46 (median)	Cohort study (MESA)	8
Formiga, 2014 (57)	Spain	CVD mortality	85 (mean)	25	312	2.8 (median)	Cohort study (Octabaix study)	4
di Giuseppe, 2014 (58)	Europe	HF	51 (mean)	221	1349	8.2 (mean)	Case-cohort study (EPIC-Potsdam)	8
Khaw, 2014 (27)	United Kingdom	CVD mortality CVD events	62 (mean)	854 4514	14,641 14,641	13 (mean)	Case-cohort study (EPIC-Norfolk)	6
Lee, 2014 (59)	United Kingdom	CVD mortality	60 (mean)	72	2452	4.3 (median)	Cohort study (EMAS)	6

(Continued)

TABLE 1 (Continued)

First author, year (ref)	Country	Endpoint	Age, y	Cases, <i>n</i>	Participants at risk, <i>n</i>	Follow-up, y	Design (study name)	Quality score
Wannamethee, 2014 (60)	United Kingdom	HF	60–79	287	3646	13 (mean)	Cohort study (British Regional Heart Study)	8
Chien, 2015 (61)	Taiwan	CVD events	60.2 (mean)	263	1816	9.6 (median)	Cohort study (CCCC)	6
Lutsey, 2015 (62)	USA	HF	56 (mean)	1763	12,215	14	Cohort study (ARIC)	7
Jassal, 2010 (63)	USA	CVD mortality	74 (mean)	111	1073	6.4 (mean)	Cohort study (Rancho Bernardo Study)	5
Anderson, 2010 (64)	USA	IHD/MI	55 (mean)	763	21,853	1.3 (mean)	Cohort study (None)	5
		HF		594	23,793			
		Stroke		197	26,025			
		CVD mortality		1193	27,686			
		Composite CVD		1304	20,069			
Afzal, 2014 (65)	Denmark	CVD mortality	58 (mean)	2877	9902	19.1 (median)	Cohort study (Copenhagen City Heart Study)	7
				317	25,432	5.8 (median)	Cohort study (Copenhagen General Population Study)	
Karakas, 2013 (66)	Germany	IHD (men)	35–74	225	964	11 (mean)	Case-cohort study	6
		IHD (women)		73	819		(MONICA/KORA Augsburg)	

¹ ABC, Aging and Body Composition; ARIC, Atherosclerosis Risk in Communities Study; CCCC, Chin-Shan Community Cardiovascular Cohort Study; CHS, Cardiovascular Health Study; CVD, cardiovascular disease; EMAS, European Male Ageing Study; EPIC, European Prospective Investigation into Cancer and Nutrition; ESTHER, Epidemiological investigations of the chances of preventing, recognizing early and optimally treating chronic diseases in an elderly population; HF, heart failure; HPFS, Health Professionals Follow-Up Study; IHD, ischemic heart disease; InCHIANTI, Invecchiare in Chianti; MESA, Multi-Ethnic Study of Atherosclerosis; MI, myocardial infarction; MONICA/KORA Augsburg, Monitoring of Trends and Determinants in Cardiovascular Disease/Cooperative Health Research in the Region of Augsburg; MrOS, Osteoporotic Fractures in Men; NHS, Nurses' Health Study; ref, reference; ULSAM, Uppsala Longitudinal Study of Adult Men; WHI, Women's Health Initiative.

previous meta-analysis reviewed the association between vitamin D status and composite CVD but did not include heart failure (HF), and relevant studies were included without regard to the source of participants (30). Because an increasing number of prospective studies have speculated on the association between vitamin D status and the risk of CVD, we performed a comprehensive dose-response meta-analysis of prospective studies to reevaluate and update the relation between baseline serum 25(OH)D concentration and the risk of total CVD events and CVD mortality in the general population.

METHODS

Literature search strategy and selection

This meta-analysis was reported in accordance with the MOOSE (Meta-analysis of Observational Studies in Epidemiology) statement (31). We undertook a systematic search of PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>) and EMBASE (<http://www.embase.com>) through December 2015 for studies related to vitamin D and CVD. In addition, we used hand-searches of the references of all identified articles and relevant reviews to identify pertinent sources. Our search terms used for CVD included cardiovascular diseases, heart failure, myocardial infarction (MI), ischemic heart disease (IHD) and stroke. Search terms used for vitamin D included the following: vitamin D, 25-hydroxyvitamin D, 25hydroxy-vitamin D, 25(OH)D, 1,25-dihydroxyvitamin D, cholecalciferol, and ergocalciferol. CVD-related terms, vitamin D-related terms, and cohort study-related terms were combined by using the Boolean operator "AND." Finally, results were further restricted to human studies,

those in English, and those in adults aged >18 y. Details on the literature search strategy are described in **Supplemental Methods**.

Study selection

The titles and abstracts of identified studies were first screened by 2 investigators (RZ and BL) for potentially relevant sources, and we retrieved the full texts of such articles to assess eligibility. Studies were included if they were of a prospective design, examined the relation between serum 25(OH)D concentration and CVD, and provided risk estimates of RRs, HRs, or ORs and CIs. The exclusion criteria were as follows: 1) case reports, editorials, letters, meeting abstracts, or review articles; 2) retrospective studies, cross-sectional studies, or case-control studies; 3) studies that did not measure serum circulating 25(OH)D at baseline; 4) studies that did not use CVD, IHD, MI, stroke, HF, or CVD mortality as an endpoint; 5) studies that reported <3 exposure categories or did not provide RRs for per-unit changes in 25(OH)D; 6) studies that did not publish the mean or median or range of each category; and 7) studies that were based on a hospital inpatient population with a specific disease. In the case of multiple reported articles from the same cohort, only those with the highest number of cases or the longest follow-up times were included in the meta-analysis. If there were discrepancies between the reviewers (RZ and BL), then another author (GL), as the third investigator, was consulted to reach a consensus.

Data extraction

We collected data from each eligible study, including the name of the first author, publication year, geographical location, source

of cohort, study design, duration of follow-up, baseline year, mean age of participants, sample size, numbers of women and men, cardiovascular outcomes, number of cases, concentration of 25(OH)D in each category, effect sizes and 95% CIs in each category, and adjusted variables. When >1 RRs were reported, we extracted the risk ratio from the model containing the largest number of adjusted variables. For the studies in which results were published for separate subgroups instead of overall, data for those subgroups were extracted individually. We extracted data on all cardiovascular outcomes if several endpoints were reported. For information that was not available or clear in the publication, we directly contacted the author to obtain related

data. We evaluated the quality of individual records by using Newcastle-Ottawa quality-assessment scale (32). Each study was assigned 0–9 “stars” (with more stars meaning higher quality) by 2 of the investigators (RZ and RT) to ensure the accuracy of extraction.

Data synthesis and analysis

The effect sizes between serum 25(OH)D and cardiovascular outcomes were reported in each included study. Because ORs in nested case-control studies and HRs were quite similar to RRs, we used RRs as common risk estimates for all references (22, 33). According to the




FIGURE 1 Forest plot showing the pooled effects of serum 25-hydroxyvitamin D on the risk of total cardiovascular events with the use of the random-effects model (RR: 0.90; 95% CI: 0.86, 0.94), which includes 32 studies (40 estimates). The pooled effects are shown for 10-ng/mL increments in serum 25-hydroxyvitamin D. Solid diamonds and horizontal lines represent RRs (95% CIs) for the outcome of interest. Solid circles and horizontal lines represent RRs (95% CIs); the gray boxes reflect the statistical weight of the study. The dotted vertical line denotes the point estimate for the pooled RRs and the solid vertical line indicates the line of no effect. The open diamond represents the pooled RR with its 95% CI. HF, heart failure; IHD, ischemic heart disease; MI, myocardial infarction.

TABLE 2Subgroup analysis of serum 25(OH)D and risk of total CVD events and CVD mortality in dose-response meta-analysis¹

Subgroup	Total CVD events					CVD mortality				
	<i>n</i>	RR (95% CI)	<i>I</i> ² , %	<i>P</i> _h ²	<i>P</i> _h ³	<i>n</i>	RR (95% CI)	<i>I</i> ² , %	<i>P</i> _h ²	<i>P</i> _h ³
Age, y					0.542					0.534
<65	31	0.91 (0.88, 0.94)	78.2	<0.001		9	0.90 (0.86, 0.93)	22.0	0.247	
≥65	9	0.87 (0.74, 1.03)	89.8	<0.001		8	0.85 (0.72, 1.00)	88.9	<0.001	
Cases, <i>n</i>					0.201					0.063
≤200	13	0.85 (0.74, 0.98)	82.6	<0.001		9	0.87 (0.74, 1.02)	84.4	<0.001	
>200	27	0.91 (0.88, 0.94)	80.1	<0.001		8	0.89 (0.85, 0.93)	35.9	0.142	
Follow-up, y					0.031					0.729
≥10	23	0.92 (0.89, 0.96)	73.7	<0.001		10	0.89 (0.86, 0.93)	32.5	0.148	
5–<10	11	0.88 (0.77, 1.01)	93.6	<0.001		5	0.82 (0.67, 1.00)	86.8	<0.001	
<5	6	0.84 (0.78, 0.90)	45.8	0.101		2	0.96 (0.77, 1.20)	0.0	0.524	
Number of 8 adjusted variables ⁴					0.106					0.013
<6	21	0.93 (0.89, 0.97)	80.8	<0.001		8	0.93 (0.89, 0.96)	0.0	0.451	
≥6	19	0.87 (0.80, 0.94)	92.4	<0.001		9	0.85 (0.76, 0.97)	92.4	<0.001	
Quality scores, stars					0.178					0.023
≤6	22	0.91 (0.87, 0.96)	79.5	<0.001		11	0.90 (0.86, 0.94)	0.0	0.655	
>6	18	0.89 (0.83, 0.96)	92.7	<0.001		6	0.83 (0.71, 0.98)	96.4	<0.001	

¹ *n* represents the number of estimates in the subgroup analysis. CVD, cardiovascular disease; *P*_h, *P*-heterogeneity; 25(OH)D, 25-hydroxyvitamin D.² *P*_h values were for heterogeneity within a subgroup.³ *P*_h values were for heterogeneity between subgroups with the use of meta-regression.⁴ The 8 adjusted variables were sex, age, smoke, blood pressure or history of hypertension, season of blood draw, diabetes, BMI, and physical activity.

Newcastle-Ottawa scale, we considered 0–3, 4–5, and 6–9 stars to be low-, middle-, and high-quality studies, respectively.

Because most of the data we examined were published with the use of different cutoffs of 25(OH)D, we performed a dose-response meta-analysis by using the method proposed by Greenland and Longnecker (34). First, specific linear trends and 95% CIs were estimated from the natural logs of RRs across categories of serum 25(OH)D by the generalized least-square models method. Then, we conducted a random-effects model analysis to examine linear trends (35). The dose-response outcomes were presented per 10-ng/mL (25 nmol/L) increment in serum 25(OH)D in the forest plots. In addition, we assessed the potential nonlinear association between serum 25(OH)D and CVD by using restricted cubic splines with 3 knots (36). For dose-response analysis, this required ≥3 quantitative categories of use. We needed specific data for each category of serum 25(OH)D in all records, including the number of cases and total participants or person-years, RRs and CIs, and median or mean doses. For the studies that did not present the median or mean doses of serum 25(OH)D, we chose the midpoint of each category as the alternative. If the category was open-ended, the midpoint of this category was calculated by assuming the interval was the same as that of the adjacent interval. When the numbers of cases or participants in each category were not available, the number was inferred on the basis of the total number and RRs of each category (37).

*I*² statistics were used to estimate the potential heterogeneity between studies (38). If there was no evidence of between-study heterogeneity, a fixed-effects model was used to recalculate the combined RRs. In addition, we explored the source of heterogeneity by subgroup and meta-regression analyses. Studies were

stratified by duration of follow-up; mean age of participants; numbers of cases; numbers of adjusted important risk factors including sex, age, smoke, blood pressure or history of hypertension, season of blood draw, diabetes, BMI, and physical activity; and study quality. We performed a sensitivity analysis by omitting 1 study at a time to assess the effectiveness of individual studies on the pooled RRs and the robustness of the pooled RRs. Heterogeneity was confirmed with a significance of *P* ≤ 0.10. The possibility of publication bias in the included studies was evaluated by using visual inspection of a funnel plot, Begg's test, and Egger's test. We also used the trim-and-fill method (39) as a sensitivity analysis to explore the potential effects of unidentified studies on the results.

The primary endpoints of the analysis were total CVD events and CVD mortality. If the same cohort presented RRs of IHD/MI, stroke, HF, and composite CVD, we used the composite CVD result in the meta-analysis of total CVD events. In the case of unreported composite results, the specific outcomes of IHD/MI, stroke, and HF were combined in the meta-analysis as separate outcomes. In cases in which a study reported incident CVD and CVD mortality, we extracted the RRs of incident CVD in the analysis of total CVD events. All of the analyses were performed by using STATA software (version 12.0; StataCorp), and significance was defined as *P* < 0.05 with the use of a 2-sided test unless otherwise specified.

RESULTS

Literature research and study characteristics

By using the previously mentioned literature search strategy, a total of 1470 records were identified after duplicates had been removed. Through screening of the title and abstract, we retrieved




FIGURE 2 Dose-response analysis between serum 25-hydroxyvitamin D and the relative risk of total cardiovascular events. The solid line represents point estimates of the association of serum 25-hydroxyvitamin D and total cardiovascular disease risk with the use of a restricted cubic splines model, and the dashed lines indicate 95% CIs.

157 records for their full text, of which 34 were ultimately included in our meta-analysis. Details of exclusions are shown in **Supplemental Figure 1**.

The total number of participants in the included studies was 180,667, which included 9170 CVD deaths, 7074 MI or IHD cases, 3127 stroke cases, and 3037 HF cases. The results of the 34 eligible articles were derived from 27 cohorts; 8 of these articles reported >1 CVD outcomes. Among the included studies, 3 used a nested case-control design, 4 studies used a case-cohort design, and the remaining studies were all prospective cohort studies. Apart from 2 studies conducted in Asia, most were conducted in Europe and United States, with 18 and 14 studies, respectively. The participants in 2 studies were white, and 2 studies were conducted in multiethnic populations. In most of the studies, the mean age of participants was >50 y. The proportion of males was 49.5%. The number of participants in the identified studies ranged from 312 to 27,686, and cases of CVD outcomes ranged from 25 to 4514. The maximal duration of follow-up was >32 y, with the minimum follow-up being 1.3 y. Eighty-five percent of the included studies were of high quality. The extracted information is summarized in **Table 1**.

Serum 25(OH)D and the risk of total CVD events

Among the included studies, 32 were publications (27 independent studies) designed to evaluate the linear dose-response




FIGURE 3 Forest plot showing the pooled effects of serum 25-hydroxyvitamin D on the risk of cardiovascular mortality with use of the random-effects model (RR: 0.87; 95% CI: 0.80, 0.95), which includes 17 studies (18 estimates). The pooled effects are shown for 10-ng/mL increments in serum 25-hydroxyvitamin D. Solid diamonds and horizontal lines represent RRs (95% CIs) for the outcome of interest. Solid circles and horizontal lines represent RRs (95% CIs); the gray boxes reflect the statistical weight of the study. The dotted vertical line denotes the point estimate for the pooled RRs and the solid vertical line indicates the line of no effect. The open diamond represents the pooled RR with its 95% CI.

relation after excluding the 2 studies that reported on CVD mortality from the same cohort (42, 54). The estimated specific linear trends of identified studies between serum 25(OH)D and total CVD events are summarized in **Figure 1**. The pooled RR per 10-ng/mL increment from the random-effects dose-response model indicated an inverse association between vitamin D status and total CVD events (RR: 0.90; 95% CI: 0.86, 0.94), with high heterogeneity ($Q = 414.95$, $I^2 = 90.6\%$; P -heterogeneity < 0.001). To address the main source of heterogeneity, we implemented subgroup analyses according to prespecified characteristics, and significant evidence of heterogeneity was shown between subsets stratified by duration of follow-up and number of adjusted variables by the method of meta-regression (**Table 2**). The relation between serum 25(OH)D and total CVD events showed nonlinearity with the use of a restricted cubic model (P -nonlinear < 0.001) (**Figure 2**). The risk of total CVD events was lower when serum 25(OH)D was > 25 ng/mL.

There was no evidence of publication bias on the basis of visual inspection by the funnel plot (**Supplemental Figure 2**) or Begg's or Egger's test ($P = 0.415$ and 0.590 , respectively). In a sensitivity test that omitted one study each time to obtain the pooled RRs from the random-effects model, the RRs all suggested an inverse association between serum 25(OH)D and total CVD events, with little variation (data not shown). Using the trim-and-fill method, the pooled RR incorporating 2 hypothetical studies was almost unchanged (RR: 0.90; 95% CI: 0.85, 0.93).

Serum 25(OH)D and CVD mortality

A total of 17 publications (17 independent studies) were published that used RRs and 95% CIs for ≥ 3 categories or with the original dose-response data. By using the method proposed by Greenland and Longnecker (34), the linear trends for each specific study were calculated. The pooled RR was $0.88 \cdot 10 \text{ ng}^{-1} \cdot \text{mL}^{-1}$ (95% CI: $0.80, 0.96 \cdot 10 \text{ ng}^{-1} \cdot \text{mL}^{-1}$) increment, with substantial heterogeneity ($I^2 = 90.8\%$). The forest plot in **Figure 3** shows that 1 estimate [Semba, 2010 (44)] deviated from the other values and had the highest weight, and we inferred that it might be the source of high heterogeneity. After removing this study (44), the I^2 dramatically decreased to 17.6% and yielded a slightly different RR (RR: 0.90; 95% CI: 0.86, 0.93). In addition, subgroup analyses showed no significant associations between serum 25(OH)D and CVD mortality in participants aged > 65 y and in studies with ≤ 200 cases and a duration of follow-up < 5 y (**Table 2**). We noted that in other subsets inverse associations were detected. **Figure 4** shows a nonlinear relation between serum 25(OH)D and CVD mortality ($P = 0.022$).

There was no indication of publication bias when evaluated by Begg's test ($P = 0.303$) and Egger's test ($P = 0.117$). The funnel plot is shown in **Supplemental Figure 3**. A sensitivity analysis was conducted, and the RRs did not show substantial changes when omitting one study at a time (data not shown) or when using the technique of trim-and-fill (RR: 0.85; 95% CI: 0.78, 0.93).

DISCUSSION

This dose-response meta-analysis, which was based on general populations, suggested a significant inverse relation between baseline serum 25(OH)D and total CVD events as well as CVD mortality. Because of the considerable heterogeneity, the results

should be interpreted with caution. Furthermore, there was evidence of a nonlinear association between serum 25(OH)D concentration and total CVD events and CVD mortality.

The results of our meta-analysis are consistent with those of a previous meta-analysis (30), which showed an inverse association between serum 25(OH)D and total CVD events. However, Wang et al. (30), in their meta-analysis of 10 prospective studies, suggested a 7% lower risk of total CVD events per 10-ng/mL increment, whereas in the present meta-analysis there was a 10% lower risk. The small differences are probably due to 2 reasons: the previous study included patients recruited from the hospital, which may have influenced participant characteristics, and the present meta-analysis identified studies related to HF. However, HF is a disease entity with substantial heterogeneity in etiology. We included HF in our meta-analysis as one of the endpoints because it is one of the major CVDs, with a relatively high prevalence (1–2%) (67) and important clinical implications (67, 68). Furthermore, HF was defined as one of the composite CVD events in several included studies (10, 26, 27, 54). Of note, the pooled RR did not materially change after we excluded HF from the analyses (RR: 0.90; 95% CI: 0.86, 0.94). For nonlinear analysis, both studies found that at a 25(OH)D concentration of ~ 25 ng/mL, the RR of total CVD events was lowest. Above that concentration, the RR did not decrease significantly. Nevertheless, serum 25(OH)D may be confounded by the type of assay. For example, immune and mass spectrometric methods give different results. Still, defining a threshold of serum 25(OH)D in relation to the risk of total CVD events was an important finding.

For CVD, the leading cause of death worldwide (69), a recent large observational study found that low and high concentrations of 25(OH)D were related to CVD mortality and that the HRs were higher at low concentrations than at high concentrations, in a reverse J-shaped manner (25). In this cohort, Durup et al. (25) suggested that the concentration of 28 ng/mL was linked to the lowest risk of CVD mortality. There was some discrepancy in the results for high concentrations of 25(OH)D, primarily because there were insufficient data for high concentrations, especially for serum 25(OH)D > 40 ng/mL, which might also reflect assay differences. Even in this large observational cohort, only 5% of




FIGURE 4 Dose-response analysis between serum 25-hydroxyvitamin D and the relative risk of cardiovascular mortality. The solid line represents point estimates of the association of serum 25-hydroxyvitamin D and cardiovascular mortality with the use of a restricted cubic splines model, and the dashed lines indicate 95% CIs.

participants had serum 25(OH)D concentrations >40 ng/mL. Contrary to the uncertainty shown for higher concentrations, the majority of related publications (15, 44, 51, 70) showed that the lower extreme concentration of serum 25(OH)D was associated with a higher risk of CVD mortality.

A protective effect of higher baseline serum 25(OH)D on total CVD events and CVD mortality was detected in our meta-analysis. Although the exact mechanism of the association between 25(OH)D and CVD is not known, experimental studies have indicated a regulatory effect of vitamin D on cardiomyocytes and vascular smooth muscle cells (71). In addition, decreased serum 25(OH)D will activate the renin angiotensin system, consequently increasing blood pressure (72). Furthermore, it was reported that vitamin D has anti-inflammatory actions, which play an important role in atherogenesis (71, 73, 74). Vitamin D also inhibits certain matrix metalloproteinases (MMPs) that are important in plaque instability and that are known to increase in MI, notably MMP-9 and MMP-2 (75). However, the majority of results from randomized vitamin D therapy trials failed to support the protective effects against CVD (76). We should note that most randomized vitamin D trials were designed to examine the effect on the skeleton and participants were mostly women and older (73). Well-designed randomized clinical trials are needed to explore the causality of the correlation between vitamin D and CVD, especially if carried out early in the natural history of CVD.

Our meta-analysis had several strengths. First, compared with previous meta-analyses, more comprehensive outcomes were evaluated in this study, including MI or IHD, stroke, HF, and CVD mortality. For total CVD events and CVD mortality, we performed separate dose-response meta-analyses. Moreover, the included studies were all population-based studies and hospital-based studies were excluded. We were therefore more confident to extend the results to the general population. In addition, serum 25(OH)D is the main biomarker for assessing vitamin D status (77). Laboratory measurements of serum 25(OH)D are more objective than measurements of dietary vitamin D intake (78). In this meta-analysis, we used serum 25(OH)D as a measure of exposure. Furthermore, to rule out recall and selection bias, case-control studies were not eligible. Finally, the sample sizes of participants and cases were large and therefore the RR data from this study were more robust.

Several potential limitations of this meta-analysis should be mentioned. First, as with any observational study, residual confounding cannot be excluded. Nevertheless, most of the results of the identified studies were adjusted for recognized risk factors of CVD (smoke, hypertension, diabetes, BMI, physical activity). In most situations, different confounders were considered in the identified studies and the results with the largest number of adjusted covariates were chosen. Second, the number of participants with high concentrations of serum 25(OH)D was small. Third, the association between 25(OH)D concentration and CVD may be confounded by assay variations across studies, especially because most of the studies did not report the use of quality assessment schemes. In consequence, the threshold of low risk should be considered with caution. Fourth, publication bias is inevitable in any meta-analysis, although it was not shown from the funnel plot. Finally, high heterogeneity was apparent in this meta-analysis, which may have been due to differences in duration of follow-up, number of adjusted risk factors, and quality of studies.

In conclusion, the present meta-analysis showed that higher serum 25(OH)D concentrations had a protective effect on total CVD events and CVD mortality. Furthermore, the dose-response analysis showed a J-shaped association between serum 25(OH)D and total CVD events. Well-designed randomized vitamin D therapy trials are needed to confirm the role of vitamin D in preventing overt CVD, as well as to define optimal vitamin D status for a reduction in overall CVD risk.

We thank Sabine Rohrmann and Karl Michaëlsson for their help with provision of extra data to complete our meta-analysis.

The authors' responsibilities were as follows—RZ and GL: designed the research; BL and RT: extracted the data; RZ, XG, and YP: analyzed and interpreted the data; RZ: drafted the manuscript; XG, YP, YJ, HG, Yilong Wang, and Yongjun Wang: revised the manuscript for important intellectual content; RZ: had primary responsibility for the final content; and all authors: read and approved the final manuscript. None of the authors reported a conflict of interest related to the study.

REFERENCES

- de Boer IH, Kestenbaum B, Siscovick DS. Vitamin D insufficiency. *N Engl J Med* 2011;364:1379–80; author reply: 80.
- Lieben L, Carmeliet G. Vitamin D signaling in osteocytes: effects on bone and mineral homeostasis. *Bone* 2013;54:237–43.
- Weaver CM, Alexander DD, Boushey CJ, Dawson-Hughes B, Lappe JM, LeBoff MS, Liu S, Looker AC, Wallace TC, Wang DD. Calcium plus vitamin D supplementation and risk of fractures: an updated meta-analysis from the National Osteoporosis Foundation. *Osteoporos Int* 2016;27:367–76.
- Avenell A, Mak JC, O'Connell D. Vitamin D and vitamin D analogues for preventing fractures in post-menopausal women and older men. *Cochrane Database Syst Rev* 2014;4:CD000227.
- Halfon M, Phan O, Teta D. Vitamin D: a review on its effects on muscle strength, the risk of fall, and frailty. *Biomed Res Int* 2015;2015: 953241.
- Ke L, Graubard BI, Albanes D, Fraser DR, Weinstein SJ, Virtamo J, Brock KE. Hypertension, pulse, and other cardiovascular risk factors and vitamin D status in Finnish men. *Am J Hypertens* 2013;26: 951–6.
- Lim S, Kim MJ, Choi SH, Shin CS, Park KS, Jang HC, Billings LK, Meigs JB. Association of vitamin D deficiency with incidence of type 2 diabetes in high-risk Asian subjects. *Am J Clin Nutr* 2013;97:524–30.
- Gagnon C, Lu ZX, Magliano DJ, Dunstan DW, Shaw JE, Zimmet PZ, Sikaris K, Grantham N, Ebeling PR, Daly RM. Serum 25-hydroxyvitamin D, calcium intake, and risk of type 2 diabetes after 5 years: results from a national, population-based prospective study (the Australian Diabetes, Obesity and Lifestyle Study). *Diabetes Care* 2011;34:1133–8.
- Pham TM, Ekwari JP, Setayeshgar S, Veugeliers PJ. The effect of changing serum 25-hydroxyvitamin D concentrations on metabolic syndrome: a longitudinal analysis of participants of a preventive health program. *Nutrients* 2015;7:7271–84.
- Wang TJ, Pencina MJ, Booth SL, Jacques PF, Ingelsson E, Lanier K, Benjamin EJ, D'Agostino RB, Wolf M, Vasan RS. Vitamin D deficiency and risk of cardiovascular disease. *Circulation* 2008;117: 503–11.
- Michos ED, Reis JP, Post WS, Lutsey PL, Gottesman RF, Mosley TH, Sharrett AR, Melamed ML. 25-Hydroxyvitamin D deficiency is associated with fatal stroke among whites but not blacks: the NHANES-III linked mortality files. *Nutrition* 2012;28:367–71.
- Chandler PD, Buring JE, Manson JE, Giovannucci EL, Moorthy MV, Zhang S, Lee IM, Lin JH. Circulating vitamin D levels and risk of colorectal cancer in women. *Cancer Prev Res (Phila)* 2015;8:675–82.
- McDonnell SL, Baggerly C, French CB, Baggerly LL, Garland CF, Gorham ED, Lappe JM, Heaney RP. Serum 25-hydroxyvitamin D concentrations ≥ 40 ng/ml are associated with $>65\%$ lower cancer risk: pooled analysis of randomized trial and prospective cohort study. *PLoS One* 2016;11:e0152441.
- Skaaby T, Husemoen LL, Pisinger C, Jorgensen T, Thuesen BH, Fenger M, Linneberg A. Vitamin D status and cause-specific mortality: a general population study. *PLoS One* 2012;7:e52423.

15. Pilz S, Dobnig H, Nijpels G, Heine RJ, Stehouwer CDA, Snijder MB, Van Dam RM, Dekker JM. Vitamin D and mortality in older men and women. *Clin Endocrinol (Oxf)* 2009;71:666–72.
16. Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, Durazo-Arvizu RA, Gallagher JC, Gallo RL, Jones G, et al. The 2011 report on Dietary Reference Intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab* 2011;96:53–8.
17. Smith SC Jr., Collins A, Ferrari R, Holmes DR Jr., Logstrup S, McGhie DV, Ralston J, Sacco RL, Stam H, Taubert K, et al. Our time: a call to save preventable death from cardiovascular disease (heart disease and stroke). *Glob Heart* 2012;7:297–305.
18. Peterlik M, Boonen S, Cross HS, Lamberg-Allardt C. Vitamin D and calcium insufficiency-related chronic diseases: an emerging world-wide public health problem. *Int J Environ Res Public Health* 2009;6:2585–607.
19. Theodoratou E, Tzoulaki I, Zgaga L, Ioannidis JP. Vitamin D and multiple health outcomes: umbrella review of systematic reviews and meta-analyses of observational studies and randomised trials. *BMJ* 2014;348:g2035.
20. Bassuk SS, Manson JE, Lee IM, Cook NR, Christen WG, Bubes VY, Gordon DS, Copeland T, Friedenberg G, D'Agostino DM, et al. Baseline characteristics of participants in the VITamin D and Omega-3 Trial (VITAL). *Contemp Clin Trials* 2016;47:235–43.
21. Brøndum-Jacobsen P, Benn M, Jensen GB, Nordestgaard BG. 25-Hydroxyvitamin D levels and risk of ischemic heart disease, myocardial infarction, and early death: population-based study and meta-analyses of 18 and 17 studies. *Arterioscler Thromb Vasc Biol* 2012;32:2794–802.
22. Sun Q, Pan A, Hu FB, Manson JE, Rexrode KM. 25-Hydroxyvitamin D levels and the risk of stroke: a prospective study and meta-analysis. *Stroke* 2012;43:1470–7.
23. Kühn T, Kaaks R, Teucher B, Hirche F, Dierkes J, Weikert C, Katzke V, Boeing H, Stangl GI, Buijsse B. Plasma 25-hydroxyvitamin D and its genetic determinants in relation to incident myocardial infarction and stroke in the European prospective investigation into cancer and nutrition (EPIC)-Germany study. *PLoS One* 2013;8:e69080.
24. Perna L, Schöttker B, Holleczeck B, Brenner H. Serum 25-hydroxyvitamin D and incidence of fatal and nonfatal cardiovascular events: a prospective study with repeated measurements. *J Clin Endocrinol Metab* 2013;98:4908–15.
25. Durup D, Jørgensen HL, Christensen J, Tjønneland A, Olsen A, Halkjær J, Lind B, Heegaard AM, Schwarz P. A reverse J-shaped association between serum 25-hydroxyvitamin D and cardiovascular disease mortality: the CopD study. *J Clin Endocrinol Metab* 2015;100:2339–46.
26. Michaëlsson K, Baron JA, Snellman G, Gedeberg R, Byberg L, Sundström J, Berglund L, Årnlöv J, Hellman P, Blomhoff R, et al. Plasma vitamin D and mortality in older men: a community-based prospective cohort study. *Am J Clin Nutr* 2010;92:841–8.
27. Khaw KT, Luben R, Wareham N. Serum 25-hydroxyvitamin D, mortality, and incident cardiovascular disease, respiratory disease, cancers, and fractures: a 13-y prospective population study. *Am J Clin Nutr* 2014;100:1361–70.
28. Drechsler C, Pilz S, Obermayer-Pietsch B, Verduijn M, Tomaschitz A, Krane V, Espe K, Dekker F, Brandenburg V, März W, et al. Vitamin D deficiency is associated with sudden cardiac death, combined cardiovascular events, and mortality in haemodialysis patients. *Eur Heart J* 2010;31:2253–61.
29. Pilz S, Dobnig H, Fischer JE, Wellnitz B, Seelhorst U, Boehm BO, März W. Low vitamin D levels predict stroke in patients referred to coronary angiography. *Stroke* 2008;39:2611–3.
30. Wang L, Song Y, Manson JE, Pilz S, März W, Michaëlsson K, Lundqvist A, Jassal SK, Barrett-Connor E, Zhang C, et al. Circulating 25-hydroxy-vitamin D and risk of cardiovascular disease: a meta-analysis of prospective studies. *Circ Cardiovasc Qual Outcomes* 2012;5:819–29.
31. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA, Thacker SB; Meta-analysis Of Observational Studies in Epidemiology (MOOSE) Group. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA* 2000;283:2008–12.
32. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses [Internet]. [cited 2017 Jan 8]. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.
33. Berger S, Raman G, Vishwanathan R, Jacques PF, Johnson EJ. Dietary cholesterol and cardiovascular disease: a systematic review and meta-analysis. *Am J Clin Nutr* 2015;102:276–94.
34. Greenland S, Longnecker MP. Methods for trend estimation from summarized dose-response data, with applications to meta-analysis. *Am J Epidemiol* 1992;135:1301–9.
35. Orsini N, Bellocco R, Greenland S. Generalized least squares for trend estimation of summarized dose-response data. *Stata J* 2006;6:40–57.
36. Durrleman S, Simon R. Flexible regression models with cubic splines. *Stat Med* 1989;8:551–61.
37. Bekkering GE, Harris RJ, Thomas S, Mayer AM, Beynon R, Ness AR, Harbord RM, Bain C, Smith GD, Sterne JA. How much of the data published in observational studies of the association between diet and prostate or bladder cancer is usable for meta-analysis? *Am J Epidemiol* 2008;167:1017–26.
38. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
39. Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;56:455–63.
40. Giovannucci E, Liu Y, Hollis BW, Rimm EB. 25-Hydroxyvitamin D and risk of myocardial infarction in men: a prospective study. *Arch Intern Med* 2008;168:1174–80.
41. Kilkkinen A, Knekt P, Aro A, Rissanen H, Marniemi J, Heliövaara M, Impivaara O, Reunanen A. Vitamin D status and the risk of cardiovascular disease death. *Am J Epidemiol* 2009;170:1032–9.
42. Cawthon PM, Parimi N, Barrett-Connor E, Laughlin GA, Ensrud KE, Hoffman AR, Shikany JM, Cauley JA, Lane NE, Bauer DC, et al. Serum 25-hydroxyvitamin D, parathyroid hormone, and mortality in older men. *J Clin Endocrinol Metab* 2010;95:4625–34.
43. Hutchinson MS, Grimnes G, Joakimsen RM, Figenschau Y, Jorde R. Low serum 25-hydroxyvitamin D levels are associated with increased all-cause mortality risk in a general population: the Tromsø study. *Eur J Endocrinol* 2010;162:935–42.
44. Semba RD, Houston DK, Bandinelli S, Sun K, Cherubini A, Cappola AR, Guralnik JM, Ferrucci L. Relationship of 25-hydroxyvitamin D with all-cause and cardiovascular disease mortality in older community-dwelling adults. *Eur J Clin Nutr* 2010;64:203–9.
45. Eaton CB, Young A, Allison MA, Robinson J, Martin LW, Kuller LH, Johnson KC, Curb JD, Van Horn L, McTiernan A, et al. Prospective association of vitamin D concentrations with mortality in postmenopausal women: results from the Women's Health Initiative (WHI). *Am J Clin Nutr* 2011;94:1471–8.
46. Kestenbaum B, Katz R, de Boer I, Hoofnagle A, Sarnak MJ, Shlipak MG, Jenny NS, Siscovick DS. Vitamin D, parathyroid hormone, and cardiovascular events among older adults. *J Am Coll Cardiol* 2011;58:1433–41.
47. Kritchevsky SB, Tooze JA, Neiberg RH, Schwartz GG, Hausman DB, Johnson MA, Bauer DC, Cauley JA, Shea MK, Cawthon PM, et al. 25-Hydroxyvitamin D, parathyroid hormone, and mortality in black and white older adults: the Health ABC Study. *J Clin Endocrinol Metab* 2012;97:4156–65.
48. Lin SW, Chen W, Fan JH, Dawsey SM, Taylor PR, Qiao YL, Abnet CC. Prospective study of serum 25-hydroxyvitamin D concentration and mortality in a Chinese population. *Am J Epidemiol* 2012;176:1043–50.
49. Messenger W, Nielson CM, Li H, Beer T, Barrett-Connor E, Stone K, Shannon J. Serum and dietary vitamin D and cardiovascular disease risk in elderly men: a prospective cohort study. *Nutr Metab Cardiovasc Dis* 2012;22:856–63.
50. Brøndum-Jacobsen P, Nordestgaard Bø G, Schnohr P, Benn M. 25-Hydroxyvitamin D and symptomatic ischemic stroke: an original study and meta-analysis. *Ann Neurol* 2013;73:38–47.
51. Deng X, Song Y, Manson JE, Signorello LB, Zhang SM, Shrubsole MJ, Ness RM, Seidner DL, Dai Q. Magnesium, vitamin D status and mortality: results from US National Health and Nutrition Examination Survey (NHANES) 2001 to 2006 and NHANES III. *BMC Med* 2013;11:187.
52. Robinson-Cohen C, Hoofnagle AN, Ix JH, Sachs MC, Tracy RP, Siscovick DS, Kestenbaum BR, de Boer IH. Racial differences in the association of serum 25-hydroxyvitamin D concentration with coronary heart disease events. *JAMA* 2013;310:179–88.

53. Rohrmann S, Braun J, Bopp M, Faeh D. Inverse association between circulating vitamin D and mortality—dependent on sex and cause of death? *Nutr Metab Cardiovasc Dis* 2013;23:960–6.
54. Schöttker B, Haug U, Schomburg L, Köhrle J, Perna L, Müller H, Holleczeck B, Brenner H. Strong associations of 25-hydroxyvitamin D concentrations with all-cause, cardiovascular, cancer, and respiratory disease mortality in a large cohort study. *Am J Clin Nutr* 2013;97:782–93.
55. Skaaby T, Husemoen LLN, Pisinger C, Jørgensen T, Thuesen BH, Fenger M, Linneberg A. Vitamin D status and incident cardiovascular disease and all-cause mortality: a general population study. *Endocrine* 2013;43:618–25.
56. Bansal N, Zelnick L, Robinson-Cohen C, Hoofnagle AN, Ix JH, Lima JA, Shoben AB, Peralta CA, Siscovick DS, Kestenbaum B, et al. Serum parathyroid hormone and 25-hydroxyvitamin D concentrations and risk of incident heart failure: the Multi-Ethnic Study of Atherosclerosis. *J Am Heart Assoc* 2014;3:e001278.
57. Formiga F, Ferrer A, Megido MJ, Boix L, Contra A, Pujol R, Almeda J, Sanz H, Badia T, Lobato A, et al. Low serum vitamin D is not associated with an increase in mortality in oldest old subjects: the Octabaix Three-Year Follow-up Study. *Gerontology* 2014;60:10–5.
58. di Giuseppe R, Buijsse B, Hirche F, Wirth J, Arregui M, Westphal S, Isermann B, Hense HW, Dierkes J, Boeing H, et al. Plasma fibroblast growth factor 23, parathyroid hormone, 25-hydroxyvitamin D3, and risk of heart failure: a prospective, case-cohort study. *J Clin Endocrinol Metab* 2014;99:947–55.
59. Lee DM, Vanderschueren D, Boonen S, O'Neill TW, Pendleton N, Pye SR, Ravindrarajah R, Gielen E, Claessens F, Bartfai G, et al. Association of 25-hydroxyvitamin D, 1,25-dihydroxyvitamin D and parathyroid hormone with mortality among middle-aged and older European men. *Age Ageing* 2014;43:528–35.
60. Wannamethee SG, Welsh P, Papacosta O, Lennon L, Whincup PH, Sattar N. Elevated parathyroid hormone, but not vitamin D deficiency, is associated with increased risk of heart failure in older men with and without cardiovascular disease. *Circ Heart Fail* 2014;7:732–9.
61. Chien KL, Hsu HC, Chen PC, Lin HJ, Su TC, Chen MF, Lee YT. Total 25-hydroxyvitamin D concentration as a predictor for all-cause death and cardiovascular event risk among ethnic Chinese adults: a cohort study in a Taiwan community. *PLoS One* 2015;10:e0123097.
62. Lutsey PL, Michos ED, Misialek JR, Pankow JS, Loehr L, Selvin E, Reis JP, Gross M, Eckfeldt JH, Folsom AR. Race and vitamin D binding protein gene polymorphisms modify the association of 25-hydroxyvitamin D and incident heart failure: the ARIC (Atherosclerosis Risk in Communities) study. *JACC Heart Fail* 2015;3:347–56.
63. Jassal SK, Chonchol M, von Muhlen D, Smits G, Barrett-Connor E. Vitamin D, parathyroid hormone, and cardiovascular mortality in older adults: the Rancho Bernardo Study. *Am J Med* 2010;123:1114–20.
64. Anderson JL, May HT, Horne BD, Bair TL, Hall NL, Carlquist JF, Lappe DL, Muhlestein JB. Relation of vitamin D deficiency to cardiovascular risk factors, disease status, and incident events in a general healthcare population. *Am J Cardiol* 2010;106:963–8.
65. Afzal S, Brøndum-Jacobsen P, Bojesen SE, Nordestgaard BG. Genetically low vitamin D concentrations and increased mortality: Mendelian randomisation analysis in three large cohorts. *BMJ* 2014;349:g6330.
66. Karakas M, Thorand B, Zierer A, Huth C, Meisinger C, Roden M, Rottbauer W, Peters A, Koenig W, Herder C. Low levels of serum 25-hydroxyvitamin D are associated with increased risk of myocardial infarction, especially in women: results from the MONICA/KORA Augsburg case-cohort study. *J Clin Endocrinol Metab* 2013;98:272–80.
67. Metra M, Carubelli V, Ravera A, Stewart Coats AJ. Heart failure 2016: still more questions than answers. *Int J Cardiol* 2017;227:766–77.
68. Krumholz HM, Chen YT, Wang Y, Vaccarino V, Radford MJ, Horwitz RJ. Predictors of readmission among elderly survivors of admission with heart failure. *Am Heart J* 2000;139:72–7.
69. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, Abraham J, Adair T, Aggarwal R, Ahn SY, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095–128. Erratum in: *Lancet* 2013;381:628.
70. Schierbeck LL, Rejnmark L, Tofteng CL, Stilgren L, Eiken P, Mosekilde L, Køber L, Jensen JEB. Vitamin D deficiency in postmenopausal, healthy women predicts increased cardiovascular events: a 16-year follow-up study. *Eur J Endocrinol* 2012;167:553–60.
71. Motiwala SR, Wang TJ. Vitamin D and cardiovascular risk. *Curr Hypertens Rep* 2012;14:209–18.
72. Ajabshir S, Asif A, Nayer A. The effects of vitamin D on the renin-angiotensin system. *J Nephropathol* 2014;3:41–3.
73. Al Mheid I, Patel RS, Tangpricha V, Quyyumi AA. Vitamin D and cardiovascular disease: is the evidence solid? *Eur Heart J* 2013;34:3691–8.
74. Jouni ZE, Winzerling JJ, McNamara DJ. 1,25-Dihydroxyvitamin D3-induced HL-60 macrophages: regulation of cholesterol and LDL metabolism. *Atherosclerosis* 1995;117:125–38.
75. Timms PM, Mannan N, Hitman GA, Noonan K, Mills PG, Syndercombe-Court D, Aganna E, Price CP, Boucher BJ. Circulating MMP9, vitamin D and variation in the TIMP-1 response with VDR genotype: mechanisms for inflammatory damage in chronic disorders? *QJM* 2002;95:787–96.
76. Mao PJ, Zhang C, Tang L, Xian YQ, Li YS, Wang WD, Zhu XH, Qiu HL, He J, Zhou YH. Effect of calcium or vitamin D supplementation on vascular outcomes: a meta-analysis of randomized controlled trials. *Int J Cardiol* 2013;169:106–11.
77. Haddad JG Jr, Hahn TJ. Natural and synthetic sources of circulating 25-hydroxyvitamin D in man. *Nature* 1973;244:515–7.
78. Scragg R. Vitamin D and public health: an overview of recent research on common diseases and mortality in adulthood. *Public Health Nutr* 2011;14:1515–32.