

No association between blood telomere length and longitudinally assessed diet or adiposity in a young adult Filipino population

Hilary J. Bethancourt^{1,2} · Mario Kratz^{3,4,5} · Shirley A. A. Beresford^{3,4} ·
M. Geoffrey Hayes^{6,7,8} · Christopher W. Kuzawa^{7,9} · Paulita L. Duazo¹⁰ ·
Judith B. Borja^{10,11} · Daniel T. A. Eisenberg^{1,2}

Received: 15 July 2015 / Accepted: 12 October 2015 / Published online: 26 October 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract

Purpose Telomeres, DNA–protein structures that cap and protect chromosomes, are thought to shorten more rapidly when exposed to chronic inflammation and oxidative stress. Diet and nutritional status may be a source of inflammation and oxidative stress. However, relationships between telomere length (TL) and diet or adiposity have primarily been studied cross-sectionally among older, overweight/obese populations and yielded inconsistent results. Little is known about the relationship between diet or body composition and TL among younger, low- to normal-weight populations. It also remains unclear how cumulative exposure to a specific diet or body composition during the years of growth and development, when telomere attrition is most rapid, may be related to TL in adulthood.

Methods In a sample of 1459 young adult Filipinos, we assessed the relationship between blood TL at ages 20.8–22.5 and measures of BMI *z*-score, waist circumference, and diet collected between the ages of 8.5 and 22.5. TL was

measured using monochrome multiplex quantitative PCR, and diet was measured using multiple 24-h recalls.

Results We found no associations between blood TL and any of the measures of adiposity or between blood TL and the seven dietary factors examined: processed meats, fried/grilled meats and fish, non-fried fish, coconut oil, fruits and vegetables, bread and bread products, and sugar-sweetened beverages.

Conclusions Considering the inconsistencies in the literature and our null results, small differences in body composition and consumption of any single pro- or anti-inflammatory dietary component may not by themselves have a meaningful impact on telomere integrity, or the impact may differ across distinct ecological circumstances.

Keywords Telomere length · Diet · Nutrition · Adiposity · Nutrition transition

✉ Hilary J. Bethancourt
hilaryjb@u.washington.edu

¹ Department of Anthropology, University of Washington, Seattle, WA, USA

² Center for Studies in Demography and Ecology, University of Washington, Seattle, WA, USA

³ Department of Epidemiology, School of Public Health, University of Washington, Seattle, WA, USA

⁴ Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA

⁵ Division of Metabolism, Endocrinology, and Nutrition, Department of Medicine, University of Washington, Seattle, WA, USA

⁶ Division of Endocrinology, Metabolism and Molecular Medicine, Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

⁷ Department of Anthropology, Northwestern University, Evanston, IL, USA

⁸ Center for Genetic Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

⁹ Institute for Policy Research, Northwestern University, Evanston, IL, USA

¹⁰ Office of Population Studies Foundation, University of San Carlos, Cebu City, Philippines

¹¹ Department of Nutrition and Dietetics, University of San Carlos, Cebu City, Philippines

Introduction

Modern industrialized diets tend to include highly processed animal products, refined grains, and added fats and sugars but often lack adequate fruits and vegetables. Such diets are thought to cause increased risk of chronic non-communicable diseases and reduced longevity [1–4]. One mechanism via which diet may influence health outcomes is through its potential effects on the length of telomeres, the repeating DNA–protein structures that protect the ends of linear chromosomes from loss of genetic information [5–7]. Telomeres of somatic cells shorten with each cellular replication due to the ‘end-replication problem’ [6, 8], but they may shorten more rapidly if exposed to chronic oxidative stress [9–11] and inflammation [12, 13]. Critically short telomere length (TL) leads to cellular senescence and apoptosis [14, 15], and TL is thought to play a causal role in aging-related disease processes [16].

Certain dietary factors may contribute to or help protect against oxidative stress and inflammation [17–23] and may thereby influence telomere integrity. For instance, diets high in processed meats, refined grains, and added sugars tend to be associated with increased inflammation, whereas diets higher in fruits, vegetables, fish, and whole grains tend to be related to reduced inflammation [17, 20, 24–27]. Saturated fats are widely thought to be pro-inflammatory [28–30], though this may only be the case for long-chain saturated fatty acids [31, 32]. Also, meats cooked at high temperatures (e.g., fried or grilled) are high in advanced glycation end products (AGEs), which are thought to contribute to oxidative stress and inflammation [33, 34].

In line with the evidence on diet and inflammation, several studies have observed longer blood TL among individuals with higher Alternative Healthy Eating Index scores or higher Mediterranean diet scores [35–38]. Further studies have reported associations between blood TL and specific foods or nutrients. For example, some studies have reported inverse associations between blood TL and the consumption of all meat [38], red and/or processed meat [39, 40], saturated fat [41, 42], refined grains [38, 43], and non-diet soda [40, 44]. Positive associations have been observed between blood TL and consumption of fruits and/or vegetables [38, 40, 41, 45, 46] and omega-3 fatty acid supplements [47]. However, findings have not been consistent; in fact, it is more common that studies report null associations between blood TL and consumption of meat [38, 43, 45, 48], saturated fat [37, 43, 49, 50], omega-3 fatty acids or fish [38–41, 43, 45, 48–50], and fruits and vegetables [39, 48, 50, 51].

Notably, many of the populations of the aforementioned studies on diet and TL were, on average, overweight or obese, with the primary exceptions being the few study

populations in China and Korea [40, 48]. Increased adiposity can be an independent contributor to oxidative stress and inflammation [52, 53]. This makes it difficult to clearly separate any direct association between diet and TL from the potential indirect association related to dietary effects on adiposity. Few studies have examined how the relationship between diet and TL may depend on adiposity measures. One study [41] found that associations between blood TL and margarine intake among men and vegetable intake among women were significant among overweight and obese individuals but not among their normal-weight counterparts. Another study [54] suggested that salt intake may be related to blood TL in overweight and obese adolescents but not in normal-weight adolescents. In other words, it may be that the body can generally prevent damage caused by the inflammatory and oxidative properties of food, except when the body is already in a pro-inflammatory state, as in the case of chronic caloric excess and overweight or obesity.

TL attrition is most rapid in the pre-adult years, and it has been suggested that these years might be an especially sensitive period in which environmental and lifestyle factors could influence lifelong TL [55]. In particular, we might expect that cumulative exposure to certain dietary factors and adiposity levels during growth and development could have an important impact on TL in adulthood. However, the majority of studies on TL are cross-sectional and conducted among middle-aged and older adult populations. Few studies have explored the relationship between measures of diet or nutritional status and TL among youth. One study reported that estimated levels of total dietary antioxidants were positively associated with blood TL in a sample of 5- to 18-year-olds in Spain, while white bread and dietary glycemic load were both inversely associated with blood TL [43]. That study did not explore whether those associations differed by body composition, though over 60 % of the adolescents in that study population was overweight or obese. Another study found that blood TL in normal-weight 3- to 9-year-old Australian children was inversely associated with plasma zinc status [56].

A few other studies have explored the relationship between TL and measures of adiposity among younger populations. One study reported an inverse association between blood TL and obesity status among 2- to 17-year-old French children [57], while another found a similar association among 5- to 12-year-old Arab boys but not girls [58]. Among a large sample of 31-year-old Finnish males and females, both body mass index (BMI) and waist-to-hip ratio were found to be inversely associated with blood TL [59], and the inverse association between BMI and blood TL was found to be stronger at younger ages (<30 years old) than at older ages in a US population aged 8–90 years old [60]. In contrast to these studies, however,

no associations were found between TL and measures of adiposity in other studies of children [61], adolescents [62], or young adults [49].

Given the inconsistent findings across the literature and the dearth of information on diet, adiposity, and TL from young, non-obese populations, the present study utilizes longitudinal data on diet and body composition measured among Filipinos during the adolescent and early adult years to test whether specific foods and overall nutritional status during years of rapid cellular turnover are associated with blood TL in early adulthood. Still in the midst of a nutrition transition, consumption of meat, soft drinks, and total fat has increased among Filipino populations in recent decades [63–65], and rates of obesity and cardiometabolic disease have, consequently, been rising [65–68]. Thus, this study setting provides a unique opportunity to explore how diet trends in the early stages of the global nutrition transition [69] may be associated with TL. Specifically, we hypothesized that consumption of fried/grilled meats, processed meats, sugar-sweetened beverages, refined bread and bread products, and coconut oil (the primary cooking oil in the Philippines and a major source of saturated fat) across the adolescent and early adult years would be inversely associated with blood TL, but that consumption of non-fried fish and fruits and vegetables would be positively associated with blood TL in young adult Filipinos. Given the dual burden of relatively high prevalence of undernutrition and increasing prevalence of overnutrition in the Philippines [64, 65, 67, 68, 70], we hypothesized that the cumulative levels of adiposity across the childhood, adolescent, and early adult years would have a curvilinear association with blood TL. We also predicted that diet–TL associations would depend on overall adiposity levels.

Materials and methods

Study participants

Data came from the Cebu Longitudinal Health and Nutrition Survey (CLHNS) [71, 72]. Briefly, this study began in 1983 with the enrolment of 3327 pregnant women from 17 urban and 16 rural neighborhoods from Cebu, the second largest metropolitan area of the Philippines. Follow-up questionnaires on diet, nutritional status, and demographics were administered in 1991–1992, 1994–1995, 1998–1999, 2002–2003, and 2005–2006 when the 1983–1984 birth cohort was on average 8.5, 11.5, 15.5, 18.7, and 21.5 years old. In 2005, venous blood samples were collected from 1779 individuals from the 1983–1984 birth cohort [73]. Only data from singletons in the cohort were used for the analyses conducted herein. All study protocols were originally reviewed and approved by the Institutional Review Board of University of

North Carolina at Chapel Hill. Telomere measurement and analysis using de-identified samples and data were not considered human subject research by Northwestern University's or University of Washington's Institutional Review Boards.

Measurement of TL

Details of the methods used for measuring blood TL can be found elsewhere [73, 74]. In brief, DNA was extracted from the blood samples of 1753 singletons from the 1983–1984 birth cohort, and TL (measured as the relative telomere to single copy gene ratio or “*T/S* ratio”) was analyzed using a modified version the monochrome multiplex quantitative polymerase chain reaction (MMqPCR) method [73–75]. The geometric mean intra-assay coefficient of variation was 5.7 %, and the correlation of TL between samples that were analyzed on two separate runs was 0.92 ($p < 0.0001$). Adjustment for well position further increased the precision of the TL measures [74]. TL measured in whole blood, as is most common in studies exploring relationships between diet or adiposity and TL, is generally considered to be exclusively of leukocyte origin. However, because of the possibility of other cellular and non-cellular sources of DNA in the blood [73, 76], we refer to TL measured in this way as blood TL rather than leukocyte TL.

Assessment of adiposity

Longitudinal assessment of adiposity was based on BMI *z*-scores and waist circumference measurements. As BMI is strongly age-dependent during the childhood and adolescent years, BMI *z*-scores were used instead of BMI to account for age differences of individuals within each survey wave. In order to obtain a more generalizable measure of body mass, World Health Organization reference values for 5- to 19-year-olds, which are used to assess childhood growth worldwide [77–79], were used to estimate BMI *z*-scores of our population sample using age (in months) and weight and height measurements from 1991, 1994, 1998, 2002, and 2005. A comparable BMI *z*-score for ages 19–23 was obtained using the 19-year-old reference values.

Waist circumference was measured in 1998, 2002, and 2005. Because no generalizable reference exists to estimate waist circumference *z*-scores, models of waist circumference adjusted for height and age at the time of each waist circumference measurement.

Assessment of diet

Diet during the adolescent and early adult years was assessed using multiple 24-h recalls administered by trained personnel in 1994, 1998, 2002, and 2005. Surveys were conducted without parental supervision after 1994.

One day of intake was recorded in the 1994–1995 survey wave, and two days of intake were recorded in subsequent survey years.

Estimates of average total energy intake during each survey year were estimated from 24-h recall data using the 1997 Philippines Food Composition Tables from the Food and Nutrition Research Institute [80], which was periodically updated to include newer food items. Coconut oil consumption was estimated by multiplying the weight of fried or sautéed foods consumed by an absorption factor ranging from 0.025 to 0.17, depending on the kind of food and cooking method [81]. Estimates of daily intake (in grams) of processed meat, fried or broiled/grilled meat, non-fried fish, soft drinks, bread and bread products, and fruits and vegetables were calculated directly from the 24-h recall data. The processed meat variable represented consumption of processed beef and pork products, including chorizo (local sausages), hot dogs, and other canned meat and meat products. The fried and broiled/grilled meats and fish variable included beef, poultry, pork, and fish products. The non-fried fish variable included intake of any fresh, smoked, dried, canned, or fermented fish and shellfish consumed raw, boiled, or sautéed; it excluded fried and broiled fish, which, like fried or broiled meat, tend to be high in AGEs [33]. Fruit and vegetable consumption was the combination of all fresh, pickled, or dried vegetables (leafy and non-leafy vegetables and tubers), cooked and raw, combined with fresh, dried, and canned fruits. Estimates of sugar-sweetened beverage intake were based on reports of consuming any kind of soda/soft drink (diet soda consumption is uncommon in Cebu) and non-100 % juices.

Statistical analyses

To obtain a measure of cumulative adiposity levels across the juvenile, adolescent, and early adult years that could be used to test linear and quadratic associations with blood TL, BMI *z*-scores were averaged across the five measurements taken from 1991 to 2005, and waist circumference was averaged across the three available measurements from 1998 to 2005. Women who were pregnant during any of the anthropometric measurement periods ($n = 113$) or who had given birth within 6 months prior to any of the survey waves ($n = 68$) were excluded.

As with the adiposity measures, kilocalories and each food group of interest (measured in grams) was averaged across the seven 24-h recalls available for the 1994–2005 years (Table 2). Females and males ($n = 25$) with average kilocalorie values below 500 and 800 or above 3500 and 4000, respectively, were excluded [82].

Multiple linear regression models with robust standard errors were used to test associations between relative blood TL (*T/S* ratio) and average BMI *z*-score; average waist

circumference; and average intake of processed meat products, grilled/fried meat products, non-fried fish, coconut oil, bread products, fruits and vegetables, and sugar-sweetened beverages. Interactions between dietary factors and average BMI *z*-score were also tested in the diet-specific models. As age and sex are thought to be the strongest, most consistent predictors of TL [16, 55, 83–85], each model initially controlled for age (in months at the time of blood collection in 2005), sex, and the interaction between age and sex. Additionally, initial models included ten principal component scores of genome-wide genetic variation that control for potential population structure [86–88]. Fully adjusted models further controlled for log-transformed average inflation-adjusted household income (collected at birth and again in the five survey waves between 1991 and 2005), smoking status in 2005 (entered as former, current, or never smoker dummy variable categories due to lack of accurate information on pack-years), and average urbanicity score (a scale from 0 to 70 derived from data on population size, population density, communication resources, transportation services, markets, education facilities, and health care services [89]). These additional lifestyle, socioeconomic, and environmental variables were added to the fully adjusted models because of evidence suggesting that TL may be shorter, on average, among smokers and among those growing up in more adverse socioeconomic circumstances [90–93], as well as evidence of strong relationships between urbanicity and energy density in the Philippines [64, 65, 89]. In addition to these variables, the models of waist circumference further adjusted for age and height in each measurement period. The diet-related models further adjusted for average kilocalorie consumption across the four survey waves 1994–2005. All age, kilocalorie, height, and waist variables were mean-centered.

Likelihood ratio tests were used to test the combined effect on model fit of adding linear and quadratic terms for average BMI *z*-score or average waist circumference (Table 5: Models 1A and 1B) and adding BMI *z*-scores and waist circumference measures from different years (Table 5: Models 2A and 2B) to the fully adjusted baseline model. The models assessing adiposity–TL associations were run initially on the entire sample and then separately on males and females; we present here just the sex-specific results for the fully adjusted models.

The models examining diet–TL associations first tested each dietary variable of interest individually in both simple (Table 6: Model 3A) and fully adjusted models (Table 6: Model 3B). Interactions between each individual dietary variable and average BMI *z*-score was then tested in the fully adjusted model (Table 6: Model 3C). Finally, as no significant interactions were found at our adjusted alpha value, all dietary variables of interest were added simultaneously to the fully adjusted model without interaction

Table 1 Sample demographics

TL and demographic characteristics	Males (<i>n</i> = 850)		Females (<i>n</i> = 609)	
	Mean (SD)	Range	Mean (SD)	Range
Relative blood TL (<i>T/S</i> ratio)	0.76 (0.16)	0.20–1.27	0.80 (0.17)	0.029–1.43
Age, 2005 (years)	21.7 (0.3)	20.9–22.5	21.7 (0.4)	20.8–22.5
Average weekly household income, 1983–2005 (Philippine Pesos, inflation-adjusted)	480.7 (366.9)	68.5–3509.3	499.2 (334.3)	70.6–3490.9
Average urbanicity score, 1983–2005 ^a	37.2 (12.6)	7.8–55.7	37.4 (12.1)	8.5–53.7
Smoking status, 2005 (%)	%		%	
Never	18.2		67.7	
Former	49.1		5.9	
Current	32.7		26.4	

^a Urbanicity score ranges from 0 to 70 with higher scores representing greater urbanization

Table 2 Anthropometric characteristics of study sample

Anthropometric measures	Males (<i>n</i> = 850)		Females (<i>n</i> = 609)	
	Mean (SD)	Range	Mean (SD)	Range
BMI <i>z</i> -score, 1991	−0.8 (0.9)	−4.9 to 4.3	−0.8 (0.9)	−4.4 to 2.6
% Underweight (<−2 SD)	9.2		7.2	
% Overweight or obese (>1 SD)	2.7		1.5	
BMI <i>z</i> -score, 1994	−1.2 (1.1)	−4.5 to 3.3	−1.0 (1.1)	−4.8 to 2.9
% Underweight (<−2 SD)	20.4		18.7	
% Overweight or obese (>1 SD)	4.5		5.1	
BMI <i>z</i> -score, 1998	−1.0 (1.1)	−4.0 to 3.0	−0.7 (1.0)	−4.1 to 2.0
% Underweight (<−2 SD)	16.5		10.2	
% Overweight or obese (>1 SD)	3.3		5.1	
BMI <i>z</i> -score, 2002	−0.9 (1.0)	−3.9 to 4.0	−0.6 (1.0)	−3.6 to 2.8
% Underweight (<−2 SD)	12.5		6.4	
% Overweight or obese (>1 SD)	4.0		6.4	
BMI <i>z</i> -score, 2005	−0.6 (1.1)	−3.8 to 3.8	−0.6 (1.1)	−3.5 to 3.8
% Underweight (<−2 SD)	6.8		8.2	
% Overweight or obese (>1 SD)	6.5		7.9	
Average BMI <i>z</i> -score, 1991–2005	−0.9 (0.9)	−3.3 to 3.7	−0.7 (0.9)	−3.2 to 2.0
BMI (kg/m ²), 2005	21.0 (3.0)	14.5 to 40.3	20.1 (3.3)	13.9 to 41.2
% <18.5 kg/m ²	14.9		33.0	
% ≥23.0 kg/m ²	19.3		14.6	
% ≥25.0 kg/m ²	9.3		8.0	
Waist circumference 1998	64.7 (5.7)	50.2 to 114.5	62.9 (5.6)	50.0 to 93.8
Waist circumference 2002	68.6 (6.4)	55.5 to 116.8	65.6 (6.3)	51.0 to 97.5
Waist circumference 2005	72.2 (7.5)	56.5 to 112.0	67.5 (7.6)	53.5 to 112.2
Average waist circumference, 1998–2005	68.5 (6.0)	57.0 to 111.5	65.3 (5.9)	54.2 to 96.9

terms (Table 6: Model 3D). In the latter model, a likelihood ratio test was used to test the combined effect on model fit of adding all dietary variables to the baseline model, and Wald's tests were used to test the combined effect of (1) processed and grilled/fried animal products; (2) non-fried fish and fruits and vegetables; and (3) bread and bread products and sugar-sweetened beverages.

All models were conducted on the 1459 (609 females and 850 males) with complete blood TL, BMI, waist circumference, dietary, income, urbanicity, and smoking data (excluding women who were pregnant during or recently prior to any survey wave). With this sample, we had at least 80 % power in all models to detect a change in R^2 as small as 0.01 at Bonferroni-adjusted alpha values of 0.0125 and

Table 3 Dietary characteristics of study sample

Dietary variables ^a	Males (<i>n</i> = 850)		Females (<i>n</i> = 609)	
	Mean (SD)	Range	Mean (SD)	Range
Kilocalories	1943 (547.5)	810–3907	1423 (439.8)	556–3396
Meat and poultry (all, g/day)	117.2 (79.7)	0–452.0	89.8 (58.1)	0–380.7
Processed red meat (g/day)	25.1 (25.3)	0–169.5	20.7 (21.2)	0–198.6
Grilled/fried meat and fish (g/day)	90.3 (64.7)	0–568.3	81.6 (54.0)	0–327.2
Fish (all, g/day)	46.8 (26.2)	0–155.3	36.5 (20.6)	0–133.0
Non-fried fish (g/day)	26.9 (20.4)	0–126.9	19.8 (15.1)	0–94.6
Coconut oil (g/day)	9.1 (5.0)	0–28.3	8.0 (4.4)	0.4–31.7
Other added oils and fat (g/day)	7.4 (6.6)	0–48.5	7.0 (5.5)	0–33.3
Fruit and vegetables (all, g/day)	55.4 (43.7)	0–325.6	53.9 (38.3)	0–229.3
Fruit (g/day)	10.0 (21.7)	0–240.4	14.4 (22.6)	0–182.1
Non-root vegetables (g/day)	26.2 (26.7)	0–208.0	22.4 (19.8)	0–119.4
Root vegetables and plantains (g/day)	19.2 (23.9)	0–217.7	17.2 (18.4)	0–103.6
Rice (g/day)	603.3 (259.8)	0–1591.1	351.4 (166.8)	0–1033.9
Bread and bread products (g/day)	52.8 (37.9)	0–252.5	48.7 (27.3)	0–165.7
Soft drinks and artificial juices (oz/day)	3.4 (3.4)	0–21.5	3.5 (3.1)	0–18.9

^a Dietary variables are the averages of intakes reported in the 1994, 1998, 2002, and 2005 survey waves

0.0071 in the adiposity and dietary models, respectively, to account for the four separate tests of linear and quadratic associations with adiposity measures and the separate tests of seven dietary factors. These adjusted alpha values were used to assess significance, but all *p* values reported herein are the raw, unadjusted *p* values.

Results

Our study population was 20.8–22.5 (mean = 21.7) years old in 2005 when blood TL was measured (Table 1). Average household income and urbanicity scores were comparable between males and females, but a substantially greater proportion of males were current or former smokers compared to females.

Overall, the prevalence of underweight was greater than the prevalence of overweight in most years among both males and females (Table 2), though this trend began to reverse in 2005 for males. Given the BMI *z*-score cut-offs for underweight and overweight of <−2 SD and >1 SD, respectively [77], up to 20.4 % of males and 18.7 % of females were considered underweight in any given year of childhood or adolescence, while no more than 6.4 % of males and females were considered in the overweight or obese categories during the younger years. By early adulthood, still 14.9 % of males and 33.0 % of females were considered underweight (BMI < 18.5 kg/m²) but up to 19.3 % of males and 14.6 % of females could be considered above “normal” weight at the suggested BMI cutoff of 23 kg/m² for Asian populations [94].

Consumption of animal products, processed bread products, and sugar-sweetened beverages remained low on average in this population (Table 3). Only about half of the population was consuming at least one serving (85 g or 3 oz) of any kind of meat per day on average. Similarly, only about 60 % the population ate the equivalent of one slice of bread on a daily basis. An even smaller proportion of the population (about 10 %) was consuming an average of one 8-oz serving of sugar-sweetened beverages on a regular basis. At the same time, consumption of more traditional Filipino foods, other than rice, was also low. A mere 7 and 12 % of this population ate an average of one serving (85 g or 3 oz) or more of fish or even a single serving (100 g or 3.5 oz) or more of fruits and vegetables, respectively.

As reported previously for this population [73] and in line with previous studies [76, 85], blood TL was, on average, shorter among males ($\beta = -0.036$, $p < 0.0001$) and slightly shorter with each additional month of age ($\beta = -0.0035$, $p = 0.015$) in the simple baseline model (Table 4). The strength of the relationship between age and blood TL was attenuated when variables for average household income, average urbanicity score, and smoking status were added to the baseline model. Average urbanicity score was significantly positively associated with blood TL ($\beta = 0.0028$, $p < 0.0001$), while average log-transformed income was, intriguingly, significantly inversely associated with blood TL ($\beta = -0.027$, $p < 0.0001$). Controlling for these additional variables almost doubled the adjusted *R*² of the baseline model.

No further substantial changes in the adjusted *R*² were observed when adding our main adiposity or dietary

Table 4 Baseline regression models ($n = 1459$)

	Simple baseline model (adjusted $R^2 = 0.04$)		Fully adjusted baseline model (adjusted $R^2 = 0.07$)	
	β (95 % CI)	p	β (95 % CI)	p
Age in 2005 (months)	−0.0035 (−0.0063, −0.00070)	0.015	−0.0021 (−0.0050, 0.00073)	0.15
Sex (male)	−0.036 (−0.053, −0.020)	<0.0001	−0.043 (−0.063, −0.023)	<0.0001
Age*sex interaction	−0.0031 (−0.0068, 0.00070)	0.11	−0.0029 (−0.0067, 0.00082)	0.13
Principal component scores (PCS)				
PCS 1	−0.60 (−0.94, −0.26)	0.00	−0.45 (−0.79, −0.11)	0.0090
PCS 2	0.071 (−0.27, 0.42)	0.69	0.088 (−0.25, 0.42)	0.61
PCS 3	0.17 (−0.18, 0.53)	0.34	0.12 (−0.23, 0.47)	0.50
PCS 4	0.088 (−0.26, 0.44)	0.62	0.033 (−0.31, 0.38)	0.85
PCS 5	−0.12 (−0.46, 0.21)	0.47	−0.11 (−0.45, 0.22)	0.50
PCS 6	−0.073 (−0.42, 0.27)	0.68	−0.040 (−0.38, 0.30)	0.82
PCS 7	−0.25 (−0.59, 0.09)	0.16	−0.18 (−0.51, 0.16)	0.30
PCS 8	0.24 (−0.098, 0.58)	0.16	0.22 (−0.10, 0.55)	0.18
PCS 9	0.018 (−0.32, 0.35)	0.92	−0.13 (−0.47, 0.21)	0.44
PCS 10	0.41 (0.065, 0.75)	0.020	0.38 (0.043, 0.71)	0.027
Ln (average income 1983–2005)			−0.027 (−0.041, −0.012)	<0.0001
Average urbanicity score (1983–2005)			0.0028 (0.0021, 0.0035)	<0.0001
Smoking status				
Current smoker			0.012 (−0.010, 0.035)	0.28
Former smoker			0.0094 (−0.011, 0.030)	0.37

predictors to any of the models (Tables 5, 6). There was no evidence in our study population of a linear or quadratic association between blood TL and BMI z -scores or waist circumference measures in any of the models tested (p values for all likelihood ratio tests ≥ 0.10 without correction for multiple testing) (Table 5). Though the linear and quadratic terms in Model 1B were suggestive of a curvilinear association between blood TL and average waist circumference among males, the likelihood ratio test indicated that the combined effect of the linear and curvilinear terms was not statistically significant, particularly after adjustment for multiple comparisons.

Likewise, there was no evidence that any of the dietary variables examined were associated with blood TL (Table 6). In the simple dietary model (Model 3A), most of the coefficients were in the opposite direction of what we hypothesized, and there were no significant associations (all p values ≥ 0.3 without correction for multiple testing). In the fully adjusted model (Model 3B), and particularly in the fully adjusted model that included interactions with average BMI z -score (Model 3C), most of the coefficients were in the hypothesized direction, but still none of the associations in these fully adjusted models were significant (all p values ≥ 0.08 without correction for multiple testing). Only the interaction between non-fried fish and average BMI z -score approached significance at an unadjusted

alpha value (β for interaction = 0.031, $p = 0.02$), but this would not be considered significant after accounting for multiple comparisons. Finally, there was no evidence of a relationship between groups of dietary variables and blood TL in the fully adjusted model that included all dietary variables together (Model 3D) (all Wald's test p values ≥ 0.68 ; likelihood ratio test p value = 0.96).

Discussion

This study assessed the association between blood TL and measures of adiposity and pro- and anti-inflammatory dietary factors among a young, non-Western population. The data used for this study provided a unique opportunity to assess how diet across years of growth, development, and rapid cellular turnover may impact TL of a generally low-to normal-weight population. Contrary to expectations, we did not find evidence to support our hypotheses that diet and adiposity levels during the childhood, adolescent, and early adult years are associated with blood TL measured in early adulthood.

The results of the present study contrast with reports of inverse associations between blood TL and obesity status or adiposity measures among children [57] and young adults [59, 60], as well as evidence for variable associations

Table 5 Relationship between adiposity measures and blood TL among males and females

	Males (<i>n</i> = 850)		Females (<i>n</i> = 609)	
	β (95 % CI)	<i>p</i> ^a	β (95 % CI)	<i>p</i> ^a
Model 1A				
Average BMI 1991–2005				
Linear term	0.0015 (−0.013, 0.016)	0.84	0.0022 (−0.019, 0.024)	0.84
Quadratic term	−0.0013 (−0.0079, 0.0053)	0.71	0.0078 (−0.0036, 0.019)	0.18
Likelihood ratio test		0.82		0.20
Model 2A				
BMI <i>z</i> -score 1991	−0.0011 (−0.017, 0.015)	0.90	−0.0025 (−0.024, 0.019)	0.82
BMI <i>z</i> -score 1994	−0.0076 (−0.023, 0.0079)	0.34	0.019 (−0.0032, 0.042)	0.09
BMI <i>z</i> -score 1998	−0.00061 (−0.021, 0.019)	0.95	−0.014 (−0.04, 0.013)	0.31
BMI <i>z</i> -score 2002	0.014 (−0.0055, 0.034)	0.16	−0.0078 (−0.036, 0.020)	0.59
BMI <i>z</i> -score 2005	−0.0012 (−0.017, 0.014)	0.88	−0.0055 (−0.029, 0.018)	0.64
Likelihood ratio test		0.63		0.35
Model 1B				
Average waist circumference 1991–2005 ^a				
Linear term	0.0029 (0.000088, 0.0056)	0.04	−0.0012 (−0.0037, 0.0013)	0.34
Quadratic term	−0.00012 (−0.00022, −0.000017)	0.02	0.00013 (−0.000073, 0.00033)	0.21
Likelihood ratio test		0.10		0.37
Model 2B				
Waist circumference 1998 ^b	0.0012 (−0.00203, 0.0044)	0.47	−0.00048 (−0.0043, 0.0034)	0.81
Waist circumference 2002 ^b	−0.00085 (−0.0045, 0.0028)	0.64	−0.00081 (−0.0046, 0.0030)	0.68
Waist circumference 2005 ^b	0.00056 (−0.0018, 0.0029)	0.64	0.00038 (−0.0025, 0.0033)	0.80
Likelihood ratio test		0.74		0.90

All models control for mean-centered age in 2005 (in months), sex, the interaction between age and sex, log-transformed average inflation-adjusted income from 1983 to 2005, average urbanicity score from 1983 to 2005, smoking status (current, former, and never smoker), and 10 principal component scores of genome-wide data

^a All *p* values are raw, unadjusted *p* values. Significance levels in these models are set at an adjusted alpha value of 0.0125

^b All models with waist circumference further controlled for mean-centered height and mean-centered age in each of the measurement periods

between adiposity and blood TL in males and females [58, 95]. This could arguably be due to the low proportion of overweight and obese individuals in our sample, at least in comparison with the Finnish sample of young adults in which 49 and 32 % of males and females, respectively, were above the 25 kg/m² BMI cutoff for normal weight [59]. Nonetheless, these results are consistent with other studies that reported no associations between adiposity measures and TL among samples of children [61], adolescents [62], and young adults [49], despite substantially higher average BMI levels compared to those in our sample. These results can also be added to the larger body of inconsistent findings between adiposity and TL among adults overall [as reviewed in 85, 96], even when considering primarily Western populations in which the prevalence of overweight and obesity is considerably higher than in the Philippines.

Our null findings from the models testing the associations between blood TL and the specific food categories of interest contrast with the one study of adolescents that

found significant inverse relationships between blood TL and white bread consumption [43]. They also contrast with several studies among adult populations that reported inverse associations between blood TL and red and/or processed meat [39, 40], and soda consumption [40, 44] and positive associations between blood TL and consumption of fruit [40, 46] and vegetables [38, 41, 45].

Nonetheless, as few studies reporting significant diet–TL relationships take into account multiple comparisons, the strength of those findings remains unclear. Overall, our null findings are actually in agreement with a greater number of studies that found no association between blood TL and fruits and vegetables assessed together [50, 51] or separately [39, 43, 48] and studies that found no association between fish consumption and blood TL in fully adjusted models [39, 40, 43, 45, 48]. No previous studies explored the relationship between coconut oil and TL. However, considering the saturated fatty acid content of coconut oil, the null results in the present study contrast with the two studies that found inverse associations between consumption

Table 6 Relationship between dietary factors and blood TL ($n = 1459$)

	Model 3A ^a		Model 3B ^b		Model 3C ^c		Model 3D ^d	
	β (95 % CI)	p^e	β (95 % CI)	p^e	β (95 % CI)	p^e	β (95 % CI)	p^e
Calories (100 kcal)								
Processed meat (50 g)	0.00048 (−0.017, 0.018)	0.96	−0.0046 (−0.022, 0.013)	0.61	−0.0072 (−0.028, 0.014)	0.51	−0.00089 (−0.0032, 0.0014)	0.45
Interaction with BMI								
Grilled/fried meat and fish (50 g)	−0.00045 (−0.0087, 0.0078)	0.91	−0.0031 (−0.011, 0.0051)	0.45	−0.0040 (−0.013, 0.0054)	0.66	−0.0032 (−0.012, 0.0057)	0.48
Interaction with BMI								
Non-fried fish (50 g)	−0.0023 (−0.026, 0.021)	0.85	0.0054 (−0.018, 0.029)	0.65	−0.0015 (−0.0077, 0.0048)	0.65		
Interaction with BMI								
Coconut oil (5 g)	0.0021 (−0.0083, 0.013)	0.69	0.0010 (−0.0095, 0.011)	0.86	0.0016 (−0.0098, 0.0131)	0.78	0.0028 (−0.0083, 0.014)	0.62
Interaction with BMI								
Fruits and vegetables (50 g)	−0.0035 (−0.013, 0.0060)	0.47	0.0041 (−0.0054, 0.014)	0.39	0.0020 (−0.0098, 0.0131)	0.77	0.0043 (−0.0053, 0.014)	0.38
Interaction with BMI								
Bread and bread products (50 g)	0.0068 (−0.0059, 0.019)	0.30	0.0019 (−0.011, 0.014)	0.77	−0.0024 (−0.00702, 0.00904)	0.66		
Interaction with BMI								
Soft drinks and juice (4 oz)	0.0011 (−0.0095, 0.012)	0.84	0.0010 (−0.0095, 0.011)	0.86	−0.0016 (−0.018, 0.015)	0.85	0.0021 (−0.011, 0.015)	0.75
Interaction with BMI								
Wald's Tests								
Processed meat, grilled/fried meat and fish					−0.0041 (−0.017, 0.0084)	0.52		0.74
Fish and vegetables					−0.00052 (−0.014, 0.013)	0.94	0.0017 (−0.0090, 0.012)	0.75
Bread and sweetened beverages					−0.0020 (−0.012, 0.0080)	0.70		0.89
Likelihood ratio test								
Combined dietary factors								0.96

^a Simple model testing each dietary variable individually only controlling for mean-centered age in 2005 (in months), sex, the interaction between age and sex, 10 principal component scores of genome-wide data, and mean-centered average kilocalories from 1994 to 2005

^b Fully adjusted model testing each dietary variable individually controlling for all in Model 3A in addition to log-transformed average inflation-adjusted income from 1983 to 2005, average urbanicity score from 1983 to 2005, and smoking status (current, former, and never smoker)

^c Fully adjusted model testing interaction between individual dietary variables and average BMI z-score from 1991 to 2005

^d Fully adjusted model testing all dietary variables together in the same model

^e All p values are raw, unadjusted p values. Significance levels in these models are set at an adjusted alpha value of 0.0071

of all or specific saturated fatty acids and blood TL among males [41] and females [42] but agree with other studies that found no association between total saturated fat intake and blood TL [37, 43, 49, 50]. There are no previous studies exploring the relationship between fried and broiled/grilled animal products and TL with which to compare the results of the present study, but there are several studies that looked at consumption of meat in general and found no statistically significant associations with blood TL [43, 45, 48] or found associations only within specific subpopulations [38].

Given that the nutrition transition was still in a relatively early stage at the time of data collection, particularly in the rural areas of the Philippines [65], it may be that the diet of our study population was not yet particularly pro-inflammatory or that consumption of the pro- and anti-inflammatory foods investigated was not variable enough across individuals to detect a marked association with blood TL. On the one hand, it was remarkable how consistently low the intake of antioxidant-rich fruits and vegetables was in this young Filipino population compared to reported intake levels of Spanish adolescents [43] and US, European, and Chinese adult populations [37, 39, 41, 45, 48, 50]. On the other hand, consumption of other common “Western” pro-inflammatory foods was also relatively low, with the exception of processed meat, which was consumed at levels comparable to a US adult population in which inverse associations between processed meat and blood TL were found [39]. Despite reports of increased soda consumption in the Philippines in recent decades [63, 65], average consumption of sugar-sweetened beverages was still only about a quarter of the levels reported in a US adult population in which a significant inverse association between non-diet soda and blood TL was found [44] and about 10–50 % lower than consumption levels of another US adult population [39] and the Spanish adolescent population [43] in which no associations with blood TL were found. Similarly, average consumption of bread and bread products was less than half that of the young Spanish population in which significant associations between white bread and blood TL were reported [43]. Though intake of added oils and fats among this Filipino population was comparable to an elderly Chinese population in which a borderline (albeit sex-specific) inverse association with blood TL was reported [48], coconut oil, the primary source of added fat in the Filipino diet, may actually be relatively anti-inflammatory compared to other saturated or unsaturated fats [31, 32]. Moreover, the overall low caloric intake in this population may be protective against inflammation and oxidative stress [97, 98]. Caloric intake in the earlier years was about half that of the Spanish adolescents, and at ages 20–22 it was lower, on average, than the energy intake of many US and European adult populations in which diet–TL

relationships have been assessed [39, 41, 45, 49, 50]. In fact, recent studies have demonstrated that this cohort of young Filipinos has, on average, lower levels of inflammatory cytokines than age-matched controls from US and European populations [99, 100].

Viewed alongside the generally negative findings in the literature, our data suggest that, at least among younger and low- to normal-weight individuals, differences in adiposity may not have a substantial impact on blood TL. Likewise, individual dietary components may not have a meaningful impact on telomere integrity, at least when caloric consumption and adiposity levels are otherwise low. Though there was little suggestion of an interaction between dietary variables and average BMI z-score, the low proportion of overweight or obese individuals may have limited the ability to detect this.

Some researchers have recently suggested that TL is not the biomarker of cumulative inflammation and oxidative stress that it was previously thought to be and that, compared with biological factors, lifestyle factors play a relatively minor role in determining TL [16, 55, 101]. However, given the characteristics of our study sample and several methodological limitations, it remains unclear whether this is an explanation for the null results of the analyses presented here. For example, given the low proportion of overweight individuals and overall low levels of inflammation among this population, any effects of adiposity on inflammation and, thereby, on blood TL may have been too small to detect with our sample size. Similarly, considering the potential synergistic and antagonistic effects of different dietary components [102] and the significant associations between blood TL and both income and level of urbanization in our study sample, it may be that the overall combination of physiological, dietary, lifestyle, and environmental factors matter more than any individual factor. It may also take longer than we originally hypothesized for the cumulative effect of diet and adiposity to become manifest as differences in blood TL. Indeed, a number of studies that found no significant association between TL and individual dietary factors among middle-aged and older adult populations did find significant associations between TL and indices of overall healthy dietary or lifestyle patterns [37, 51, 103].

Furthermore, given the known potential for measurement error when using qPCR methods [85, 101] or attempting to measure the usual diet of either juvenile, adolescent, or adult populations [82, 104], we cannot rule out the possibility that statistical noise combined with limited power may have precluded our ability to detect associations in this sample. However, our MMqPCR measure of blood TL is well validated and shows the expected magnitude of associations with blood TL measured using the gold standard Southern Blot method, as well as with age, sex, maternal

blood TL, and paternal age [73, 74]. Our sample size is also larger than some other studies that reported associations between TL and diet or adiposity [39, 43, 45, 57, 58]. Moreover, the multiple 24-h recalls across time should have increased the reliability of our dietary measures [105] and should have provided more accurate measures of average dietary intake than the one-time food frequency questionnaires used in almost every other known study on diet and TL.

Additional sources of variability and residual confounding may have come from factors either not measured in this study or not controlled for in the analyses. Had we been able to obtain measures of blood TL at two different time points, we would have been better equipped to distinguish between variability in blood TL caused by inherited and early biological factors versus variability in the actual rate of telomere shortening. Also, it is intriguing that average household income was negatively associated with blood TL in our study sample. These results are not necessarily contradictory to the mixed findings of other studies [106]. However, these findings may indicate that other unmeasured socioeconomic, lifestyle, environmental, and early development factors could be moderating the effect of adiposity and diet on blood TL in our study population, and the effect of those factors may differ across varying ecological circumstances.

In summary, this study found no evidence that individual dietary characteristics or measures of adiposity across the childhood, adolescent, and early adult years are associated with inter-individual variability in the blood TL of low- to normal-weight young adult Filipino males and females. These results contribute to the existing body of mixed evidence for TL–diet and TL–adiposity relationships and provide new information regarding specific food groups (e.g., grilled/fried meats and coconut oil) and adiposity measures across childhood and adolescent years among a non-Western population. The null results of this and other studies suggest that small differences in body composition and consumption of any single pro- or anti-inflammatory dietary component may not have a meaningful impact on TL, or the impact may differ depending on the broader set of nutritional, lifestyle, and environmental circumstances. Diet and adiposity still undoubtedly play an important role in long-term health and longevity, but telomere attrition may play a relatively minor role, if any, in those specific causal pathways.

Acknowledgments Partial support for this research came from a National Science Foundation Graduate Research Fellowship (DGE-0718145 and DGE-1256082) to HJB and *Shanahan Endowment Fellowship* and *Eunice Kennedy Shriver* National Institute of Child Health and Human Development research infrastructure and training grants (R24 HD042828 and T32 HD007543) to the Center for Studies

in Demography and Ecology at the University of Washington. The qPCR laboratory analysis was funded by National Science Foundation (Doctoral Dissertation Improvement Grant BCS-0962282), the Wenner-Gren Foundation (Gr. 8111), and institutional support from Northwestern University. Data and sample collection was funded by the National Institute of Health (Grants TW05596, DK078150, RR20649, ES10126, and DK056350). Importantly, we would like to acknowledge the participants and staff of the Cebu Longitudinal Health and Nutrition Survey who made this study possible.

Compliance with ethical standards

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

1. Heidemann C, Schulze MB, Franco OH, van Dam RM, Mantzoros CS, Hu FB (2008) Dietary patterns and risk of mortality from cardiovascular disease, cancer, and all causes in a prospective cohort of women. *Circulation* 118:230–237
2. Schulze MB, Fung TT, Manson JE, Willett WC, Hu FB (2006) Dietary patterns and changes in body weight in women. *Obesity* 14:1444–1453
3. Popkin BM (2006) Global nutrition dynamics: the world is shifting rapidly toward a diet linked with noncommunicable diseases. *Am J Clin Nutr* 84:289–298
4. Hu FB, Rimm EB, Stampfer MJ, Ascherio A, Spiegelman D, Willett WC (2000) Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr* 72:912–921
5. Blackburn EH (2005) Telomeres and telomerase: their mechanisms of action and the effects of altering their functions. *FEBS Lett* 579:859–862
6. de Lange T (2009) How telomeres solve the end-protection problem. *Science* 326:948–952
7. O'Sullivan RJ, Karlseder J (2010) Telomeres: protecting chromosomes against genome instability. *Nat Rev Mol Cell Biol* 11:171–181
8. Harley CB, Vaziri H, Counter CM, Allsopp RC (1992) The telomere hypothesis of cellular aging. *Exp Gerontol* 27:375–382
9. Von Zglinicki T (2002) Oxidative stress shortens telomeres. *Trends Biochem Sci* 27:339–344
10. Von Zglinicki T (2000) Role of oxidative stress in telomere length regulation and replicative senescence. *Ann NY Acad Sci* 908:99–110
11. Kawanishi S, Oikawa S (2004) Mechanism of telomere shortening by oxidative stress. *Ann NY Acad Sci* 1019:278–284
12. Aviv A (2004) Telomeres and human aging: facts and fids. *Sci Aging Knowl Environ* 2004:pe43
13. Houben MJ, Moonen HJJ, van Schooten FJ, Hageman GJ (2008) Telomere length assessment: biomarker of chronic oxidative stress? *Free Radic Biol Med* 44:235–246
14. Blackburn EH (2001) Switching and signaling at the telomere. *Cell* 106:661–673
15. Blackburn EH (2000) Telomere states and cell fates. *Nature* 408:53–56
16. Aviv A, Kark JD, Susser E (2015) Telomeres, atherosclerosis, and human longevity: a causal hypothesis. *Epidemiology* 26:295–299
17. Lopez-Garcia E, Schulze MB, Fung TT, Meigs JB, Rifai N, Manson JE, Hu FB (2004) Major dietary patterns are related to

- plasma concentrations of markers of inflammation and endothelial dysfunction. *Am J Clin Nutr* 80:1029–1035
18. Chrysoshoou C, Panagiotakos DB, Pitsavos C, Das UN, Stefanadis C (2004) Adherence to the Mediterranean diet attenuates inflammation and coagulation process in healthy adults: the Attica study. *J Am Coll Cardiol* 44:152–158
 19. Fitó M, Guxens M, Corella D et al (2007) Effect of a traditional Mediterranean diet on lipoprotein oxidation: a randomized controlled trial. *Arch Intern Med* 167:1195–1203
 20. Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, D'Armiento M, D'Andrea F, Giugliano D (2004) Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA* 292:1440–1446
 21. Margioris AN (2009) Fatty acids and postprandial inflammation. *Curr Opin Clin Nutr Metab Care* 12:129–137
 22. Kallio P, Kolehmainen M, Laaksonen DE, Pulkkinen L, Atalay M, Mykkänen H, Uusitupa M, Poutanen K, Niskanen L (2008) Inflammation markers are modulated by responses to diets differing in postprandial insulin responses in individuals with the metabolic syndrome. *Am J Clin Nutr* 87:1497–1503
 23. Galland L (2010) Diet and inflammation. *Nutr Clin Pract* 25:634–640
 24. Schulze MB, Hoffmann K, Manson JE, Willett WC, Meigs JB, Weikert C, Heidemann C, Colditz GA, Hu FB (2005) Dietary pattern, inflammation, and incidence of type 2 diabetes in women. *Am J Clin Nutr* 82:675–684
 25. Nettleton JA, Steffen LM, Mayer-Davis EJ, Jenny NS, Jiang R, Herrington DM, Jacobs DR (2006) Dietary patterns are associated with biochemical markers of inflammation and endothelial activation in the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr* 83:1369–1379
 26. Gao X, Bermudez OI, Tucker KL (2004) Plasma C-reactive protein and homocysteine concentrations are related to frequent fruit and vegetable intake in Hispanic and non-Hispanic white elders. *J Nutr* 134:913–918
 27. Holt EM, Steffen LM, Moran A, Basu S, Steinberger J, Ross JA, Hong C-P, Sinaiko AR (2009) Fruit and vegetable consumption and its relation to markers of inflammation and oxidative stress in adolescents. *J Am Diet Assoc* 109:414–421
 28. Kennedy A, Martinez K, Chuang C-C, LaPoint K, McIntosh M (2009) Saturated fatty acid-mediated inflammation and insulin resistance in adipose tissue: mechanisms of action and implications. *J Nutr* 139:1–4
 29. Muñoz A, Costa M (2013) Nutritionally mediated oxidative stress and inflammation. *Oxidative Med Cellular Longev* 2013:610950. doi:[10.1155/2013/610950](https://doi.org/10.1155/2013/610950)
 30. Nicholls SJ, Lundman P, Harmer JA, Cutri B, Griffiths KA, Rye K-A, Barter PJ, Celermajer DS (2006) Consumption of saturated fat impairs the anti-inflammatory properties of high-density lipoproteins and endothelial function. *J Am Coll Cardiol* 48:715–720
 31. Intahphuak S, Khonsung P, Panthong A (2010) Anti-inflammatory, analgesic, and antipyretic activities of virgin coconut oil. *Pharm Biol* 48:151–157
 32. Marten B, Pfeuffer M, Schrezenmeier J (2006) Medium-chain triglycerides. *Int Dairy J* 16:1374–1382
 33. Uribarri J, Woodruff S, Goodman S, Cai W, Chen X, Pyzik R, Yong A, Striker GE, Vlassara H (2010) Advanced glycation end products in foods and a practical guide to their reduction in the diet. *J Am Diet Assoc* 110(911–916):e912
 34. Vlassara H, Striker GE (2011) AGE restriction in diabetes mellitus: a paradigm shift. *Nat Rev Endocrinol* 7:526–539
 35. Sun Q, Shi L, Prescott J, Chiuve SE, Hu FB, De Vivo I, Stampfer MJ, Franks PW, Manson JE, Rexrode KM (2012) Healthy lifestyle and leukocyte telomere length in U.S. women. *PLoS One* 7:e38374
 36. Boccardi V, Esposito A, Rizzo MR, Marfella R, Barbieri M, Paolisso G (2013) Mediterranean diet, telomere maintenance and health status among elderly. *PLoS One* 8:e62781
 37. Crous-Bou M, Fung TT, Prescott J, Julin B, Du M, Sun Q, Rexrode KM, Hu FB, De Vivo I (2014) Mediterranean diet and telomere length in Nurses' Health Study: population based cohort study. *BMJ* 349:g6674
 38. Gu Y, Honig L, Schupf N, Lee J, Luchsinger J, Stern Y, Scarmeas N (2015) Mediterranean diet and leukocyte telomere length in a multi-ethnic elderly population. *AGE* 37:1–13
 39. Nettleton JA, Diez-Roux A, Jenny NS, Fitzpatrick AL, Jacobs DR (2008) Dietary patterns, food groups, and telomere length in the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr* 88:1405–1412
 40. Lee JY, Jun NR, Yoon D, Shin C, Baik I (2015) Association between dietary patterns in the remote past and telomere length. *Eur J Clin Nutr* 69:1048–1052
 41. Tiainen AM, Männistö S, Blomstedt PA, Moltchanova E, Perälä MM, Kaartinen NE, Kajantie E, Kananen L, Hovatta I, Eriksson JG (2012) Leukocyte telomere length and its relation to food and nutrient intake in an elderly population. *Eur J Clin Nutr* 66:1290–1294
 42. Song Y, You N-CY, Song Y, Kang MK, Hou L, Wallace R, Eaton CB, Tinker LF, Liu S (2013) Intake of small-to-medium-chain saturated fatty acids is associated with peripheral leukocyte telomere length in postmenopausal women. *J Nutr* 143:907–914
 43. García-Calzón S, Moleres A, Martínez-González MA, Martínez JA, Zalba G, Martí A (2015) Dietary total antioxidant capacity is associated with leukocyte telomere length in a children and adolescent population. *Clin Nutr* 34:694–699
 44. Leung CW, Laraia BA, Needham BL, Rehkopf DH, Adler NE, Lin J, Blackburn EH, Epel ES (2014) Soda and cell aging: associations between sugar-sweetened beverage consumption and leukocyte telomere length in healthy adults from the National Health and Nutrition Examination Surveys. *Am J Public Health* 104:2425–2431
 45. Marcon F, Siniscalchi E, Crebelli R, Saieva C, Sera F, Fortini P, Simonelli V, Palli D (2012) Diet-related telomere shortening and chromosome stability. *Mutagenesis* 27:49–57
 46. Hou L, Savage SA, Blaser MJ, Perez-Perez G, Hoxha M, Dioni L, Pegoraro V, Dong LM, Zatonski W, Lissowska J, Chow W-H, Baccarelli A (2009) Telomere length in peripheral leukocyte DNA and gastric cancer risk. *Cancer Epidemiol Biomark Prev* 18:3103–3109
 47. O'Callaghan N, Parletta N, Milte CM, Benassi-Evans B, Fenech M, Howe PRC (2014) Telomere shortening in elderly individuals with mild cognitive impairment may be attenuated with ω -3 fatty acid supplementation: a randomized controlled pilot study. *Nutrition* 30:489–491
 48. Chan R, Woo J, Suen E, Leung J, Tang N (2010) Chinese tea consumption is associated with longer telomere length in elderly Chinese men. *Br J Nutr* 103:107–113
 49. Kark JD, Goldberger N, Kimura M, Sinnreich R, Aviv A (2012) Energy intake and leukocyte telomere length in young adults. *Am J Clin Nutr* 95:479–487
 50. Cassidy A, De Vivo I, Liu Y, Han J, Prescott J, Hunter DJ, Rimm EB (2010) Associations between diet, lifestyle factors, and telomere length in women. *Am J Clin Nutr* 91:1273–1280
 51. Bekaert S, De Meyer T, Rietzschel ER, De Buyzere ML, De Bacquer D, Langlois M, Segers P, Cooman L, Van Damme P, Cassiman P, Van Crielinge W, Verdonck P, De Backer GG, Gillebert TC, Van Oostveldt P (2007) Telomere length and

- cardiovascular risk factors in a middle-aged population free of overt cardiovascular disease. *Aging Cell* 6:639–647
52. Dandona P, Aljada A, Bandyopadhyay A (2004) Inflammation: the link between insulin resistance, obesity and diabetes. *Trends Immunol* 25:4–7
 53. Keaney JF, Larson MG, Vasan RS, Wilson PWF, Lipinska I, Corey D, Massaro JM, Sutherland P, Vita JA, Benjamin EJ (2003) Obesity and systemic oxidative stress: clinical correlates of oxidative stress in the Framingham Study. *Arterioscler Thromb Vasc Biol* 23:434–439
 54. Zhu H, Bhagatwala J, Pollock NK, Parikh S, Gutin B, Stallmann-Jorgensen I, Thomas J, Harshfield GA, Dong Y (2015) High sodium intake is associated with short leukocyte telomere length in overweight and obese adolescents. *Int J Obes* 39:1249–1253
 55. Benetos A, Kark JD, Susser E, Kimura M, Sinnreich R, Chen W, Steenstrup T, Christensen K, Herbig U, von Bornemann Hjelmberg J, Srinivasan SR, Berenson GS, Labat C, Aviv A (2013) Tracking and fixed ranking of leukocyte telomere length across the adult life course. *Aging Cell* 12:615–621
 56. Milne E, O'Callaghan N, Ramankutty P, de Klerk NH, Greenop KR, Armstrong BK, Miller M, Fenech M (2015) Plasma micronutrient levels and telomere length in children. *Nutrition* 31:331–336
 57. Buxton JL, Walters RG, Visvikis-Siest S, Meyre D, Froguel P, Blakemore AI (2011) Childhood obesity is associated with shorter leukocyte telomere length. *J Clin Endocrinol Metab* 96:1500–1505
 58. Al-Attas OS, Al-Daghri N, Bamakhramah A, Shaun Sabico S, McTernan P, Huang T-K (2010) Telomere length in relation to insulin resistance, inflammation and obesity among Arab youth. *Acta Paediatr* 99:896–899
 59. Buxton JL, Das S, Rodriguez A, Kaakinen M, Couto Alves A, Sebert S, Millwood IY, Laitinen J, O'Reilly PF, Jarvelin M-R, Blakemore AIF (2014) Multiple measures of adiposity are associated with mean leukocyte telomere length in the northern Finland birth cohort 1966. *PLoS One* 9:e99133
 60. Lee M, Martin H, Firpo MA, Demerath EW (2011) Inverse association between adiposity and telomere length: the Fels Longitudinal Study. *Am J Hum Biol* 23:100–106
 61. Zannolli R, Mohn A, Buoni S, Pietrobelli A, Messina M, Chiarelli F, Miracco C (2008) Telomere length and obesity. *Acta Paediatr* 97:952–954
 62. Zhu H, Wang X, Gutin B, Davis CL, Keeton D, Thomas J, Stallmann-Jorgensen I, Mookken G, Bundy V, Snieder H (2011) Leukocyte telomere length in healthy Caucasian and African-American adolescents: relationships with race, sex, adiposity, adipokines, and physical activity. *J Pediatr* 158:215–220
 63. Adair LS, Popkin BM (2005) Are child eating patterns being transformed globally? *Obes Res* 13:1281–1299
 64. Kelles A, Adair L (2009) Offspring consume a more obesogenic diet than mothers in response to changing socioeconomic status and urbanization in Cebu, Philippines. *Int J Behav Nutr Phys Act* 6:47
 65. Pedro MRA, Barba CV, Benavides-de Leon R (2008) Nutrition transition in the Philippines. *Philippine Population Review* 6:1–19
 66. Adair LS, Gultiano S, Suchindran C (2011) 20-year trends in Filipino women's weight reflect substantial secular and age effects. *J Nutr* 141:667–673
 67. Food and Nutrition Research Institute (2012) Philippine nutrition: facts and figures 2011. Food and Nutrition Research Institute – Department of Science and Technology, Taguig City
 68. Adair LS (2004) Dramatic rise in overweight and obesity in adult Filipino women and risk of hypertension. *Obes Res* 12:1335–1341
 69. Drewnowski A, Popkin BM (1997) The nutrition transition: new trends in the global diet. *Nutr Rev* 55:31–43
 70. Ke-You G, Da-Wei F (2001) The magnitude and trends of under- and over-nutrition in Asian countries. *Biomed Environ Sci* 14:53–60
 71. Adair LS, Popkin BM, Akin JS, Guilkey DK, Gultiano S, Borja J, Perez L, Kuzawa CW, McDade T, Hindin MJ (2011) Cohort profile: the Cebu Longitudinal Health and Nutrition Survey. *Int J Epidemiol* 40:619–625
 72. Feranil A, Gultiano S, Adair L (2008) The Cebu longitudinal health and nutrition survey: two decades later. *Asia Pac Popul J* 23:39–54
 73. Eisenberg DTA, Hayes MG, Kuzawa CW (2012) Delayed paternal age of reproduction in humans is associated with longer telomeres across two generations of descendants. *Proc Natl Acad Sci* 109:10251–10256
 74. Eisenberg DTA, Kuzawa CW, Hayes MG (2015) Improving qPCR telomere length assays: Controlling for well position effects increases statistical power. *Am J Hum Biol* 27(4):570–575. doi:10.1002/ajhb.22690
 75. Cawthon RM (2009) Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Res* 37:e21
 76. Eisenberg DTA (2011) An evolutionary review of human telomere biology: the thrifty telomere hypothesis and notes on potential adaptive paternal effects. *Am J Hum Biol* 23:149–167
 77. World Health Organization (2007) Growth reference 5–19 years. World Health Organization, Geneva
 78. de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 85:660–667
 79. de Onis M, Onyango A, Borghi E, Siyam A, Blossner M, Lutter C (2012) Worldwide implementation of the WHO child growth standards. *Public Health Nutr* 15:1603–1610
 80. Food and Nutrition Research Institute (1997) Food composition tables recommended for use in the Philippines. Manila, Philippines
 81. Feranil AB, Duazo PL, Kuzawa CW, Adair LS (2011) Coconut oil predicts a beneficial lipid profile in pre-menopausal women in the Philippines. *Asia Pac J Clin Nutr* 20:190
 82. Willett W (2013) Nutritional epidemiology. Oxford University Press, New York
 83. Farzaneh-Far R, Lin J, Epel E, Lapham K, Blackburn E, Whooley MA (2010) Telomere length trajectory and its determinants in persons with coronary artery disease: longitudinal findings from the Heart and Soul Study. *PLoS One* 5:e8612
 84. Okuda K, Bardeguet A, Gardner JP, Rodriguez P, Ganesh V, Kimura M, Skurnick J, Awad G, Aviv A (2002) Telomere length in the newborn. *Pediatr Res* 52:377–381
 85. Sanders JL, Newman AB (2013) Telomere length in epidemiology: a biomarker of aging, age-related disease, both, or neither? *Epidemiol Rev* 35:112–131
 86. Croteau-Chonka DC, Marvelle AF, Lange EM, Lee NR, Adair LS, Lange LA, Mohlke KL (2011) Genome-wide association study of anthropometric traits and evidence of interactions with age and study year in Filipino women. *Obesity* 19:1019–1027
 87. Croteau-Chonka DC, Wu Y, Li Y, Fogarty MP, Lange LA, Kuzawa CW, McDade TW, Borja JB, Luo J, AbdelBaky O, Combs TP, Adair LS, Lange EM, Mohlke KL (2012) Population-specific coding variant underlies genome-wide association with adiponectin level. *Hum Mol Genet* 21:463–471
 88. Wu Y, McDade T, Kuzawa C, Borja J, Li Y, Adair L, Mohlke K, Lange L (2012) Genome-wide association with C-reactive protein levels in CLHNS: evidence for the CRP and HNF1A

- loci and their interaction with exposure to a pathogenic environment. *Inflammation* 35:574–583
89. Dahly DL, Adair LS (2007) Quantifying the urban environment: a scale measure of urbanicity outperforms the urban–rural dichotomy. *Soc Sci Med* 64:1407–1419
 90. Valdes AM, Andrew T, Gardner JP, Kimura M, Oelsner E, Cherkas LF, Aviv A, Spector TD (2005) Obesity, cigarette smoking, and telomere length in women. *Lancet* 366:662–664
 91. Cherkas LF, Aviv A, Valdes AM, Hunkin JL, Gardner JP, Surdulescu GL, Kimura M, Spector TD (2006) The effects of social status on biological aging as measured by white-blood-cell telomere length. *Aging Cell* 5:361–365
 92. Needham BL, Adler N, Gregorich S, Rehkopf D, Lin J, Blackburn EH, Epel ES (2013) Socioeconomic status, health behavior, and leukocyte telomere length in the National Health and Nutrition Examination Survey, 1999–2002. *Soc Sci Med* 85:1–8
 93. Epel ES, Blackburn EH, Lin J, Dhabhar FS, Adler NE, Morrow JD, Cawthon RM (2004) Accelerated telomere shortening in response to life stress. *Proc Natl Acad Sci USA* 101:17312–17315
 94. World Health Organization Expert Consultation (2004) Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 363:157
 95. Nordfjäll K, Eliasson M, Stegmayr B, Melander O, Nilsson P, Roos G (2008) Telomere length is associated with obesity parameters but with a gender difference. *Obesity* 16:2682–2689
 96. Müezziner A, Zaineddin AK, Brenner H (2014) Body mass index and leukocyte telomere length in adults: a systematic review and meta-analysis. *Obes Rev* 15:192–201
 97. Dirks AJ, Leeuwenburgh C (2006) Caloric restriction in humans: potential pitfalls and health concerns. *Mech Ageing Dev* 127:1–7
 98. Chung HY, Cesari M, Anton S, Marzetti E, Giovannini S, Seo AY, Carter C, Yu BP, Leeuwenburgh C (2009) Molecular inflammation: underpinnings of aging and age-related diseases. *Ageing Res Rev* 8:18–30
 99. McDade TW, Rutherford JN, Adair L, Kuzawa C (2009) Population differences in associations between C-reactive protein concentration and adiposity: comparison of young adults in the Philippines and the United States. *Am J Clin Nutr* 89:1237–1245
 100. McDade TW, Tallman PS, Adair LS, Borja J, Kuzawa CW (2011) Comparative insights into the regulation of inflammation: levels and predictors of interleukin 6 and interleukin 10 in young adults in the Philippines. *Am J Phys Anthropol* 146:373–384
 101. Hunt SC, Kark JD, Aviv A (2015) Association between shortened leukocyte telomere length and cardio-metabolic outcomes. *Circ Cardiovasc Genet* 8:4–7
 102. Wang S, Meckling KA, Marcone MF, Kakuda Y, Tsao R (2011) Synergistic, additive, and antagonistic effects of food mixtures on total antioxidant capacities. *J Agric Food Chem* 59:960–968
 103. Mirabello L, Huang W-Y, Wong JYY, Chatterjee N, Reding D, David Crawford E, De Vivo I, Hayes RB, Savage SA (2009) The association between leukocyte telomere length and cigarette smoking, dietary and physical variables, and risk of prostate cancer. *Aging Cell* 8:405–413
 104. Livingstone M, Robson P, Wallace J (2004) Issues in dietary intake assessment of children and adolescents. *Br J Nutr* 92:S213–S222
 105. Kristal AR, Peters U, Potter JD (2005) Is it time to abandon the food frequency questionnaire? *Cancer Epidemiol Biomark Prev* 14:2826–2828
 106. Robertson T, Batty GD, Der G, Fenton C, Shiels PG, Benzeval M (2013) Is socioeconomic status associated with biological aging as measured by telomere length? *Epidemiol Rev* 35:98–111