

Is peroxisome proliferator-activated receptor gamma (PPAR γ) a therapeutic target for the treatment of pulmonary hypertension?

David E. Green¹, Roy L. Sutliff¹, C. Michael Hart¹

¹Department of Medicine, Emory University, Atlanta Veterans Affairs Medical Center, Decatur, Georgia, USA

ABSTRACT

Pulmonary hypertension (PH), a progressive disorder associated with significant morbidity and mortality, is caused by complex pathways that culminate in structural and functional alterations of the pulmonary circulation and increases in pulmonary vascular resistance and pressure. Diverse genetic, pathological, or environmental triggers stimulate PH pathogenesis culminating in vasoconstriction, cell proliferation, vascular remodeling, and thrombosis. We conducted a thorough literature review by performing MEDLINE searches via PubMed to identify articles pertaining to PPAR γ as a therapeutic target for the treatment of PH. This review examines basic and preclinical studies that explore PPAR γ and its ability to regulate PH pathogenesis. Despite the current therapies that target specific pathways in PH pathogenesis, including prostacyclin derivatives, endothelin-receptor antagonists, and phosphodiesterase type 5 inhibitors, morbidity and mortality related to PH remain unacceptably high, indicating the need for novel therapeutic approaches. Consequently, therapeutic targets that simultaneously regulate multiple pathways involved in PH pathogenesis have gained attention. This review focuses on peroxisome proliferator-activated receptor gamma (PPAR γ), a member of the nuclear hormone receptor superfamily of ligand-activated transcription factors. While the PPAR γ receptor is best known as a master regulator of lipid and glucose metabolism, a growing body of literature demonstrates that activation of PPAR γ exerts antiproliferative, antithrombotic, and vasodilatory effects on the vasculature, suggesting its potential efficacy as a PH therapeutic target.

Key Words: Peroxisome proliferator-activated receptor gamma, pulmonary hypertension, therapy

INTRODUCTION

Pulmonary hypertension (PH) is a rare and progressive disorder with a prevalence of 15 cases per million.^[1] The hemodynamic definition of PH requires elevation of the mean pulmonary artery pressure >25 mm Hg at rest or 30 mm Hg with exercise and a mean pulmonary-capillary wedge pressure or left ventricular end-diastolic pressure \leq 15 mm Hg. The World Health Organization (WHO) now classifies PH into five groups with similar disease mechanisms, histopathologic features and responses to treatment.^[2] Current concepts suggest that PH pathogenesis involves three primary processes: vasoconstriction, cellular proliferation/vascular remodeling, and thrombosis.^[3] Evolving evidence

suggests that peroxisome proliferator-activated receptor gamma (PPAR γ), a member of the nuclear hormone receptor superfamily of ligand-activated transcription factors, can favorably modulate cellular proliferation, vascular tone and coagulation. This review provides an overview of PPAR γ and considers how PPAR γ impacts these primary processes involved in PH pathogenesis.

A thorough literature review was conducted to identify articles pertaining to PPAR as a therapeutic target for the treatment of PH. We performed MEDLINE searches through PubMed in the National Library of Medicine to identify relevant articles. Articles accepted for review

Address correspondence to:

Dr. C. Michael Hart
Atlanta VA Medical Center (151),
1670 Clairmont Road,
Decatur, Georgia 30033, USA
E-mail: michael.hart3@va.gov

Access this article online

Quick Response Code:



Website: www.pulmonarycirculation.org

DOI: 10.4103/2045-8932.78101

Pulm Circ 2011;1:33-47

included full text journal articles, clinical trials, reviews, guidelines, and randomized controlled trials. Articles not accepted for review included letters, editorials and correspondence, and manuscripts published before 1985.

PPAR γ AND PH

Overview of PPAR γ biology

Peroxisome proliferator-activated receptors (PPARs) belong to subfamily 1 of the nuclear hormone receptor superfamily (Nuclear Receptors Nomenclature Committee, 1999). There are three distinct isoforms of PPARs (α , β/δ , and γ) that are expressed throughout the body and possess similar structural and functional features. PPARs are responsive to the lipid status of the cell and act as ligand-activated transcription factors. PPAR isoforms are distinguished by their tissue distribution, ligand specificity and target genes. PPAR α is most highly expressed in tissues with high catabolic rates such as the liver, kidney, heart, brown adipose tissue, and the intestine.^[4,5] PPAR β/δ has the broadest expression pattern with the expression in many cases dependent on the degree of cell proliferation and differentiation.^[4,6,7] PPAR γ exists as two isoforms that differ in their N terminus. PPAR γ 2 is found in adipose and PPAR γ 1 is more widely expressed in tissues such as the brain, vascular tissues, small intestine, and lymphatic tissues.^[4,8] Each PPAR isoform is preferentially activated by a wide range of naturally occurring or metabolized lipids derived from the diet or from intracellular signaling pathways, which include n-3 and n-6 family polyunsaturated fatty acids (PUFAs) and eicosanoid products of cyclooxygenase and lipoxygenase.

As illustrated in Figure 1, transcriptional regulation by PPARs requires the formation of heterodimers with retinoid X receptor (RXR) isoforms. PPARs typically form heterodimers with RXR α (RXR α). PPAR:RXR heterodimers bind to DNA at sites composed of the hexameric direct repeat sequence, AGGTCA, separated by a single nucleotide. Gene regulation involves ligand-induced conformational changes in the PPAR receptor which mediate interaction with specific coactivator (e.g. steroid receptor coactivator-1 and p300) and corepressor molecules. Coactivator proteins possess either histone acetyltransferase activity or recruit other proteins with this activity to the transcription start site. Acetylation of histone proteins alters chromatin structure, facilitating the binding of RNA polymerase and the initiation of transcription.^[9] PPAR γ can also repress gene expression by interfering with the clearance of corepressors from selected promoters.^[10] Due to their numerous metabolic and therapeutic actions, PPARs have become pharmacological targets. For example, PPAR α agonists

such as fibrate medications are used in the treatment of dyslipidemia and may have anti-inflammatory effects. PPAR β/δ selective agonists such as GW0742 augment high density lipoprotein (HDL) cholesterol and have been implicated in promoting tissue repair.^[9-12] On the other hand, PPAR γ agonists such as rosiglitazone and pioglitazone, used to treat type 2 diabetes, have beneficial effects in vascular disease in animal models.^[11,12] These reports emphasize that ligands for PPAR receptors can exert diverse biological effects in numerous organs and tissues. Among these PPAR receptors, PPAR γ is of particular interest in the pulmonary circulation.

The role of altered PPAR γ expression in PH

PPAR γ is abundantly expressed in many cell types in the lung, including those of the pulmonary vascular wall such as endothelial and smooth muscle cells. Several studies indicate that expression of PPAR γ is reduced by conditions associated with PH. Ameshima and colleagues were the first to demonstrate that compared to normal or chronic obstructive pulmonary disease (COPD) patients, patients with PH had significantly reduced or absent expression of PPAR γ in precapillary arteriolar plexiform lesions.^[13] PPAR γ expression was markedly reduced in endothelial cells isolated from PH patients when compared to normal patients (unpublished observation). These clinical findings have also been corroborated using *in vivo* experimental models of PH. For example, PPAR γ expression was reduced in pulmonary vascular lesions in the rat model of hypoxia-induced PH.^[13,14] Similarly, using *in vitro* cell culture models, increased shear




Figure 1: Schematic illustration of PPAR γ -mediated gene regulation. PPAR γ ligands originating from outside or inside the cell bind to the PPAR γ receptor, stimulating formation of a heterodimer with the retinoid X receptor (RXR). The PPAR γ -RXR heterodimer binds to PPAR response elements (PPRE) comprising hexameric repeats of the sequence, AGGTCA-X-AGGTCA. This active heterodimer recruits coactivators to and/or derecruits corepressor molecules from the transcriptional start site, leading to increased transcription of selected genes. Activation of PPAR γ can also inhibit the expression of genes regulated by specific proinflammatory transcription factors such as NF- κ B, AP-1, and STAT

stress or hypoxia was demonstrated to directly alter PPAR γ expression. Exposure of ECV304 endothelial cells to increased fluid shear stress decreased PPAR γ expression.^[13] Similarly, exposure of endothelial cells to 1% hypoxia decreased expression of PPAR γ .^[15] Collectively, these findings suggest that PPAR γ expression is reduced in PH and that cells exposed to conditions that promote PH have decreased PPAR γ expression. These reductions in PPAR γ could contribute to an abnormal, proliferative, and apoptosis-resistant endothelial cell phenotype.

To further examine the role of PPAR γ in pulmonary vascular biology, more recent studies have employed PPAR γ knockout mice. Because global deletion of PPAR γ results in embryonic lethality,^[16] investigators have examined experimental animals with tissue-targeted deletion of PPAR γ . For example, Guignabert and co-workers reported that targeted deletion of PPAR γ in the vascular endothelium of mice (ePPAR $\gamma^{-/-}$) results in spontaneous PH with right ventricular hypertrophy and muscularization of small distal pulmonary arteries.^[17] The ePPAR $\gamma^{-/-}$ mice exposed to chronic hypoxia (10% O $_2$) for 3 weeks developed a similar degree of PH as wild-type control mice. However, following cessation of hypoxia, PH persisted longer in the ePPAR $\gamma^{-/-}$ mice compared to wild-type mice exposed to hypoxia, suggesting that reduced endothelial PPAR γ signaling is sufficient to cause mild PH and impair recovery from chronic hypoxia exposure.^[17] Targeted deletion of PPAR γ from smooth muscle (smPPAR $\gamma^{-/-}$) also resulted in spontaneous PH in mice.^[18] Microarray analysis of bovine pulmonary artery endothelial cells following treatment with a PPAR γ antagonist revealed alterations in the expression of numerous genes including those that might stimulate cell cycle progression and proliferation.^[19] Taken together, these reports suggest that loss of PPAR γ function in pulmonary vascular wall cells stimulates PH pathogenesis.

PPAR γ activation ameliorates experimental PH

Mounting experimental evidence indicates that PPAR γ stimulation ameliorates PH development in animal models of PH. Monocrotaline (MCT)-induced PH and vascular remodeling in the rat were attenuated by treatment with the PPAR γ ligands, pioglitazone or troglitazone.^[20] Interestingly, PPAR γ ligands also inhibited MCT-induced vascular wall thickening and staining for proliferating cell nuclear antigen, suggesting that PPAR γ ligands suppressed cell proliferation and vascular remodeling.^[20] In Wistar-Kyoto rats exposed to continuous hypobaric hypoxia for 3 weeks, treatment with rosiglitazone attenuated hypoxia-induced right ventricular hypertrophy and vascular smooth muscle cell (VSMC) proliferation, as well as pulmonary vascular collagen and elastin deposition, infiltration of c-Kit-positive cells into the adventitia, and matrix metalloproteinase-2 (MMP-2) activity. In this

study, rosiglitazone failed to attenuate hypoxia-induced increases in pulmonary artery pressure, an observation attributed to the inability of PPAR γ ligands to modulate Rho kinase signaling, a critical mediator of pulmonary vasoconstriction.^[21] Hansmann and colleagues reported that ApoE knockout mice fed high fat diets developed significant increases in right ventricular systolic pressure, pulmonary vascular remodeling and right ventricular hypertrophy and that administration of PPAR γ ligands in this model attenuated PH.^[22] An elegant series of experiments in this model provided evidence that PPAR γ ligands attenuated PH by inhibiting platelet derived growth factor (PDGF) signaling.

Male C57Bl/6 mice exposed to chronic hypoxia (10% O $_2$) for 3 weeks developed PH that was attenuated by treatment with the PPAR γ agonist, rosiglitazone (10 mg/kg/day by gavage) during the final 10 days of hypoxia exposure.^[15] Rosiglitazone treatment also reduced hypoxia-induced right ventricular hypertrophy and muscularization of small pulmonary arterioles. From a therapeutic perspective, this study also demonstrated that rosiglitazone could reverse the established PH by introducing rosiglitazone treatment only after animals developed PH.^[15] The mechanisms of these therapeutic effects were attributed to PPAR γ -mediated reductions in Nox4 expression, oxidative stress, and PDGF signaling in the lung. Collectively, these reports indicate that PPAR γ ligands attenuated pulmonary vascular remodeling and hypertension caused by a variety of stimuli in experimental models. The effect of alterations in PPAR γ expression and activation on PH in various experimental models is summarized in Table 1. These studies have begun to identify specific pathways modulated by PPAR γ in the pathogenesis of PH. Additional evidence, reviewed below, suggests that PPAR γ has the potential to regulate a diverse spectrum of pathways and mediators implicated in PH. The ensuing section will consider how targeting PPAR γ can potentially modulate additional pathways fundamental to the pathogenesis of PH.

ASPECTS OF PH PATHOGENESIS THAT ARE POTENTIALLY REGULATED BY PPAR γ

Vascular tone in PH

Common mechanisms in the pathogenesis of PH are endothelial dysfunction, reduced endothelial production of vasodilators, and increased production of vasoconstrictors leading to vasoconstriction and increased pulmonary vascular resistance. Recent advances in PH therapeutics have sought to promote vascular function by restoring vasodilator levels. For example, circulating levels of the

Table 1: Current evidence linking PPAR γ and pulmonary hypertension

Author	Year	Ref. No.	Observation
Decreased PPAR γ expression is associated with PH			
Ameshima	2003	13	PPAR γ expression reduced in lung tissue from PH patients
Hansmann	2008	18	Shear stress decreased PPAR γ expression in ECV304 cells
Guignabert	2009	17	Targeted smooth muscle PPAR γ deletion in mice caused PH
Nisbet	2010	15	Targeted endothelial PPAR γ deletion in mice caused PH
Kim	2010	14	Hypoxia decreased PPAR γ in HPASMC and HPAEC
PPAR γ activation reduced PH			
Matsuda	2005	19	Hypoxia decreased lung PPAR γ in rat model
Crossno	2007	20	MCT-induced PH in rats attenuated by PPAR γ ligands
Hansmann	2008	22	Hypoxia-induced vascular remodeling in rats attenuated by PPAR γ ligands
Nisbet	2010	15	High fat diet-induced PH in Apo-E deficient mice attenuated by PPAR γ ligands
Kim	2010	14	Hypoxia-induced PH and vascular remodeling in mice attenuated by PPAR γ ligands
Kim	2010	14	Hypoxia-induced PH in rats attenuated by PPAR γ ligands

vasodilator, prostacyclin, are decreased in PH,^[23] and the administration of prostacyclin or its analogues represents a significant advance in PH therapy.^[24] Nitric oxide (NO)-mediated vasorelaxation is also impaired in PH.^[25] Phosphodiesterase type 5 inhibitors which prolong NO-mediated increases in cGMP are now employed in selected patients with PH.^[24] Conversely, levels of vasoconstrictors such as endothelin-1 (ET-1) and thromboxane (TXA₂) are increased in PH patients.^[26] By blocking these vasoconstrictive effects, both endothelin receptor antagonists and calcium channel blockers comprise additional strategies in the current PH therapeutic armamentarium. In addition to altering vascular tone, these agents may also modulate structural changes in the pulmonary vasculature and vascular remodeling, sequelae of vascular injury and increased intraluminal pressure or flow. While the current therapies target individual mediators or mechanisms in PH pathogenesis, PPAR γ may simultaneously modulate several of these pathways involved in PH pathogenesis as described below.

Prostacyclin

As a critical regulator of pulmonary vascular function, the endothelial-derived mediator, prostacyclin, is a potent vasodilator that inhibits platelet aggregation and exerts anti-inflammatory, anti-thrombotic, and anti-proliferative vascular effects.^[27] Overexpression of prostacyclin synthase protected mice from chronic hypoxia-induced PH, whereas prostacyclin-receptor deficient mice were sensitized to hypoxia-induced PH.^[28] Prostacyclin synthase expression was reduced in the pulmonary arteries of patients with severe PH compared to normal subjects, and the vascular endothelium was found to be the major site of lung vascular prostacyclin synthase expression.^[29] In patients with PH, prostacyclin derivatives decreased urinary isoprostane metabolites, an index

of oxidative stress, without altering TXA₂.^[30] Currently, augmenting prostacyclin levels constitutes a therapeutic strategy in PH, but the precise cellular mechanisms responsible for prostacyclin-mediated benefits remain to be defined. The classical signaling pathway activated by prostacyclin involves binding the G-protein coupled cell surface prostacyclin receptor (IP), which when activated, stimulates adenyl cyclase and increases cellular cAMP content. However, prostacyclin and its analogues can also activate PPAR receptors including PPAR δ and PPAR γ to mediate biological effects.^[31,32] Activation of the PPAR γ receptor with thiazolidinedione also reduced systemic vascular production of the potent vasoconstrictor, thromboxane,^[33] and attenuated iNOS and Cox-2 upregulation.^[34] These reports emphasize that PPAR γ can regulate prostanoid production and that additional studies will be required to determine if PPAR γ can regulate prostacyclin and its metabolites in the pulmonary circulation.

Nitric oxide

In addition to prostacyclin, NO represents an additional endothelium-derived vasodilator whose bioavailability is reduced in PH.^[35,36] Downregulation of the enzyme that produces nitric oxide, endothelial nitric oxide synthase (eNOS), has been described in PH in some studies,^[37] whereas others report unchanged or increased levels of the enzyme. (More consistent evidence demonstrates that endothelium-derived, NO-mediated vasodilation is impaired in models of PH.) Thus, it is likely that reductions in NO bioavailability in PH are more closely related to post-translational alterations in eNOS regulation and/or enhanced NO degradation rather than reduced eNOS expression. The critical role of NO bioavailability in PH is supported by the evidence that genetic deletion of eNOS enhanced susceptibility to hypoxia-induced PH,^[38]

a defect reversed by adenoviral-mediated transfection of the pulmonary vasculature with eNOS.^[39] Furthermore, overexpression of eNOS attenuated hypoxia-induced PH.^[40] In addition, NO inhalation improves pulmonary hemodynamics and quality of life in a subset of patients with PH,^[41] and recent advances in cell-based eNOS gene transfer to the lung have demonstrated that eNOS can reverse the established PH in animal models and facilitate restoration of the pulmonary microvasculature.^[42]

NO is produced constitutively in vascular endothelial cells from the amino acid, L-arginine, by the Type III, eNOS isoform. Enzyme activity is largely regulated by intracellular Ca²⁺, cofactor availability, and eNOS post-translational modifications including phosphorylation and interaction with other proteins such as caveolin and heat shock protein 90 (hsp90).^[43] Derangements in eNOS phosphorylation and interactions between eNOS and caveolin have been reported in PH. PPAR γ ligands can affect these post-translational pathways.^[44,45] PPAR γ increased endothelial NO release by reducing the inhibitory interaction between eNOS and caveolin, increasing interactions between eNOS and the molecular chaperone, hsp90, and enhancing phosphorylation of eNOS on serine 1177, all post-translational modifications associated with enhanced eNOS activity.^[46,47] PPAR γ agonists can increase NO bioavailability by blunting the degradation of inducible nitric oxide synthase (iNOS) and by decreasing serum levels of asymmetric dimethylarginine (ADMA), the endogenous inhibitor of NOS.^[48-50]

Once produced, endothelial-derived NO reduces: 1) vascular tone,^[51] 2) platelet activation and aggregation,^[52] 3) stimulated vascular smooth muscle proliferation,^[53] and 4) leukocyte adherence.^[54] To exert these biological effects, NO must diffuse into the vascular wall where its activity may be limited by local concentrations of superoxide. Superoxide combines at diffusion-limited rates with NO, forming the potent oxidant, peroxynitrite, thereby reducing the vascular protective effects of NO and enhancing oxidative stress. Peroxynitrite also oxidizes the NOS cofactor, tetrahydrobiopterin.^[55] Deficiency of tetrahydrobiopterin alters electron flow through eNOS to molecular oxygen rather than arginine, producing superoxide rather than NO, a condition referred to as eNOS uncoupling. Enhanced superoxide production in the vascular wall may, therefore, reduce NO bioavailability through multiple mechanisms. As a result, NO bioavailability can be regulated not only by the rate of NO formation, but also by the rate of NO degradation. Thus, increased superoxide generation constitutes an important mechanism of NO inactivation and endothelial dysfunction in the vascular wall. Current evidence indicates that PPAR γ can stimulate endothelial NO release and simultaneously

reduce superoxide generation in vascular endothelial cells, suggesting that PPAR γ ligands could enhance NO bioavailability and the vascular protective effects of NO to favorably modulate vasoconstriction and PH.^[46,47]

Endothelin-1

The potent vasoconstricting polypeptide, ET-1, has been implicated in PH pathogenesis. ET-1 receptors are upregulated in the lung in both animal models and patients with PH.^[25] ET-1, as well as endothelium-derived reactive oxygen species (ROS), attenuated NO-dependent pulmonary vasodilation, following exposure to chronic hypoxia in isolated rat lungs.^[56] ET-1-induced pulmonary vasoconstriction was markedly reduced by administration of Cu/Zn superoxide dismutase and was completely attenuated in gp91phox deficient mice.^[57] These findings suggest that NADPH oxidase and superoxide play an important role in pulmonary vascular effects of ET-1. ET-1 receptor antagonists have been employed in patients with PH to improve functional status and other indices of PH related morbidity,^[56] further suggesting that ET-1 is an important mediator of pulmonary vascular dysregulation. Emerging evidence in several disease states indicates that PPAR γ activation attenuates ET-1 signaling. PPAR ligands inhibited ET-1 secretion by vascular endothelial cells *in vitro*.^[57-60] Similarly, in non-diabetic patients with metabolic syndrome, treatment with rosiglitazone reduced several markers of vascular inflammation including plasma levels of ET-1 and improved markers of metabolic control, while lowering blood pressure and improving flow-mediated vasodilation.^[61] PPAR γ activation also reduced hypertrophy and anti-apoptotic effects caused by ET-1 in cardiac myocytes *in vitro* through altered nuclear factor of activated T cells (NFAT) signaling. Treatment with PPAR γ ligands in several rat models of hypertension reduced ET-1 expression in cardiac and vascular tissues.^[62] Collectively, these findings suggest that PPAR γ activation can attenuate expression of ET-1 in cardiovascular tissues in response to a variety of stimuli and can attenuate ET-1-mediated signaling in selected models. Potential pathways by which PPAR γ regulates vascular tone are illustrated in Figure 2.

Abnormal vascular remodeling, inflammation, and cell proliferation in PH

Remodeling of the pulmonary vasculature can reduce its cross-sectional diameter and compliance, increase pulmonary vascular resistance, and contribute to sustained PH. Studies examining the molecular mechanisms underlying pulmonary vascular remodeling have implicated growth factor pathways as well as matrix remodeling in the development and progression of PH.^[63] Inflammation also plays a significant role in altered pulmonary vascular function during the development

of PH.^[64] Inflammatory markers are elevated in PH, and the plexiform lesions that characterize severe PH are surrounded by macrophages, T and B lymphocytes, and dendritic cells.^[65] These cells may exacerbate PH by releasing growth factors, ROS and additional cytokines.^[66] Chemokines such as CX3CL1, CCL5 and MCP-1 which recruit inflammatory cells are elevated in PH patients.^[67-69] Therefore, agents that target the generation of these oxidative and inflammatory stimuli in the pulmonary vascular wall may reduce vascular dysfunction and attenuate the development or progression of PH. Relationships between PPAR γ and these complex pathways implicated in PH pathogenesis are illustrated in Figure 3.

NADPH oxidases

NADPH oxidases, a major source of superoxide production in the vasculature, have been implicated in PH and contribute to endothelial dysfunction and vascular cell proliferation.^[70,71] Originally described in phagocytic cells, the gp91phox-based NADPH oxidase is a multicomponent, membrane-associated, enzyme that catalyzes the one electron reduction of oxygen to superoxide, using NADH or NADPH as the electron donor.^[70] The classical phagocytic NADPH oxidase is composed of several components or subunits including the membrane-bound gp91phox (Nox2) and p22phox subunits as well as the cytosolic p47phox and p67phox subunits that, when stimulated, combine with the small G-protein, rac, and translocate to the membrane to activate the enzyme complex. On the other hand, in nonphagocytic cells, the catalytic moiety of NADPH oxidases is composed of one or more Nox2 homologues, Nox 1, 3, 4, 5, Duox1 or Duox2.^[72] These Nox homologues associate with the membrane-bound p22phox subunit and are differentially regulated and targeted to distinct subcellular loci, suggesting that these oxidases serve unique roles in cell function. Nox1 and 3 are activated through interactions with rac and the p47phox and p67phox homologues, NOXA1 and NOXO1.

Current evidence indicates that Nox4 expression is increased in hypoxia-induced PH in the mouse and in the pulmonary vasculature of patients with PH.^[15,73] Nox4 is highly expressed in vascular wall cells including smooth muscle and endothelial cells where it is constitutively active.^[74] Furthermore, hypoxia increased Nox4 expression and pulmonary artery smooth muscle cell (PASMC) proliferation,^[73] and Nox4 stimulated the proliferation of endothelial and smooth muscle cells.^[75] Hypoxia stimulated Nox4 expression and cell proliferation in the mouse lung *in vivo* and in human pulmonary artery endothelial cell (HPAEC) and human pulmonary artery smooth muscle cell (HPASMC) *in vitro*.^[15] Treatment with rosiglitazone during the last 10 days of hypoxia exposure reduced Nox4 levels and ROS production and attenuated

hypoxia-induced PH, right ventricular hypertrophy and vascular remodeling. Similarly, rosiglitazone attenuated hypoxia-induced Nox4 expression and proliferation in human PAEC and PASMC *in vitro*. Taken together, these findings suggest that NADPH oxidases are important mediators of cell proliferation and vasoconstriction in PH that can be regulated by PPAR γ [summarized in Figure 4].




Figure 2: Potential pathways by which PPAR γ regulates vascular tone. Activation of PPAR γ potentially modulates several pathways involved in the regulation of vascular tone. In addition, prostacyclin or its analogues may activate PPAR γ . Arrowheads at the end of lines denote stimulatory effects, whereas perpendicular lines denote inhibitory effects. (ADMA – Asymmetric dimethyl arginine; cAMP – Cyclic adenosine nucleotide monophosphate; cGMP – Cyclic guanosine nucleotide monophosphate; eNOS – Endothelial nitric oxide synthase; ET-1 – Endothelin-1; ET-R – Endothelin-1 receptor; NO – Nitric oxide; IP receptor – Prostacyclin receptor)




Figure 3: Potential pathways by which PPAR γ regulates vascular remodeling, cell proliferation and hypertrophy in PH. PPAR γ has the potential to regulate a complex variety of pathways that are involved in remodeling of the pulmonary vasculature during the development of PH. Arrowheads at the end of lines denote stimulatory effects, whereas perpendicular lines denote inhibitory effects. *Pathway components that are inhibited by PPAR γ , *pathway components that are stimulated by PPAR γ . (BMP-2 – Bone morphogenetic protein-2; BMPR-2 – BMP-2 receptor; EPC – endothelial progenitor cell; ET-1 – Endothelin-1; ET-R – Endothelin-1 receptor; ROS – Reactive oxygen species; TF – Inflammatory transcription factors (e.g. NF- κ B); TGF- β – Transforming growth factor beta; TGF β R – TGF- β receptor)

Although increases in Nox4 mRNA levels were associated with increased Nox4 activity,^[76] detailed understanding of Nox4 transcriptional regulation remains to be established. Nox4 induction has been reported in response to diverse stimuli including hypoxia in kidney and ischemia in brain.^[71] In smooth muscle cells, activators of Nox4 transcription include urokinase, plasminogen activator, angiotensin II, transforming growth factor-beta 1 (TGF)- β 1, and tumor necrosis factor (TNF)- α .^[77] In contrast, in endothelial cells, oscillatory shear stress^[78] and PPAR γ activation^[79] suppressed Nox4 mRNA levels. However, few studies have examined regulatory elements in the Nox4 promoter. Hypoxia stimulated activation of the Nox4 promoter in part through nuclear factor (NF)- κ B-mediated signaling and enhanced p65 binding to the Nox4 promoter which increased Nox4 expression and activity.^[80] Furthermore, treatment with rosiglitazone inhibited hypoxia-induced Nox4 expression and activity, proliferation, and p65–Nox4 promoter interaction.^[80] HIF-1 α and the E2F family transcription factors also activated the Nox4 promoter although their regulation by PPAR γ at this site has yet to be confirmed.^[81] Based on evidence that Nox4 stimulates smooth muscle and endothelial cell proliferation, these findings suggest that activation of PPAR γ can attenuate the proliferation of pulmonary vascular wall cells that may participate in the pathogenesis of PH.^[82] Coupled with the ability of NADPH oxidases to attenuate NO bioavailability, Nox4, in particular, and NADPH oxidases, in general, may be an important target of PPAR γ amelioration of PH.

PDGF signaling

PDGF participates in PH pathogenesis. Two genes (A and




Figure 4: The effects of PPAR γ activation on chronic hypoxia-induced PH and PDGF signaling. PPAR γ activation decreases oxidative stress and hypoxic vasoconstriction by blunting hypoxia-induced increases NADPH oxidase expression and decreases in NO, respectively. PPAR γ stimulated PTEN expression blocks PDGF-mediated vascular smooth muscle cell proliferation. Arrowheads at the end of lines denote stimulatory effects, whereas perpendicular lines denote inhibitory effects

B) produce three biologically active forms of PDGF protein (AA, AB, and BB).^[83,84] These proteins activate one or more PDGF receptors ($\alpha\alpha$, $\alpha\beta$, or $\beta\beta$) to stimulate cell migration and survival. Ligand binding promotes PDGF receptor tyrosine autophosphorylation and subsequent activation of several downstream signaling pathways including Src, phosphatidylinositol 3 kinase (PI3K), phospholipase C γ , and Ras. These signaling pathways are largely activated by recruitment of these enzymes to PDGF-R SH-2 domains. The composition of the downstream signaling pathways activated by PDGF and their integration into specific cellular responses continue to be defined.^[85] Although the expression of PDGF and its receptors is limited in the vascular wall at baseline, several pathological stimuli, including alterations in blood pressure and shear stress, induce peptide and receptor expression.^[84] PDGF receptor expression was increased in the lungs of patients with PH, and the PDGF receptor antagonist, imatinib, reversed MCT- or hypoxia-induced PH in rodents and improved pulmonary vascular resistance and exercise capacity in a patient with severe idiopathic PH.^[86-88] PPAR γ ligands attenuated hypoxia-induced PDGF activation in a mouse model of PH *in vivo*.^[15] Coupled with reports that NO inhibits PDGF signaling and that PPAR γ ligands inhibit PDGF-stimulated VSMC migration *in vitro*,^[89-91] these studies suggest that PPAR γ can regulate important proliferative signaling pathways in experimental PH, including those activated by PDGF.

PDGF receptor activity can also be regulated by phosphatase and tensin homologue deleted on chromosome 10 (PTEN), a dual specificity phosphatase that dephosphorylates both lipid and protein substrates.^[92] PTEN catalyzes the removal of the phosphate moiety from the 3-position of the phosphatidylinositol ring, converting the second messenger, PI(3,4,5)P $_3$, to PI(4,5)P $_2$.^[93] PTEN and PI3K thereby have opposing actions on cellular levels of PI(3,4,5)P $_3$. PTEN also dephosphorylates the PDGF receptor. Because PDGF receptor activation mediates cell proliferation and migration in part through stimulation of PI3K, PTEN can inhibit PDGF signaling both at the receptor and through lowering of PI(3,4,5)P $_3$ levels, thereby lowering the activity of PI3K-related downstream mediators such as the protein kinase B, Akt, which mediates survival, growth, and proliferative signals by inhibiting apoptosis.

PTEN activity is regulated at the transcriptional and post-translational level. At the transcriptional level, pathological stimuli such as ischemia reduced PTEN expression and promoted hypertrophy and remodeling.^[94,95] Limited evidence suggests that NF- κ B activation may lead to suppression of PTEN expression.^[96,97] On the other hand, the PTEN promoter contains two PPAR response elements, and several studies demonstrated that PPAR γ ligands

stimulated PTEN expression.^[98,99] Furthermore, PTEN overexpression reduced VSMC proliferation and migration and inhibited injury-induced vascular remodeling *in vivo*.^[100,101] PTEN activity is inhibited by ROS which reversibly oxidize cysteine residues in the phosphatase active site.^[102,103] In fact, NADPH oxidase-derived ROS facilitated PDGF signaling by inhibiting PTEN.^[104] These reports suggested that chronic hypoxia caused sustained PI3K/Akt activation, in part, through the generation of NADPH oxidase-derived ROS that stimulated PDGF and inhibited PTEN signaling pathways. A recent report confirmed that chronic hypoxia increased PDGF receptor activation and reduced PTEN expression in the lung.^[15] Furthermore, treatment with rosiglitazone attenuated hypoxia-induced PDGF receptor activation and restored PTEN levels in hypoxic mice to levels comparable to control animals. The ability of PPAR γ ligands to simultaneously stimulate PTEN expression and lower oxidative stress-induced PTEN inactivation while attenuating PH provides a unique strategy to lower PDGF receptor phosphorylation and activation and reduce cellular PI(3,4,5)P $_3$ levels. These integrated effects may contribute to the ability of PPAR γ to attenuate pulmonary VSMC proliferation and hypertrophy as well as vascular remodeling caused by chronic hypoxia. A hypothetical schema depicting relationships between PPAR γ , NADPH oxidase, PDGF, and PTEN is provided in Figure 4.

ALTERED BONE MORPHOGENETIC PROTEIN (BMP) AND TGF- β 1 SIGNALING IN PH

BMP and TGF- β 1 belong to the TGF beta superfamily of growth factors which are involved in multiple cellular processes including proliferation, differentiation, inflammation, and immunity.^[105,106] All TGF- β superfamily ligands are generated as inactive dimeric precursor proteins that are subsequently cleaved by proteases, activated and secreted.^[105] The ligands bind to one of two types of serine/threonine kinase receptors (type I and II) causing Smad substrate recruitment, phosphorylation, transduction of intracellular signals and nuclear translocation.^[106-108] Approximately 70% of patients with familial pulmonary arterial hypertension and 11–40% with idiopathic pulmonary arterial hypertension (IPAH) have germline mutations in bone morphogenetic protein receptor-2 (BMPR-2), a receptor needed for normal vascular development.^[109] BMPR-2 expression is decreased in some PH cases without identified BMPR mutation.^[110] In comparison to unaffected patients, IPAH patients have altered cellular growth responses to TGF- β 1 and BMP signals which favor VSMC proliferation.^[106,111] Specifically, in IPAH patients, TGF- β 1 induces a heightened SMC

proliferative response and BMP signaling fails to confer an expected antiproliferative and proapoptotic effect.^[49] TGF- β 1 signaling is responsible for pulmonary artery remodeling in rat models as evidenced by the attenuation of SMC migration and muscularization of small pulmonary arteries that occurred upon the inhibition of the TGF- β 1 receptor activin like kinase-5 (ALK-5).^[112]

Cell culture experiments reveal that TGF- β 1 promotes smooth muscle proliferation via an autocrine induction of PDGF and Nox4.^[113] Mechanistically, Nox4-derived ROS may cause transient oxidative inactivation of counterbalancing phosphatases involved in kinase-based cell growth cascades, effectively promoting cell cycle transition and proliferation. Nox4 and TGF- β 1 signaling are closely linked and both increase in response to hypoxia and mediate proliferation of HPASMC *in vitro*.^[15,114] Specifically, TGF- β 1 was found to be the proximal mediator of HPASMC proliferation through a cascade involving sequential signaling of phosphatidylinositol 3-kinase (PI3K) and serine/threonine kinase (Akt) phosphorylation, insulin-like growth factor binding protein-3 (IGFBP-3), and Nox 4.^[114] Dominant negative mutations of the TGF- β type II receptor ameliorated hypoxia-induced pulmonary vascular remodeling in mice, supporting the role of TGF- β 1 in promoting PH through enhanced cellular proliferation.^[115,116]

Several lines of evidence suggest that PPAR γ activation can modulate TGF- β 1 signaling through Smad-dependent and Smad-independent mechanisms. For example, both natural (15d-PGJ $_2$) and synthetic (ciglitazone and rosiglitazone) PPAR γ ligands inhibited the profibrotic effects of TGF- β 1 on human lung fibroblasts in a Smad-independent manner as evidenced by blunted myofibroblast differentiation and collagen I synthesis.^[117] Adenoviral PPAR γ gene overexpression similarly decreased fibroblast to myofibroblast differentiation, collagen I and III production in alkali burned mouse corneas.^[118] In this study, PPAR γ overexpression also reduced TGF- β 1 mRNA transcription in fibroblasts, cultured macrophages, and epithelial cells. In other studies, PPAR γ agonists prevented TGF- β 1 induced mesangial and hepatic stellate cell activation and extracellular matrix secretion.^[119,120] Finally, PPAR γ activation inhibited TGF- β 1-mediated Smad 3 phosphorylation and induction of connective tissue growth factor expression, a key regulator of extracellular matrix production and neointima formation following vascular injury.^[121] These findings emphasize that PPAR γ modulation of TGF- β 1 signaling is cell type and context specific and that PPAR γ agonists may blunt abnormal TGF- β 1/BMP-2 signaling and resultant pulmonary vascular remodeling in PH. Additional pathways by which PPAR γ regulates cellular proliferation and vascular remodeling are summarized in Figure 4.

Matrix alterations

PH involves remodeling of pulmonary vessels with muscularization of nonmuscular distal arterioles, proliferation and migration of VSMC, and increased production of extracellular matrix proteins including fibronectin, collagen, and elastin.^[122,123] Alterations in matrix composition may be related to increased matrix degradation resulting from an imbalance in the matrix metalloproteinases-tissue inhibitor of metalloproteinases system (MMP-TIMP) with deposition of collagen, elastin, fibronectin, and tenascin-C.^[122] Inhibition of MMPs and elastase prevented the progression and actually induced regression of vascular remodeling in an experimental model of PH.^[122] In addition, the elastase inhibitor, elafin, protected mice from chronic hypoxia-induced PH.^[124] Recent evidence indicates that the PPAR γ ligand, rosiglitazone, attenuated and reversed vascular remodeling in rat and mouse models of chronic hypoxia-induced PH.^[15,21] Rosiglitazone decreased collagen production and elastin deposition and increased MMP-2 activity. Treatment with PPAR γ ligands has also been associated with attenuation of matrix deposition in disorders other than PH.^[125-127] Taken together, these reports suggest that therapeutic mechanisms of PPAR γ activation in PH could involve attenuation of matrix deposition and remodeling in the pulmonary vasculature.

Inflammation

Several lines of evidence implicate inflammation in the pathobiology of PH. Despite the heterogeneity of disease conditions that lead to PH, similar inflammatory cells types can be found in the plexiform vascular lesions in patients with IPAH and PH associated with connective tissue disease and HIV. Perivascular mononuclear cell infiltrates comprising macrophages, T and B lymphocytes, and dendritic cells are reported in patients with PH.^[64,66,68] Circulating markers of inflammation are also increased in patients with PH.^[128] Proinflammatory cytokines and growth factors such as IL-1, IL-6, PDGF, epidermal growth factor (EGF), and vascular endothelial growth factor (VEGF) are increased and contribute to mitogenic and chemoattractant events in the vascular wall in PH.^[64] Chemokines that mediate inflammation by recruiting leukocytes, monocytes, and T cells to the vascular wall are increased in patients with PH. Chemokines, including fractalkine (CX3CL1), regulated upon activation, normal T-cell expressed and secreted (RANTES), and monocyte chemoattractant protein-1 (MCP-1), also promote pulmonary vascular remodeling in PH.^[64,68] Fractalkine is directly associated with pulmonary artery SMC proliferation in MCT-induced PH in the rat,^[129] and both RANTES^[68] and MCP-1^[130] may indirectly contribute to mitogenesis and vasoconstriction by inducing endothelin converting enzyme-1 (ECE-1) and ET-1.

The stimulation of these inflammatory pathways and their modulation by PPAR γ likely occurs at the level of transcriptional regulation. The transcription factor, NFAT, is upregulated in the pulmonary artery walls and circulating inflammatory cells of PH patients. NFAT, a master activator of T cells, regulates the expression of many inflammatory genes. In PAH, increases in NFAT confer a proliferative and apoptosis resistant phenotype on PASMC by increasing bcl-2, mitochondrial hyperpolarization, and downregulating voltage-gated potassium channels (Kv1.5).^[131] Common stimuli of PH such as hypoxia cause inflammatory responses in the pulmonary vasculature. PPAR γ is expressed in T cells, macrophages, leukocytes, and dendritic cells, suggesting that its activation could modulate inflammation in the pulmonary vasculature that contributes to the development of PH. Studies demonstrate that PPAR γ ligands attenuate inflammation in numerous models.^[132] PPAR γ activation reduced macrophage recruitment and inflammatory mediator production, impaired dendritic cell priming of T cells, as well as T lymphocyte proliferation and viability. PPAR γ ligands also induced regulatory T cells which downregulate immune responses.^[133,134] Current evidence indicates that these anti-inflammatory effects of PPAR γ are mediated not through transactivation of specific target genes, rather through physical binding to other proinflammatory transcription factors such as NF- κ B, AP-1, and STAT, leading to suppression of these proinflammatory transcriptional pathways. The precise mechanisms of these transrepression effects remain to be completely defined but may involve ligand-dependent SUMOylation of PPAR γ which targets the receptor to corepressor complexes on the promoters of inflammatory genes preventing proteasomal degradation of the corepressors and thereby causing inhibition of inflammatory gene expression.^[10]

Apoptosis

Apoptosis is an important process whereby selected cells are programmed for death and removal.^[135] Apoptosis counterbalances cell proliferation and mitotic division in a cascade of events involving caspase activation and protein cleavage, nuclear DNA fragmentation, cytoplasmic condensation, cell fragment sequestration and formation of apoptotic bodies.^[136] Ineffective apoptosis contributes to the pathology of diseases including some malignancies, autoimmunity and persistent infections. Deregulated apoptosis, on the other hand, may be implicated in atherosclerosis and other cardiovascular diseases.^[136] Triggers for apoptosis include, but are not limited to Fas death receptor stimulation by Fas ligand, recognition of cells with DNA damage, abnormal cellular migration and proliferation, ROS, angiotensin type 2, and BMP-2.^[135]

The pulmonary artery SMCs of patients with PAH demonstrated increased tendency to proliferate and exhibited resistance to normal apoptotic signals such as bone morphogenetic protein-2 (BMP-2).^[137] Accordingly, germline mutations in BMPR-2 significantly contribute to disease pathogenesis in 70% of patients with familial PAH.^[109] Impaired apoptosis and proliferation of PASMC may result from increases in the apoptosis inhibitor, survivin, and the sequential occurrences: hyperpolarization of mitochondria, increased HIF-1 α transcription and nuclear translocation, and decreased Kv1.5 voltage-gated potassium channel expression.^[131] These derangements in the regulation of apoptosis in pulmonary vascular wall cells likely contribute to the increased cell proliferation and migration characteristic of the remodeled pulmonary vasculature in PH.^[135] PPAR γ agonists promote apoptosis of proliferating VSMCs in an extracellular signal-related kinase 1/2 (ERK 1/2) - independent manner that may involve interferon regulatory factor-1 (IRF-1) or activation of the proapoptotic genes, p53 and Gadd 45.^[138,139] PPAR γ agonists also induce apoptosis in endothelial cells in a PPAR γ -dependent manner.^[140] Similarly, pulmonary artery endothelial cells derived from patients with IPAH displayed a proliferative, apoptosis-resistant phenotype associated with enhanced STAT 3 signaling, and PPAR ligands reduced STAT 3 signaling.^[141,142] Taken together, these studies indicate that multiple pathways can lead to an apoptosis-resistant phenotype in pulmonary vascular wall cells during the development of PH and that PPAR γ may counterbalance these proliferative pathways by stimulating apoptosis.

Progenitor cell recruitment

Endothelial progenitor cells (EPCs) are bone marrow-derived cells that are mobilized into the systemic circulation in response to ischemia or vascular injury.^[143] The role of EPCs in vascular repair, vascular homeostasis, and formation of new blood vessels continues to be defined.^[144] Clinical trials suggest that EPCs may serve as markers of vascular dysfunction and cardiovascular disease.^[145,146] Because endothelial dysfunction participates in the pathogenesis of PH, EPCs may play a role in pulmonary vascular disease.^[147] The endogenous erythropoietin system recruits EPCs to the lung in experimental PH in mice.^[148] Furthermore, mesenchymal stem cells overexpressing eNOS or EPCs expressing adrenomedullin attenuated MCT-induced PH in rats.^[149] These reports suggest that migration of EPCs to the pulmonary vasculature during experimental PH can exert beneficial effects. PPAR γ agonists facilitate the differentiation of angiogenic progenitor cells into EPC.^[150] PPAR γ agonists also increased the number and migratory activity of EPC in patients with type 2 diabetes and impaired endothelial function and increased EPC migratory activity and reduced EPC apoptosis in

mice.^[151] Collectively, these reports suggest that PPAR γ ligands might stimulate EPC to reduce pulmonary vascular dysfunction and reduce PH.

Thrombosis in PH

Pulmonary artery thrombosis is a common pathologic finding occurring in 48–56% of patients with IPAH.^[152,153] Thrombosis can also be found in other forms of PH associated with collagen vascular disease, HIV, portal hypertension, drugs and anorexigens. Thrombotic lesions in IPAH can be eccentric or concentric and usually occur *in situ* in peripheral muscular arteries where they form lesions that result from mural organization of thrombi.^[49,153] Thrombosis is in large part due to abnormal platelet activation and endothelial dysfunction which results in coagulation and loss of counterbalancing antifibrinolytic mechanisms.^[152,153] Shear stress from elevated pulmonary pressures may contribute to endothelial injury and release of mediators of coagulation.^[49] Accordingly, patients with IPAH have increased serum levels of coagulation mediators including plasminogen activator inhibitor type-1 (PAI-1),^[154] Von Willebrand factor (vWF),^[155] fibrinopeptide A,^[156] and factor VIIIc.^[157] The abnormal platelet activation in PH can be attributed to increases in platelet-derived thromboxane (TXA₂) and decreased levels of endothelial derived prostacyclin (PGI₂) and nitric oxide synthase (eNOS).^[158] Collectively, these imbalances favor coagulation and platelet aggregation which ultimately predisposes patients with PH to the development of thrombosis.

PPAR γ agonists modulate platelets' immunoregulatory and proinflammatory functions and favorably affect vascular patency. Platelets release a variety of proinflammatory mediators and cytokines including prostaglandin E1, TGF- β , IL-1B, PAI-1, and CD 40.^[48] For example, the CD 40 ligand is a transmembrane protein expressed on stimulated CD4+ T cells and platelets,^[159,160] whose expression can be reduced by PPAR γ ligands.^[48,161] CD 40 ligand is associated with platelet activation and increased risk for cardiovascular disease.^[48,159,160] Upon platelet activation, the CD 40 ligand is released and promotes the expression of vasoactive, inflammatory, and thrombotic mediators including cyclooxygenase-2 (COX-2), prostaglandins, TNF- α , IFN- γ , tissue factor, MMPs, selected interleukins, chemokines, and multiple adhesion molecules.^[48,159-161] Thus, CD 40 receptor–ligand binding links platelet activation and vascular inflammation to hemostasis and increased risk for intravascular thrombosis. *In vitro*, thiazolidinediones decreased platelet CD 40 ligand expression and release.^[159] *In vivo* experiments confirmed that diabetic mice treated with pioglitazone had decreased CD 40 expression after platelet activation.^[162] These studies demonstrate that PPAR γ agonists modulate platelet immunoregulatory and hemostatic function by attenuating CD 40 expression.

Since PPAR γ belongs to a nuclear receptor superfamily of transcription factors, historically PPAR γ expression was thought to be limited to nucleated cells. However, Akbiyik and colleagues reported that platelets contain PPAR γ .^[159] Given that platelets have no nucleus and express no PPAR γ mRNA, PPAR γ agonists may affect platelet function through non-transcriptional pathways such as modulation of intracellular signaling or platelet–protein interactions. For example, two studies reported that pioglitazone delayed the onset of iatrogenic arterial thrombosis in animal models.^[163,164] The mechanisms for these PPAR γ effects may include attenuation of platelet activation as evidenced by reductions in soluble and platelet bound P-selectin,^[48,162] CD 40 ligand,^[49,159] and TXA₂,^[49,159] or to direct effects of PPAR γ ligands on vascular endothelium and its production of platelet regulators. PPAR γ agonists also decreased adenosine diphosphate and arachidonic acid-induced platelet aggregation,^[159,163] ATP production,^[163] and increased nitric oxide-mediated activation of fibrinolysis and inhibition of coagulation.^[163,165] Other proposed anti-thrombotic mechanisms involve PPAR γ -induced increases in prostacyclin,^[164] thrombomodulin,^[163] as well as decreases in PAI-1 and fibrinogen.^[166,167] In summary, PPAR γ ligands likely decrease platelet activation and aggregation through direct effects on platelets and cells of the vascular wall that stimulate endothelial and platelet-derived vasodilatory mediators to maintain vascular patency and blood flow.

CONCLUSIONS AND FUTURE DIRECTIONS

The evidence reviewed above suggests that PPAR γ can participate in the regulation of numerous pathways implicated in PH pathogenesis. Activation of PPAR γ can reduce vasoconstriction, vascular remodeling and inflammation, and thrombosis that contribute to the generation and progression of PH. Studies employing animal models of PH have demonstrated that tissue-targeted deletion of PPAR γ in vascular wall cells can promote PH while activation of PPAR γ with exogenous ligands attenuated PH or vascular remodeling in MCT- or hypoxia-induced rodent models of PH.^[14,15,17,20–22] The limitations of these animal models have been recently reviewed emphasizing that the performance of pharmacological tools in the treatment of common experimental models of PH in rodents may not accurately translate to the treatment of human PAH.^[168] The investigation of PPAR γ ligands in recently reported rodent models which more closely reproduce pathological changes seen in the pulmonary arteries of patients with advanced PAH provides an experimental strategy that might circumvent some of the limitations of common existing models.^[169]

Based on the findings in this review, the next step in the evaluation of PPAR γ as a therapeutic target in PH would appear to be clinical trials employing PPAR γ ligands in patients with PH. The abundant evidence reviewed above that PPAR γ ligands can favorably modulate many pathological pathways and mediators involved in PH would support this position. In addition, the current availability of synthetic thiazolidinedione PPAR γ ligands (rosiglitazone and pioglitazone) for the treatment of type 2 diabetes in the United States suggests that clinical trials could be expedited. However, recent findings surrounding potential adverse cardiovascular events in diabetic patients taking rosiglitazone emphasize the need for caution before employing this agent in PH patients with preexisting cardiopulmonary disease.^[170,171] In contrast, clinical studies with pioglitazone in diabetic patients have reported a lowered risk for adverse cardiovascular endpoints.^[172–175] These reports emphasize that individual PPAR γ ligands may regulate unique patterns of gene expression that differentially modulate cell and tissue function. Thus, the therapeutic potential of strategies targeting PPAR γ will be optimized by future studies that must determine not only the relevant molecular pathways that are altered by PPAR γ in a ligand-specific manner, but also the cellular site of action of any ligand and its relative dependence on the PPAR γ receptor. Evidence supporting successful therapy of PH with existing PPAR γ ligands could also stimulate the development of novel pharmacological PPAR γ ligands with enhanced therapeutic efficacy and/or reduced side effects.

ACKNOWLEDGMENTS

The authors acknowledge grant support from the Research Service of the Atlanta Veterans Affairs Medical Center and the National Institutes of Health (R01 DK 074518). Dr. Green is supported by an NHLBI T32 training grant (HL076118-06).

REFERENCES

1. Humbert M, Sitbon O, Chaouat A, Bertocchi M, Habib G, Gressin V, *et al.* Pulmonary arterial hypertension in France: Results from a national registry. *Am J Respir Crit Care Med* 2006;173:1023–30.
2. Simonneau G, Robbins IM, Beghetti M, Channick RN, Delcroix M, Denton CP, *et al.* Updated clinical classification of pulmonary hypertension. *J Am Coll Cardiol* 2009;54:S43–54.
3. McLaughlin VV, Archer SL, Badesch DB, Barst RJ, Farber HW, Lindner JR, *et al.* ACCF/AHA 2009 expert consensus document on pulmonary hypertension: A report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association: Developed in collaboration with the American College of Chest Physicians, American Thoracic Society, Inc., and the Pulmonary Hypertension Association. *Circulation* 2009;119:2250–94.
4. Desvergne B, Wahli W. Peroxisome proliferator-activated receptors: Nuclear control of metabolism. *Endocr Rev* 1999;20:649–88.
5. Mandard S, Müller M, Kersten S. Peroxisome proliferator-activated receptor alpha target genes. *Cell Mol Life Sci* 2004;61:393–416.
6. Barak Y, Liao D, He W, Ong ES, Nelson MC, Olefsky JM, *et al.* Effects of peroxisome proliferator-activated receptor delta on placentalation,

- adiposity, and colorectal cancer. *Proc Natl Acad Sci U S A* 2002;99:303-8.
7. Peters JM, Lee SS, Li W, Ward JM, Gavrilova O, Everett C, *et al.* Growth, adipose, brain, and skin alterations resulting from targeted disruption of the mouse peroxisome proliferator-activated receptor beta(delta). *Mol Cell Biol* 2000;20:5119-28.
8. Zhu Y, Qi C, Korenberg JR, Chen XN, Noya D, Rao MS, *et al.* Structural organization of mouse peroxisome proliferator-activated receptor gamma (mPPAR gamma) gene: Alternative promoter use and different splicing yield two mPPAR gamma isoforms. *Proc Natl Acad Sci U S A* 1995;92:7921-5.
9. Rosen ED, Sarraf P, Troy AE, Bradwin G, Moore K, Milstone DS, *et al.* PPAR gamma is required for the differentiation of adipose tissue *in vivo* and *in vitro*. *Mol Cell* 1999;4:611-7.
10. Pascual G, Fong AL, Ogawa S, Gamliel A, Li AC, Perissi V, *et al.* A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR-gamma. *Nature* 2005;437:759-63.
11. Li AC, Brown KK, Silvestre MJ, Willson TM, Palinski W, Glass CK. Peroxisome proliferator-activated receptor gamma ligands inhibit development of atherosclerosis in LDL receptor-deficient mice. *J Clin Invest* 2000;106:523-31.
12. Xu HE, Lambert MH, Montana VG, Parks DJ, Blanchard SG, Brown PJ, *et al.* Molecular recognition of fatty acids by peroxisome proliferator-activated receptors. *Mol Cell* 1999;3:397-403.
13. Ameshima S, Golpon H, Cool CD, Chan D, Vandivier RW, Gardai SJ, *et al.* Peroxisome proliferator-activated receptor gamma (PPARgamma) expression is decreased in pulmonary hypertension and affects endothelial cell growth. *Circ Res* 2003;92:1162-9.
14. Kim EK, Lee JH, Oh YM, Lee YS, Lee SD. Rosiglitazone attenuates hypoxia-induced pulmonary arterial hypertension in rats. *Respirology* 2010;15:659-68.
15. Nisbet RE, Bland JM, Kleinhenz DJ, Mitchell PO, Walp ER, Sutliff RL, *et al.* Rosiglitazone attenuates chronic hypoxia-induced pulmonary hypertension in a mouse model. *Am J Respir Cell Mol Biol* 2010;42:482-90.
16. Barak Y, Nelson MC, Ong ES, Jones YZ, Ruiz-Lozano P, Chien KR, *et al.* PPAR gamma is required for placental, cardiac, and adipose tissue development. *Mol Cell* 1999;4:585-95.
17. Guignabert C, Alvira CM, Alastalo TP, Sawada H, Hansmann G, Zhao M, *et al.* Tie2-Mediated loss of peroxisome proliferator-activated receptor-[gamma] in mice causes PDGF-receptor {beta}-dependant pulmonary arterial muscularization. *Am J Physiol Lung Cell Mol Physiol* 2009;297:L1082-90.
18. Hansmann L, Groeger S, von Wulffen W, Bein G, Hackstein H. Human monocytes represent a competitive source of interferon-alpha in peripheral blood. *Clin Immunol* 2008;127:252-64.
19. Tian J, Smith A, Nechtman J, Podolsky R, Aggarwal S, Snead C, *et al.* Effect of PPARgamma inhibition on pulmonary endothelial cell gene expression: Gene profiling in pulmonary hypertension. *Physiol Genomics* 2009;40:48-60.
20. Matsuda Y, Hoshikawa Y, Ameshima S, Suzuki S, Okada Y, Tabata T, *et al.* [Effects of peroxisome proliferator-activated receptor gamma ligands on monocrotaline-induced pulmonary hypertension in rats]. *Nihon Kokyuki Gakkai Zasshi* 2005;43:283-8.
21. Crossno JT Jr, Garat CV, Reusch JE, Morris KG, Dempsey EC, McMurtry IF, *et al.* Rosiglitazone attenuates hypoxia-induced pulmonary arterial remodeling. *Am J Physiol Lung Cell Mol Physiol* 2007;292:L885-97.
22. Hansmann G, de Jesus Perez VA, Alastalo TP, Alvira CM, Guignabert C, Bekker JM, *et al.* An antiproliferative BMP-2/PPARgamma/apoE axis in human and murine SMCs and its role in pulmonary hypertension. *J Clin Invest* 2008;118:1846-57.
23. Christman BW, McPherson CD, Newman JH, King GA, Bernard GR, Groves BM, *et al.* An imbalance between the excretion of thromboxane and prostacyclin metabolites in pulmonary hypertension. *N Engl J Med* 1992;327:70-5.
24. Farber HW, Loscalzo J. Pulmonary arterial hypertension. *N Engl J Med* 2004;351:1655-65.
25. Giaid A, Saleh D. Reduced expression of endothelial nitric oxide synthase in the lungs of patients with pulmonary hypertension. *N Engl J Med* 1995;333:214-21.
26. Li H, Chen SJ, Chen YF, Meng QC, Durand J, Oparil S, *et al.* Enhanced endothelin-1 and endothelin receptor gene expression in chronic hypoxia. *J Appl Physiol* 1994;77:1451-9.
27. Olschewski H, Rose F, Schermuly R, Ghofrani HA, Enke B, Olschewski A, *et al.* Prostacyclin and its analogues in the treatment of pulmonary hypertension. *Pharmacol Ther* 2004;102:139-53.
28. Hoshikawa Y, Voelkel NF, Gesell TL, Moore MD, Morris KG, Alger LA, *et al.* Prostacyclin receptor-dependent modulation of pulmonary vascular remodeling. *Am J Respir Crit Care Med* 2001;164:314-8.
29. Tuder RM, Cool CD, Geraci MW, Wang J, Abman SH, Wright L, *et al.* Prostacyclin synthase expression is decreased in lungs from patients with severe pulmonary hypertension. *Am J Respir Crit Care Med* 1999;159:1925-32.
30. Robbins IM, Morrow JD, Christman BW. Oxidant stress but not thromboxane decreases with epoprostenol therapy. *Free Radic Biol Med* 2005;38:568-74.
31. Gupta RA, Tan J, Krause WF, Geraci MW, Willson TM, Dey SK, *et al.* Prostacyclin-mediated activation of peroxisome proliferator-activated receptor delta in colorectal cancer. *Proc Natl Acad Sci U S A* 2000;97:13275-80.
32. Lim H, Dey SK. A novel pathway of prostacyclin signaling-hanging out with nuclear receptors. *Endocrinology* 2002;143:3207-10.
33. Peredo HA, Mayer MA, Carranza A, Puyó AM. Pioglitazone and losartan modify hemodynamic and metabolic parameters and vascular prostanooids in fructose-overloaded rats. *Clin Exp Hypertens* 2008;30:159-69.
34. Tesse A, Al-Massarani G, Wangenstein R, Reitenbach S, Martínez MC, Andriantsitohaina R. Rosiglitazone, a peroxisome proliferator-activated receptor-gamma agonist, prevents microparticle-induced vascular hyporeactivity through the regulation of proinflammatory proteins. *J Pharmacol Exp Ther* 2008;324:539-47.
35. Adnot S, Raffestin B, Eddahibi S, Braquet P, Chabrier PE. Loss of endothelium-dependent relaxant activity in the pulmonary circulation of rats exposed to chronic hypoxia. *J Clin Invest* 1991;87:155-62.
36. Steinhorn RH, Russell JA, Lakshminrusimha S, Gugino SF, Black SM, Fineman JR. Altered endothelium-dependent relaxations in lambs with high pulmonary blood flow and pulmonary hypertension. *Am J Physiol Heart Circ Physiol* 2001;280:H311-7.
37. McQuillan LP, Leung GK, Marsden PA, Kostyk SK, Kourembanas S. Hypoxia inhibits expression of eNOS via transcriptional and posttranscriptional mechanisms. *Am J Physiol* 1994;267:H1921-7.
38. Fagan KA, Fouty BW, Tyler RC, Morris KG Jr, Hepler LK, Sato K, *et al.* The pulmonary circulation of homozygous or heterozygous eNOS-null mice is hyperresponsive to mild hypoxia. *J Clin Invest* 1999;103:291-9.
39. Champion HC, Bivalacqua TJ, Greenberg SS, Giles TD, Hyman AL, Kadowitz PJ. Adenoviral gene transfer of endothelial nitric-oxide synthase (eNOS) partially restores normal pulmonary arterial pressure in eNOS-deficient mice. *Proc Natl Acad Sci U S A* 2002;99:13248-53.
40. Ozaki M, Kawashima S, Yamashita T, Ohashi Y, Rikitake Y, Inoue N, *et al.* Reduced hypoxic pulmonary vascular remodeling by nitric oxide from the endothelium. *Hypertension* 2001;37:322-7.
41. Channick RN, Newhart JW, Johnson FW, Williams PJ, Auger WR, Fedullo PF, *et al.* Pulsed delivery of inhaled nitric oxide to patients with primary pulmonary hypertension: An ambulatory delivery system and initial clinical tests. *Chest* 1996;109:1545-9.
42. Zhao YD, Courtman DW, Ng DS, Robb MJ, Deng YP, Trogadis J, *et al.* Microvascular regeneration in established pulmonary hypertension by angiogenic gene transfer. *Am J Respir Cell Mol Biol* 2006;35:182-9.
43. Pritchard KA Jr, Ackerman AW, Gross ER, Stepp DW, Shi Y, Fontana JT, *et al.* Heat shock protein 90 mediates the balance of nitric oxide and superoxide anion from endothelial nitric-oxide synthase. *J Biol Chem* 2001;276:17621-4.
44. Konduri GG, Ou J, Shi Y, Pritchard KA Jr. Decreased association of HSP90 impairs endothelial nitric oxide synthase in fetal lambs with persistent pulmonary hypertension. *Am J Physiol Heart Circ Physiol* 2003;285:H204-11.
45. Murata T, Sato K, Hori M, Ozaki H, Karaki H. Decreased endothelial nitric-oxide synthase (eNOS) activity resulting from abnormal interaction between eNOS and its regulatory proteins in hypoxia-induced pulmonary hypertension. *J Biol Chem* 2002;277:44085-92.
46. Calnek DS, Mazzella L, Roser S, Roman J, Hart CM. Peroxisome proliferator-activated receptor gamma ligands increase release of nitric oxide from endothelial cells. *Arterioscler Thromb Vasc Biol* 2003;23:52-7.
47. Polikandriotis JA, Mazzella LJ, Rupnow HL, Hart CM. Peroxisome proliferator-activated receptor gamma ligands stimulate endothelial nitric oxide production through distinct peroxisome proliferator-activated receptor gamma-dependent mechanisms. *Arterioscler Thromb Vasc Biol* 2005;25:1810-6.
48. Borchert M, Schöndorf T, Lübken G, Forst T, Pfützner A. Review of the pleiotropic effects of peroxisome proliferator-activated receptor gamma agonists on platelet function. *Diabetes Technol Ther* 2007;9:410-20.
49. Morrell NW, Adnot S, Archer SL, Dupuis J, Jones PL, MacLean MR, *et al.*

- Cellular and molecular basis of pulmonary arterial hypertension. *J Am Coll Cardiol* 2009;54:520-31.
50. Wakino S, Hayashi K, Tatematsu S, Hasegawa K, Takamatsu I, Kanda T, *et al.* Pioglitazone lowers systemic asymmetric dimethylarginine by inducing dimethylarginine dimethylaminohydrolase in rats. *Hypertens Res* 2005;28:255-62.
 51. Palmer RM, Ferrige AG, Moncada S. Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 1987;327:524-6.
 52. Azuma H, Ishikawa M, Sekizaki S. Endothelium-dependent inhibition of platelet aggregation. *Br J Pharmacol* 1986;88:411-5.
 53. Garg UC, Hassid A. Nitric oxide-generating vasodilators and 8-bromocyclic guanosine monophosphate inhibit mitogenesis and proliferation of cultured rat vascular smooth muscle cells. *J Clin Invest* 1989;83:1774-7.
 54. Bath PM, Hassall DG, Gladwin AM, Palmer RM, Martin JF. Nitric oxide and prostacyclin. Divergence of inhibitory effects on monocyte chemotaxis and adhesion to endothelium *in vitro*. *Arterioscler Thromb* 1991;11:254-60.
 55. Kuzkaya N, Weissmann N, Harrison DG, Dikalov S. Interactions of peroxynitrite, tetrahydrobiopterin, ascorbic acid, and thiols: Implications for uncoupling endothelial nitric-oxide synthase. *J Biol Chem* 2003;278:22546-54.
 56. Jernigan NL, Walker BR, Resta TC. Endothelium-derived reactive oxygen species and endothelin-1 attenuate NO-dependent pulmonary vasodilation following chronic hypoxia. *Am J Physiol Lung Cell Mol Physiol* 2004;287:L801-8.
 57. Liu JQ, Zelko IN, Erbynn EM, Sham JS, Folz RJ. Hypoxic pulmonary hypertension: Role of superoxide and NADPH oxidase (gp91phox). *Am J Physiol Lung Cell Mol Physiol* 2006;290:L2-10.
 58. Delerive P, Martin-Nizard F, Chinetti G, Trottein F, Fruchart JC, Najib J, *et al.* Peroxisome proliferator-activated receptor activators inhibit thrombin-induced endothelin-1 production in human vascular endothelial cells by inhibiting the activator protein-1 signaling pathway. *Circ Res* 1999;85:394-402.
 59. Martin-Nizard F, Furman C, Delerive P, Kandoussi A, Fruchart JC, Staels B, *et al.* Peroxisome proliferator-activated receptor activators inhibit oxidized low-density lipoprotein-induced endothelin-1 secretion in endothelial cells. *J Cardiovasc Pharmacol* 2002;40:822-31.
 60. Fukunaga Y, Itoh H, Doi K, Tanaka T, Yamashita J, Chun TH, *et al.* Thiazolidinediones, peroxisome proliferator-activated receptor gamma agonists, regulate endothelial cell growth and secretion of vasoactive peptides. *Atherosclerosis* 2001;158:113-9.
 61. Wang TD, Chen WJ, Cheng WC, Lin JW, Chen MF, Lee YT. Relation of improvement in endothelium-dependent flow-mediated vasodilation after rosiglitazone to changes in asymmetric dimethylarginine, endothelin-1, and C-reactive protein in nondiabetic patients with the metabolic syndrome. *Am J Cardiol* 2006;98:1057-62.
 62. Iglarz M, Touyz RM, Amiri F, Lavoie MF, Diep QN, Schiffrin EL. Effect of peroxisome proliferator-activated receptor-alpha and -gamma activators on vascular remodeling in endothelin-dependent hypertension. *Arterioscler Thromb Vasc Biol* 2003;23:45-51.
 63. Jeffery TK, Morrell NW. Molecular and cellular basis of pulmonary vascular remodeling in pulmonary hypertension. *Prog Cardiovasc Dis* 2002;45:173-202.
 64. Hassoun PM, Mouthon L, Barberà JA, Eddahibi S, Flores SC, Grimminger F, *et al.* Inflammation, growth factors, and pulmonary vascular remodeling. *J Am Coll Cardiol* 2009;54:S10-9.
 65. Tudor RM, Groves B, Badesch DB, Voelkel NF. Exuberant endothelial cell growth and elements of inflammation are present in plexiform lesions of pulmonary hypertension. *Am J Pathol* 1994;144:275-85.
 66. Tudor RM, Voelkel NF. Pulmonary hypertension and inflammation. *J Lab Clin Med* 1998;132:16-24.
 67. Balabanian K, Foussat A, Dorfmueller P, Durand-Gasselin I, Capel F, Bouchet-Delbos L, *et al.* CX(3)C chemokine fractalkine in pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2002;165:1419-25.
 68. Dorfmueller P, Zarka V, Durand-Gasselin I, Monti G, Balabanian K, Garcia G, *et al.* Chemokine RANTES in severe pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2002;165:534-9.
 69. Sanchez O, Marcos E, Perros F, Fadel E, Tu L, Humbert M, *et al.* Role of endothelium-derived CC chemokine ligand 2 in idiopathic pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2007;176:1041-7.
 70. Griendling KK, Sorescu D, Ushio-Fukai M. NAD(P)H oxidase: Role in cardiovascular biology and disease. *Circ Res* 2000;86:494-501.
 71. Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: Physiology and pathophysiology. *Physiol Rev* 2007;87:245-313.
 72. Lambeth JD. Nox/Duox family of nicotinamide adenine dinucleotide (phosphate) oxidases. *Curr Opin Hematol* 2002;9:11-7.
 73. Mittal M, Roth M, König P, Hofmann S, Dony E, Goyal P, *et al.* Hypoxia-dependent regulation of nonphagocytic NADPH oxidase subunit NOX4 in the pulmonary vasculature. *Circ Res* 2007;101:258-67.
 74. Ambasta RK, Kumar P, Griendling KK, Schmidt HH, Busse R, Brandes RP. Direct interaction of the novel Nox proteins with p22phox is required for the formation of a functionally active NADPH oxidase. *J Biol Chem* 2004;279:45935-41.
 75. Djordjevic T, BelAiba RS, Bonello S, Pfeilschifter J, Hess J, Görlach A. Human urotensin II is a novel activator of NADPH oxidase in human pulmonary artery smooth muscle cells. *Arterioscler Thromb Vasc Biol* 2005;25:519-25.
 76. Serrander L, Cartier L, Bedard K, Banfi B, Lardy B, Plastre O, *et al.* NOX4 activity is determined by mRNA levels and reveals a unique pattern of ROS generation. *Biochem J* 2007;406:105-14.
 77. Lambeth JD, Kawahara T, Diebold B. Regulation of Nox and Duox enzymatic activity and expression. *Free Radic Biol Med* 2007;43:319-31.
 78. Sorescu GP, Song H, Tressel SL, Hwang J, Dikalov S, Smith DA, *et al.* Bone morphogenic protein 4 produced in endothelial cells by oscillatory shear stress induces monocyte adhesion by stimulating reactive oxygen species production from a nox1-based NADPH oxidase. *Circ Res* 2004;95:773-9.
 79. Hwang J, Kleinhenz DJ, Lassègue B, Griendling KK, Dikalov S, Hart CM. Peroxisome proliferator-activated receptor-gamma ligands regulate endothelial membrane superoxide production. *Am J Physiol Cell Physiol* 2005;288:C899-905.
 80. Lu X, Murphy TC, Nanes MS, Hart CM. PPAR[gamma] regulates hypoxia-induced Nox4 expression in human pulmonary artery smooth muscle cells through NF-[kappa]B. *Am J Physiol Lung Cell Mol Physiol* 2010;299:L559-66.
 81. Zhang L, Sheppard OR, Shah AM, Brewer AC. Positive regulation of the NADPH oxidase NOX4 promoter in vascular smooth muscle cells by E2F. *Free Radic Biol Med* 2008;45:679-85.
 82. Chen K, Kirber MT, Xiao H, Yang Y, Keaney JF Jr. Regulation of ROS signal transduction by NADPH oxidase 4 localization. *J Cell Biol* 2008;181:1129-39.
 83. Fredriksson L, Li H, Eriksson U. The PDGF family: Four gene products form five dimeric isoforms. *Cytokine Growth Factor Rev* 2004;15:197-204.
 84. Raines EW. PDGF and cardiovascular disease. *Cytokine Growth Factor Rev* 2004;15:237-54.
 85. Tallquist M, Kazlauskas A. PDGF signaling in cells and mice. *Cytokine Growth Factor Rev* 2004;15:205-13.
 86. Balasubramaniam V, Le Cras TD, Ivy DD, Grover TR, Kinsella JP, Abman SH. Role of platelet-derived growth factor in vascular remodeling during pulmonary hypertension in the ovine fetus. *Am J Physiol Lung Cell Mol Physiol* 2003;284:L826-33.
 87. Ghofrani HA, Seeger W, Grimminger F. Imatinib for the treatment of pulmonary arterial hypertension. *N Engl J Med* 2005;353:1412-3.
 88. Schermuly RT, Dony E, Ghofrani HA, Pullamsetti S, Savai R, Roth M, *et al.* Reversal of experimental pulmonary hypertension by PDGF inhibition. *J Clin Invest* 2005;115:2811-21.
 89. Goetze S, Xi XP, Kawano H, Gotlibowski T, Fleck E, Hsueh WA, *et al.* PPAR gamma-ligands inhibit migration mediated by multiple chemoattractants in vascular smooth muscle cells. *J Cardiovasc Pharmacol* 1999;33:798-806.
 90. Marx N, Schönbeck U, Lazar MA, Libby P, Plutzky J. Peroxisome proliferator-activated receptor gamma activators inhibit gene expression and migration in human vascular smooth muscle cells. *Circ Res* 1998;83:1097-103.
 91. Law RE, Goetze S, Xi XP, Jackson S, Kawano Y, Demer L, *et al.* Expression and function of PPARgamma in rat and human vascular smooth muscle cells. *Circulation* 2000;101:1311-8.
 92. Gericke A, Munson M, Ross AH. Regulation of the PTEN phosphatase. *Gene* 2006;374:1-9.
 93. Maehama T, Dixon JE. The tumor suppressor, PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-trisphosphate. *J Biol Chem* 1998;273:13375-8.
 94. Schwartzbauer G, Robbins J. The tumor suppressor gene PTEN can regulate cardiac hypertrophy and survival. *J Biol Chem* 2001;276:35786-93.
 95. Tong H, Chen W, Steenbergen C, Murphy E. Ischemic preconditioning activates phosphatidylinositol-3-kinase upstream of protein kinase C. *Circ Res* 2000;87:309-15.
 96. Han SW, Roman J. Fibronectin induces cell proliferation and inhibits apoptosis in human bronchial epithelial cells: Pro-oncogenic effects mediated by PI3-kinase and NF-kappa B. *Oncogene* 2006;25:4341-9.

97. Wang Q, Zhou Y, Wang X, Chung DH, Evers BM. Regulation of PTEN expression in intestinal epithelial cells by c-Jun NH2-terminal kinase activation and nuclear factor-kappaB inhibition. *Cancer Res* 2007;67:7773-81.
98. Farrow B, Evers BM. Activation of PPARgamma increases PTEN expression in pancreatic cancer cells. *Biochem Biophys Res Commun* 2003;301:50-3.
99. Patel L, Pass I, Coxon P, Downes CP, Smith SA, Macphee CH. Tumor suppressor and anti-inflammatory actions of PPARgamma agonists are mediated via upregulation of PTEN. *Curr Biol* 2001;11:764-8.
100. Huang J, Kontos CD. Inhibition of vascular smooth muscle cell proliferation, migration, and survival by the tumor suppressor protein PTEN. *Arterioscler Thromb Vasc Biol* 2002;22:745-51.
101. Huang J, Niu XL, Phippen AM, Annex BH, Kontos CD. Adenovirus-mediated intraarterial delivery of PTEN inhibits neointimal hyperplasia. *Arterioscler Thromb Vasc Biol* 2005;25:354-8.
102. Lee SR, Yang KS, Kwon J, Lee C, Jeong W, Rhee SG. Reversible inactivation of the tumor suppressor PTEN by H₂O₂. *J Biol Chem* 2002;277:20336-42.
103. Leslie NR, Bennett D, Lindsay YE, Stewart H, Gray A, Downes CP. Redox regulation of PI 3-kinase signalling via inactivation of PTEN. *EMBO J* 2003;22:5501-10.
104. Kwon J, Lee SR, Yang KS, Ahn Y, Kim YJ, Stadtman ER, *et al.* Reversible oxidation and inactivation of the tumor suppressor PTEN in cells stimulated with peptide growth factors. *Proc Natl Acad Sci U S A* 2004;101:16419-24.
105. Goumans MJ, Liu Z, ten Dijke P. TGF-beta signaling in vascular biology and dysfunction. *Cell Res* 2009;19:116-27.
106. Upton PD, Morrell NW. TGF-beta and BMPR-II pharmacology--implications for pulmonary vascular diseases. *Curr Opin Pharmacol* 2009;9:274-80.
107. Miyazono K. Transforming growth factor-beta signaling in epithelial-mesenchymal transition and progression of cancer. *Proc Jpn Acad Ser B Phys Biol Sci* 2009;85:314-23.
108. Richter A, Yeager ME, Zaiman A, Cool CD, Voelkel NF, Tudor RM. Impaired transforming growth factor-beta signaling in idiopathic pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2004;170:1340-8.
109. Cogan JD, Vnencak-Jones CL, Phillips JA 3rd, Lane KB, Wheeler LA, Robbins IM, *et al.* Gross BMPR2 gene rearrangements constitute a new cause for primary pulmonary hypertension. *Genet Med* 2005;7:169-74.
110. Du L, Sullivan CC, Chu D, Cho AJ, Kido M, Wolf PL, *et al.* Signaling molecules in nonfamilial pulmonary hypertension. *N Engl J Med* 2003;348:500-9.
111. Humbert M, Morrell NW, Archer SL, Stenmark KR, MacLean MR, Lang IM, *et al.* Cellular and molecular pathobiology of pulmonary arterial hypertension. *J Am Coll Cardiol* 2004;43:135-245.
112. Long L, Crosby A, Yang X, Southwood M, Upton PD, Kim DK, *et al.* Altered bone morphogenetic protein and transforming growth factor-beta signaling in rat models of pulmonary hypertension: Potential for activin receptor-like kinase-5 inhibition in prevention and progression of disease. *Circulation* 2009;119:566-76.
113. Sturrock A, Cahill B, Norman K, Huecksteadt TP, Hill K, Sanders K, *et al.* Transforming growth factor-beta1 induces Nox4 NAD(P)H oxidase and reactive oxygen species-dependent proliferation in human pulmonary artery smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 2006;290:L661-73.
114. Ismail S, Sturrock A, Wu P, Cahill B, Norman K, Huecksteadt T, *et al.* NOX4 mediates hypoxia-induced proliferation of human pulmonary artery smooth muscle cells: The role of autocrine production of transforming growth factor-beta1 and insulin-like growth factor binding protein-3. *Am J Physiol Lung Cell Mol Physiol* 2009;296:L489-99.
115. Ambalavanan N, Nicola T, Hagood J, Bulger A, Serra R, Murphy-Ullrich J, *et al.* Transforming growth factor-beta signaling mediates hypoxia-induced pulmonary arterial remodeling and inhibition of alveolar development in newborn mouse lung. *Am J Physiol Lung Cell Mol Physiol* 2008;295:L86-95.
116. Chen G, Khalil N. TGF-beta1 increases proliferation of airway smooth muscle cells by phosphorylation of map kinases. *Respir Res* 2006;7:2.
117. Burgess HA, Daugherty LE, Thatcher TH, Lakatos HF, Ray DM, Redonnet M, *et al.* PPARgamma agonists inhibit TGF-beta induced pulmonary myofibroblast differentiation and collagen production: Implications for therapy of lung fibrosis. *Am J Physiol Lung Cell Mol Physiol* 2005;288:L1146-53.
118. Saika S, Yamanaka O, Nishikawa-Ishida I, Kitano A, Flanders KC, Okada Y, *et al.* Effect of Smad7 gene overexpression on transforming growth factor beta-induced retinal pigment fibrosis in a proliferative vitreoretinopathy mouse model. *Arch Ophthalmol* 2007;125:647-54.
119. Zou R, Xu G, Liu XC, Han M, Jiang JJ, Huang Q, *et al.* PPARgamma agonists inhibit TGF-beta-PKA signaling in glomerulosclerosis. *Acta Pharmacol Sin* 2010;31:43-50.
120. Zhao C, Chen W, Yang L, Chen L, Stimpson SA, Diehl AM. PPARgamma agonists prevent TGFbeta1/Smad3-signaling in human hepatic stellate cells. *Biochem Biophys Res Commun* 2006;350:385-91.
121. Fu M, Zhang J, Zhu X, Myles DE, Willson TM, Liu X, *et al.* Peroxisome proliferator-activated receptor gamma inhibits transforming growth factor beta-induced connective tissue growth factor expression in human aortic smooth muscle cells by interfering with Smad3. *J Biol Chem* 2001;276:45888-94.
122. Cowan KN, Jones PL, Rabinovitch M. Elastase and matrix metalloproteinase inhibitors induce regression, and tenascin-C antisense prevents progression, of vascular disease. *J Clin Invest* 2000;105:21-34.
123. Stenmark KR, Fagan KA, Frid MG. Hypoxia-induced pulmonary vascular remodeling: Cellular and molecular mechanisms. *Circ Res* 2006;99:675-91.
124. Zaidi SH, You XM, Ciura S, Husain M, Rabinovitch M. Overexpression of the serine elastase inhibitor elafin protects transgenic mice from hypoxic pulmonary hypertension. *Circulation* 2002;105:516-21.
125. Hao GH, Niu XL, Gao DF, Wei J, Wang NP. Agonists at PPAR-gamma suppress angiotensin II-induced production of plasminogen activator inhibitor-1 and extracellular matrix in rat cardiac fibroblasts. *Br J Pharmacol* 2008;153:1409-19.
126. Makino N, Sugano M, Satoh S, Oyama J, Maeda T. Peroxisome proliferator-activated receptor-gamma ligands attenuate brain natriuretic peptide production and affect remodeling in cardiac fibroblasts in reoxygenation after hypoxia. *Cell Biochem Biophys* 2006;44:65-71.
127. Peng Y, Liu H, Liu F, Liu Y, Li J, Chen X. Troglitazone inhibits synthesis of transforming growth factor-beta1 and reduces matrix production in human peritoneal mesothelial cells. *Nephrology (Carlton)* 2006;11:516-23.
128. Bakouboula B, Morel O, Faure A, Zobairi F, Jesel L, Trinh A, *et al.* Procoagulant membrane microparticles correlate with the severity of pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2008;177:536-43.
129. Perros F, Dorfmueller P, Souza R, Durand-Gasselin I, Godot V, Capel F, *et al.* Fractalkine-induced smooth muscle cell proliferation in pulmonary hypertension. *Eur Respir J* 2007;29:937-43.
130. Molet S, Furukawa K, Maghazachi A, Hamid Q, Giaid A. Chemokine- and cytokine-induced expression of endothelin 1 and endothelin-converting enzyme 1 in endothelial cells. *J Allergy Clin Immunol* 2000;105:333-8.
131. Bonnet S, Rochefort G, Sutendra G, Archer SL, Haromy A, Webster L, *et al.* The nuclear factor of activated T cells in pulmonary arterial hypertension can be therapeutically targeted. *Proc Natl Acad Sci U S A* 2007;104:11418-23.
132. Szanto A, Nagy L. The many faces of PPARgamma: Anti-inflammatory by any means? *Immunobiology* 2008;213:789-803.
133. Hontecillas R, Bassaganya-Riera J. Peroxisome proliferator-activated receptor gamma is required for regulatory CD4+ T cell-mediated protection against colitis. *J Immunol* 2007;178:2940-9.
134. Wohlfert EA, Nichols FC, Nevius E, Clark RB. Peroxisome proliferator-activated receptor gamma (PPARgamma) and immunoregulation: Enhancement of regulatory T cells through PPARgamma-dependent and -independent mechanisms. *J Immunol* 2007;178:4129-35.
135. Gurbanov E, Shiliang X. The key role of apoptosis in the pathogenesis and treatment of pulmonary hypertension. *Eur J Cardiothorac Surg* 2006;30:499-507.
136. Haunstetter A, Izumo S. Apoptosis: Basic mechanisms and implications for cardiovascular disease. *Circ Res* 1998;82:1111-29.
137. Zhang S, Fantozzi I, Tigno DD, Yi ES, Platoshyn O, Thistlethwaite PA, *et al.* Bone morphogenetic proteins induce apoptosis in human pulmonary vascular smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L740-54.
138. Okura T, Nakamura M, Takata Y, Watanabe S, Kitami Y, Hiwada K. Troglitazone induces apoptosis via the p53 and Gadd45 pathway in vascular smooth muscle cells. *Eur J Pharmacol* 2000;407:227-35.
139. Aizawa Y, Kawabe J, Hasebe N, Takekura N, Kikuchi K. Pioglitazone enhances cytokine-induced apoptosis in vascular smooth muscle cells and reduces intimal hyperplasia. *Circulation* 2001;104:455-60.
140. Bishop-Bailey D, Hla T. Endothelial cell apoptosis induced by the peroxisome proliferator-activated receptor (PPAR) ligand 15-deoxy-Delta12, 14-prostaglandin J2. *J Biol Chem* 1999;274:17042-8.
141. Masri FA, Xu W, Comhair SA, Asosingh K, Koo M, Vasanthi A, *et al.*

- Hyperproliferative apoptosis-resistant endothelial cells in idiopathic pulmonary arterial hypertension. *Am J Physiol Lung Cell Mol Physiol* 2007;293:L548-54.
142. Park EJ, Park SY, Joe EH, Jou I. 15d-PGJ2 and rosiglitazone suppress Janus kinase-STAT inflammatory signaling through induction of suppressor of cytokine signaling 1 (SOCS1) and SOCS3 in glia. *J Biol Chem* 2003;278:14747-52.
 143. Asahara T, Murohara T, Sullivan A, Silver M, van der Zee R, Li T, *et al.* Isolation of putative progenitor endothelial cells for angiogenesis. *Science* 1997;275:964-7.
 144. Rosenzweig A. Circulating endothelial progenitors--cells as biomarkers. *N Engl J Med* 2005;353:1055-7.
 145. Hill JM, Zalos G, Halcox JP, Schenke WH, Waclawiw MA, Quyyumi AA, *et al.* Circulating endothelial progenitor cells, vascular function, and cardiovascular risk. *N Engl J Med* 2003;348:593-600.
 146. Werner N, Kosiol S, Schiegl T, Ahlers P, Walenta K, Link A, *et al.* Circulating endothelial progenitor cells and cardiovascular outcomes. *N Engl J Med* 2005;353:999-1007.
 147. Sata M. Role of circulating vascular progenitors in angiogenesis, vascular healing, and pulmonary hypertension: Lessons from animal models. *Arterioscler Thromb Vasc Biol* 2006;26:1008-14.
 148. Satoh K, Kagaya Y, Nakano M, Ito Y, Ohta J, Tada H, *et al.* Important role of endogenous erythropoietin system in recruitment of endothelial progenitor cells in hypoxia-induced pulmonary hypertension in mice. *Circulation* 2006;113:1442-50.
 149. Nagaya N, Kangawa K, Kanda M, Uematsu M, Horio T, Fukuyama N, *et al.* Hybrid cell-gene therapy for pulmonary hypertension based on phagocytosing action of endothelial progenitor cells. *Circulation* 2003;108:889-95.
 150. Wang CH, Ciliberti N, Li SH, Szmítko PE, Weisel RD, Fedak PW, *et al.* Rosiglitazone facilitates angiogenic progenitor cell differentiation toward endothelial lineage: A new paradigm in glitazone pleiotropy. *Circulation* 2004;109:1392-400.
 151. Gensch C, Clever YP, Werner C, Hanhoun M, Böhm M, Laufs U. The PPAR-gamma agonist pioglitazone increases neoangiogenesis and prevents apoptosis of endothelial progenitor cells. *Atherosclerosis* 2007;192:67-74.
 152. Chaouat A, Weitzenblum E, Higenbottam T. The role of thrombosis in severe pulmonary hypertension. *Eur Respir J* 1996;9:356-63.
 153. Herve P, Humbert M, Sitbon O, Parent F, Nunes H, Legal C, *et al.* Pathobiology of pulmonary hypertension. The role of platelets and thrombosis. *Clin Chest Med* 2001;22:451-8.
 154. Welsh CH, Hassell KL, Badesch DB, Kressin DC, Marlar RA. Coagulation and fibrinolytic profiles in patients with severe pulmonary hypertension. *Chest* 1996;110:710-7.
 155. Hoepfer MM, Sosada M, Fabel H. Plasma coagulation profiles in patients with severe primary pulmonary hypertension. *Eur Respir J* 1998;12:1446-9.
 156. Eisenberg PR, Lucore C, Kaufman L, Sobel BE, Jaffe AS, Rich S. Fibrinogen levels indicative of pulmonary vascular thrombosis in patients with primary pulmonary hypertension. *Circulation* 1990;82:841-7.
 157. Altman R, Scaccia A, Rouvier J, Gurfinkel E, Favaloro R, Perrone S, *et al.* Coagulation and fibrinolytic parameters in patients with pulmonary hypertension. *Clin Cardiol* 1996;19:549-54.
 158. Tudor RM, Chacon M, Alger L, Wang J, Taraseviciene-Stewart L, Kasahara Y, *et al.* Expression of angiogenesis-related molecules in plexiform lesions in severe pulmonary hypertension: Evidence for a process of disordered angiogenesis. *J Pathol* 2001;195:367-74.
 159. Akbiyik F, Ray DM, Gettings KF, Blumberg N, Francis CW, Phipps RP. Human bone marrow megakaryocytes and platelets express PPARgamma, and PPARgamma agonists blunt platelet release of CD40 ligand and thromboxanes. *Blood* 2004;104:1361-8.
 160. Schönbeck U, Libby P. CD40 signaling and plaque instability. *Circ Res* 2001;89:1092-103.
 161. Henn V, Slupsky JR, Gräfe M, Anagnostopoulos I, Förster R, Müller-Berghaus G, *et al.* CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells. *Nature* 1998;391:591-4.
 162. Bodary PF, Vargas FB, King SA, Jongeward KL, Wickenheiser KJ, Eitzman DT. Pioglitazone protects against thrombosis in a mouse model of obesity and insulin resistance. *J Thromb Haemost* 2005;3:2149-53.
 163. Li D, Chen K, Sinha N, Zhang X, Wang Y, Sinha AK, *et al.* The effects of PPAR-gamma ligand pioglitazone on platelet aggregation and arterial thrombus formation. *Cardiovasc Res* 2005;65:907-12.
 164. Smyth SS, Jennings JL. PPARgamma agonists: A new strategy for antithrombotic therapy. *J Thromb Haemost* 2005;3:2147-8.
 165. Cylwik D, Mogiński A, Buczek W. L-arginine and cardiovascular system. *Pharmacol Rep* 2005;57:14-22.
 166. Fonseca VA, Reynolds T, Hemphill D, Randolph C, Wall J, Valiquet TR, *et al.* Effect of troglitazone on fibrinolysis and activated coagulation in patients with non-insulin-dependent diabetes mellitus. *J Diabetes Complications* 1998;12:181-6.
 167. Kato K, Yamada D, Midorikawa S, Sato W, Watanabe T. Improvement by the insulin-sensitizing agent, troglitazone, of abnormal fibrinolysis in type 2 diabetes mellitus. *Metabolism* 2000;49:662-5.
 168. Stenmark KR, Meyrick B, Galie N, Mooi WJ, McMurtry IF. Animal models of pulmonary arterial hypertension: The hope for etiological discovery and pharmacological cure. *Am J Physiol Lung Cell Mol Physiol* 2009;297:L1013-32.
 169. Abe K, Toba M, Alzoubi A, Ito M, Fagan KA, Cool CD, *et al.* Formation of plexiform lesions in experimental severe pulmonary arterial hypertension. *Circulation* 2010;121:2747-54.
 170. Nissen SE, Wolski K. Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes. *N Engl J Med* 2007;356:2457-71.
 171. Rosen CJ. The rosiglitazone story--lessons from an FDA Advisory Committee meeting. *N Engl J Med* 2007;357:844-6.
 172. Dormandy JA, Charbonnel B, Eckland DJ, Erdmann E, Massi-Benedetti M, Moules IK, *et al.* Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitAzone Clinical Trial In macroVascular Events): A randomised controlled trial. *Lancet* 2005;366:1279-89.
 173. Erdmann E, Dormandy JA, Charbonnel B, Massi-Benedetti M, Moules IK, Skene AM, *et al.* The effect of pioglitazone on recurrent myocardial infarction in 2,445 patients with type 2 diabetes and previous myocardial infarction: Results from the PROactive (PROactive 05) Study. *J Am Coll Cardiol* 2007;49:1772-80.
 174. Lincoff AM, Wolski K, Nicholls SJ, Nissen SE. Pioglitazone and risk of cardiovascular events in patients with type 2 diabetes mellitus: A meta-analysis of randomized trials. *JAMA* 2007;298:1180-8.
 175. Wilcox R, Bousser MG, Betteridge DJ, Scherthaner G, Piras V, Kupfer S, *et al.* Effects of pioglitazone in patients with type 2 diabetes with or without previous stroke: Results from PROactive (PROspective pioglitAzone Clinical Trial In macroVascular Events 04). *Stroke* 2007;38:865-73.

Source of Support: Atlanta Veterans Affairs Medical Center and the National Institutes of Health (R01 DK 074518). NHLBI T32 training grant (HL076118-06). **Conflict of Interest:** None declared.

Staying in touch with the Journal

1) Table of Contents (TOC) email alert

Receive an email alert containing the TOC when a new complete issue of the journal is made available online. To register for TOC alerts go to www.pulmonarycirculation.org/signup.asp.

2) RSS feeds

Really Simple Syndication (RSS) helps you to get alerts on new publication right on your desktop without going to the journal's website. You need a software (e.g. RSSReader, Feed Demon, FeedReader, My Yahoo!, NewsGator and NewzCrawler) to get advantage of this tool. RSS feeds can also be read through FireFox or Microsoft Outlook 2007. Once any of these small (and mostly free) software is installed, add www.pulmonarycirculation.org/rssfeed.asp as one of the feeds.