

# Traditional approach versus Stewart approach for acid–base disorders: Inconsistent evidence

SAGE Open Medicine  
Volume 6: 1–9  
© The Author(s) 2018  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/2050312118801255  
journals.sagepub.com/home/smo



Satoshi Kimura<sup>1</sup> , Muhammad Shabsigh<sup>2</sup>  
and Hiroshi Morimatsu<sup>1</sup> 

## Abstract

**Purpose:** The traditional approach and the Stewart approach have been developed for evaluating acid–base phenomena. While some experts have suggested that the two approaches are essentially identical, clinical researches have still been conducted on the superiority of one approach over the other one. In this review, we summarize the concepts of each approach and investigate the reasons of the discrepancy, based on current evidence from the literature search.

**Methods:** In the literature search, we completed a database search and reviewed articles comparing the Stewart approach with the traditional, bicarbonate-centered approach to November 2016.

**Results:** Our literature review included 17 relevant articles, 5 of which compared their diagnostic abilities, 9 articles compared their prognostic performances, and 3 articles compared both diagnostic abilities and prognostic performances. These articles show a discrepancy over the abilities to detect acid–base disturbances and to predict patients' outcomes. There are many limitations that could yield this discrepancy, including differences in calculation of the variables, technological differences or errors in measuring variables, incongruences of reference value, normal range of the variables, differences in studied populations, and confounders of prognostic strength such as lactate.

**Conclusion:** In conclusion, despite the proposed equivalence between the traditional approach and the Stewart approach, our literature search shows inconsistent results on the comparison between the two approaches for diagnostic and prognostic performance. We found crucial limitations in those studies, which could lead to the reasons of the discrepancy.

## Keywords

Henderson–Hasselbalch, Stewart, anion gap, strong ion difference, strong ion gap

Date received: 27 October 2017; accepted: 27 August 2018

## Introduction

Originally, Henderson<sup>1</sup> recognized that carbon dioxide and bicarbonate were key elements of carbonate mass action. Hasselbalch<sup>2</sup> developed it into the negative logarithmic pH notation. Henderson–Hasselbalch equation considers bicarbonate one of the strongest buffers and determinants of pH in our physiologic system. In order to separate metabolic and respiratory components in acid–base disorders, the concept of base excess (BE) was first introduced by Siggaard-Andersen et al.<sup>3</sup> and became the head of the Copenhagen school. On the other hand, exploiting the flaw of using in vitro concept of BE in a living organism, Schwartz and Relman<sup>4</sup> developed the bicarbonate-centered approach setting out the relationship between partial pressure of carbon dioxide (pCO<sub>2</sub>) and bicarbonate ion (HCO<sub>3</sub><sup>-</sup>) in vivo, which became the center of the Boston school. The difference of the two approaches for

metabolic components generated the “great trans-Atlantic acid-base debate” between the Boston school and the Copenhagen school.<sup>5</sup>

In the late 1900s, Peter Stewart questioned the bicarbonate-centered approach and the base excess method for acid-base phenomenon.<sup>6–8</sup> In his concept, each variable is classified as a dependent or independent factor in determining the H<sup>+</sup>

<sup>1</sup>Department of Anesthesiology and Resuscitation, Okayama University Hospital, Okayama, Japan

<sup>2</sup>Department of Anesthesiology, The Ohio State University Wexner Medical Center, Columbus, OH, USA

### Corresponding author:

Satoshi Kimura, Department of Anesthesiology and Resuscitation, Okayama University Hospital, 2-5-1, Shikata-cho, Kita-ku, Okayama 700-8558, Japan.

Email: kimsato1034@hotmail.co.jp



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

concentration of a solution, resulting in pH through the dissociation of water, in order to maintain electrical neutrality.<sup>9,10</sup> Although both the BE approach and the Stewart approach were developed in physio-chemical terms, the Stewart approach is sometimes called “physicochemical,” “modern,” or strong ion approach.<sup>6,7</sup> In contrast, the bicarbonate-centered approach and the base excess approach are called “traditional approach.”<sup>11</sup> Currently, most of the modern blood gas analyzers report both  $\text{HCO}_3^-$  and BE for many clinicians to use the traditional approach.

Since then, both the traditional and the Stewart approaches have been relevant subjects for clinical and research discussions. While some experts have suggested that the two approaches are essentially identical,<sup>12,13</sup> clinical researches have still been conducted and discussed which approach has a better performance as a diagnostic or prognostic tool. In this review, we summarize the concepts of each approach and investigate the reasons of this discrepancy, based on current evidence from the literature search, despite the proposed identity.

## The traditional approach

### Bicarbonate-centered (“Boston”) approach

In the early 1900s, an acid was defined as a substance that is capable of donating hydrogen to a solution, and a base was defined as a substance capable of accepting hydrogen from a solution. Henderson<sup>2</sup> first recognized that bicarbonate is a unique and important buffer, which has the ability to bind or release hydrogen ions in a solution to keep the pH relatively constant, in a physiologic system at constant  $\text{pCO}_2$ . Henderson–Hasselbalch equation provides a simple relationship among the respiratory parameter ( $\text{pCO}_2$ ), the non-respiratory parameter bicarbonate ( $\text{HCO}_3^-$ ), and the overall acidity parameter (pH).<sup>14</sup>

Based on the equation, Schwartz and Relman<sup>4</sup> developed the  $\text{CO}_2/\text{HCO}_3^-$  approach predicting the nature of acid–base disorders. Although it is relatively easy to understand and to apply in clinical settings, there are some weaknesses we need to consider. Since there are non-bicarbonate buffers such as albumin and hemoglobin, a change in bicarbonate concentration does not always reflect the total amount of non-respiratory acids or bases.<sup>15</sup> Furthermore, the equation listing  $\text{pCO}_2$  and bicarbonate as determinants of pH can mislead their interdependence.

### BE and standard BE (“Copenhagen”) approach

In 1948, Singer and Hastings<sup>16</sup> introduced the concept of the buffer base, which is the sum of all plasma buffer anions and is composed of bicarbonate ion and non-volatile, weak acid buffers (mainly albumin and phosphate). It is shown that a change in a buffer base corresponds to a change in the metabolic component of acid–base balance and develops into the BE methodology.<sup>17,18</sup>

In 1960, Siggaard-Andersen et al.<sup>3,19</sup> measured the plasma bicarbonate concentration at a fixed temperature and partial  $\text{pCO}_2$  and compared the difference between their results and a reference value. When corrected by a constant, this difference yields the BE, which represents the amount of acid or alkali that must be added to 1 L of oxygenated blood, exposed in vitro to a  $\text{pCO}_2$  of 40 mmHg to achieve the average normal pH of 7.40.<sup>19,20</sup>

Blood BE measures the metabolic component that is independent from the respiratory component and incorporates the effect of hemoglobin as a buffer.<sup>19,20</sup> The most commonly used formula for calculating the BE is the Van Slyke equation, developed by Siggaard-Andersen<sup>19</sup>

$$\text{BE} = \left\{ \begin{array}{l} \left[ \text{HCO}_3^- \right] - 24.4 + \left( 2.3 * \left[ \text{Hemoglobin (Hb)} \right] + 7.7 \right) \\ * (\text{pH} - 7.4) \\ * (1 - 0.023 * [\text{Hb}]) \end{array} \right\}$$

The BE equation suffers from inaccuracy in vivo with changes in  $\text{pCO}_2$ , possibly due to equilibration across the entire extracellular fluid space, which is composed of whole blood and interstitial fluid.<sup>7,21,22</sup> Therefore, the equation was modified to “Standardize” the effect of hemoglobin on  $\text{CO}_2$  titration in order to improve the accuracy in vivo<sup>7</sup>

$$\text{Standard base excess (SBE)} = 0.9287 * \left\{ \begin{array}{l} \left[ \text{HCO}_3^- \right] \\ - 24.4 + 14.83 \\ * (\text{pH} - 7.4) \end{array} \right\}$$

However, the standard base excess (SBE) is still slightly subject to  $\text{pCO}_2$  change.<sup>7</sup> Furthermore, this equation assumes normal non-buffer ion levels; however, a decrease in albumin or phosphate, which is commonly encountered in intensive care unit (ICU), results in more unstable SBE.<sup>7,8</sup> In addition, the BE and SBE methods are unable to detect complicated acid–base disorders or identify different types of metabolic acidosis.

### Anion gap

The anion gap (AG), the difference between unmeasured plasma anions and the unmeasured plasma cations,<sup>8</sup> is an additional diagnostic tool to assess the metabolic components of the acid–base equilibrium. Albumin and phosphate, one of the circulatory proteins, mainly account for the AG under normal conditions. The rest of the possible candidates are composed of urate, lactate, ketone bodies, sulfate, salicylates, penicillins, citrate, pyruvate, and acetate.<sup>23,24</sup>

This additional diagnostic tool provides new insight to the traditional approach, classifying metabolic acidosis into normal AG acidosis and high AG acidosis. However, severe pH disturbances and changes in the concentration of serum

albumin, which behaves as an anion, have a significant impact on the AG.<sup>25,26</sup> Those disadvantages lower the sensitivity and specificity of this diagnostic tool to detect metabolic acidosis.

A noticeable attempt to improve the practical AG was the introduction of the corrected anion gap (AGc). The most popular AGc is “albumin-corrected” AG. For each 10 g/L decrement in the serum albumin concentration, the AG is expected to decrease by 2.5 mmol/L and needs to be corrected to compensate for abnormality of serum albumin concentration.<sup>8</sup> However, this AGc attributes a fixed negative charge to albumin, taking no consideration for pH effects on the imidazole groups of albumin.<sup>7</sup> In addition, this AGc ignores the phosphate contribution to all of the weak acids that might need to be considered.<sup>27–29</sup>

## Stewart approach

### Concept of the Stewart approach

Stewart<sup>9,10</sup> questioned the traditional approach for acid–base equilibrium evaluation. He modeled a solution that contained a complex mixture of ions of constant charge over the physiological pH range (strong ions), non-volatile proton donor/acceptors which transfer H<sup>+</sup> within the physiological pH range (weak acid/base), and the volatile bicarbonate–CO<sub>2</sub> buffer system.<sup>8</sup> Key aspect of Stewart’s concept was the classification of each variable as dependent or independent in determining the H<sup>+</sup> concentration of the solution. In his theory, there are three responsible variables to independently determine the dissociation of water, and consequently the hydrogen ion concentration, in order to maintain electrical neutrality: (1) strong ion difference (SID), (2) total concentration of weak acids (A<sub>TOT</sub>), and (3) partial pCO<sub>2</sub> of the solution.<sup>8,30</sup> Thus, in the Stewart’s approach, metabolic disorders are the results of changes in SID or A<sub>TOT</sub>.<sup>7,31</sup>

### Apparent SID and effective SID

Apparent SID (SIDa) represents the difference between measured strong cations and strong anions.<sup>7</sup> With the development of devices capable of detecting “unmeasured” ions (which we could not measure routinely), current calculation of the SIDa contains the following ions<sup>7</sup>

$$\text{SIDa} = \left( [\text{Na}^+] + [\text{K}^+] + [\text{Ca}^{2+}] + [\text{Mg}^{2+}] \right) - \left( [\text{Cl}^-] + [\text{L-lactate}^-] \right)$$

where Na denotes sodium, K denotes potassium, Ca denotes calcium, Mg denotes magnesium, and Cl denotes chloride.

On the other hand, SID calculated to account for electrical neutrality is viewed as the effective SID (SIDE).<sup>7</sup> The SIDE can be calculated as the sum of bicarbonate and weak acids ([A<sup>-</sup>]), mainly albumin and phosphate<sup>8</sup>

$$\text{SIDE} = [\text{HCO}_3^-] + [\text{Alb}^-] + [\text{Pi}^-]$$

where Alb denotes albumin and Pi denotes inorganic phosphate

### SIG

Although the law of electrical neutrality in the body requires SIDA and SIDE to be equal, failure to measure the concentration of all strong and weak ions in plasma yields a gap between the two. Thus, SIG, the difference of SIDA and SIDE, quantifies [unmeasured anions]–[unmeasured cations] of both strong and weak ions.<sup>7</sup>

One of the theoretical advantages of SIG over AG is the pure representation of unmeasured ions. Although both AG and SIG represent unmeasured ions, the “unmeasured” ions derived from AG are composed of [Mg<sup>2+</sup>], [Ca<sup>2+</sup>], [A<sup>-</sup>] (mainly albumin and phosphate), [Lactate<sup>-</sup>], and [other ions] clinicians do not routinely measure, whereas the unmeasured ions expressed by the SIG are composed of just [other ions]. While normal AG ranges from 7 to 17 mEq/L when using [K<sup>+</sup>] for the calculation, SIG is close to zero in normal situations.<sup>8</sup> Although the albumin-corrected AG eliminates the effect of hypo/hyper albuminemia, the gap still persists.

### A<sub>TOT</sub>

Consideration of A<sub>TOT</sub> alternations for acid–base disorders is another key aspect of this approach compared to the traditional one.<sup>7,31</sup> A<sub>TOT</sub>, representing all non-bicarbonate buffers, is made up of mainly serum albumin and other minor charges such as phosphate and globulins.<sup>7,31</sup> In the Stewart approach, an increase in A<sub>TOT</sub> would result in metabolic acidosis and a decrease would result in metabolic alkalosis.<sup>7</sup>

There is a controversy over the existence of A<sub>TOT</sub> acidosis/alkalosis.<sup>32,33</sup> Although observations in vitro show that alterations in albumin concentration can affect acidity, there is no credible demonstration that the living organism, especially the liver, regulates albumin to maintain acid–base homeostasis.<sup>30</sup> One of the explanations is that the theoretical slight weak acid loss secondary to hypoproteinemia is compensated for by a decrease in SID (adjusted SID) without changes in pH, HCO<sub>3</sub><sup>-</sup>, and BE as commonly seen in ICU, rather than a complex acid–base disorder such as a mixed metabolic acidosis/hypoalbuminemic alkalosis.<sup>20,34,35</sup>

Although the traditional approach and the physicochemical approach originated from different concepts as mentioned above, their mathematical comparison showed very few differences once model coefficients are estimated in the consistent manner.<sup>12</sup> Representation of the bicarbonate buffers is almost identical, and representation of non-bicarbonate buffering in the van Slyke equation can be derived from the equations of Stewart. Representation of electrical neutrality comes from the preservation of charge equation described by

Singer and Hastings.<sup>16</sup> For both approaches, measurement of plasma protein concentration is essential if unmeasured anions are to be distinguished from protein buffers.<sup>12</sup> However, many clinical researches have still been conducted on which method is more informative and useful in clinical situations, and there has been no consistent conclusion. In order to find the reasons of the consistency, we conducted a literature search focusing on two main comparisons: diagnostic and prognostic performance of those approaches.

## Literature search

The PubMed Database was initially searched from inception to 15 November 2016 to compare the physicochemical approach with the traditional approach. The search was performed with the relevant medical subject heading terms and strategies: ((SID) OR (strong ion gap)) AND ((AG) OR (BE)). References of selected publications were individually inspected for additional articles that might have been omitted or overlooked in the electronic database search.

The inclusion criteria for the review were (a) studies using both approaches for the same population and (b) studies comparing the diagnostic and/or predictive abilities directly or indirectly. Studies using the traditional methods with AG but without AGc were excluded because non-corrected AG lacks consideration of abnormal albumin concentration commonly seen in the ICU, and many studies already have shown that the simple AG cannot detect acid–base disorders that the Stewart method can identify.<sup>36–39</sup> Nonhuman studies, case reports, abstracts, and unpublished or any studies in which full text was not available were excluded.

## Results

Our electrical literature search revealed 192 studies. One hundred and five nonhuman studies, case reports, abstracts, or otherwise irrelevant studies were excluded. Among 87 potentially relevant articles, we exclude 41 studies that did not compare the two approaches as for diagnostic and/or predictive performance and 29 studies that did not calculate corrected AG for the comparison. Thus, the remaining 17 articles were included in this review. Eight studies compared their diagnostic abilities and 12 articles compared their prognostic performances (Table 1).

### *Inconsistent results on the superiority of one approach over the other approach*

While 10 studies have shown the potential superiority of the Stewart approach,<sup>6,27–29,40,44,46,48–50</sup> four articles failed to show the superiority of the physicochemical approach over the traditional one,<sup>33,41–43</sup> and three articles even showed greater strength of the traditional method than the modern one.<sup>24,45,47</sup>

## Discussion

### *Reasons for inconsistent results on diagnostic performance*

Our literature search shows a discrepancy over the ability to detect acid–base disturbances on diagnostic performance of the two approaches. There are several possible explanations for the discordance. The first thing to be mentioned is the calculation of each variable in both approaches. Table 1 shows there are many differences in inclusive ions, especially lactate, phosphate, and magnesium ion, of each calculation of AGc and SIG among the studies. In addition, cumulative differences or errors in each variable should be considered. As each mathematical equation contains more measurement, there could be greater variability in the parameters, such as SIDa, SIDe, and SIG in Stewart approach, because the differences are exaggerated via complicated mathematical calculations.<sup>51</sup> As shown by Matousek et al.,<sup>12</sup> there should be no difference between the approaches from a mathematical perspective. However, it is true only when the same ions are measured and taken into account and each measurement is accurate. Those differences of each calculation and potential cumulative errors could lead to the discordance about the usefulness as a diagnostic tool between the two approaches.

Another potential reason is technological differences or errors in measuring each variable. Morimatsu et al.<sup>52</sup> showed that chloride measurements, made with point-of-care blood gas and electrolyte analyzers, differed significantly from those made using central laboratory biochemistry analyzers, resulting in different SID values and assessments of the acid–base status. Nguyen et al.<sup>51</sup> compared two central laboratory analyzers for electrolyte measurement and reported that the biochemistry laboratory analyzers have large differences from each other. It should be noted that 12 of 17 articles in our review measured electrolytes using central laboratory analyzers, many of which are currently using diluted blood sample and indirect ion selective electrodes in order to measure the electrolytes, rather than blood gas analyzers (Table 1). Measurements by this method are affected by hypoalbuminemia and could be inaccurate compared with the ones measured by blood gas analyzers.<sup>53</sup> Studies that used indirect ion selective electrodes could lead to wrong calculation and acid–base interpretation, which could make an implausible conclusion. Thus, interpretation of the results in papers comparing these approaches needs attention on the analyzer that each study used. We found a wide variety of machines and technologies used to measure pH, pCO<sub>2</sub>, and electrolytes in those articles, which could be one of the reasons for the inconsistent results on this topic.

Reference value of each parameter is another problem. The dependency on site recommends reference value should be determined in each institution.<sup>46</sup> However, our review showed that while only five studies collected healthy controls for the reference,<sup>33,40,46,49,50</sup> other studies

**Table 1.** Differences in studied population, measured ions, calculation of variables, and references among articles comparing the two approaches.

Author	Studied population	n	Measurements of electrolytes	What is AGc corrected for?	Reference of AGc (mmol/L)	Calculation of SIG	Reference of SIG	Main results and comments
Fencel et al. <sup>40</sup>	ICU	152	Blood gas analyzer	Alb	>21 (based on healthy subjects)	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	>14 (based on healthy subjects)	While unmeasured strong anions represented by SIG detected 35% of patients with normal BE, AGc found 59% of hidden metabolic acid–base disturbances
Cusack et al. <sup>41</sup>	ICU	100	Blood gas analyzer	Alb	>12	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	>0	SIG and SDe in Stewart principle appear to offer no advantage in prediction of outcome
Rocktaeschel et al. <sup>42</sup>	ICU	300	Central laboratory	Alb	N/A	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	N/A	AUROC curves of AGc, SDe, and SIG for mortality prediction were relatively small
Hucker et al. <sup>43</sup>	Accident and emergency department	1424	Central laboratory	Alb	N/A	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	N/A	All of each single variable in both approach have similar and unreliable predictive value
Martin et al. <sup>24</sup>	Surgical ICU, trauma	2152	Central laboratory	Alb, Lac	N/A	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	>0	AUROC for mortality was strong for AGc with AUROC values of 0.68 compared with that for SIG (0.54)
Gunnerson et al. <sup>27</sup>	ICU	9799	Central laboratory	Alb, Pi, Lac	N/A	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	>50% of SBE	AGc identified only 84% of patients classified as SIG acidosis
Dubin et al. <sup>33</sup>	ICU	935	Central laboratory	Alb	3 SD above or below the mean of 7 normal volunteers	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	3 SD above or below the mean of 7 normal volunteers	SIG, not AGc, was an independent predictors of mortality (OR 1.065; 95% CI 1.03–1.10; p = 0.001) When AGc was included in acid–base analysis, the Stewart approach did not offer any diagnostic or prognostic advantages
Kaplan and Kellum <sup>28</sup>	ICU, major trauma	78	Central laboratory	Alb, Pi, Lac	N/A	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	N/A	Although AGc had acceptable ROC curves (0.86) for 28-day mortality, it was significantly inferior to SIG (0.96) (p = 0.018)
Boniatti et al. <sup>29</sup>	ICU (medical and surgical)	175	Central laboratory	Alb, Pi, Lac	>=17	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	>2	There was significant difference between survivors and non-survivors in SIG (p = 0.01), but not in AGc (p = 0.11)
Abdulraof Menesi et al. <sup>44</sup>	Kidney transplant	83	Central laboratory	Alb	>16	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	>3	A greater percentage of patients presented with an increase in unexplained anions by SIG than by AGc (42 vs 32%, respectively) (p value: N/A)
Ratanarat et al. <sup>45</sup>	Medical and surgical ICU	410	Blood gas analyzer	Alb	>12	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	>0	According to ROC curves, the predictive ability to discriminate between survivors and non-survivors of AGc and SIG were 0.72 and 0.67, respectively

(Continued)

Table 1. (Continued)

Author	Studied population	n	Measurements of electrolytes	What is AGc corrected for?	Reference of AGc (mmol/L)	Calculation of SIG	Reference of SIG	Main results and comments
Zheng et al. <sup>6</sup>	Nephrology ICU, metabolic acidosis	78	Central laboratory	Alb	N/A	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	N/A	SIG value was associated with mortality at 24 h, 72 h, 1 week, 1 month, and 3 months after acute kidney injury, whereas AGc was not associated with mortality at each follow-up
Antonogiannaki et al. <sup>46</sup>	Emergency department	365	Central laboratory	Alb	> 17 (based on healthy volunteers)	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	> 6 (based on healthy volunteers)	Significantly fewer patients with unmeasured anions acidosis were identified with AGc than those with SIGc (p=0.001)
Ho et al. <sup>47</sup>	ICU	6878	Blood gas analyzer	Alb	N/A	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Alb <sup>-</sup> , Pi <sup>-</sup> (Lac <sup>-</sup> )	N/A	The abilities to predict hospital mortality in SIG (AUROC 0.52) and SDe (0.63) are modest, whereas AGc (0.67) and BE (0.69) has stronger ability to differentiate between survivors and non-survivors
Morgan et al. <sup>48</sup>	CPB	60	Blood gas analyzer	Alb	> 20	Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	> 4	AUROC of SIG for detecting "unmeasured anions" was significantly higher than that of AGc (0.81 vs 0.79; p = 0.048)
Guérin et al. <sup>49</sup>	Chronic respiratory failure	128	Central laboratory	Alb	8 healthy volunteers	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	8 healthy volunteers	The Stewart approach detected high SDe in 13% of normal SBE and in 20% of normal AGc, and low SDe in 22% of non-elevated HCO <sub>3</sub> <sup>-</sup> , providing better diagnostic performance
Shen et al. <sup>50</sup>	Acute pancreatitis	186	Central laboratory	Alb, Lac	13 health volunteers	Mg <sup>2+</sup> , Ca <sup>2+</sup> , Lac <sup>-</sup> , Alb <sup>-</sup> , Pi <sup>-</sup>	13 health volunteers	SIG, but not AGc, had significant independent correlations with disease severity

ICU: intensive care unit; AGc: corrected anion gap; BE: base excess; SBE: standard base excess; SID: strong ion difference; SDe: effective strong ion difference; SIG: strong ion gap; SIGc: corrected strong ion gap; Mg: magnesium; Ca: calcium; Alb: albumin; Pi: inorganic phosphate; Lac: lactate; HCO<sub>3</sub>: bicarbonate; AUROC: area under receiver operating characteristic curve; ROC: receiver operating characteristic; N/A: not applicable; OR: odds ratio; CI: confidential interval; SD: standard deviation.

used pre-determined numbers,<sup>29,41,44,45,48</sup> and the method for reference value selection in those studies was not specified.<sup>6,24,27,28,42,43</sup> These incongruences of reference value due to arbitrary choice may cause a variety of discordant results. As there is no consensus on the normal range of each variable, especially in the Stewart approach, we recommend that future researchers collect healthy controls for reference in each research institute.

We also need to pay attention to the differences in the normal result range between the two approaches in these studies, since more than one parameter in each method aims to represent the same concept. For example, Boniatti et al. defined normal SBE as  $-5$  to  $+5$  and normal SIDe as  $38-42$  mEq/L. Since changes in BE represent changes in SIDe if  $A_{TOT}$  is normal,<sup>8</sup> the large difference of normal range ( $10$  vs  $4$ ) could mislead the interpretation. This sort of “unfair” comparison might be one reason of the inconsistent results.

Studied populations need another consideration. Several studies showed that patients with renal failure,<sup>54</sup> liver diseases,<sup>55</sup> sepsis and trauma<sup>56</sup> often have accumulations of unmeasured anions. However, Dubin et al.<sup>33</sup> and Ho et al.<sup>47</sup> reported patient demographics in their studies; the percentage of patients with shock, acute renal failure, and hepatic failure was only 13%, 13%, and 4%, respectively, in one study, and liver diseases were only 2% in the other. A study by Cusack et al.<sup>41</sup> included a high proportion of post-elective surgery patients, who generally have low severity of illness and low mortality. Hucker et al.<sup>43</sup> did not provide details about reasons for admission, patients’ illness severity, or the underlying medical conditions of patients in their accident and emergency department. For patients with severe illness, measuring more ions and involving them into variables such as AG and SIG could demonstrate their potential ability to detect unmeasured anions, revealing more detailed acid–base disturbances, no matter which approach is used. For populations with a small number of severe patients, measuring more ions would not be needed for detailed analysis of acid–base disorders. Thus, the combination of variety of the studied populations and the aforementioned differences of calculation in each variable among those studies could be one of the reasons of their inconsistent results.

### *Reasons for inconsistent results on prognostic performance*

Those factors as the potential reasons for inconsistent results on diagnostic performance could also yield a controversy about the prognostic performance of the two approaches. Some authors have investigated the predictive value of the traditional approach and the physicochemical Stewart approach, mainly, AGc versus SIG. One of their questions is “Is there any association between AGc or SIG and outcomes?” or “Can AGc or SIG levels be used as a marker of poor outcomes?” Here the difference of measured and involved ions in each calculation could again mislead the

conclusion. Simple comparison of AGc with SIG does not always answer these particular questions. Although both parameters represent unmeasured ions, consideration of lactate for AGc and SIG depends on each individual study. A bulk of evidence has shown that the level of lactate is associated with poor prognosis.<sup>57,58</sup> If we would like to simply compare the prognostic abilities of the two approaches, the contribution of lactate should be removed from their equations. Only five of all 17 articles remove the effect of lactate from their calculations of AGc and SIG.

It is not only lactate but also other ions, such as magnesium, that need to be considered when comparing the two methods. The changes in magnesium concentration are usually so small that they may usually be neglected, but this simplification is not applicable if the changes are significant. Theoretically, an increased level of magnesium reduces the AG, increases SID, and does not change SIG. There are no studies so far that compare these approaches for patients with abnormal serum magnesium concentrations. Thus, radical question of the comparison should not be “Is there any association between AGc or SIG and outcomes?” but “Is there any association between unmeasured anions that we does not measure in clinical practice and outcomes?” In order to answer this question directly, we need to exclude the contribution of lactate and other measured ion.

Finally, we cannot forget the effect of fluids used for resuscitation, which lead to iatrogenic acidosis. Hayhoe et al.<sup>59</sup> found 40% of acidosis were attributed to the use of polygeline, which acts as an acid resulting in increased unmeasured circulating anions. Similarly, gelatin-derived colloids have also been found to iatrogenically increase the SIG due to increased unmeasured anions.<sup>59</sup> None of the studies included in our review provided detailed information about the type and volume of administered resuscitation fluids. This iatrogenically fluid-induced increment of SIG and metabolic acidosis in less critical patients is not expected to have many adverse outcomes, and therefore, the prognostic value of these indices of the Stewart approach could be wrongly affected.

## **Conclusion**

Although the traditional approach and the Stewart approach are seen as complementary giving the same information about the acid–base phenomena despite their different concepts, our literature search shows inconsistent results on the comparison between the traditional approach and the physicochemical approach for their diagnostic and prognostic performance. Many studies to date have crucial limitations in comparing these approaches. Those limitations are considered the reasons for the discrepancy in clinical researches.

## **Declaration of conflicting interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## ORCID iDs

Satoshi Kimura  <https://orcid.org/0000-0002-0408-5836>

Hiroshi Morimatsu  <https://orcid.org/0000-0002-6744-0508>

## References

- Henderson LJ. The theory of neutrality regulation in the animal organism. *Am J Physiol* 1907; 18: 427–448.
- Hasselbalch KA. Die Berechnung der Wasserstoffzahl des blutes auf der freien und gebundenen Kohlensäure desselben, und die Sauerstoffbindung des Blutes als Funktion der Wasserstoffzahl. *Biochem Z* 1916; 78: 112–144.
- Siggaard-Andersen O, Engel K, Jorgensen K, et al. A micro method for determination of pH, carbon dioxide tension, base excess and standard bicarbonate in capillary blood. *Scand J Clin Lab Invest* 1960; 12: 172–176.
- Schwartz WB and Relman AS. A critique of the parameters used in the evaluation of acid-base disorders. “Whole-blood buffer base” and “standard bicarbonate” compared with blood pH and plasma bicarbonate concentration. *N Engl J Med* 1963; 268: 1382–1388.
- Bunker JP. The great trans-atlantic acid-base debate. *Anesthesiology* 1965; 26: 591–594.
- Stewart PA. Independent and dependent variables of acid-base control. *Respir Physiol* 1978; 33(1): 9–26.
- Stewart PA. *How to understand acid-base: a quantitative acid-base primer for biology and medicine*. New York: Elsevier North Holland Inc, 1981.
- Siggaard-Andersen O. *The acid-base status of the blood*. 4th ed. Copenhagen: Munksgaard, 1974, and Baltimore: William & Wilkins Company, 1974.
- Kellum JA and Elbers PWG. *Stewart’s textbook of acid-base* Lulu Enterprises, UK Ltd, 2009.
- Todorović J, Nešović-Ostojić J, Milovanović A, et al. The assessment of acid-base analysis: comparison of the “traditional” and the “modern” approaches. *Med Glas (Zenica)* 2015; 12(1): 7–18.
- Kishen R, Honoré PM, Jacobs R, et al. Facing acid–base disorders in the third millennium—the Stewart approach revisited. *Int J Nephrol Renovasc Dis* 2014; 7: 209–217.
- Matousek S, Handy J and Rees SE. Acid-base chemistry of plasma: consolidation of the traditional and modern approaches from a mathematical and clinical perspective. *J Clin Monit Comput* 2011; 25(1): 57–70.
- Wooten EW. Calculation of physiological acid-base parameters in multicompartments systems with application to human blood. *J Appl Physiol (1985)* 2003; 95(6): 2333–2344.
- Van Slyke DD. Studies of acidosis I. The bicarbonate concentration of the blood plasma; its significance and its determination as a measure of acidosis. *J Biol Chem* 1917; 30: 289–246.
- Hughes R and Brain MJ. A simplified bedside approach to acid–base: fluid physiology utilizing classical and physicochemical approaches. *Anaesth Intens Care Med* 2013; 14(10): 445–452.
- Singer RB and Hastings AB. An improved clinical method for the estimation of disturbances of the acid-base balance of human blood. *Medicine (Baltimore)* 1948; 27(2): 223–242.
- Astrup P, Jorgensen K, Siggaard-Andersen O, et al. The acid-base metabolism. A new approach. *Lancet* 1960; 1(7133): 1035–1039.
- Severinghaus JW. Acid-base balance nomogram: a Boston-Copenhagen detente. *Anesthesiology* 1976; 45(5): 539–541.
- Siggaard-Andersen O. The van Slyke equation. *Scand J Clin Lab Invest Suppl* 1977; 146: 15–20.
- Kellum JA. Determinants of blood pH in health and disease. *Crit Care* 2000; 4(1): 6–14.
- Brackett NC, Cohen JJ and Schwartz WB. Carbon dioxide titration curve of normal man. Effect of increasing degrees of acute hypercapnia on acid-base equilibrium. *N Engl J Med* 1965; 272: 6–12.
- Prys-Roberts C, Kelman GR and Nunn JF. Determination of the in vivo carbon dioxide titration curve of anaesthetized man. *Br J Anaesth* 1966; 38(7): 500–509.
- Staempfli HR and Constable PD. Experimental determination of net protein charge and A(tot) and K(a) of nonvolatile buffers in human plasma. *J Appl Physiol (1985)* 2003; 95(2): 620–630.
- Martin M, Murray J, Berne T, et al. Diagnosis of acid-base derangements and mortality prediction in the trauma intensive care unit: the physicochemical approach. *J Trauma* 2005; 58(2): 238–243.
- Morgan TJ, Cowley DM, Weier SL, et al. Stability of the strong ion gap versus the anion gap over extremes of pCO<sub>2</sub> and pH. *Anaesth Intensive Care* 2007; 35(3): 370–373.
- Figge J, Rossing TH and Fencl V. The role of serum proteins in acid-base equilibria. *J Lab Clin Med* 1991; 117(6): 453–467.
- Gunnerson KJ, Saul M, He S, et al. Lactate versus non-lactate metabolic acidosis: a retrospective outcome evaluation of critically ill patients. *Crit Care* 2006; 10(1): R22.
- Kaplan LJ and Kellum JA. Comparison of acid-base models for prediction of hospital mortality after trauma. *Shock* 2008; 29(6): 662–666.
- Boniatti MM, Cardoso PR, Castilho RK, et al. Acid-base disorders evaluation in critically ill patients: we can improve our diagnostic ability. *Intensive Care Med* 2009; 35(8): 1377–1382.
- Adrogué HJ and Madias NE. Assessing acid-base status: physiologic versus physicochemical approach. *Am J Kidney Dis* 2016; 68(5): 793–802.
- Rastegar A. Clinical utility of Stewart’s method in diagnosis and management of acid-base disorders. *Clin J Am Soc Nephrol* 2009; 4(7): 1267–1274.
- Balasubramanian N, Havens PL and Hoffman GM. Unmeasured anions identified by the Fencl-Stewart method predict mortality better than base excess, anion gap, and lactate in patients in the pediatric intensive care unit. *Crit Care Med* 1999; 27(8): 1577–1581.
- Dubin A, Meneses MM, Masevicius FD, et al. Comparison of three different methods of evaluation of metabolic acid-base disorders. *Crit Care Med* 2007; 35(5): 1264–1270.

34. Wilkes P. Hypoproteinemia, strong-ion difference, and acid-base status in critically ill patients. *J Appl Physiol* (1985) 1998; 84(5): 1740–1748.
35. Siggaard-Andersen O and Fogh-Andersen N. Base excess or buffer base (strong ion difference) as measure of a non-respiratory acid-base disturbance. *Acta Anaesthesiol Scand Suppl* 1995; 107: 123–128.
36. Miller AC, Subramanian RA, Safi F, et al. Influenza A 2009 (H1N1) virus in admitted and critically ill patients. *J Intensive Care Med* 2012; 27(1): 25–31.
37. Kaplan LJ and Kellum JA. Initial pH, base deficit, lactate, anion gap, strong ion difference, and strong ion gap predict outcome from major vascular injury. *Crit Care Med* 2004; 32(5): 1120–1124.
38. Antonini B, Piva S, Paltenghi M, et al. The early phase of critical illness is a progressive acidic state due to unmeasured anions. *Eur J Anaesthesiol* 2008; 25(7): 566–571.
39. Kaplan LJ, Cheung NH, Maerz L, et al. A physicochemical approach to acid-base balance in critically ill trauma patients minimizes errors and reduces inappropriate plasma volume expansion. *J Trauma* 2009; 66(4): 1045–1051.
40. Fencel V, Jabor A, Kazda A, et al. Diagnosis of metabolic acid-base disturbances in critically ill patients. *Am J Respir Crit Care Med* 2000; 162(6): 2246–2251.
41. Cusack RJ, Rhodes A, Lochhead P, et al. The strong ion gap does not have prognostic value in critically ill patients in a mixed medical/surgical adult ICU. *Intensive Care Med* 2002; 28(7): 864–869.
42. Rocktaeschel J, Morimatsu H, Uchino S, et al. Unmeasured anions in critically ill patients: can they predict mortality? *Crit Care Med* 2003; 31(8): 2131–2136.
43. Hucker TR, Mitchell GP, Blake LD, et al. Identifying the sick: can biochemical measurements be used to aid decision making on presentation to the accident and emergency department. *Br J Anaesth* 2005; 94(6): 735–741.
44. Abdulraof Menesi F, Verzola D, Villaggio B, et al. Evaluation of metabolic acidosis in patients with a kidney graft: comparison of the bicarbonate-based and strong ion-based methods. *Transplant Proc* 2011; 43(4): 1055–1062.
45. Ratanarat R, Sodapak C, Poompichet A, et al. Use of different approaches of acid-base derangement to predict mortality in critically ill patients. *J Med Assoc Thai* 2013; 96(Suppl. 2): S216–S223.
46. Antonogiannaki EM, Mitrouska I, Amargianitakis V, et al. Evaluation of acid-base status in patients admitted to ED-physicochemical vs traditional approaches. *Am J Emerg Med* 2015; 33(3): 378–382.
47. Ho KM, Lan NS, Williams TA, et al. A comparison of prognostic significance of strong ion gap (SIG) with other acid-base markers in the critically ill: a cohort study. *J Intensive Care* 2016; 4: 43.
48. Morgan TJ, Anstey CM and Wolf MB. A head to head evaluation of 8 biochemical scanning tools for unmeasured ions. *J Clin Monit Comput* 2017; 31(2): 449–457.
49. Guérin C, Nesme P, Leray V, et al. Quantitative analysis of acid-base disorders in patients with chronic respiratory failure in stable or unstable respiratory condition. *Respir Care* 2010; 55(11): 1453–1463.
50. Shen X, Ke L, Yang D, et al. The prognostic value of the strong ion gap in acute pancreatitis. *J Crit Care* 2016; 36: 140–145.
51. Nguyen BV, Vincent JL, Hamm JB, et al. The reproducibility of Stewart parameters for acid-base diagnosis using two central laboratory analyzers. *Anesth Analg* 2009; 109(5): 1517–1523.
52. Morimatsu H, Rocktäschel J, Bellomo R, et al. Comparison of point-of-care versus central laboratory measurement of electrolyte concentrations on calculations of the anion gap and the strong ion difference. *Anesthesiology* 2003; 98(5): 1077–1084.
53. Van den Ancker W, Haagen IA and van der Voort PH. Direct sodium measurement prevents underestimation of hyponatremia in critically ill patients. *Intensive Care Med* 2015; 41(3): 553–554.
54. Naka T, Bellomo R, Morimatsu H, et al. Acid-base balance in combined severe hepatic and renal failure: a quantitative analysis. *Int J Artif Organs* 2008; 31(4): 288–294.
55. Kellum JA, Kramer DJ and Pinsky MR. Strong ion gap: a methodology for exploring unexplained anions. *J Crit Care* 1995; 10(2): 51–55.
56. Clowes GH, George BC, Villee CA, et al. Muscle proteolysis induced by a circulating peptide in patients with sepsis or trauma. *N Engl J Med* 1983; 308(10): 545–552.
57. Bakker J, Coffernils M, Leon M, et al. Blood lactate levels are superior to oxygen-derived variables in predicting outcome in human septic shock. *Chest* 1991; 99(4): 956–962.
58. Blow O, Magliore L, Claridge JA, et al. The golden hour and the silver day: detection and correction of occult hypoperfusion within 24 hours improves outcome from major trauma. *J Trauma* 1999; 47(5): 964–969.
59. Hayhoe M, Bellomo R, Liu G, et al. The aetiology and pathogenesis of cardiopulmonary bypass-associated metabolic acidosis using polygeline pump prime. *Intensive Care Med* 1999; 25(7): 680–685.