

How Well Do Survey Studies Capture Alcohol's Harm to Others?

Ingeborg Rossow

Norwegian Institute for Alcohol and Drug Research, Oslo, Norway.

Supplementary Issue: Harm to Others from Substance Use and Abuse

ABSTRACT: Empirical studies assessing alcohol's harm to others very often rely on population survey data. This study addresses some of the problems and challenges in using survey data for this purpose. Such problems include the limited capacity of population surveys in identifying infrequent harm and long-term consequences of drinking. Moreover, the drinker may report the alcohol-related harm or the person being harmed may report the damage. However, irrespective of who reports the harm, causal attribution to drinking is problematic. Challenges for future population surveys to address alcohol's harm to others include the need for improved models and understanding of complex mechanisms to guide empirical studies within the broad range of harm. Study designs other than cross-sectional surveys, such as longitudinal study designs and combinations of population surveys and other data sources, are likely to overcome some of the identified problems in current population surveys of alcohol's harm to others.

KEYWORDS: alcohol, harm to others, population surveys, methodological issues, causal attribution

SUPPLEMENT: Harm to Others from Substance Use and Abuse

CITATION: Rossow. How Well Do Survey Studies Capture Alcohol's Harm to Others?. *Substance Abuse: Research and Treatment* 2015;9(S2) 99–106 doi: 10.4137/SART.S23503.

TYPE: Opinion

RECEIVED: July 06, 2015. **RESUBMITTED:** September 21, 2015. **ACCEPTED FOR PUBLICATION:** September 23, 2015.

ACADEMIC EDITOR: Gregory Stuart, Editor in Chief

PEER REVIEW: Five peer reviewers contributed to the peer review report. Reviewers' reports totaled 1696 words, excluding any confidential comments to the academic editor.

FUNDING: Author discloses no funding sources.

COMPETING INTERESTS: Author discloses no potential conflicts of interest.

CORRESPONDENCE: ir@sirus.no

COPYRIGHT: © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

Paper subject to independent expert blind peer review. All editorial decisions made by independent academic editor. Upon submission manuscript was subject to anti-plagiarism scanning. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties. This journal is a member of the Committee on Publication Ethics (COPE).

Published by Libertas Academica. Learn more about this journal.

Background

While alcohol's harm to the health of the drinker is well described, much less is known about the social harm to drinkers and the negative effects on people other than the drinker (ie, alcohol's harm to others).^{1,2} Indeed, some particular areas of alcohol's harm to others have been studied quite extensively, including fetal alcohol syndrome and other consequences of prenatal alcohol exposure,^{3,4} drunk driving and alcohol-related traffic accidents,⁵ and alcohol-related violence.^{6,7} However, the broad range of other health and social harms that people may experience as a consequence of others' drinking seems to have been less extensively examined, and the total burden to the society because of drinking is still incomplete. In recent years, a renewed and growing interest in measuring the extent and types of alcohol's harm to others (also termed "externalities," "collateral damage," or "second-hand effects of alcohol use") has developed.^{1,8–13} While a comprehensive review of the literature on alcohol's harm to others is still warranted,¹³ there is an impression that many of the studies in this area are based on cross-sectional population surveys.¹³

In this study, the articles that were retrieved from a search in MEDLINE (June 2015, applying the search terms "alcohol" and "harm to others") serve as examples of literature on alcohol's harm to others. Out of 34 journal articles

that solely or in part addressed alcohol's harm to others, two-thirds were published during the past 5 years (since 2010), which may indicate a renewed interest. Moreover, the majority of empirical studies (18 of 25) employed cross-sectional general population survey data to estimate the prevalence of alcohol's harm to others. The remaining articles were reviews, conceptual framework papers, commentaries, or the like,^{1,2,14–20} or they were empirical studies, employing other research designs and methods or partly addressing issues other than the prevalence of harm.^{21–27} Notably, this fairly simple literature search does not provide a complete retrieval of relevant publications on alcohol's harm to others, but it may be suggestive of some characteristics of this literature. Thus, it seems likely that cross-sectional survey data from the general population are frequently used to estimate the extent of various types of harm attributable to others' drinking. While this approach is important, at least as a first step to illustrate that such harm is commonly experienced, this approach also represents a number of challenges and inherent problems. Using the selected cross-sectional survey studies as examples of the epidemiological literature on alcohol's harm to others, this study addresses some of these problems and suggests additional research designs, and methods to further enhance our understanding of the extent and distribution of alcohol's harm to others.



Use of Cross-Sectional Survey Data in Estimating Alcohol's Harm to Others

Whose perspective and what kind of harm? While alcohol's harm to others inherently involves at least two parties (the drinker and the person(s)/company/community being harmed), cross-sectional survey data typically reflect the perspective of only one of these parties, most often the adult victim of others' actions. Thus, among 18 cross-sectional population surveys identified in the recent MEDLINE search and presented in Table 1, 16 studies investigated harm from others' drinking as experienced and reported by the victims. One study investigated harm to children as reported by their parent,²⁸ and in one study, the respondents were asked to report on harm from their own drinking, in addition to the harm they had experienced due to others' drinking.²⁹ As noted in later sections, a one-sided perspective on alcohol's harm to others carries a range of potential problems.

Surveys in this area appear to have been mostly conducted in the general adult population, or they were student samples that illustrated harm to adult persons. Thus, harm to children is often not addressed in these studies, although there are exceptions, for instance, parents reporting harm to their children from others' drinking²⁸ and adults reporting retrospectively on both exposure to alcoholic parents and harm during their childhood.³⁰ Studies of psychosocial adjustment in adult children of alcoholics^{31,32} also rely on retrospective assessments of exposure to parental heavy drinking; however, these studies often address concurrent problems and possible long-term effects of parents' drinking behavior. Moreover, a broad range of harm from someone's drinking is not (solely) experienced by individual persons, for instance, harm to work places (eg, lost productivity) or to public properties (eg, vandalism). These types of harm are difficult to capture with this study design, and are therefore easily neglected in such studies.

Another problem with the victim's perspective is that the respondent is asked either to assess whether an assailant who had been drinking committed the harm or to attribute the harm as a consequence of someone's drinking. Both types of assessments are frequently used in surveys (Table 1), and both carry a significant risk of bias. In many cases, victims may not know whether the assailant had been drinking or not.³³ While the respondent may be better informed when the assailant is well known (eg, a household member), judgements about whether the assailant had been drinking may be quite difficult when the assailant is a stranger or less well known.²⁹ This makes judgements about causal attribution even more difficult and less reliable.

Furthermore, the perpetrator and the victim's perspectives may not deviate only with respect to judgements of drinking and attribution of harm to drinking but also with respect to what is perceived as harmful¹ and the respondent's willingness to report harm. Asking for both perpetrator and victim experiences within the same survey sample, Callinan and Room²⁹ found that the proportions of those

who reported harm from others' drinking (verbal or physical aggression) were much higher than the proportions of those who reported having committed alcohol-related harm. Thus, it seems that at least some instances of alcohol-related harm are less likely to be perceived and reported as harm by the perpetrator compared to the victim. This may reflect unawareness of consequences of one's own drinking or an unwillingness to report harm (or both), suggesting that by using the perpetrator's perspective, the prevalence of harm is likely to be underestimated.

The types of harm typically examined in cross-sectional population surveys are in the form of events (eg, been assaulted) or conditions (eg, financial troubles).¹ These instances of harm often include "social victimization,"³⁴ such as having been insulted, disturbed by loud people, involved in a quarrel, lost friendship, experienced family problems, had property vandalized, been assaulted (or pushed or hit), and had financial troubles. These types of harm have been examined in the studies summarized in Table 1, and they have also been reported in previous empirical studies^{34,35} and reviews.^{1,16} Surveys of social harm, as reported by the drinker, also often include several of these types of harm (eg, fights, quarrels, family, and friendship problems).¹⁶ They may also include other types of harm (or increasing costs) to a company or to society, such as lost working time or poor work performance and being arrested by (or having been in trouble with) the police.¹⁶

Thus, survey approaches seem to cover harm to others according to all of the main types of relationships (ie, the drinker's relationship with family/household, with workplace, and with friends and strangers).¹ It seems, however, that harm to the workplace is more likely to be covered when applying the perpetrator's perspective. Obviously, surveys tend to address harm that is quite frequently experienced by victims. These instances of harm tend to be less severe and they may, in many cases, be characterized as a nuisance (eg, kept awake at night). Correspondingly, highly severe forms of harm with long-lasting effects on the individual and carrying significant costs to society (eg, severe physical injuries and severe child abuse and neglect) are less frequently experienced and they are rarely covered in surveys. This is not to say that the issue of severity of harm has been neglected in survey studies. Indeed, several literature reviews and empirical studies have noted the large variability in the severity of these instances of harm and, in part, have offered approaches to address this aspect of harm.^{1,10,16,34,36} Yet, it may also be noted that certain items used to capture harm from others' drinking also include events that do not necessarily imply harm to any person or belonging other than the drinker. For instance, being a passenger with a driver who had drunk too much¹⁶ will most likely not lead to any harm to the passenger.

Another problem is that several harm measures often used within this survey tradition are rather unspecific and vague with respect to the nature, severity, and extent/amount of harm. Examples of harm measures of this kind include

**Table 1.** Overview of 18 selected cross-sectional survey studies on alcohol's harm to others.

AUTHOR, YEAR, STUDY	SAMPLING METHOD, SAMPLE SIZE	WHOSE PERSPECTIVE	TYPES OF HARM	REFERENCE PERIOD AND QUANTIFICATION OF HARM	TYPES OF DRINKING INVOLVED	RELATIONSHIP OFFENDER-VICTIM	ATTRIBUTION TO OTHER'S DRINKING	METHODOLOGICAL CONCERNS
Diep, 2015 ³⁷	Multi-stage sampling of students from 12 universities in Vietnam, n = 6011	Victim	7 items (eg, insulted, kept awake, physically hurt)	Last year. Frequency of harm (2 cat)	Not specified	Not specified	Not specified	Cross-sectional design hampers causal inference of association with drinking in co-habitants
Dussaillant, 2015 ³⁸	Probabilistic sampling of adults in 13 Chilean regions, n = 1500	Victim	Well-being, health problems	Current? Graded measures	Exposure to heavy drinkers	2 cat (inside or outside household)	Statistical modeling of association between exposure to heavy drinkers and outcomes	Small sample and less precise estimates
Jiang, 2015 ⁴⁵	Random sampling of adults from phone number list in Australia, n = 2649	Victim	4 items on additional tasks and spending time (eg, on cleaning up, transport)	Last year. No quantification	Drinking	6 cat	Attributed harm to most harmful drinker's drinking	Low response rate, too few observations to describe relationship types between harmful drinker and respondent
Mugavin, 2014 ³⁹	Random sampling of adults from phone number list in Australia, n = 2649	Victim	Calling police, seeking health-related services	Last year. Some grading of extent	Drinking	2 categories	Attributed harm to other's drinking	More detail on event warranted, amount of harm experienced left to respondent's judgement
Karriker-Jaffe, 2014 ⁴⁶	Random sampling adults in the USA, n = 10121	Victim	Family problems, crime (property damage or physically hurt)	Last year. No quantification	Drinking	Not specified	Family problems attributed to other's drinking, crime assailant assessed as had been drinking	Cross-sectional design hampers causal attribution of association with neighborhood disadvantage, assessment of other's drinking may be misclassified, assailants not identified
Callinan, 2014 ⁴⁹	Random digit dialing sampling of adults in Australia, n = 5001	Victim and perpetrator	Victim: 8 items (eg, fear, physical abuse, verbal abuse). Perpetrator: verbal abuse, physical abuse	Last year. No quantification	Drinking	Not specified	Some victimization items attributed to other's drinking, other victimization items and perpetrator items related to assailant's drinking	More difficult to assess influence of drinking in strangers than in family/friends
Huhtanen, 2012 ⁴⁴	Random sampling from census records of adults in Finland, n = 4657	Victim	6 items, eg, physically hurt, property damage, insulted	Last year. No quantification	Intoxication	Not specified	Assessed assailant as intoxicated	Gender differences in perception of alcohol related harm
Laslett, 2012 ²⁸	Random sampling of adults from phone number list in Australia, n = 2649	Victim's parent	Been unsupervised, criticized, physically hurt, witness serious violence, call protection agency	Last year. Some grading of severity	Drinking	Not specified	Attributed harm to other's drinking	Harm rate likely underestimated (due to low response rate, uncertainty about harm from respondent's drinking) response bias in attribution of other's drinking, not clear whether one or more children harmed per family

(Continued)

Table 1. (Continued)

AUTHOR, YEAR, STUDY	SAMPLING METHOD, SAMPLE SIZE	WHOSE PERSPECTIVE	TYPES OF HARM	REFERENCE PERIOD AND QUANTIFICATION OF HARM	TYPES OF DRINKING INVOLVED	RELATIONSHIP OFFENDER-VICTIM	ATTRIBUTION TO OTHER'S DRINKING	METHODOLOGICAL CONCERNS
Wilkinson, 2012 ⁵⁸	Random sampling of adults from phone number list in Australia, n = 2649	Victim	5 amenity problems (eg, kept awake, felt unsafe, property damage)	Last year. No quantification	Drinking	Strangers	Attributed harm to other's drinking	
Casswell, 2011 ⁴⁷	Random sampling of 12–80 year olds with landline telephone in New Zealand, n = 3068	Victim	24 items, eg, feel threatened, physically hurt, property damage	Last year. No quantification	Drinking	5 cat	Attributed harm to drinking by the drinker who most negatively affected respondent	Difficult to establish causal relationship in cross-sectional survey
Ferris, 2011 ⁴³	Random sampling of adults from phone number list in Australia, n = 2649	Victim	Mental health, depression or anxiety	Current (?). No quantification	Drinking	7 cat	Statistical modelling exposure to heavy drinkers	Not validated outcome measure
Laslett, 2011 ¹⁰	Random sampling of adults from phone number list in Australia, n = 2649	Victim	18 items, eg, emotionally hurt, feel threatened, physically hurt	Last year. No quantification per harm type	Drinking	7 cat	Attributed harm to heavy drinker's drinking	Population survey not best for studying severe effects
Casswell, 2011 ⁴⁰	Random sampling of 12–80 year olds with landline telephone in New Zealand, n = 3068	Victim	Well-being, health problems	Current (?). Graded measures	Exposure to heavy drinkers	8 cat	Statistical modelling exposure and outcome	Cross-sectional design and insufficient control for confounders
Connor, 2010 ⁵⁹	Random sampling of students from 6 universities in New Zealand, n = 4071	Victim	Unwanted sexual advance, sexual assault	Past 4 weeks (?). No quantification	Drinking	Other student	Harm assessed as effect of other's drinking	Differential willingness (drinkers vs victim) to attribute events to drinking
Connor, 2009 ⁴¹	Stratified sample of adults in New Zealand, n = 16480	Victim	Physical assault, sexual assault	Past year. Number of events	Alcohol use	7 cat	Assessed assailant as affected by alcohol	Victim's report implies misclassification
Rossow, 2004 ⁴²	Three stage (municipality, household, individual) sampling of adults in Norway, n = 2170	Victim	7 items, eg, physically hurt, property damage, insulted, kept awake	Past year. Frq, 3 cat	Intoxication	Not specified	Assessed assailant as intoxicated	Low response rate and under-estimation of harms, not sufficiently distinct harm items
Kellner, 1996 ⁶⁰	Random sampling of adults in Yukon from telephone list and household list, n = 1348	Victim	7 items, eg, insulted, disturbed, passenger of drunk driver, family problems, physically hurt	Past year. No quantification per item but summary score of number of harm items	Drinking	Not specified	Harm assessed as effect of other's drinking	Not discussed
Wechsler, 1994 ⁶¹	Two-stage sampling (university, individuals) of students in the USA, n = 17592	Victim	8 items, eg, insulted, physically hurt, property damage, unwanted sexual advance	Since beginning of school year. No quantification	Drinking	Another student	Harm assessed as effect of other's drinking	Not discussed



“family problems” and “financial troubles.” As noted by Greenfield et al.¹², family problems may be of any severity and duration, may affect few or many children, and may or may not engage social agencies. Similarly, financial troubles may also be of any duration and severity, and they may affect few or many people in a household or family. Other examples of vague harm items used in surveys include “child neglect” and “emotional abuse.”

How is harm quantified? Many survey studies have applied “the last year”/“the past 12 months” as a reference period for reporting harm (Table 1). Within the survey tradition, many types of harm from others’ drinking are damaging events, such as disturbances, property damage, insults, quarrels, and assault (Table 1). In some studies, the frequency (or some other graded measure) of such harmful events is reported,^{9,37–41} allowing for – but not necessarily leading to – the calculation of an (crude) incidence rate. More common, however, is to report the number of persons (proportion) being harmed in a year (Table 1). As some persons experience the same type of harm several times a year,^{40,41} the prevalence figures are likely to underestimate the amount of harmful events.

Some types of harm from others’ drinking are not events, but rather are lasting states, for instance, poor mental health, family troubles, or financial problems.^{9,16,42} Some quantification of these instances of harm at the individual level is seen in studies applying scales for measuring well-being or mental health,^{9,42} whereas there seems to be no tradition in surveys for applying quantifiable measures of other types of nonevent harm, such as family troubles or financial problems.

How is harm attributed to others’ drinking? Survey studies employ various approaches to the attribution of harm to another person’s drinking. Often the respondent (the drinker or the harmed person) is asked to assess whether the harm was caused by drinking (Table 1). Thus, the drinker is asked whether he/she caused the harm to someone else due to his/her own drinking. Correspondingly, the harmed person is asked whether another person’s drinking caused the experienced harm. The validity of such causal attribution is obviously problematic and more so when victims report harm from strangers. Consequently, the harm estimate is likely biased.

Another survey tradition is to ask whether the assailant had been drinking/was intoxicated/was a heavy drinker (Table 1). In studies applying such alcohol-related harm measures, the author(s) often make the implicit assumption that the reported harm is attributable to drinking. For instance, the authors interpret harm by other persons who were intoxicated as social or negative “consequences from other people’s drinking.”^{41,43} Such causal attribution may inflate estimates of harm caused by others’ drinking.

A third approach taken in survey studies is to ask respondents about the exposure to heavy drinkers and about harm victimization and then perform statistical modeling of these exposure and harm measures (Table 1). With this approach, adequate identification of – and control for – all relevant

confounding factors is essential for inferring causality. In reality, however, such control is probably limited to the available covariates, such as sociodemographic characteristics,³⁸ and may thus be insufficient for drawing causal inferences,⁹ and the estimate of harm attributable to others’ drinking is likely biased.

Further Limitations and Gaps

As noted earlier, several characteristics of cross-sectional surveys of alcohol’s harm to others imply that many types of harm are not well covered (mainly, the more severe forms of harm) or they are imprecisely assessed and/or the estimates are biased. Further limitations that apply to cross-sectional studies have also been noted by previous authors (Table 1). These include low response rates, which are likely to cause a downward biased estimation of harm,^{28,41,44} and a lack of time ordering of exposure and outcome measures with a cross-sectional design, thereby hampering the inference of causality.^{37,45,46}

Harm to a person due to another person’s drinking is often interactional.^{1,16} Moreover, persons who drink more frequently and who drink more heavily per occasion compared to others are most at risk, both of causing – at least some types of – harm to others due to their own drinking^{2,6,47} and of experiencing harm from others’ drinking.^{10,12,36,41,43} This may suggest that these types of harm often occur in contexts where both the assailant and the victim have been drinking. In line with this, Cherpitel et al.²³ found that in two-thirds of violent injuries, either the assailant or the victim had been drinking, and in 39% of all injury cases, drinking by someone was seen as causal. In almost a third of the violent injuries, both the assailant and the victim had been drinking, and in these cases, causal attribution to someone’s drinking occurred more often compared to cases where only one of the assailant or the victim had been drinking. This probably applies to a wider range of harm resulting from the social interaction between two or several intoxicated persons.

As population surveys generally interview one person per household or family (or other relevant social networks in this context), a simultaneous perspective of both the drinker and those harmed by his/her drinking is rarely seen. In 1985, Kaye Fillmore published a study on the extent and characteristics of social harm from others’ drinking,³⁴ and she concluded that the “results suggest the need for research which more boldly takes the interaction of perpetrator and victim into account in alcohol-related problems.” In recent years, while many studies have addressed the extent and characteristics of the social victimization of drinking, her conclusion still applies, and the need for better understanding of the social interaction that underlies alcohol’s harm to others is still significant.

Another limitation to most survey studies of alcohol’s harm to others pertains to the assessment of possible long-term effects of drinking and effects of long-term exposure. Indeed, when asking about the exposure to heavy drinkers and the experience of harm, as has been done in several studies,^{9,10,31,32,36,38} it



is possible that long-term effects of exposure to heavy drinkers (eg, in childhood) are captured as part of the outcome measures, but they can hardly be disentangled from short-term effects. Similarly, it is possible that long-term exposure to a heavy drinker (eg, a spouse) has a stronger impact on harm outcomes than does the short-term exposure. However, this may prove difficult to examine with a cross-sectional survey design due to the reversed causality (ie, harm experience may affect length of exposure). Thus, cross-sectional survey studies seem more appropriate for capturing immediate and simple or direct effects, whereas long-term effects, effects of long-term exposure, indirect effects, and interaction effects probably are obtained more adequately by other study designs.

Finally, we may assume that observed correlations between exposure to someone else's drinking and risk of harm are moderated and mediated by various factors. Therefore, the use of theories regarding mechanisms of effect is crucial for identifying important confounding factors and for conducting adequate analyses of causal effects. However, theory-driven analyses of harm attributable to others' drinking seem to be far between. For instance, in a systematic review of prospective studies, when examining the influence of parental drinking on their children, only a smaller fraction of the included studies had employed theories to guide the analyses and assessment of causality.⁴⁸

Challenges for Future Population Surveys Addressing Alcohol's Harm to Others

While the tradition of using surveys to assess social harm from (heavy) drinking and victimization from others' drinking dates back to several decades,^{16,34} this research area may still be regarded as being in its infancy or early years, and thus, the potential for improvements seems considerable. The following are a few suggestions of possible directions for further development in this area.

As noted earlier, several harm measures that are often used are vague and imprecise in various respects. Within the cross-sectional survey tradition, the development of more precise measures of several types of harm is therefore important. Better precision is needed in several respects. The content of the term "harm" should be fairly specific. Thus, a broad concept (for instance, "family problems") may be more precisely covered by specifically asking about the various types of problems and harm that subsume under this category (eg, physical fights, breakup of marriage, and poor economy). Furthermore, some quantification of the harm in question is needed to obtain better precision in the overall account of alcohol's harm to others. For some types of harm, frequency of harmful events (such as fights) may be useful as a quantity indicator, whereas for other types of harm (eg, poor economy), some quantitative indicator is better sought in terms of duration and severity. Development of better precision in harm measurement is likely to include multiple strategies, including qualitative methods⁴⁹ and stages of piloting and validation.

Study designs other than the conventional cross-sectional survey design may offer further potential for improving our assessment of the types, magnitude, and nature of alcohol's harm to others. First, survey studies that sample the involvement of both the assailant and the victim can provide relevant data from the perspectives of both the drinker and the harmed person and thereby can obtain more reliable measures on both exposure and harm. For example, samples of pairs of spouses are used in studies of drinking behavior and intimate partner violence,⁵⁰ and samples of parent-child dyads or entire households have been used in studies of parental drinking and adverse outcomes in children.⁵¹⁻⁵³ These types of survey samples also allow for graded measurements of exposure and measuring of harm without asking about the attribution to drinking, which facilitates the assessment of risk function and provides another approach to the assessment of causality.

Second, among observational studies, prospective cohort studies offer the best capacity for the assessment of causality. This research design is particularly relevant for studies of long-term effects, the effects of long-term exposure, or effects other than immediate effects, for example, the effects of parental drinking on their offspring. A recent scoping review of prospective cohort studies of parental drinking⁵⁴ showed that there is a significantly rich body of literature in this regard; however, few studies addressed adverse outcomes in offspring other than alcohol or other substance use. Thus, there is clearly a need for further prospective cohort studies examining various possible adverse psychosocial consequences for children of parental drinking. When applying prospective cohort studies, there is also clearly a need for theory-driven analyses to enhance the capacity for causal inferences.⁴⁸

Relatedly, there is a potential for applying survey data that are linked with various types of register data. Thus, the exposure measure (eg, parents' or intimate partner's drinking) can be obtained from the survey data, and the outcomes measure (eg, harm to children or spouse) can be obtained from registers. In particular, harm that is not frequently experienced, for instance, severe health-related harm, may be measured more adequately through the use of register data. Moreover, combining survey data on exposure and long-term follow-up data from registers on outcomes may be a valuable approach for the assessment of long-term harmful effects of others' drinking. It is also likely that the good coverage of national registers reduces the attrition bias that is often expected in longitudinal studies. Britt af Klinteberg et al.⁵⁵ study provides one example of using this kind of research design to address the possible effects of paternal drinking on criminal behavior and mortality in the offspring.

Another example of using combined data to study the possible long-term effects of others' drinking is Peter Nilsson's investigation of alcohol exposure during pregnancy and the harmful consequences for the offspring in adulthood.⁵⁶ This study applies a natural experiment of increased

availability of alcohol for exposure measures and register data on education and welfare payments as outcome measures, and the combination of data sources and analytic strategies provides a good basis for inferring a causal effect of prenatal alcohol exposure on adverse outcomes in adulthood. Admittedly, this type of study design does not allow for the estimation of risk curves or alcohol-attributable fraction, but, on the other hand, it suggests that alcohol policies may have long-term consequences borne by individuals other than the drinkers.

The importance of demonstrating harmful effects on third persons and to inform policy-making in a health area is well demonstrated with respect to tobacco. The growing evidence of the effects of passive smoking, a form of harm to others, was indeed a key component in changing policies and practices to denormalize the tobacco use.⁵⁷ In the same way, we may assume that further assessments of how we can improve survey studies of alcohol's harm to others benefit not only the scientific understanding but also the policy's significance.

Acknowledgments

The author is grateful for the valuable comments and suggestions by anonymous reviewers on a previous version.

Author Contributions

Conceived the concepts: IR. Analyzed the data: IR. Wrote the first draft of the manuscript: IR. Agree with manuscript results and conclusions: IR. Developed the structure and arguments for the paper: IR. Made critical revisions and approved final version: IR. The author reviewed and approved of the final manuscript.

REFERENCES

- Room R, Ferris J, Laslett A-M, Livingston M, Mugavin J, Wilkinson C. The drinker's effect on the social environment: a conceptual framework for studying alcohol's harm to others. *Int J Environ Res Public Health*. 2010;7:1855–71.
- Gmel G, Rehm J. Harmful alcohol use. *Alcohol Res Health*. 2003;27(1):52–62.
- Guerri C, Bazinet A, Riley EP. Foetal alcohol spectrum disorders and alterations in brain and behaviour. *Alcohol Alcohol*. 2009;44(2):108–14.
- Streissguth AP, Bookstein FL, Barr HM, Sampson PD, O'malley K, Young JK. Risk factors for adverse life outcomes in fetal alcohol syndrome and fetal alcohol effects. *J Dev Behav Pediatr*. 2004;25(4):228–38.
- Taylor B, Irving HM, Kanteres F, et al. The more you drink, the harder you fall: a systematic review and meta-analysis of how acute alcohol consumption and injury or collision risk increase together. *Drug Alcohol Depend*. 2010;110(1):108–16.
- Room R, Rossow I. Share of violence attributable to drinking. *J Subst Use*. 2001;6:218–28.
- Rossow I, Bye EK. The problem of alcohol-related violence: an epidemiological and public health perspective. In: McMurran M, ed. *Alcohol-Related Violence: Prevention and treatment*. Chichester: Wiley; 2013:3–18.
- Giesbrecht N, Cukier S, Steeves DAN. Collateral damage from alcohol: implications of 'second-hand effects of drinking' for populations and health priorities. *Addiction*. 2010;105(8):1323–5.
- Casswell S, You RQ, Huckle T. Alcohol's harm to others: reduced wellbeing and health status for those with heavy drinkers in their lives. *Addiction*. 2011;106(6):1087–94.
- Laslett A-M, Room R, Ferris J, Wilkinson C, Livingston M, Mugavin J. Surveying the range and magnitude of alcohol's harm to others in Australia. *Addiction*. 2011;106(9):1603–11.
- Babor TF. Commentary on Laslett et al. (2011): alcohol-related collateral damage and the broader issue of alcohol's social costs. *Addiction*. 2011;106(9):1612–3.
- Greenfield TK, Ye Y, Kerr W, Bond J, Rehm J, Giesbrecht N. Externalities from alcohol consumption in the 2005 US National Alcohol Survey: implications for policy. *Int J Environ Res Public Health*. 2009;6(12):3205–24.
- Laslett A-M, Callinan S, Pennay A. The increasing significance of alcohol's harm to others research. *Drugs Alcohol Today*. 2013;13(3):163–72.
- Room R. Alcohol, the individual and society: what history teaches us. *Addiction*. 1997;92(3s1):7–12.
- Paton A. Reflections on alcohol and the young. *Alcohol Alcohol*. 1999;34(4):502–5.
- Room R. Concepts and items in measuring social harm from drinking. *J Subst Abuse*. 2000;12(1):93–111.
- Room R, Graham K, Rehm J, Jernigan D, Monteiro M. Drinking and its burden in a global perspective: policy considerations and options. *Eur Addict Res*. 2003;9(4):165–75.
- Navarro HJ, Doran CM, Shakeshaft AP. Measuring costs of alcohol harm to others: a review of the literature. *Drug Alcohol Depend*. 2011;114(2):87–99.
- Rehm J. The risks associated with alcohol use and alcoholism. *Alcohol Res Health*. 2011;34(2):135.
- Leon DA, Keenan K. Commentary: alcohol, child development and harm to others: a 'hard' problem. *Int J Epidemiol*. 2012;41(4):1097–100.
- Weitzman ER, Chen Y-Y. Risk modifying effect of social capital on measures of heavy alcohol consumption, alcohol abuse, harms, and secondhand effects: national survey findings. *J Epidemiol Community Health*. 2005;59(4):303–9.
- Nutt DJ, King LA, Phillips LD. Drug harms in the UK: a multicriteria decision analysis. *Lancet*. 2010;376(9752):1558–65.
- Cherpitel CJ, Ye Y, Bond J, Room R, Borges G. Attribution of alcohol to violence-related injury: self and other's drinking in the event. *J Stud Alcohol Drugs*. 2012;73(2):277–84.
- Connor J, Casswell S. Alcohol-related harm to others in New Zealand: evidence of the burden and gaps in knowledge. *New Zeal Med J*. 2012;125(1360):11–27.
- Taylor M, Mackay K, Murphy J, et al. Quantifying the RR of harm to self and others from substance misuse: results from a survey of clinical experts across Scotland. *BMJ Open*. 2012;2(4):e000774.
- Yang S, Kramer MS. Paternal alcohol consumption, family transition and child development in a former Soviet country. *Int J Epidemiol*. 2012;41(4):1086–96.
- Wood K, Patterson C, Katikireddi SV, Hilton S. Harms to 'others' from alcohol consumption in the minimum unit pricing policy debate: a qualitative content analysis of UK newspapers (2005–12). *Addiction*. 2014;109(4):578–84.
- Laslett A-M, Ferris J, Dietze P, Room R. Social demography of alcohol – related harm to children in Australia. *Addiction*. 2012;107(6):1082–9.
- Callinan S, Room R. Harm, tangible or feared: young Victorians' adverse experiences from others' drinking or drug use. *Int J Drug Policy*. 2014;25(3):401–6.
- Dube SR, Anda RF, Felitti VJ, Croft JB, Edwards VJ, Giles WH. Growing up with parental alcohol abuse: exposure to childhood abuse, neglect, and household dysfunction. *Child Abuse Negl*. 2001;25(12):1627–40.
- Weitzman ER, Wechsler H. Alcohol use, abuse, and related problems among children of problem drinkers: findings from a national survey of college alcohol use. *J Nerv Ment Dis*. 2000;188(3):148–54.
- Harter SL. Psychosocial adjustment of adult children of alcoholics: a review of the recent empirical literature. *Clin Psychol Rev*. 2000;20(3):311–37.
- Greenfield LA. *Alcohol and Crime: An Analysis of National Data on the Prevalence of Alcohol Involvement in Crime*. Washington, DC: US Department of Justice, Office of Justice Programs, Bureau of Justice Statistics; 1998.
- Fillmore KM. The social victims of drinking. *Br J Addict*. 1985;80(3):307–14.
- Mäkelä P, Fonager K, Hibell B, Nordlund S, Sabroe S, Simpura J. *Drinking Habits in the Nordic Countries*. Oslo: National Institute for Alcohol and Drug Research; 1999.
- Laslett A-M, Catalano P, Chikritzhs Y, et al. *The Range and Magnitude of Alcohol's Harm to Others*. Fitzroy, Vic: AER Centre for Alcohol Policy Research, Turning Point Alcohol and Drug Centre; 2010.
- Diep PB, Knibbe RA, Giang KB, De Vries N. Secondhand effects of alcohol use among students in Vietnam. *Global Health Action*. 2015;8:25848.
- Dussailant F, Fernandez M. Alcohol's harm to others' well-being and health: a comparison between Chile and Australia. *Alcohol Alcohol*. 2015.
- Mugavin J, Livingston M, Laslett A-M. Seeking help because of others' drinking. *Drug Alcohol Rev*. 2014;33(2):161–8.
- Connor J, You R, Casswell S. Alcohol-related harm to others: a survey of physical and sexual assault in New Zealand. *New Zeal Med J*. 2009;122:10–20.
- Rossow I, Hauge R. Who pays for the drinking? Characteristics of the extent and distribution of social harms from others' drinking. *Addiction*. 2004;99(9):1094–102.
- Ferris JA, Laslett A-M, Livingston M, Room R, Wilkinson C. The impacts of others' drinking on mental health. *Med J Aust*. 2011;195(3):22–6.
- Huhtanen P, Tigerstedt C. Women and young adults suffer most from other people's drinking. *Drug Alcohol Rev*. 2012;31(7):841–6.
- Jiang H, Callinan S, Laslett AM, Room R. Correlates of caring for the drinkers and others among those harmed by another's drinking. *Drug Alcohol Rev*. 2015;34(2):162–9.



45. Karriker-Jaffe KJ, Greenfield TK. Gender differences in associations of neighbourhood disadvantage with alcohol's harms to others: a cross-sectional study from the USA. *Drug Alcohol Rev.* 2014;33(3):296–303.
46. Casswell S, Harding JF, You RQ, Huckle T. Alcohol's harm to others: self-reports from a representative sample of New Zealanders. *New Zeal Med J.* 2011;124(1336):75–84.
47. Rossow I, Keating P, Felix L, McCambridge J. Does parental drinking influence children's drinking? A systematic review of prospective cohort studies. *Addiction.* 2015.
48. Rossow I, Felix L, Keating P, McCambridge J. Parental drinking and adverse outcomes in children – a scoping review of cohort studies. *Drug Alcohol Rev.* 2015.
49. Manton E, MacLean S, Laslett A-M, Room R. Alcohol's harm to others: using qualitative research to complement survey findings. *Int J Alcohol Drug Res.* 2014; 3(2):143–8.
50. Caetano R, Cunradi CB, Schafer J, Clark CL. Intimate partner violence and drinking patterns among white, black, and Hispanic couples in the US. *J Subst Abuse.* 2000;11(2):123–38.
51. Alati R, Kinner SA, Hayatbakhsh MR, Mamun AA, Najman JM, Williams GM. Pathways to ecstasy use in young adults: anxiety, depression or behavioural deviance? *Drug Alcohol Depend.* 2008;92(1):108–15.
52. Fergusson DM, Lynskey MT, Horwood LJ. Childhood exposure to alcohol and adolescent drinking patterns. *Addiction.* 1994;89(8):1007–16.
53. Van Der Vorst H, Vermulst AA, Meeus WH, Deković M, Engels RC. Identification and prediction of drinking trajectories in early and mid-adolescence. *J Clin Child Adolesc Psychol.* 2009;38(3):329–41.
54. Britt af Klinteberg B, Almquist Y, Beijer U, Rydelius P-A. Family psychosocial characteristics influencing criminal behaviour and mortality-possible mediating factors: a longitudinal study of male and female subjects in the Stockholm Birth Cohort. *BMC Public Health.* 2011;11(1):756.
55. Nilsson P. *Does A Pint A Day Affect Your Child's Pay? The Effect of Prenatal Alcohol Exposure on Adult Outcomes.* Uppsala University. Cemmap Working Paper; No. CWP22/08; 2008. Available at: <http://dx.doi.org/10.1920/wp.cem>.
56. International Agency for Research on Cancer (IARC). *Evaluating the Effectiveness of Smoke-Free Policies.* Vol 2. Geneva: World Health Organization; 2009.
57. Wilkinson C, Livingston M. Distances to on-and off-premise alcohol outlets and experiences of alcohol-related amenity problems. *Drug Alcohol Rev.* 2012;31(4):394–401.
58. Connor J, Gray A, Kypri K. Drinking history, current drinking and problematic sexual experiences among university students. *Aust N Z J Public Health.* 2010;34(5):487–94.
59. Kellner F, Webster I, Chanteloup F. Describing and predicting alcohol use-related harm: an analysis of the Yukon alcohol and drug survey. *Subst Use Misuse.* 1996;31(11–12):1619–38.
60. Wechsler H, Davenport A, Dowdall G, Moeykens B, Castillo S. Health and behavioral consequences of binge drinking in college: a national survey of students at 140 campuses. *JAMA.* 1994;272(21):1672–7.