

RESEARCH

Open Access



Contributions to the fixed point theory of diagonal operators

Adrian Petruşel^{1,2*}, Ioan A Rus¹ and Marcel-Adrian Şerban¹

*Correspondence:

petrusel@math.ubbcluj.ro

¹Faculty of Mathematics and
Computer Science, Babeş-Bolyai
University Cluj-Napoca,
Kogălniceanu Street, No. 1,
Cluj-Napoca, Romania

²Academy of Romanian Scientists,
Splaiul Independentei no. 54,
Bucharest, 050094, Romania

Abstract

In this paper, we introduce the notion of diagonal operator, we present the historical roots of diagonal operators and we give some fixed point theorems for this class of operators. Our approaches are based on the weakly Picard operator technique, difference equation techniques, and some fixed point theorems for multi-valued operators. Some applications to differential and integral equations are given. We also present some research directions.

MSC: 47H10; 54H25; 47J25; 65J15; 47H09

Keywords: diagonal operator; fixed point; fixed point structure; coupled fixed point; weakly Picard operator; difference equation; multi-valued operator; differential equation; integral equation; research direction

1 Introduction and preliminary notions and results

In this section we will present some useful notions and results concerning diagonal operators, coupled fixed point operators, and iterations of some operators generated by the above concepts.

1.1 Diagonal operators

Let X be a nonempty set and $V : X \times X \rightarrow X$ be an operator. By definition, the operator $U_V : X \rightarrow X$, defined by

$$U_V(x) := V(x, x), \quad \text{for all } x \in X,$$

is called the diagonal operator corresponding to the operator V .

We also consider the following operators generated by an operator $V : X \times X \rightarrow X$:

(1) The operator $C_V : X \times X \rightarrow X \times X$ defined by

$$C_V(x, y) := (V(x, y), V(y, x)).$$

By definition an element

$$(x, y) \in F_{C_V} := \{(u, v) \in X \times X \mid C_V(u, v) = (u, v)\}$$

is called a coupled fixed point of V (see [1]; see also [2, 3]). We remark that

$$(x, x) \in F_{C_V} \iff x \in F_{U_V}.$$

(2) The operator $D_V : X \times X \rightarrow X \times X$ defined by $D_V(x, y) := (y, V(x, y))$.

We have $(x, y) \in F_{D_V} \iff x = y$ and $x \in F_{U_V}$.

(3) The operator $T_V : X \rightarrow \mathcal{P}(X)$ is defined by

$$T_V(x) := F_{V(\cdot, x)}.$$

It is clear that $F_{U_V} = F_{T_V}$.

The aim of this paper is to present some historical roots of the diagonal operators, to study the fixed points of this class of operators, and to give some applications. Some new research directions are also presented.

More precisely, the plan of the paper is the following:

1. Introduction and preliminary notions and results.
2. Historical roots of the diagonal operators.
3. Iterations of the operators C_V and U_V .
4. Iterations of the operator D_V and the difference equation

$$x_{n+2} = V(x_n, x_{n+1}), \quad n \in \mathbb{N}, x_0, x_1 \in X.$$

5. Fixed point results for the operator T_V .

6. Applications.

7. Research directions.

References.

1.2 L -Spaces [4–6]

Following Fréchet [4], we present now the concept of L -space.

Definition 1.1 Let X be a nonempty set. Let

$$s(X) := \{(x_n)_{n \in \mathbb{N}} \mid x_n \in X, n \in \mathbb{N}\}.$$

Let $c(X)$ be a subset of $s(X)$ and $\text{Lim} : c(X) \rightarrow X$ be an operator. By definition, the triple $(X, c(X), \text{Lim})$ is called an L -space (denoted by (X, \rightarrow)) if the following conditions are satisfied:

- (i) if $x_n = x$, for all $n \in \mathbb{N}$, then $(x_n)_{n \in \mathbb{N}} \in c(X)$ and $\text{Lim}(x_n)_{n \in \mathbb{N}} = x$;
- (ii) if $(x_n)_{n \in \mathbb{N}} \in c(X)$ and $\text{Lim}(x_n)_{n \in \mathbb{N}} = x$, then, for all subsequences $(x_{n_i})_{i \in \mathbb{N}}$ of $(x_n)_{n \in \mathbb{N}}$, we have $(x_{n_i})_{i \in \mathbb{N}} \in c(X)$ and

$$\text{Lim}(x_{n_i})_{i \in \mathbb{N}} = x.$$

By definition, an element of $c(X)$ is said to be a convergent sequence and $\text{Lim}(x_n)_{n \in \mathbb{N}}$ is the limit of this sequence. If $\text{Lim}(x_n)_{n \in \mathbb{N}} = x$, then we will write

$$x_n \rightarrow x \quad \text{as } n \rightarrow \infty.$$

Remark 1.1 An L -space is any set endowed with a structure implying a notion of convergence for sequences. As examples of L -spaces we mention the following:

- (1) If (X, τ) is a Hausdorff topological space, then $(X, \xrightarrow{\tau})$ is an L -space.
- (2) If (X, d) is a metric space then (X, \xrightarrow{d}) is an L -space.
- (3) If $(X, \|\cdot\|)$ is a normed space then $(X, \xrightarrow{\|\cdot\|})$ and (X, \rightharpoonup) are L -spaces.

1.3 Weakly Picard operators [5, 7–11], etc.

Let (X, \rightharpoonup) be an L -space. By definition, $f : X \rightarrow X$ is said to be a weakly Picard operator (WPO) if the sequence $(f^n(x))_{n \in \mathbb{N}}$ of successive approximations converges for all $x \in X$ and the limit (which may depend on x) is a fixed point of f . If f is weakly Picard operator and $F_f = \{x^*\}$, then, by definition, f is called a Picard operator (PO).

If $f : X \rightarrow X$ is WPO, then we define the operator $f^\infty : X \rightarrow X$ by

$$f^\infty(x) := \lim_{n \rightarrow \infty} f^n(x).$$

Now let (X, d) be a metric space. By definition, a WPO $f : X \rightarrow X$ is called ψ -WPO if $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is increasing, continuous in 0 with $\psi(0) = 0$, and

$$d(x, f^\infty(x)) \leq \psi(d(x, f(x))), \quad \text{for all } x \in X.$$

1.4 Measures of noncompactness [12–17], etc.

Let X be a Banach space. We will denote by $P_b(X)$ the family of all nonempty bounded subsets of S .

We will use the symbol $\alpha_K : P_b(X) \rightarrow \mathbb{R}_+$, for the Kuratowski measure of noncompactness on X , while $\alpha_H : P_b(x) \rightarrow \mathbb{R}_+$ will denote the Hausdorff measure of noncompactness on X . The following results are well known.

Darbo’s theorem *Let X be a Banach space, $Y \in P_{b,cl,cv}(X)$ and $f : Y \rightarrow Y$ be an operator. We suppose that:*

- (i) f is continuous;
- (ii) there exists $l \in [0, 1[$ such that:

$$\alpha_K(f(A)) \leq l \cdot \alpha_K(A), \quad \text{for all } A \in P(Y).$$

Then:

- (a) $F_f \neq \emptyset$;
- (b) F_f is a compact subset of Y .

Sadovskii’s theorem *Let X be a Banach space, $Y \in P_{b,cl,cv}(X)$ and $f : Y \rightarrow Y$ be an operator. We suppose that:*

- (i) f is continuous;
- (ii) $\alpha_H(f(A)) < \alpha_H(A)$, for all $A \in P(Y)$ with $\alpha_H(A) \neq 0$.

Then:

- (a) $F_f \neq \emptyset$;
- (b) F_f is a compact subset of Y .

1.5 Fixed point structures [14]

Let \mathcal{C} be a class of structured sets (ordered sets, topological spaces, metric spaces, Banach spaces, Hilbert spaces, ...). Let us denote by Set^* the class of all nonempty sets. For $X \in \text{Set}^*$ we shall use the notations:

$$P(X) := \{Y \subset X \mid Y \neq \emptyset\} \quad \text{and} \quad P(\mathcal{C}) := \{A \in P(X) \mid X \in \mathcal{C}\},$$

$$\mathbb{M}(A, B) := \{f : A \rightarrow B \mid f \text{ is an operator}\} \quad \text{and} \quad \mathbb{M}(A) := \mathbb{M}(A, A).$$

Now we consider the following multi-valued operators:

$$S : \mathcal{C} \rightarrow \text{Set}^*, X \mapsto S(X) \subset P(X),$$

$$M : D_M \subset P(\mathcal{C}) \times P(\mathcal{C}) \rightarrow \mathbb{M}(P(\mathcal{C}), P(\mathcal{C})), (A, B) \mapsto M(A, B) \subset \mathbb{M}(A, B).$$

By a fixed point structure on $X \in \mathcal{C}$, we understand a triple $(X, S(X), M)$ with the following properties:

- (i) $A \in S(X) \Rightarrow (A, A) \in D_M$;
- (ii) $A \in S(X), f \in M(A) \Rightarrow F_f \neq \emptyset$.

The following examples illustrate this notion.

(1) *The fixed point structure (f.p.s.) of Tarski*

Let \mathcal{C} be the class of complete lattices. If (X, \preceq) is a complete lattice,

$$S(X) := \{A \in P(X) \mid (A, \preceq) \text{ is a complete lattice}\}$$

and

$$M(A) := \{f : A \rightarrow A \mid f \text{ is increasing}\},$$

then, by Tarski's fixed point theorem, we see that $(X, S(X), M)$ is a f.p.s.

(2) *The f.p.s. of contractions*

Let \mathcal{C} be the class of complete metric spaces. For a complete metric space (X, d) , we consider

$$S(X) := P_{cl}(X)$$

and

$$M(A) := \{f : A \rightarrow A \mid f \text{ is a contraction}\}.$$

Then the Banach contraction principle implies that $(X, S(X), M)$ is a f.p.s.

(3) *The f.p.s. of Schauder*

Let \mathcal{C} be the class of Banach spaces. For a Banach space X , if we consider

$$S(X) := P_{cp,cv}(X)$$

and

$$M(A) = M(A, A) := C(A, A),$$

then, by Schauder's theorem, we see that $(X, S(X), M)$ is a f.p.s.

A similar notion of fixed point structure can be defined for multi-valued operators (see [14], pp.139-142).

A triple $(X, S(X), M^0)$ is a multi-valued fixed point structure (m.f.p.s.) if the following properties hold:

- (i) $S(X) \subset P(X)$ and $S(X) \neq \emptyset$;
- (ii) $M^0 : P(X) \multimap \bigcup_{Y \in P(X)} M^0(Y), Y \multimap M^0(Y) \subset \mathbb{M}(Y)$, where $\mathbb{M}(Y)$ is the set of all self multi-valued operators on Y ;
- (iii) $Y \in S(X), T \in M^0(Y) \Rightarrow F_T \neq \emptyset$.

1.6 Acyclic topological spaces [15, 18]

Let X be a compact metric space and $H_q(x)$ be the q -dimensional Čech homology on \mathbb{Q} of X . By definition, X is called acyclic if $H_q(X) = 0$ for $q \geq 1$ and $H_0(X) \approx \mathbb{Q}$.

The following result is a particular case of the Eilenberg-Montgomery theorem (see [15, 17, 18]).

Theorem 1.1 *Let Y be a compact convex subset of a Banach space E and $T : Y \rightarrow P(Y)$ be an upper semi-continuous multi-valued operator with acyclic values. Then $F_T \neq \emptyset$.*

2 Historical roots of the diagonal operators

There are some roots of the diagonal operators as the following examples reveal.

Example 2.1 (Difference equations [19–22]) Let (X, \rightarrow) be an L -space and $V : X \times X \rightarrow X$ be an operator. We consider the following difference equation:

$$x_{n+2} = V(x_n, x_{n+1}), \quad n \in \mathbb{N}, x_0, x_1 \in X.$$

Let us suppose that $(x_n)_{n \in \mathbb{N}}$ is a solution of this equation with the property that

$$x_n \rightarrow x^* \quad \text{as } n \rightarrow \infty.$$

If the function V is continuous, then we have $x^* = V(x^*, x^*)$. Thus, x^* is a fixed point of the diagonal operator corresponding to V .

Example 2.2 (Krasnoselskii (1955) [23]) Let X be a Banach space, $Y \in P_{b,cl,cv}(X)$ and $f, g : Y \rightarrow Y$ be two operators. We suppose that:

- (i) f is a contraction;
- (ii) g is complete continuous;
- (iii) $f(x) + g(y) \in Y$, for all $x, y \in Y$.

Under these conditions, Krasnoselskii proved that the operator $f + g : Y \rightarrow Y$ has at least a fixed point.

If we consider the operator $V : Y \times Y \rightarrow Y$, defined by

$$V(x, y) := f(x) + g(y),$$

then $f + g : Y \rightarrow Y, x \mapsto f(x) + g(x)$ is the diagonal operator corresponding to V .

Example 2.3 (Browder (1966; [24]; see also [12, 13, 16, 25–28], etc.) Let X be a Banach space, $Y \in P_{b,op}(X)$ and $V : X \times X \rightarrow X$ be a continuous operator. Then Browder considered the operator $U : \bar{Y} \rightarrow X$ defined by $U(x) := V(x, x)$.

Moreover, Browder introduced the following notions:

- (1) U is strictly semicontractive if, for each fixed x in X , $V(\cdot, x)$ is Lipschitzian with constant $l < 1$ and $V(x, \cdot)$ is compact.
- (2) U is weakly semicontractive if, for each x in X , the operator $V(\cdot, x)$ is nonexpansive and $V(x, \cdot)$ is compact.

Example 2.4 (Ziebur (1962 and 1965); [29, 30]) Let $b \in \mathbb{R}^m$ and $h, k \in \mathbb{R}_+^*$. One consider a set $\Omega := \prod_{i=1}^m [b_i - k, b_i + k]$ and a function $f \in C([0, h] \times \Omega, \mathbb{R}^m)$. Let us consider the Cauchy problem

$$(C_1) \quad \begin{cases} x' = f(t, x), \\ x(0) = x_0. \end{cases}$$

Then Ziebur introduced a function $F \in C([0, h] \times \Omega \times \Omega, \mathbb{R}^m)$ with the following property:

- (a) $F(t, x, x) = f(t, x)$, for all $t \in [0, h], x \in \mathbb{R}^m$;
- (b) $F(t, \cdot, x)$ is increasing;
- (c) $F(t, x, \cdot)$ is decreasing.

The following Cauchy problem was also considered:

$$(C_2) \quad \begin{cases} x' = F(t, x, y), \\ y' = F(t, y, x), \end{cases} \quad (x, y)(0) = (x_0, x_0)$$

and it is proved that if the Cauchy problem (C_2) has a unique solution then the problem (C_1) has a unique solution too and the Picard sequence converges to that solution.

Example 2.5 (Amann (1973, 1977), Opoitsev (1975; [2, 31–33]))

In [2] the author presents the following result ‘concerning so-called intervined’:

Let X be a chain complete ordered set possessing a least and a greatest element. Let $g : X \times X \rightarrow X$ be a mapping such that:

- (i) $g(\cdot, y) : X \rightarrow X$ is increasing for every $y \in X$;
- (ii) $g(x, \cdot) : X \rightarrow X$ is decreasing for every $x \in X$.

Then there exist two points $\bar{x}, \hat{x} \in X$ such that $\bar{x} \leq \hat{x}$, $g(\bar{x}, \hat{x}) = \bar{x}$, and $g(\hat{x}, \bar{x}) = \hat{x}$. Moreover, if $f(x) := g(x, x)$ for all $x \in X$, then

$$F_f \subset [\bar{x}, \hat{x}].$$

Remark 2.1 For other examples on this topic see [1, 33–37], etc.

Example 2.6 (Quasilinear differential equations; see [17, 34, 35, 38], etc.) Diagonal operators also appears by the linearization of a quasilinear differential equations. For example, let us consider the Cauchy problem:

$$\begin{cases} x'(t) = A(t)x(t) + f(t, x(t)), & t \in [a, b], \\ x(a) = x_0, \end{cases} \tag{2.1}$$

where $A \in C([a, b], \mathbb{R}^{m \times m})$ and $f \in C([a, b] \times \mathbb{R}^m, \mathbb{R}^m)$.

Then, for each $u \in C([a, b], \mathbb{R}^m)$ with $u(a) = x_0$ we consider the linearized Cauchy problem

$$\begin{cases} x'(t) = A(t)x(t) + f(t, u(t)), & t \in [a, b], \\ x(a) = x_0, \end{cases} \tag{2.2}$$

Let $S \subset C^1([a, b], \mathbb{R}^m)$ be the solution set of the problem (2.1) and let $T(u)$ the solution set of the problem (2.2). Then $S = F_T$.

3 Iterations of C_V and U_V

The following result is the starting point for this section.

Lemma 3.1 *Let (X, \rightarrow) be an L -space and $V : X \times X \rightarrow X$ be an operator. We suppose that the operator C_V is WPO. Then we have:*

- (a) U_V is WPO;
- (b) $C_V^\infty(x, x) = (U_V^\infty(x), U_V^\infty(x))$, for all $x \in X$;
- (c) if C_V is a PO, then:
 - (1) $F_{C_V} = \{(x^*, x^*)\}$;
 - (2) U_V is PO and $F_{U_V} = \{x^*\}$.

Proof (a) + (b). We remark that $C_V(x, x) = (U_V(x), U_V(x))$. From this we have

$$C_V^n(x, x) = (U_V^n(x), U_V^n(x))$$

and

$$C_V^n(x, x) \rightarrow C^\infty(x, x) = (U_V^\infty(x), U_V^\infty(x)) \quad \text{as } n \rightarrow \infty, \text{ for all } x \in X.$$

(c). Follows from the definition of PO and from (a) + (b).

Now we consider instead of the L -space (X, \rightarrow) a metric space (X, d) .

We will consider, on $X \times X$, the following metrics:

$$d_1((x_1, y_1), (x_2, y_2)) := d(x_1, x_2) + d(y_1, y_2), \tag{3.1}$$

$$d_2((x_1, y_1), (x_2, y_2)) := [(d(x_1, x_2))^2 + (d(y_1, y_2))^2]^{\frac{1}{2}}, \tag{3.2}$$

and

$$d_\infty((x_1, y_1), (x_2, y_2)) := \max(d(x_1, x_2), d(y_1, y_2)). \tag{3.3}$$

□

In the case of metric spaces we have the following result.

Theorem 3.1 *Let (X, d) be a metric space and $V : X \times X \rightarrow X$ be an operator. Then:*

- (a) If C_V is ψ – WPO with respect to the metric d_1 , then U_V is a θ – WPO where

$$\theta(r) := \frac{1}{2}\psi(2r), \quad r \in \mathbb{R}_+.$$

(b) If C_V is ψ – WPO with respect to the metric d_2 , then U_V is a θ – WPO where

$$\theta(r) := \frac{1}{\sqrt{2}}\psi(\sqrt{2}r), \quad r \in \mathbb{R}_+.$$

(c) If C_V is ψ – WPO with respect to the metric d_∞ , then U_V is a ψ – WPO.

Proof (a). From

$$d_1((x, y), C_V^\infty(x, y)) \leq \psi(d_1((x, y), C_V(x, y))), \quad \text{for all } (x, y) \in X \times X,$$

it follows that

$$d_1((x, x), (U_V^\infty(x), U_V^\infty(x))) \leq \psi(d_1((x, x), (V(x, x), V(x, x)))), \quad \text{for all } x \in X,$$

so

$$d(x, U_V^\infty(x)) \leq \frac{1}{2}\psi(2d(x, U_V(x))), \quad \text{for all } x \in X.$$

(b). If C_V is ψ – WPO with respect to the metric d_2 then

$$d_2((x, x), C_V^\infty(x, x)) \leq \psi(d_2((x, x), (V(x, x), V(x, x)))), \quad \text{for all } x \in X,$$

which means that

$$\sqrt{2}d(x, U_V^\infty(x)) \leq \psi(\sqrt{2}d(x, U_V(x))), \quad \text{for all } x \in X,$$

so we get the conclusion.

(c). If C_V is ψ – WPO with respect to the metric d_∞ then

$$d_\infty((x, x), C_V^\infty(x, x)) \leq \psi(d_\infty((x, x), (V(x, x), V(x, x)))), \quad \text{for all } x \in X,$$

so

$$d(x, U_V^\infty(x)) \leq \psi(d(x, U_V(x))), \quad \text{for all } x \in X,$$

which proves that U_V is a ψ – WPO. □

The following result is a coupled fixed point theorem in a complete b -metric space, which has as an additional conclusion the fact that the operator C_V is a Picard operator.

Theorem 3.2 ([39]) *Let (X, d) be a complete b -metric space with constant $s \geq 1$. Let $V : X \times X \rightarrow X$ be an operator. Assume that there exists $k \in (0, 1)$ such that, for all $(x, y), (u, v) \in X \times X$, we have*

$$d(V(x, y), V(u, v)) + d(V(y, x), V(v, u)) \leq k[d(x, u) + d(y, v)].$$

Then there exists a unique solution $(x^*, y^*) \in X \times X$ of the following coupled fixed point problem :

$$\begin{cases} x = V(x, y), \\ y = V(y, x). \end{cases} \tag{3.4}$$

and, for any initial point $(x_0, y_0) \in X \times X$, the sequences $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}$ defined, for $n \in \mathbb{N}$, by

$$\begin{cases} x_{n+1} = V(x_n, y_n), \\ y_{n+1} = V(y_n, x_n), \end{cases} \tag{3.5}$$

converge to x^* and, respectively, to y^* as $n \rightarrow \infty$.

In particular, the operator $C_V : X \times X \rightarrow X \times X$ given by $C_V(x, y) := (V(x, y), V(y, x))$ is a Picard operator.

Proof For the sake of completeness we present here the sketch of the proof. We introduce on $Z := X \times X$ the functional $\tilde{d} : Z \times Z \rightarrow \mathbb{R}_+$ defined by

$$\tilde{d}((x, y), (u, v)) := d(x, u) + d(y, v).$$

Notice that, as before, \tilde{d} is a b -metric on Z with the same constant $s \geq 1$ and, if the space (X, d) is complete, then (Z, \tilde{d}) is complete too.

We consider now the operator $F : Z \rightarrow Z$ given by

$$F(x, y) := (V(x, y), V(y, x)).$$

It is easy to prove now that F is a contraction in (Z, \tilde{d}) with constant $k \in (0, 1)$, i.e.,

$$\tilde{d}(F(z), F(w)) \leq k\tilde{d}(z, w), \quad \text{for all } z, w \in Z.$$

Thus, we can apply for F the b -metric space version of the contraction principle given by Czerwik (see, for example, Theorem 12.2, p.115 in [40]) and we get the conclusion. \square

Another result involves the coupled fixed point problem in a complete metric space under a contraction condition on the graphic of the operator. In this case, we will see that C_V is a weakly Picard operator. Let us also point out that we denote $V^2(x, y) := V(V(x, y), V(y, x))$ and $V^2(y, x) := V(V(y, x), V(x, y))$, while the graphic of an operator $U : X \rightarrow X$ is denoted by $\text{Graph}(U) := \{(x, y) \in X \times X : y = U(x)\}$.

Theorem 3.3 *Let (X, d) be a complete metric space and $V : X \times X \rightarrow X$ be an operator. Assume that there exists $k \in (0, 1)$ such that, for all $(x, y) \in X \times X$, we have*

$$d(V(x, y), V^2(x, y)) + d(V(y, x), V^2(y, x)) \leq k[d(x, V(x, y)) + d(y, V(y, x))].$$

Then there exists at least one solution $(x^*, y^*) \in X \times X$ of the coupled fixed point problem (3.4) and, for any initial point $(x_0, y_0) \in X \times X$, the sequences $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}$ defined by (3.5) converge to x^* and, respectively, to y^* as $n \rightarrow \infty$.

In particular, $C_V : X \times X \rightarrow X \times X$ given by $C_V(x, y) := (V(x, y), V(y, x))$ is a weakly Picard operator.

Proof We consider again on $Z := X \times X$ the functional $\tilde{d} : Z \times Z \rightarrow \mathbb{R}_+$ defined by

$$\tilde{d}((x, y), (u, v)) := d(x, u) + d(y, v).$$

As before, (Z, \tilde{d}) is a complete metric space.

We consider now the operator $F : Z \rightarrow Z$ given by

$$F(x, y) := (V(x, y), V(y, x)).$$

It is easy to prove now that F is a graphic contraction in (Z, \tilde{d}) with constant $k \in (0, 1)$, i.e.,

$$\tilde{d}(F(z), F^2(z)) \leq k\tilde{d}(z, F(z)), \quad \text{for all } z = (x, y) \in Z.$$

Thus, the conclusion follows by the graphic contraction principle; see, for example, [8] or [10]. □

Remark 3.1 It is worth to mention that the above results can easily be considered in the framework of an ordered metric space X , under contraction type conditions imposed for comparable elements (with respect to a partial order relation \leq on X); see for example [3, 39, 41–43], etc.

We will consider now some qualitative properties concerning the behavior of an operator $A : X \rightarrow X$, where (X, d) is a metric space. More precisely, we consider the following notions:

(i) *the fixed point equation*

$$x = A(x), \quad x \in X$$

is called *well-posed* if $F_A = \{x_A^*\}$ and for any $x_n \in X$, $n \in \mathbb{N}$ a sequence in X such that

$$d(x_n, A(x_n)) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

we have

$$x_n \rightarrow x_A^* \quad \text{as } n \rightarrow \infty.$$

(ii) The operator A has the *Ostrowski property* (or the operator A has the limit shadowing property) if $F_A = \{x^*\}$ and for any $x_n \in X$, $n \in \mathbb{N}$ a sequence in X such that

$$d(x_{n+1}, A(x_n)) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

we have

$$x_n \rightarrow x^* \quad \text{as } n \rightarrow \infty.$$

(iii) *The fixed point equation*

$$x = A(x), \quad x \in X$$

is *generalized Ulam-Hyers stable* if there exists an increasing function $\theta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, continuous in 0 with $\theta(0) = 0$, and for each $\varepsilon > 0$ and for each solution y^* of the inequality

$$d(y, A(y)) \leq \varepsilon$$

there exists a solution x^* of the fixed point equation with

$$d(x^*, y^*) \leq \theta(\varepsilon).$$

By the above notions, we have the following result.

Theorem 3.4 *Let (X, d) be a metric space and $V : X \times X \rightarrow X$ be an operator. Then:*

- (a) *If the fixed point equations for C_V is well-posed and $F_{C_V} = \{(x^*, x^*)\}$, then the fixed point equations for U_V is well-posed.*
- (b) *If the operator C_V has the Ostrowski property and $F_{C_V} = \{(x^*, x^*)\}$, then the operator U_V has the Ostrowski property.*
- (c) *If the fixed point equations for C_V is generalized Ulam-Hyers stable and all the fixed points of C_V are of the form (x^*, x^*) , then the fixed point equations for U_V is generalized Ulam-Hyers stable.*

Proof (a). If $F_{C_V} = \{(x^*, x^*)\}$ then $F_{U_V} = \{x^*\}$. Let $(x_n)_{n \in \mathbb{N}} \subset X$ such that $d(x_n, U_V(x_n)) \rightarrow 0$ as $n \rightarrow +\infty$. Then

$$d_*((x_n, x_n), C_V(x_n, x_n)) \rightarrow 0, \quad n \rightarrow +\infty,$$

where d_* is one of the metrics on $X \times X$ defined by (3.1)-(3.3). Since the fixed point equation for C_V is well-posed

$$d_*((x_n, x_n), (x^*, x^*)) \rightarrow 0, \quad n \rightarrow +\infty,$$

therefore

$$d(x_n, x^*) \rightarrow 0, \quad n \rightarrow +\infty,$$

so the fixed point equation for U_V is well-posed.

- (b). Let $(x_n)_{n \in \mathbb{N}} \subset X$ such that $d(x_n, U_V(x_{n+1})) \rightarrow 0$ as $n \rightarrow +\infty$. Then

$$d_*((x_n, x_n), C_V(x_{n+1}, x_{n+1})) \rightarrow 0, \quad n \rightarrow +\infty,$$

where d_* is one of the metrics on $X \times X$ defined by (3.1)-(3.3). Since the operator C_V has the Ostrowski property

$$d_*((x_n, x_n), (x^*, x^*)) \rightarrow 0, \quad n \rightarrow +\infty,$$

therefore

$$d(x_n, x^*) \rightarrow 0, \quad n \rightarrow +\infty,$$

so the operator U_V has the Ostrowski property.

(c). If $(x^*, x^*) \in F_{C_V}$ then $x^* \in F_{U_V}$. Let $\varepsilon > 0$ and $y^* \in X$ be a solution of the inequality

$$d(y, U_V(y)) \leq \varepsilon,$$

then

$$d_1((y^*, y^*), C_V(y^*, y^*)) = 2d(y^*, U_V(y^*)) \leq 2\varepsilon.$$

From the Ulam-Hyers stability of the fixed point equation for C_V we see that there exists $(x^*, x^*) \in F_{C_V}$ such that

$$d_1((y^*, y^*), (x^*, x^*)) \leq \theta(2\varepsilon) \iff 2d(y^*, x^*) \leq \theta(2\varepsilon),$$

so

$$d(y^*, x^*) \leq \theta_1(\varepsilon),$$

where $\theta_1(t) = \frac{1}{2}\theta(2t)$ which proves that the fixed point equations for U_V is generalized Ulam-Hyers stable.

If we replace the metric d_1 on $X \times X$ defined by (3.1) with the metric d_2 or d_∞ on $X \times X$ defined by (3.2), respectively, by (3.3), we get the same conclusion but instead of $\theta_1(t)$ we have a different function, in the case of d_2 we have $\theta_2(t) = \frac{1}{\sqrt{2}}\theta(\sqrt{2}t)$, and in the case of d_∞ we have $\theta_\infty(t) = \theta(t)$. □

Remark 3.2 For other considerations on the operator C_V see [39, 44–56], etc.

4 Iterations of the operator D_V and the difference equation $x_{n+2} = V(x_n, x_{n+1})$, $n \in \mathbb{N}, x_0, x_1 \in X$

Let X be a nonempty set and $V : X \times X \rightarrow X$. Let $(x_n, y_n) := D_V^n(x_0, y_0), (x_0, y_0) \in X \times X$. We remark that $(x_n)_{n \in \mathbb{N}}$ is a solution of the difference equation

$$x_{n+2} = V(x_n, x_{n+1}), \quad n \in \mathbb{N}, x_0, x_1 = y_0 \in X. \tag{4.1}$$

Moreover, we have the following.

Lemma 4.1 *Let (X, \rightarrow) be an L-space and $V : X \times X \rightarrow X$ a continuous operator. Then the following statements are equivalent:*

- (i) D_V is Picard with $F_{D_V} = \{(x^*, x^*)\}$;
- (ii) D_V^2 is Picard operator with $F_{D_V^2} = \{(x^*, x^*)\}$;

(iii) x^* is globally asymptotically stable solution of the difference equation

$$x_{n+2} = V(x_n, x_{n+1}), \quad n \in \mathbb{N}.$$

Proof (i) \Leftrightarrow (ii). See Lemma 2.2 in [57]. (i) \Leftrightarrow (iii). See [20, 21]. □

Theorem 4.1 *Let (X, d) be a complete metric space, $V : X \times X \rightarrow X$ and $\varphi : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$. We suppose that:*

- (i) φ is increasing;
- (ii) $\sum_{k=0}^\infty \phi^k(r) < +\infty$, where $\phi(r) = \varphi(r, r)$, $r \in \mathbb{R}_+$;
- (iii) $\varphi(r, 0) + \varphi(0, r) \leq \phi(r)$, $r \in \mathbb{R}_+$;
- (iv) $d(V(x_0, x_1), V(x_1, x_2)) \leq \varphi(d(x_0, x_1), d(x_1, x_2))$, for all $x_0, x_1, x_2 \in X$.

Then

- (a) $F_{U_V} = \{x^*\}$.
- (b) If $(x_n)_{n \in \mathbb{N}}$ is a solution of the difference equation (4.1) then $x_n \rightarrow x^*$ as $n \rightarrow +\infty$.

Proof Let $x_0, x_1 \in X$ and $(x_n)_{n \in \mathbb{N}}$ be defined by the difference equation (4.1). We have

$$\begin{aligned} d(x_2, x_3) &= d(V(x_0, x_1), V(x_1, x_2)) \leq \varphi(d(x_0, x_1), d(x_1, x_2)) \\ &\leq \phi(d_\infty((x_0, x_1), (x_1, x_2))), \\ d(x_3, x_4) &= d(V(x_1, x_2), V(x_2, x_3)) \leq \varphi(d(x_1, x_2), d(x_2, x_3)) \\ &\leq \varphi(d_\infty((x_0, x_1), (x_1, x_2)), \phi(d_\infty((x_0, x_1), (x_1, x_2)))) \\ &\leq \phi(d_\infty((x_0, x_1), (x_1, x_2))), \\ d(x_4, x_5) &= d(V(x_2, x_3), V(x_3, x_4)) \leq \varphi(d(x_2, x_3), d(x_3, x_4)) \\ &\leq \varphi(\phi(d_\infty((x_0, x_1), (x_1, x_2))), \phi(d_\infty((x_0, x_1), (x_1, x_2)))) \\ &= \phi^2(d_\infty((x_0, x_1), (x_1, x_2))). \end{aligned}$$

By induction we get

$$d(x_n, x_{n+1}) \leq \phi^{\lfloor \frac{n}{2} \rfloor}(d_\infty((x_0, x_1), (x_1, x_2))),$$

thus

$$\begin{aligned} d(x_n, x_{n+p}) &\leq \sum_{i=0}^{p-1} d(x_{n+i}, x_{n+i+1}) \leq \sum_{i=0}^{p-1} \phi^{\lfloor \frac{n+i}{2} \rfloor}(d_\infty((x_0, x_1), (x_1, x_2))) \\ &\leq 2 \sum_{j=\lfloor \frac{n}{2} \rfloor}^{\lfloor \frac{n+p-1}{2} \rfloor} \phi^j(d_\infty((x_0, x_1), (x_1, x_2))) \rightarrow 0 \quad \text{as } n, p \rightarrow +\infty, \end{aligned}$$

so $(x_n)_{n \in \mathbb{N}}$ is fundamental, therefore there exists $x^* \in X$ such that $x_n \rightarrow x^*$ as $n \rightarrow +\infty$.

By the continuity assumption on φ in 0 we have

$$\begin{aligned} d(x^*, V(x^*, x^*)) &\leq d(x^*, x_n) + d(x_n, V(x_{n-1}, x^*)) + d(V(x_{n-1}, x^*), V(x^*, x^*)) \\ &\leq d(x^*, x_n) + \varphi(d(x_{n-2}, x_{n-1}), d(x_{n-1}, x^*)) + \varphi(d(x_{n-1}, x^*), d(x^*, x^*)), \end{aligned}$$

thus $x^* = V(x^*, x^*)$, which means that $x^* \in F_{U_V}$.

If there exist $x^*, y^* \in F_{U_V}$ then

$$\begin{aligned} d(x^*, y^*) &= d(V(x^*, x^*), V(y^*, y^*)) \\ &\leq d(V(x^*, x^*), V(x^*, y^*)) + d(V(x^*, y^*), V(y^*, y^*)) \\ &\leq \varphi(0, d(x^*, y^*)) + \varphi(d(x^*, y^*), 0) \leq \phi(d(x^*, y^*)), \end{aligned}$$

but $\phi(r) < r$ for all $r \in \mathbb{R}_+$, so $d(x^*, y^*) = 0$. □

Theorem 4.2 *Let (X, d) be a complete metric space and $V : X \times X \rightarrow X$. We suppose that there exist $l_1, l_2 \in \mathbb{R}_+$, $l_1 + l_2 < 1$ such that*

$$d(V(x_1, y_1), V(x_2, y_2)) \leq l_1 d(x_1, x_2) + l_2 d(y_1, y_2), \quad \text{for all } x_i, y_i \in X, i = 1, 2.$$

Then

- (a) D_V^2 is a $(l_1 + l_2)$ -contraction;
- (b) $F_{D_V} = \{(x^*, x^*)\}$ and $F_{U_V} = \{x^*\}$;
- (c) if $(x_n)_{n \in \mathbb{N}}$ is a solution of the difference equation

$$x_{n+2} = V(x_n, x_{n+1}), \quad n \in \mathbb{N},$$

then $x_n \rightarrow x^*$ as $n \rightarrow \infty$.

Proof (a). Let us consider the complete metric space $(X \times X, d_\infty)$ where d_∞ is defined by (3.3). We have

$$\begin{aligned} d_\infty(D_V^2(x_1, y_1), D_V^2(x_2, y_2)) &= \max\{d(V(x_1, y_1), V(x_2, y_2)), d(V(y_1, V(x_1, y_1)), V(y_2, V(x_2, y_2)))\} \end{aligned}$$

and

$$\begin{aligned} d(V(x_1, y_1), V(x_2, y_2)) &\leq (l_1 + l_2) \cdot d_\infty((x_1, y_1), (x_2, y_2)), \\ d(V(y_1, V(x_1, y_1)), V(y_2, V(x_2, y_2))) &\leq l_1 d(y_1, y_2) + l_2 d(V(x_1, y_1), V(x_2, y_2)) \\ &\leq (l_1 + l_2) \cdot d_\infty((x_1, y_1), (x_2, y_2)). \end{aligned}$$

Thus

$$d_\infty(D_V^2(x_1, y_1), D_V^2(x_2, y_2)) \leq (l_1 + l_2) \cdot d_\infty((x_1, y_1), (x_2, y_2)),$$

so D_V^2 is an $(l_1 + l_2)$ -contraction.

From Lemma 4.1 we get (b) and (c). □

For related results concerning the difference equation (4.1) see [19–22, 57, 58], etc.

5 Fixed point results for the operator T_V

A possible approach for the study of the fixed points of the operator T_V is given by the following general result.

Lemma 5.1 *Let X be a nonempty set, $(X, S_1(X), M_1)$ be a f.p.s. on X and $(X, S_2(X), M_2^0)$ be a m.f.p.s. on X . Let $Y \in S_1(X) \cap S_2(X)$ and $V : Y \times Y \rightarrow Y$. We suppose that:*

- (i) $S_1(X) \cap S_2(X) \neq \emptyset$;
- (ii) $V(\cdot, x) \in M_1(Y)$, for each $x \in Y$;
- (iii) $T_V \in M_1^0(Y)$.

Then $F_{T_V} \neq \emptyset$ and $F_{U_V} = F_{T_V}$.

Proof Since $Y \in S_1(X) \cap S_2(X)$ and using (ii) we obtain $T_V(x) \neq \emptyset$, for each $x \in Y$. Since $Y \in S_2(X)$ and using (iii) we get $F_{T_V} \neq \emptyset$. On the other hand, $F_{U_V} = F_{T_V}$. □

In particular, we have the following consequences of the above approach.

Theorem 5.1 *Let X be a Banach space and $Y \in P_{cp,cv}(X)$. Let $V : Y \times Y \rightarrow Y$ be an operator such that:*

- (i) $V : Y \times Y \rightarrow Y$ is continuous;
 - (ii) the set $\{u \in Y \mid u = V(u, x)\}$ is convex, for each $x \in Y$.
- Then $F_{T_V} \neq \emptyset$, i.e., there exists $x^* \in Y$ such that $x^* = V(x^*, x^*)$.*

Proof Since $V(\cdot, x) : Y \rightarrow Y$ is continuous and $Y \in P_{cp,cv}(X)$, by Schauder’s fixed point theorem, we get $F_{V(\cdot, x)} \neq \emptyset$, for each $x \in Y$. Moreover, by (ii), the set $F_{V(\cdot, x)}$ is convex, for each $x \in Y$. On the other hand, by the continuity of V , we see that the set $\{(x, u) \in Y \times Y \mid u = V(u, x)\}$ is closed in $Y \times Y$. Thus, the multi-valued operator $T_V : Y \rightarrow P(Y)$ given by $T_V(x) := F_{V(\cdot, x)}$ has a closed graphic. Since the co-domain Y is compact, T_V is upper semi-continuous on Y . Hence, we get $T_V : Y \rightarrow P_{cp,cv}(Y)$ and it is upper semi-continuous. By Bohnenblust-Karlin’s fixed point theorem we get $F_{T_V} \neq \emptyset$. □

Theorem 5.2 *Let X be a uniformly convex Banach space and $Y \in P_{cp,cv}(X)$. Let $V : Y \times Y \rightarrow Y$ be an operator such that:*

- (i) $V : Y \times Y \rightarrow Y$ is continuous;
- (ii) $V(\cdot, x) : Y \rightarrow Y$ is nonexpansive, for each $x \in Y$.

Then $F_{T_V} \neq \emptyset$.

Proof Since $V(\cdot, x) : Y \rightarrow Y$ is nonexpansive and $Y \in P_{cp,cv}(X)$, by Browder-Ghōde-Kirk’s fixed point theorem, we see that the set $F_{V(\cdot, x)}$ is nonempty and convex, for each $x \in Y$. On

the other hand, by the continuity of V , we see that the set $\{(x, u) \in Y \times Y \mid u = V(u, x)\}$ is closed in $Y \times Y$. Thus, the multi-valued operator $T_V : Y \rightarrow P(Y)$ given by $T_V(x) := F_{V(\cdot, x)}$ has a closed graphic. Since the co-domain Y is compact, T_V is upper semi-continuous on Y . Hence, we get $T_V : Y \rightarrow P_{cp,cv}(Y)$ and it is upper semi-continuous. Our conclusion follows by Bohnenblust-Karlin's fixed point theorem. \square

Theorem 5.3 *Let $(X, \|\cdot\|)$ be a Banach space and $Y \in P_{cl,cv}(X)$. Let $V : Y \times Y \rightarrow Y$ be an operator such that:*

(i) *there exists $\alpha \in (0, 1)$ such that, for each $x \in X$, we have*

$$\|V(u, x) - V(v, x)\| \leq \alpha \|u - v\|, \quad \text{for all } u, v \in Y;$$

(ii) *for each $u \in Y$ the operator $V(u, \cdot) : Y \rightarrow Y$ is continuous;*

(iii) *for each $u \in Y$ the set $V(u, Y)$ is relatively compact.*

Then $F_{T_V} \neq \emptyset$.

Proof Since, for every $x \in Y$, the operator $V(\cdot, x) : Y \rightarrow Y$ is a contraction, for each $x \in Y$ there exists a unique $u^* = u^*(x) \in Y$ such that $V(u^*, x) = u^*$. Thus, the operator $T_V : Y \rightarrow Y$ given by $T_V(x) := u^*(x)$ is a self single-valued operator on Y . Notice that

$$T_V(x) = V(u^*(x), x) \quad \text{for each } x \in Y.$$

Moreover, T_V is continuous since

$$\begin{aligned} & \|T_V(x_1) - T_V(x_2)\| \\ &= \|V(u^*(x_1), x_1) - V(u^*(x_2), x_2)\| \\ &\leq \|V(u^*(x_1), x_1) - V(u^*(x_2), x_1)\| + \|V(u^*(x_2), x_1) - V(u^*(x_2), x_2)\| \\ &\leq \alpha \|u^*(x_1) - u^*(x_2)\| + \|V(u^*(x_2), x_1) - V(u^*(x_2), x_2)\| \\ &= \alpha \|T_V(x_1) - T_V(x_2)\| + \|V(u^*(x_2), x_1) - V(u^*(x_2), x_2)\|. \end{aligned}$$

As a consequence, we get the continuity of the single-valued operator T_V :

$$\|T_V(x_1) - T_V(x_2)\| \leq \frac{1}{1-\alpha} \cdot \|V(u^*(x_2), x_1) - V(u^*(x_2), x_2)\| \rightarrow 0 \quad \text{as } x_1 \rightarrow x_2.$$

By (iii), $T_V(Y)$ is relatively compact. Thus, by Schauder's fixed point theorem, there exists $x^* \in Y$ such that $x^* = T_V(x^*)$. As a consequence, $x^* = u^* = V(u^*, x^*) = V(x^*, x^*)$. \square

Theorem 5.4 *Let $(X, \|\cdot\|)$ be a Banach space and $Y \in P_{cp,cv}(X)$. Let $V : Y \times Y \rightarrow Y$ be an operator such that:*

(i) *there exists $\alpha \in (0, 1)$ such that, for each $x \in X$, we have*

$$\|V(u, x) - V(V(u, x), x)\| \leq \alpha \|u - V(u, x)\|, \quad \text{for all } x, u \in Y;$$

(ii) *$V : Y \times Y \rightarrow Y$ is continuous;*

(iii) *the set $\{u \in Y \mid u = V(u, x)\}$ is convex, for each $x \in Y$.*

Then $F_{T_V} \neq \emptyset$.

Proof Notice first that, for every $x \in Y$, the operator $V(\cdot, x) : Y \rightarrow Y$ is a graphic contraction. Thus, for each $x \in Y$, the set $F_{V(\cdot, x)}$ is nonempty. Moreover, by the continuity of V , the set $F_{V(\cdot, x)}$ is closed. Thus, the operator $T_V : Y \rightarrow P(Y)$ given by $T_V(x) := F_{V(\cdot, x)}$ is a multi-valued operator with closed graph. Since Y is compact, we see that T_V is upper semi-continuous on Y with compact and (by (iii)) convex values. The conclusion follows by Bohnenblust-Karlin’s fixed point theorem. \square

Another result of this type can be reached using the above mentioned particular variant of the Eilenberg-Montgomery theorem; see Theorem 1.1.

Theorem 5.5 *Let $(X, \|\cdot\|)$ be a Banach space and $Y \in P_{ep,cv}(X)$. Let $V : Y \times Y \rightarrow Y$ be an operator such that:*

- (i) *for each $x \in X$ the operator $V(\cdot, x)$ is nonexpansive and compact;*
- (ii) *the operator $V : Y \times Y \rightarrow Y$ is continuous.*

Then $F_{T_V} \neq \emptyset$.

Proof Notice first that, by Theorem 1.63 in [59], the set $F_{V(\cdot, x)}$ is nonempty and acyclic, for each $x \in Y$. On the other hand, by the continuity of V , we see that the set $\{(x, u) \in Y \times Y \mid u = V(u, x)\}$ is closed in $Y \times Y$. Thus, the multi-valued operator $T_V : Y \rightarrow P(Y)$ given by $T_V(x) := F_{V(\cdot, x)}$ has a closed graphic. Since the co-domain Y is compact, T_V is upper semi-continuous on Y . Hence, we see that $T_V : Y \rightarrow P(Y)$ has acyclic values and it is upper semi-continuous. The conclusion follows by Theorem 1.1. \square

Remark 5.1 For the case of the Eilenberg-Montgomery fixed point theorem see [15, 34, 35], etc.

Remark 5.2 For the fixed point theory of multivalued operators see [14, 15, 17, 18], etc.

6 Applications

6.1 Fredholm type integral equations

Let $\Omega \subset \mathbb{R}^m$ be a bounded domain and $C(\overline{\Omega})$ be the Banach space with

$$\|x\|_\infty := \max_{t \in \overline{\Omega}} |x(t)|.$$

We consider the integral equation, in $C(\overline{\Omega})$,

$$x(t) = \int_\Omega K(t, s, x(s), x(s)) ds, \quad t \in \overline{\Omega}, \tag{6.1}$$

where $K \in C(\overline{\Omega}, \overline{\Omega} \times \mathbb{R}^2)$.

Let us consider now the operator $V : C(\overline{\Omega}) \times C(\overline{\Omega}) \rightarrow C(\overline{\Omega})$ defined by

$$V(x, y)(t) := \int_\Omega K(t, s, x(s), y(s)) ds.$$

By Theorem 4.1 we have the following.

Theorem 6.1 *We suppose that:*

- (i) $\varphi : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ satisfies conditions (i)-(iii) in Theorem 4.1;

(ii) $|K(t, s, u, v) - K(t, s, v, w)| \leq \frac{1}{\text{mes}(\Omega)} \varphi(|u - v|, |v - w|)$ for all $t, s \in \overline{\Omega}$, $u, v, w \in \mathbb{R}$.

Then:

- (a) Equation (6.1) has a unique solution, $x^* \in C(\overline{\Omega})$.
- (b) The sequence $(x_n)_{n \in \mathbb{N}}$, defined by

$$x_{n+2}(t) = \int_{\Omega} K(t, s, x_n(s), x_{n+1}(s)) ds, \quad t \in \overline{\Omega},$$

converges to x^* for all $x_0, x_1 \in C(\overline{\Omega})$.

6.2 A periodic boundary value problem

We will consider now a periodic boundary value problem of the following type:

$$\begin{cases} -x'' = f(t, x, x), & t \in [a, b], \\ x(a) = 0, & x(b) = 0, \end{cases} \tag{6.2}$$

where $f : [a, b] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is a given continuous function.

This problem is equivalent to a Fredholm type integral equation of the following form:

$$x(t) = \int_a^b G(t, s) f(s, x(s), x(s)) ds, \quad t \in [a, b],$$

where $G : [a, b] \times [a, b] \rightarrow \mathbb{R}_+$ is the corresponding Green function.

Let us define now the operator $V : C[a, b] \times C[a, b] \rightarrow C[a, b]$ defined by

$$V(x, y)(t) := \int_a^b G(t, s) f(s, x(s), y(s)) ds.$$

By Theorem 4.1 we have the following.

Theorem 6.2 *We suppose that:*

- (i) $\varphi : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ satisfies conditions (i)-(iii) in Theorem 4.1;
- (ii) $|f(s, u, v) - f(s, v, w)| \leq \frac{8}{(b-a)^2} \cdot \varphi(|u - v|, |v - w|)$ for all $t, s \in [a, b]$, $u, v, w \in \mathbb{R}$.

Then:

- (a) The boundary value problem (6.2) has a unique solution, x^* .
- (b) The sequence $(x_n)_{n \in \mathbb{N}}$, defined by

$$x_{n+2}(t) = \int_a^b f(s, x_n(s), x_{n+1}(s)) ds, \quad t \in [a, b]$$

converges to x^* for all $x_0, x_1 \in C[a, b]$.

6.3 Other applications

Other applications of the abstract results given in this paper can be obtained for the case of functional-differential equations and functional-integral equations (or inclusions) which appear in [6, 21, 36, 44, 53, 60], etc.

7 Research directions

7.1 Mixed monotone operators

Let (X, \leq) be an ordered set and $U : X \rightarrow X$. Under these conditions there exists $V : X \times X \rightarrow X$ such that:

- (i) $V(\cdot, x)$ is increasing;
- (ii) $V(x, \cdot)$ is decreasing;
- (iii) $U = U_V$.

References: [2, 29, 30], etc.

7.2 Difference equations for diagonal operators

Let (X, \rightarrow) be an L -space and $U : X \rightarrow X$ an operator. Under which conditions is U a diagonal operator with respect to some $V : X \times X \rightarrow X$ such that each solution $(x_n)_{n \in \mathbb{N}}$ of the difference equation

$$x_{n+1} = V(x_n, x_{n+1}), \quad n \in \mathbb{N},$$

converges to a fixed point of U ?

References: [19, 20, 22, 57, 61, 62], etc.

7.3 Fixed point structures approach to diagonal operators

Let $(X, S(X), M)$ be a fixed point structure on X , $Y \in S(X)$ and $V : Y \times Y \rightarrow Y$. Under which conditions on V do we have $U_V \in M(Y)$?

Commentaries Let X be a Banach space, $(X, P_{b,cl,cv}(X), M)$ the fixed point structure of Darbo. Let $Y \in P_{b,cl,cv}(X)$ and $V : Y \times Y \rightarrow Y$. We suppose that:

- (i) V is continuous;
- (ii) $V(\cdot, y)$ is a l -contraction, for all $y \in Y$;
- (iii) $V(x, \cdot)$ is compact, for all $x \in Y$.

Then $U_V \in M(Y)$ (see [12, 13, 27]).

References: [12, 24–27], etc.

Competing interests

All the authors have read Springer Open's guidance on competing interests and we confirm that none of the authors have any competing interests in the manuscript.

Authors' contributions

All the authors have made equal contributions to this work. All authors read and approved the final manuscript.

Acknowledgements

The authors are thankful to the anonymous reviewer(s) for his (her, their) useful remarks and constructive suggestions.

Received: 12 August 2016 Accepted: 22 September 2016 Published online: 03 October 2016

References

1. Guo, D, Lakshmikantham, V: Coupled fixed points of nonlinear operators with applications. *Nonlinear Anal.* **11**, 623-632 (1987)
2. Amann, H: Order Structures and Fixed Points. SAFA, vol. 2, pp. 1-51, University of Calabria (1977)
3. Bota, M-F, Petruşel, A, Petruşel, G, Samet, B: Coupled fixed point theorems for singlevalued operators in b -metric spaces. *Fixed Point Theory Appl.* **2015**, 231 (2015)
4. Fréchet, M: Les espaces abstraits. Gauthier-Villars, Paris (1928)
5. Rus, IA: Picard operators and applications. *Sci. Math. Jpn.* **58**, 191-219 (2003)
6. Rus, IA: Some nonlinear functional differential and integral equations, via weakly Picard operator theory: a survey. *Carpath. J. Math.* **26**(2), 230-258 (2010)

7. Berinde, V, Păcurar, M, Rus, IA: From a Dieudonné theorem concerning the Cauchy problem to an open problem in the theory of weakly Picard operators. *Carpath. J. Math.* **30**, 283-292 (2014)
8. Rus, IA: Weakly Picard mappings. *Comment. Math. Univ. Carol.* **34**(4), 769-773 (1993)
9. Rus, IA, Petruşel, A, Şerban, M-A: Weakly Picard operators: equivalent definitions, applications and open problems. *Fixed Point Theory* **7**(1), 3-22 (2006)
10. Rus, IA: *Generalized Contractions and Applications*. Transilvania Press, Cluj-Napoca (2001)
11. Rus, IA: A Müller's example to Cauchy problem: an operatorial point of view. *Creative Math. Inform.* **21**(2), 215-220 (2012)
12. Nussbaum, RD: The fixed point index and fixed point theorems for k -set-contractions. Ph.D. Dissertation, University of Chicago (1969)
13. Petryshyn, WV: Fixed point theorems for various classes of 1-set-contractive and 1-ball-contractive mappings in Banach spaces. *Trans. Am. Math. Soc.* **182**, 323-352 (1973)
14. Rus, IA: *Fixed Point Structure Theory*. Cluj University Press, Cluj-Napoca (2006)
15. Andres, J, Górniewicz, L: *Topological Fixed Point Principles for Boundary Value Problems*. Kluwer Academic, Dordrecht (2003)
16. Browder, FE, Nussbaum, RD: The topological degree for noncompact nonlinear mappings in Banach spaces. *Bull. Am. Math. Soc.* **74**, 671-676 (1968)
17. Granas, A, Dugundji, J: *Fixed Point Theory*. Springer, Berlin (2003)
18. Brown, RF, Furi, M, Górniewicz, L, Jiang, B (eds.): *Handbook of Topological Fixed Point Theory*. Springer, Berlin (2005)
19. Ortega, J, Rheinboldt, WC: *Iterative Solution of Nonlinear Equations in Several Variables*. Academic Press, New York (1960)
20. Rus, IA: An abstract point of view in the nonlinear difference equations. In: *Conference on Analysis, Functional Equations, Approximation and Convexity, in Honour of Professor Elena Popoviciu, Cluj-Napoca*, pp. 272-276 (1999)
21. Şerban, MA: *Fixed Point Theory for Operators on Cartesian Product Spaces*. Cluj University Press, Cluj-Napoca (2002) (in Romanian)
22. Rus, IA: An iterative method for the solution of the equation $x = f(x, \dots, x)$. *Rev. Anal. Numér. Théor. Approx.* **10**(1), 95-100 (1981)
23. Krasnoselskii, MA: Two remarks on the method of successive approximations. *Usp. Mat. Nauk* **10**(1), 123-127 (1955)
24. Browder, FE: Fixed point theorems for nonlinear semicontractive mappings in Banach spaces. *Arch. Ration. Mech. Anal.* **21**(4), 259-269 (1966)
25. Browder, FE: Semicontractive and semiaccretive nonlinear mappings in Banach spaces. *Bull. Am. Math. Soc.* **74**, 660-665 (1968)
26. Kirk, WA: On nonlinear mappings of strongly semicontractive type. *J. Math. Anal. Appl.* **27**, 409-412 (1969)
27. Webb, JRL: Fixed point theorems for nonlinear semicontractive operators in Banach spaces. *J. Lond. Math. Soc.* **1**, 683-688 (1969)
28. Soardi, P: Punti uniti di semicontrazioni. *Atti Accad. Sci. Torino, Cl. Sci. Fis. Mat. Nat.* **108**, 51-57 (1974)
29. Ziebur, AD: Uniqueness and the convergence of successive approximations. *Proc. Am. Math. Soc.* **13**, 899-903 (1962)
30. Ziebur, AD: Uniqueness and the convergence of successive approximations. II. *Proc. Am. Math. Soc.* **16**, 335-340 (1965)
31. Opoitsev, VI: Heterogenic and combined-concave operators. *Sib. Math. J.* **16**, 781-792 (1975) (in Russian)
32. Opoitsev, VI: A generalization of the theory of monotone and concave operators. *Trans. Mosc. Math. Soc.* **36**, 243-280 (1979)
33. Diamond, P, Opoitsev, VI: Stability of nonlinear difference inclusions. *Dyn. Contin. Discrete Impuls. Syst., Ser. A Math. Anal.* **8**, 353-371 (2001)
34. Anichini, G: Nonlinear problems for systems of differential equations. *Nonlinear Anal.* **1**(6), 691-699 (1977)
35. Anichini, G, Conti, G: How to make use of the solution set to solve boundary value problems. *Prog. Nonlinear Differ. Equ. Appl.* **40**, 15-25 (2000)
36. Deimling, K, Lakshmikantham, V: Quasi-solutions and their role in the qualitative theory of differential equations. *Nonlinear Anal.* **4**, 657-663 (1980)
37. Opoitsev, VI, Khurodze, TA: *Nonlinear Operators in Spaces with a Cone*. Tbilis Gos. University, Tbilisi (1984) (in Russian), 271 pp.
38. Cronin, J: *Fixed Points and Topological Degree in Nonlinear Analysis*. Amer. Math. Soc. Math. Survey, vol. 11 (1964)
39. Petruşel, A, Petruşel, G, Samet, B, Yao, J-C: Coupled fixed point theorems for symmetric contractions in b -metric spaces with applications to operator equation systems. *Fixed Point Theory* **17**(2), 457-476 (2016)
40. Kirk, WA, Shahzad, N: *Fixed Point Theory in Distance Spaces*. Springer, Heidelberg (2014)
41. Petruşel, A, Petruşel, G, Samet, B, Yao, J-C: A study of the coupled fixed point problem for operators satisfying a max-symmetric condition in b -metric spaces with applications to a boundary value problem. *Miskolc Math. Notes* **17**(1), 501-516 (2016)
42. Radenović, S: Bhaskar-Lakshmikantham type results for monotone mappings in partially ordered metric spaces. *Int. J. Nonlinear Anal. Appl.* **5**(2), 37-49 (2014)
43. Radenović, S: Coupled fixed point theorems for monotone mappings in partially ordered metric spaces. *Kragujev. J. Math.* **38**(2), 249-257 (2014)
44. Bainov, D, Hristova, S: Monotone-iterative techniques of V. Lakshmikantham for a boundary value problem for systems of integro-differential equations. *Math. J. Toyama Univ.* **18**, 169-178 (1995)
45. Berinde, V: Generalized coupled fixed point theorems for mixed monotone mappings in partially ordered metric spaces. *Nonlinear Anal.* **74**, 7347-7355 (2011)
46. Berinde, V: Coupled fixed point theorems for φ -contractive mixed monotone mappings in partially ordered metric spaces. *Nonlinear Anal.* **75**(6), 3128-3228 (2012)
47. Chen, Y-Z: Existence theorems of coupled fixed points. *J. Math. Anal. Appl.* **154**, 142-150 (1991)
48. Gana Bhaskar, T, Lakshmikantham, V: Fixed point theorems in partially ordered metric spaces and applications. *Nonlinear Anal.* **65**, 1379-1393 (2006)
49. Guo, D, Lakshmikantham, V: *Nonlinear Problems in Abstract Cones*. Academic Press, New York (1988)

50. Guo, D, Cho, YJ, Zhu, J: Partial Ordering Methods in Nonlinear Problems. Nova Science Publishers, Hauppauge (2004)
51. Krasnoselskii, MA, Zabreiko, PP: Geometrical Methods of Nonlinear Analysis. Springer, Berlin (1984)
52. Lakshmikantham, V, Ćirić, L: Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces. *Nonlinear Anal.* **70**, 4341-4349 (2009)
53. Lishan, L: Iterative method for solutions and coupled quasi-solutions of nonlinear Fredholm integral equations in ordered Banach spaces. *Indian J. Pure Appl. Math.* **27**(10), 959-972 (1996)
54. Xiao, J-Z, Zhu, X-H, Shen, Z-M: Common coupled fixed point results for hybrid nonlinear contractions in metric spaces. *Fixed Point Theory* **14**, 235-250 (2013)
55. Popa, D, Lungu, N: On an exponential inequality. *Demonstr. Math.* **38**(3), 667-674 (2005)
56. Soleimani Rad, G, Shukla, S, Rahimi, H: Some relations between n -tuple fixed point and fixed point results. *Rev. R. Acad. Cienc. Exactas Fis. Nat., Ser. A Mat.* **109**(2), 471-481 (2015)
57. Rus, IA: Remarks on a LaSalle conjecture on global asymptotic stability. *Fixed Point Theory* **17**(1), 159-172 (2016)
58. Presić, SB: Sur une classe d'inequations aux differences finies et sur la convergence de certaines suites. *Publ. Inst. Math.* **5**(19), 75-78 (1965)
59. Djebali, S, Gorniewicz, L, Ouahab, A: Solution Sets for Differential Equations and Inclusions. de Gruyter, Berlin (2012)
60. Heikkilä, S, Lakshmikantham, V: Monotone Iterative Techniques for Discontinuous Nonlinear Differential Equations. Dekker, New York (1991)
61. Rus, MD: Fixed point theorems for generalized contractions in partially ordered metric spaces, with semi-monotone metric. *Nonlinear Anal.* **74**, 1804-1813 (2011)
62. Rus, MD: The method of monotone iterations for mixed monotone operators in partially ordered sets and order-attractive fixed points. *Fixed Point Theory* **15**, 579-594 (2014)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
