

Circulating Neutrophil MicroRNAs as Biomarkers for the Detection of Lung Cancer

Jie Ma^{1,2,†}, Ning Li^{1,3,†}, Yanli Lin¹, Chhavi Gupta¹ and Feng Jiang¹

¹Department of Pathology, University of Maryland School of Medicine, Baltimore, MD, USA. ²Department of Clinical Biochemistry, Jiangsu University School of Medicine, Zhenjiang, Jiangsu Province, China. ³College of Chinese Traditional Materia Medica, Shenyang Pharmaceutical University, Shenyang, Liaoning Province, China. [†]These authors contributed equally to this work.

ABSTRACT: Peripheral neutrophils are the predominant circulating leukocytes and an important component of innate and adaptive immune systems, which is a primary defense against cancer. MicroRNAs (miRNAs) can modulate neutrophil functions and play important roles in cancer pathogenesis by regulating neutrophil gene expression. To investigate if assessment of differential miRNA levels of peripheral neutrophils has the potential for diagnosis of non-small-cell lung cancer (NSCLC), we examine neutrophils of 15 patients with stage I NSCLC and 15 smokers without cancer. We identify five neutrophil miRNAs that have an abnormal level in patients with NSCLC versus smokers without cancer. In a training set of 82 patients with lung cancer and 73 controls, a set of two genes (miRs-26a-2-3p and 574-3p) are developed, producing 77.8% sensitivity and 78.1% specificity for NSCLC detection. Furthermore, in a testing set of 60 patients with lung cancer and 58 smokers, the performance of analyzing the two miRNAs for lung cancer detection is confirmed. This study for the first time shows that a neutrophil miRNA profile may serve as a new category of circulating biomarkers for the detection of NSCLC.

KEYWORDS: microRNA, peripheral neutrophils, biomarkers, lung cancer, diagnosis

CITATION: Ma et al. Circulating Neutrophil MicroRNAs as Biomarkers for the Detection of Lung Cancer. *Biomarkers in Cancer* 2016;8 1–7 doi:10.4137/BIC.S37333.

TYPE: Original Research

RECEIVED: November 10, 2015. **RESUBMITTED:** December 13, 2015. **ACCEPTED FOR PUBLICATION:** December 22, 2015.

ACADEMIC EDITOR: Barbara Guinn, Editor in Chief

PEER REVIEW: Three peer reviewers contributed to the peer review report. Reviewers' reports totaled 1010 words, excluding any confidential comments to the academic editor.

FUNDING: This work was supported in part by NCI R01CA161837, VA merit Award I01 CX000512, LUNGevity/Upstage Foundation Early Detection Award, a grant support from DeCesaris/Prout Cancer Foundation, and The University of Maryland Research and Innovation Seed Program (FJ). The authors confirm that the funder had no influence over the study design, content of the article, or selection of this journal.

COMPETING INTERESTS: Authors disclose no potential conflicts of interest.

COPYRIGHT: © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

CORRESPONDENCE: fjiang@som.umaryland.edu

Paper subject to independent expert blind peer review. All editorial decisions made by independent academic editor. Upon submission manuscript was subject to anti-plagiarism scanning. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties. This journal is a member of the Committee on Publication Ethics (COPE).

Published by Libertas Academica. Learn more about this journal.

Introduction

The early diagnosis of lung cancer by low-dose computed tomography (LDCT) can significantly reduce the mortality.¹ However, >25% of smokers screened by LDCT have indeterminate pulmonary nodules (PNs), of which 96% are finally diagnosed to be false-positives.¹ Circulating biomarkers that can be used alone or complement CT for more sensitively and specifically detecting NSCLC are urgently needed.

Neutrophils are the predominant circulating leukocyte population, accounting for approximately 70% of peripheral leukocytes.² Neutrophils have emerged as an important component of effector and regulatory circuits in the innate and adaptive immune systems,³ which are essential for immune reaction as a primary defense against cancer. Furthermore, the integration of immune and tumor cells promotes various molecular and cellular processes underlying multiple interrelated steps that define cancer initiation, development, and progression.⁴ In addition, neutrophils contribute to the multiple steps of tumorigenesis by mediating cytotoxicity, tumor cell apoptosis, immunologically mediated tumor rejection, and antitumoral immune memory.⁵ Previous studies have provided strong evidence for the existence of N1 (anti-tumoral) and N2 (protumoral) tumor-associated neutrophils (TANs).^{2,6–9} N1 neutrophils can kill tumor cells and thus play a beneficial and protective role for the host. Conversely,

N2 neutrophils promote carcinogenesis in certain situations and are more closely associated with the invasive and metastatic effects in tumorigenesis. The assessment of TANs, particularly in surgically resected tumor tissue specimens, could predict disease-specific and overall survival in patients with metastatic and localized tumors.² However, there have been no reports regarding the use of peripheral neutrophils or the related molecular changes as potential circulating biomarkers for the detection of cancer at the early stage.

De novo induction of microRNAs (miRNAs) is part of the crucial regulatory circuits that control neutrophil gene expression³ and hence modulates neutrophil functions.^{10–13} For instance, miR-223 can regulate innate immune genes whose functions are related to neutrophil biology.¹⁴ Furthermore, the effect of miRNAs (miRs-491-3p, 34b, 595, 328, 1281, and 483-3p) has an important impact on neutrophil-related pathways.¹⁵ miR-17 can regulate interleukin-8, a member of the cysteine-X-cysteine-chemokine, and has critical roles in neutrophil recruitment.¹⁶ In addition, miRNAs (miRs-15b, 26a/b, 29a, 30b, 106b, and 93) govern the outcome of the neutrophil response to specific bacterial infections by mediating cytokine production.¹⁵ miR-130a modulates transforming growth factor β 1 control of neutrophil differentiation.¹⁷ Moreover, both miR-17 and miR-31 can control neutrophil adherence on endothelial cells.¹⁸ Therefore, the investigation

of the neutrophil-related miRNAs could help understand neutrophil function and tumor biology. More importantly, characterizing differential miRNA expression profiles of peripheral neutrophils might provide an immunomonitoring approach for tumorigenesis. Since neutrophils are the largest circulating leukocyte population in blood, hereby we investigate if miRNAs of peripheral neutrophils could be used as new circulation biomarkers for lung cancer.

Materials and Methods

Patient cohorts. Our study was approved by the Institutional Review Boards of the Baltimore VA Medical Center and University of Maryland Medical Center. The research complied with the principles of the Declaration of Helsinki, and patients gave their written, informed consent to participate in this research. We recruited 157 patients with NSCLC (45 stage I, 35 II, 37 III, and 40 IV) and 146 smokers without any cancer, from whom peripheral blood was drawn as previously described.¹⁹ The NSCLC cases consisted of two major types: squamous cell carcinoma and adenocarcinoma. These smokers without cancer had granulomatous inflammation ($n = 72$), nonspecific inflammatory changes ($n = 53$), or lung infections ($n = 21$). To delineate miRNA expression profiles in neutrophils of patients with lung cancer by an array platform, we used peripheral neutrophil specimens of 15 patients with stage I NSCLC and 15 smokers as an exploratory set. Furthermore, we randomly selected 82 patients with lung cancer with different stages and 73 smokers as a training set (Table 1). We used the training cohort to develop a panel

of lung cancer biomarkers. We also selected 60 patients with stage I NSCLC and 58 cancer-free subjects as a testing set of cohort to confirm the performance of the biomarkers in the detection of NSCLC (Table 2).

Neutrophil isolation and processing. Peripheral neutrophils were isolated from blood as previously described.^{19–22} Briefly, a nucleated cell suspension was prepared using lymphocyte separation medium (MP Biomedicals, LLC). The cell pellet was lysed with ammonium chloride solution (Becton Dickinson and Company BD Biosciences). Neutrophils were isolated from the cells by using the Neutrophil Enrichment Kit (Stem Cell Technologies). The purity of the isolated cell populations was assessed by flow cytometry as previously described.²² RNA was extracted from the cells as described in our earlier studies.^{19,23,24} Furthermore, we determined the purity, concentration, and integrity of RNA as described in our previously published articles.^{19,23,24} Briefly, the purity and concentration of RNA were determined by OD260/280 readings using a dual beam UV spectrophotometer (Eppendorf AG). A clean sample should have a 260/280 nm OD ratio of 1.8–2.0. RNA integrity was determined by capillary electrophoresis using the RNA 6000 Nano Lab-on-a-Chip kit and the Bioanalyzer 2100 (Agilent Technologies). A sample that had RNA integrity number of ≥ 7 was considered as the one with very high quality. Only RNA extracts with RNA integrity number values >7 underwent in further analysis.

Real-time polymerase chain reaction-based microarray analysis of miRNAs. The analysis of miRNA expression profiles in neutrophils was performed by *Exiqon Services* (Exiqon)

Table 1. Characteristics of patients with lung cancer and cancer-free controls in a training set.

	NSCLC CASES (n = 82)	CONTROLS (n = 73)	P-VALUE
Age	66.27 (SD 10.16)	65.68 (SD 12.38)	0.39
Sex			0.32
Female	26	27	
Male	56	46	
Race			0.09
White	61	56	
African American	21	17	
Pack-years	43.68 (Range, 0–169)	8.45 (Range, 0–97)	<0.01
Nodule size (cm)	5.23 (Range, 95)	0.78 (Range, 53.56)	<0.01
Stage			
I	19		
II	21		
III	21		
IV	21		
Histological type			
Adenocarcinoma	43		
Squamous cell carcinoma	39		

Abbreviation: NSCLC, non-small cell lung cancer.

Table 2. Characteristics of patients with stage I NSCLC and cancer-free controls in a testing set.

	NSCLC CASES (n = 60)	CONTROLS (n = 58)	P-VALUE
Age	67.79 (SD 8.23)	64.68 (SD 10.37)	0.36
Sex			0.46
Female	21	20	
Male	39	38	
Race			0.36
White	36	36	
African American	24	22	
Pack-years	49.34 (Range, 0–187)	10.25 (Range, 0–86)	<0.01
Nodule size (cm)	5.85 (Range, 99)	0.84 (Range, 59.37)	<0.01
Stage			
I	11		
II	14		
III	16		
IV	19		
Histological type			
Adenocarcinoma	31		
Squamous cell carcinoma	29		

Abbreviation: NSCLC, non-small cell lung cancer.

as described in our previous report.²¹ Briefly, 6 μ L RNA was reversely transcribed in 30 μ L reactions using the miRCURY LNATM Universal RT miRNA PCR, Polyadenylation and cDNA synthesis kit (Exiqon). Complementary DNA (cDNA) was diluted 100 \times and assayed in 10 μ L polymerase chain reactions (PCRs) according to the protocol for miRCURY LNATM Universal RT miRNA PCR; each miRNA was assayed once by quantitative PCR (qPCR) on the miRNA Ready-to-Use PCR, Haman Panel I using ExiLent SYBR[®] Green master mix. Negative controls excluding template from the reverse transcription reaction were performed and profiled like the samples. The amplification was performed in a LightCycler[®] 480 Real-Time PCR System (Roche) in 384-well plates. The amplification curves were made by using quantification cycle (C_q), which was used as a relative value for further quantification of the tested genes. We normalized the resulted data by using the average of assays detected in the samples (average assay C_q). We used a *t*-test with a cutoff of *P*-value <0.05 to discover differentially expressed genes between patients with NSCLC and cancer-free smokers.

Quantitative reverse transcriptase PCR. Quantitative reverse transcriptase (qRT)-PCR was carried out using a protocol that was developed in our laboratory.^{19,20} Briefly, 100 ng RNA was reversely transcribed by a T100 thermal cycler (Applied Biosystems) using miRCURY LNATM Universal cDNA Synthesis Kit (Exiqon). The thermocycler parameters were as follows: hold for 60 minutes at 42°C and for five minutes at 95°C. For qPCR analysis, after 40 \times dilution, 4 μ L of cDNA was combined with 5 μ L of miRCURY LNATM Universal RT miRNA PCR ExiLent SYBR Green master

mix (Exiqon) and 1 μ L PCR primers set to produce a PCR reaction in a total volume of 10.0 μ L. qPCR was carried out on an CFX96 thermocycler (Bio-Rad) at 95°C for 10 minutes, followed by 40 cycles of 95°C for 15 seconds and 60°C for one minute. qPCR data were analyzed by using the Manager software (Bio-Rad) with an automatic C_t setting for assigning the baseline and threshold for C_t determination. All tests were performed in triplicates. For the normalization of the miRNA qPCR data, we screened the expression of all miRNAs represented in the above array-based panel analysis to find the miRNAs with the minimal variation between the two groups. miR-484 was selected to be a control gene because the expression was stable in all neutrophil samples without difference between the cases and controls. We calculated miRNA expression using the equation $2 - \Delta C_t$, where $\Delta C_t = C_t$ (targeted miRNA) – C_t (miR-484). C_t means that in a PCR assay, a positive reaction is detected by accumulation of a fluorescent signal. Red blood cell (RBC)-related miRNAs (mir-451 and mir-92a) were analyzed using qRT-PCR.

Statistical analysis. We used Kruskal–Wallis test for comparing continuous variables or significance test for Spearman's rank correlation coefficient to determine the relations of the miRNAs and clinicopathologic and demographic characteristics of the subjects. We created receiver operating characteristic (ROC) curve and calculated the area under the ROC (AUC) value of each miRNA using numerical integration of the ROC curve. We utilized AUC to decide the accuracy of a biomarker on its capacity to differentiate cases from controls. We used Pearson's correlation analysis to evaluate correlation between the miRNAs.

Results

Identifying miRNAs that showed a differential expression level in peripheral neutrophils of lung cancer patients versus cancer-free subjects. Purity of the isolated peripheral neutrophils was greater than 95% as assessed by flow cytometry. Extracted RNA from the neutrophils was very pure as demonstrated by a 260/280 ratio of 1.8–2.0 and had very high quality as demonstrated by a RNA integrity number of ≥ 7 . The measured miRNA expressions by using miRNA array in the replicates of each sample were highly correlated (all $P < 0.0001$). Of the 372 miRNAs embodied on the miRNA array, 141 were measurable in all the neutrophil specimens of cancer cases and controls, as they showed a $< 37 C_q$ value. Among the 141 miRNAs, five (miRs-423-3p, 26a-2-3p, 148a-3p, 18a-3p, and 574-3p) exhibited > 2.0 fold-changes (P -value < 0.05) in patients with NSCLC versus smokers (Table 3). Four of the five genes (miRs-423-3p, 148a-3p, 18a-3p, and 574-3p) had a high expression level, whereas one miRNA (miR-26a-2-3p) displayed a low level in neutrophils of patients with NSCLC versus controls.

Developing a panel of neutrophil miRNA biomarkers. Using qRT-PCR, we validated the results produced from the above miRNA array in a training set of 82 patients with lung cancer and 73 controls. All targeted five miRNAs showed $\leq 35 C_t$ values in each neutrophil specimen and thus could be consistently measurable using a different technique, qRT-PCR. Furthermore, we did not find the RBC-related miRNAs in the neutrophil specimens. Of the five miRNAs, two (miRs-26a-2-3p and 574-3p) showed a significantly different level in neutrophil samples of the patients with lung cancer versus control subjects (All $P < 0.05$) (Table 4). As shown in Figure 1, miR-26a-2-3p displayed a lower expression level, whereas miR-574-3p exhibited a higher level in neutrophils of patients with lung cancer versus control subjects. Therefore, the qRT-PCR analysis showed that the two miRNAs had changes consistent with miRNA array data in the same statistically significant direction.

The two neutrophil genes displayed AUC values of 0.71–0.74 in differentiating patients with lung cancer from

controls. Interestingly, the use of the two neutrophil miRNAs in combination created an AUC value of 0.81 (Fig. 2), being considerably higher than that of any single one used alone. There was no significant correlation between the two miRNAs' level ($r = 0.02$, $P = 0.84$), supporting a combined effort of using the two miRNAs as a panel of biomarkers. The optimal cutoff for the two biomarkers used together was $U = 0.658$, where $U = -8.369 + 3.876 \times \log(\text{miR-26a-2-3p}) - 3.4679 \times \log(\text{miR-574-3p})$. One with $U \geq 0.658$ was categorized as a cancer case. Consequently, combined use of the two miRNAs generated 77.8% sensitivity and 78.1% specificity for the detection of all-stage NSCLC. In addition, miR-26a-2-3p expression level was negatively associated with stages of NSCLC ($P = 0.04$). However, miR-574-3p did not show differences between stages of the disease ($P = 0.13$) (Supplementary Table 1). Subsequently, the examination of the two genes created a higher sensitivity for the detection of advanced-stage (III–IV) NSCLC compared with stage I–II NSCLC (83.3% vs 70.0%, $P < 0.05$), while maintaining the same specificity (78.08%). The panel of the two biomarkers did not exhibit significant relationship with age, ethnicity, and smoking history of the subjects (all $P > 0.05$) (Supplementary Table 1). The level of miR-574-3p rather than miR-26a-2-3p was associated with PN size ($P = 0.04$ and $P = 0.12$, respectively).

Validating the neutrophil miRNA biomarkers in a testing cohort. The use of the two neutrophil miRNAs in combination generated 78.3% sensitivity and 77.6% specificity in differentiating patients with all-stage lung cancer from cancer-free subjects (Supplementary Table 2). Combined use of the two genes created a higher sensitivity for the detection of advanced-stage (III–IV) NSCLC compared with stage I–II disease (82.9% vs 68.0% sensitivity, $P < 0.05$), while having the same specificity (77.6%). The validation experiment confirmed the potential of the neutrophil miRNAs as biomarkers in the detection of lung cancer.

Discussion

miRNAs have crucial functions in tumorigenesis. With the objective of developing circulating cancer biomarkers, abundant studies have been done by detecting cell-free miRNAs that are released from tumor cells in blood.^{19,20,25–30} Though displaying promise, the analysis of the extracellular genes in serum or plasma has some difficulties, limiting its use in the clinical practice. These mainly consist of 1) low recovery of miRNAs in plasma or serum,³¹ 2) the discharge of miRNAs in plasma or serum by hemolysis of RBCs, creating nonspecific consequences, and 3) sources of inconsistency of the cell-free miRNAs may cause varying or even opposing results for the identification of the same type of cancer.³² The difficulty turn out to be more complex by the fact that in blood, miRNAs are either associated with proteins, such as argonaute^{32,33} and lipoproteins,³⁴ or enclosed within cellular fragments designated as exosomes, microparticles, microvesicles, or extracellular vesicles.^{35,36}

Table 3. Five neutrophil miRNAs displayed ≥ 2.0 fold-changes ($P < 0.05$) in patients with NSCLC versus the cancer-free smokers by using Exiqon miRNA array.

miRNAs	FOLD CHANGE (NSCLC PATIENTS/CONTROLS)	P-VALUE
hsa-miR-574-3p	3.46	0.006
hsa-miR-423-3p	3.56	0.003
hsa-miR-148a-3p	2.53	0.017
hsa-miR-18a-3p	2.38	0.026
hsa-miR-26a-2-3p	–2.83	0.023

Abbreviation: NSCLC, non-small cell lung cancer.

Table 4. Expression levels of the miRNAs in neutrophils of patients with NSCLC versus cancer-free smokers by using qRT-PCR.

GENES	MEAN \pm SEM IN CANCER-FREE CONTROLS	MEAN \pm SEM IN NSCLC PATIENTS	P-VALUE	AUC
miR-574-3p	0.129 \pm 0.004	0.164 \pm 0.005	<0.001	0.739 (Std. Error, 0.041; 95% CI, 0.659 to 0.820)
miR-26a-2-3p	0.004 \pm 0.0003	0.002 \pm 0.0002	<0.001	0.708 (Std. Error, 0.042; 95% CI, 0.624 to 0.792)
miR-423-3p	0.678 \pm 0.027	0.699 \pm 0.028	0.595	0.522 (Std. Error, 0.066; 95% CI, 0.394 to 0.654)
miR-148a-3p	1.972 \pm 0.204	2.150 \pm 0.224	0.559	0.528 (Std. Error, 0.067; 95% CI, 0.395 to 0.661)
miR-18a-3p	0.035 \pm 0.001	0.038 \pm 0.001	0.213	0.567 (Std. Error, 0.065; 95% CI, 0.4393 to 0.695)

Abbreviations: qRT-PCR, quantitative reverse transcriptase PCR; NSCLC, non-small cell lung cancer; SEM, standard error of the mean; CI, confidence interval; AUC, the area under ROC curve receiver-operator characteristic curve.

The analysis and development of circuiting neutrophil miRNAs as blood-based biomarkers may address the above challenges in the development of the cell-free miRNAs because it is a cell-based approach. Furthermore, peripheral neutrophils make up 70% of white blood cells and thus can produce large amounts of RNA for reliable analysis of miRNAs. In addition, our results show that extracted RNA from the neutrophils has very high quality. Moreover, purity of neutrophils was greater than 95%, from which RNA is specifically isolated and may not contain RBC miRNAs and other blood components. Therefore, neutrophil miRNA profile may serve as a new category of circulating biomarkers for the detection of NSCLC.

Of the two miRNAs, only miR-26a-2-3p was previously found to be associated with lung cancer and had an elevated expression level in lung tumor tissues.^{19,37–39} However, our present study showed that miR-26a-2-3p exhibited a low level in neutrophils of patients with NSCLC compared with cancer-free controls. The contradictory observations suggest that the miRNA expression change in neutrophils may not simply be a reflection of that in the lung cancer tissues or mimic that in the primary tumors. Furthermore, previous reports including our own have found that numerous cell-free miRNAs released from primary cancer sites have the potential for lung cancer diagnosis.²⁰ Yet, no overlap is observed between the

neutrophil miRNAs of lung cancer versus the plasma or serum miRNAs from lung cancers.^{20,40} The neutrophil miRNAs may not be influenced by circulating cancer cells. Therefore, the assessment of the neutrophil miRNAs could present a different strategy for lung cancer diagnosis. In addition, differing from the analysis of circulating extracellular molecules that are dependent on substantial cancer burden, the determination of neutrophil miRNA expressions may function as a better surrogate window into cancer status, as the dysregulation of neutrophil miRNAs is involved in every step of tumorigenesis, such as cytotoxicity, tumor cell apoptosis, immunologically mediated tumor rejection, and antitumoral immune memory.^{3,10–13} Nevertheless, carrying out a new study to compare miRNAs of matched neutrophils, plasma, and cancer tissues of the same patients is necessary to investigate if the neutrophil miRNAs are independent from those in blood and cancers of the patients. Furthermore, analyzing the potential neutrophil miRNA biomarkers in different types of malignancy to determine if they are specific to lung cancer is also required.

Neutrophils play a dual role in tumorigenesis through innate and adaptive immune systems that can recognize and remove malignant cells, a process called *immune surveillance*. On the one hand, tumor-mediated signals can induce the formation of a pro-tumorigenic (N2) phenotype capable of supporting tumor growth and suppressing the antitumor




Figure 1. Expression levels of two miRNAs in neutrophil samples of 82 cancer-free smokers and 73 patients with NSCLC. The two miRNAs (A and B) have statistically significantly different levels in the patients with NSCLC versus cancer-free smokers (all $P < 0.05$). The inside line denotes the median.




Figure 2. ROC curve analysis of expression levels of two miRNAs (miRs-26a-2-3p and -574-3p) in neutrophils of 82 patients diagnosed with NSCLC and 73 cancer-free individuals. miRs-26a-2-3p and -574-3p produce 0.71–0.74 AUC values (**A** and **B**), being significantly lower than 0.81 AUC created from the combined use of the two miRNAs (**C**) ($P < 0.05$).

immune response. Therefore, the neutrophils can promote malignancy in certain situations. Grounded on the protumoral role, previous studies have mainly focused on the analysis of surgically resected tumor tissues to determine the association of neutrophils with outcome of cancers.^{2,15} On the other hand, serving as the immune system's front line of defense against cancer, neutrophils can also identify and destroy tumor cells by induction of cytotoxicity and tumor rejection and antitumoral immune memory (N1 phenotype). The dysfunction of the neutrophils could take place as an early event in immunogenicity or immune evasion of cancer. Our present study for the first time demonstrates that peripheral neutrophil miRNAs may service as new circulating biomarkers that could be useful for the detection of cancers at the early stage.

miR-26a family has been suggested to play an important function in modulating neutrophil biology. For example, miR-26a was one of the abundant miRNAs in neutrophils.^{41,42} miR-26a expression could be preferentially induced by *Escherichia coli* and hence target the interferon- γ transcript.⁴³ The induction of miR-26a in neutrophils had a strong impact on the protein translation. In addition, ectopic expression of miR-26a dramatically enhanced lung cancer cell migration and invasion abilities by regulating matrix metalloproteinase-2,⁴⁴ which could prevent apoptosis of tumor cells in the lungs.^{2,45} miR-26a augmented lung cancer cell metastasis potential via modulation of metastasis-related gene expression and activation of AKT pathway by phosphatase and tensin homolog suppression.⁴⁴ Moreover, dysregulation of miR-26a enhanced tumorigenic and metastatic potential by a miR26a/-26b-COX-2-MIP-2 loop.⁴⁶ Our present finding that downregulation of neutrophil miR-26a-2-3p is associated with advanced-stage NSCLC is consistent with the previous discoveries that the miRNA may play an important role in the N2 neutrophils associated with cancer invasive and metastatic effects. However, dysregulation of miR-574-3p was not reported in the regulation of immune system and neutrophil biology. We are investigating how the two miRNAs involve in the cross talk between immune and

tumor cells that could change the phenotype of tumor biology. We are also exploring which molecular signaling pathways the two miRNAs may involve in carcinogenesis and if the miRNA aberrations are drivers in multiple oncogenes.

The study has some limitations. First, we use an array that only analyzes 372 miRNA genes to delineate the neutrophil miRNAs whose changes are associated with lung cancer. Only 141 miRNAs could be detectable in the neutrophil specimens. In addition, from the 141 genes, only two are finally found to have the potential as biomarkers. The two miRNAs used together has 77.78% sensitivity and 78.08% specificity for the detection of all-stage NSCLC. However, the efficiency of the biomarkers is not enough to be used in laboratory settings. Particularly, the diagnostic performance of the two biomarkers for the early-stage NSCLC is not sufficient because the sensitivity for stage I–II NSCLC is only 68.0%. To address the concern, we are using whole-genome next-generation sequencing to identify new neutrophil miRNAs of lung cancer that can be added in the assay for precisely diagnosing lung cancer. Second, the neutrophil samples are acquired from the hospital-based patients with clinical diagnosis. Future validation of the biomarkers in a prospective and multisite LDCT lung cancer screening trial is needed.

Conclusion

In conclusion, we find a differential miRNA expression profile in peripheral neutrophils of patients with lung cancer. Furthermore, two circulating neutrophil miRNA biomarkers are developed that have the potential to be used for the detection of lung cancer. Nevertheless, finding additional neutrophil miRNA biomarkers that can be added to the two biomarkers and incorporating other types of biomarkers with the neutrophil miRNA biomarkers for lung cancer early detection are required. Moreover, given the contribution of neutrophil miRNAs to both the initiation and the progression of tumors, we believe that different miRNA profiles of neutrophils could be also used for the development of new circulating biomarkers in other cancer diseases.

Acknowledgments

The authors thank the Biostatistics Shared Service of University of Maryland School of Medicine for statistically analyzing the data.

Author Contributions

Conducted the experiments and participated in data acquisition and interpretation: JM, NL, YL, and CG. Conducted the study design, coordination, and prepared the manuscript: FJ. All authors read and approved the final manuscript.

Supplementary Materials

Supplementary table 1. Associations of patient characteristics with miRNA expression levels in a training set of cases and controls.

Supplementary table 2. A panel of the two neutrophil miRNA (miRs-26a-2-3p and 574-3p) biomarkers in a testing set of specimens.

REFERENCES

- Aberle DR, Adams AM, Berg CD, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med*. 2011;365:395–409.
- Fridlender ZG, Albelda SM. Tumor-associated neutrophils: friend or foe? *Carcinogenesis*. 2012;33:949–955.
- Mantovani A, Cassatella MA, Costantini C, Jaillon S. Neutrophils in the activation and regulation of innate and adaptive immunity. *Nat Rev Immunol*. 2011;11:519–531.
- Schreiber RD, Old LJ, Smyth MJ. Cancer immunoediting: integrating immunity's roles in cancer suppression and promotion. *Science*. 2011;331:1565–1570.
- Piccardi H, Muschel RJ, Opendakker G. On the dual roles and polarized phenotypes of neutrophils in tumor development and progression. *Crit Rev Oncol Hematol*. 2012;82:296–309.
- Sionov RV, Assi S, Gershkovitz M, et al. Isolation and characterization of neutrophils with anti-tumor properties. *J Vis Exp*. 2015;19:e52933.
- Sionov RV, Fridlender ZG, Granot Z. The multifaceted roles neutrophils play in the tumor microenvironment. *Cancer Microenviron*. 2014;4. [Epub ahead of print].
- Mishalian I, Bayuh R, Levy L, Zolotarov L, Michaeli J, Fridlender ZG. Tumor-associated neutrophils (TAN) develop pro-tumorigenic properties during tumor progression. *Cancer Immunol Immunother*. 2013;62:1745–1756.
- Fridlender ZG, Sun J, Mishalian I, et al. Transcriptomic analysis comparing tumor-associated neutrophils with granulocytic myeloid-derived suppressor cells and normal neutrophils. *PLoS One*. 2012;7:e31524.
- Tili E, Michaille JJ, Croce CM. MicroRNAs play a central role in molecular dysfunctions linking inflammation with cancer. *Immunol Rev*. 2013;253:167–184.
- Tili E, Michaille JJ. Resveratrol, microRNAs, inflammation, and cancer. *J Nuc Acids*. 2011;2011:102431.
- Tili E, Croce CM, Michaille JJ. miR-155: on the crosstalk between inflammation and cancer. *Int Rev Immunol*. 2009;28:264–284.
- Tili E, Michaille JJ, Wernicke D, et al. Mutator activity induced by microRNA-155 (miR-155) links inflammation and cancer. *Proc Natl Acad Sci U S A*. 2011;108:4908–4913.
- Gilicze AB, Wiener Z, Toth S, et al. Myeloid-derived microRNAs, miR-223, miR27a, and miR-652, are dominant players in myeloid regulation. *Biomed Res Int*. 2014;2014:870267.
- Gantier MP. The not-so-neutral role of microRNAs in neutrophil biology. *J Leukoc Biol*. 2013;94:575–583.
- Oglesby IK, Vencken SF, Agrawal R, et al. miR-17 overexpression in cystic fibrosis airway epithelial cells decreases interleukin-8 production. *Eur Respir J*. 2015;46(5):1350–1360.
- Johnnidis JB, Harris MH, Wheeler RT, et al. Regulation of progenitor cell proliferation and granulocyte function by microRNA-223. *Nature*. 2008;451:1125–1129.
- Suarez Y, Wang C, Manes TD, Pober JS. Cutting edge: TNF-induced microRNAs regulate TNF-induced expression of E-selectin and intercellular adhesion molecule-1 on human endothelial cells: feedback control of inflammation. *J Immunol*. 2010;184:21–25.
- Shen J, Todd NW, Zhang H, et al. Plasma microRNAs as potential biomarkers for non-small-cell lung cancer. *Lab Invest*. 2011;91:579–587.
- Shen J, Liu Z, Todd NW, et al. Diagnosis of lung cancer in individuals with solitary pulmonary nodules by plasma microRNA biomarkers. *BMC Cancer*. 2011;11:374.
- Ma J, Lin Y, Zhan M, Mann DL, Stass SA, Jiang F. Differential miRNA expressions in peripheral blood mononuclear cells for diagnosis of lung cancer. *Lab Invest*. 2015;95:1197–1206.
- Allantaz F, Cheng DT, Bergauer T, et al. Expression profiling of human immune cell subsets identifies miRNA-mRNA regulatory relationships correlated with cell type specific expression. *PLoS One*. 2012;7:e29979.
- Ma J, Li N, Guarnera M, Jiang F. Quantification of plasma miRNAs by digital PCR for cancer diagnosis. *Biomark Insights*. 2013;8:127–136.
- Shen J, Liao J, Guarnera MA, et al. Analysis of microRNAs in sputum to improve computed tomography for lung cancer diagnosis. *J Thorac Oncol*. 2014;9:33–40.
- Cortez MA, Bueso-Ramos C, Ferdin J, Lopez-Berestein G, Sood AK, Calin GA. MicroRNAs in body fluids—the mix of hormones and biomarkers. *Nat Rev Clin Oncol*. 2011;8:467–477.
- Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol*. 2007;9:654–659.
- Pegtel DM, Cosmopoulos K, Thorley-Lawson DA, et al. Functional delivery of viral miRNAs via exosomes. *Proc Natl Acad Sci U S A*. 2010;107:6328–6333.
- Wang K, Zhang S, Weber J, Baxter D, Galas DJ. Export of microRNAs and microRNA-protective protein by mammalian cells. *Nucleic Acids Res*. 2010;38:7248–7259.
- Vickers KC, Remaley AT. Lipid-based carriers of microRNAs and intercellular communication. *Curr Opin Lipidol*. 2012;23:91–97.
- Jarry J, Schadendorf D, Greenwood C, Spatz A, van Kempen LC. The validity of circulating microRNAs in oncology: five years of challenges and contradictions. *Mol Oncol*. 2014;8:819–829.
- Fichtlscherer S, De Rosa S, Fox H, et al. Circulating microRNAs in patients with coronary artery disease. *Circ Res*. 2010;107:677–684.
- Moldovan L, Batte KE, Trgovcich J, Wisler J, Marsh CB, Piper M. Methodological challenges in utilizing miRNAs as circulating biomarkers. *J Cell Mol Med*. 2014;18:371–390.
- Arroyo JD, Chevillet JR, Kroh EM, et al. Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. *Proc Natl Acad Sci U S A*. 2011;108:5003–5008.
- Vickers KC, Palmisano BT, Shoucri BM, Shamburek RD, Remaley AT. MicroRNAs are transported in plasma and delivered to recipient cells by high-density lipoproteins. *Nat Cell Biol*. 2011;13:423–433.
- Hunter MP, Ismail N, Zhang X, et al. Detection of microRNA expression in human peripheral blood microvesicles. *PLoS One*. 2008;3:e3694.
- Mitchell PS, Parkin RK, Kroh EM, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci U S A*. 2008;105:10513–10518.
- Jiang DS, Wang YW, Jiang J, Li SM, Liang SZ, Fang HY. MicroRNA-26a involved in toll-like receptor 9-mediated lung cancer growth and migration. *Int J Mol Med*. 2014;34:307–312.
- Shen J, Jiang F. Applications of microRNAs in the diagnosis and prognosis of lung cancer. *Expert Opin Med Diagn*. 2012;6:197–207.
- Boeri M, Verri C, Conte D, et al. MicroRNA signatures in tissues and plasma predict development and prognosis of computed tomography detected lung cancer. *Proc Natl Acad Sci U S A*. 2011;108:3713–3718.
- Sozzi G, Boeri M, Rossi M, et al. Clinical utility of a plasma-based miRNA signature classifier within computed tomography lung cancer screening: a correlative MILD trial study. *J Clin Oncol*. 2014;32:768–773.
- Ward JR, Heath PR, Catto JW, Whyte MK, Milo M, Renshaw SA. Regulation of neutrophil senescence by microRNAs. *PLoS One*. 2011;6:e15810.
- Landgraf P, Rusu M, Sheridan R, et al. A mammalian microRNA expression atlas based on small RNA library sequencing. *Cell*. 2007;129:1401–1414.
- Ma F, Xu S, Liu X, et al. The microRNA miR-29 controls innate and adaptive immune responses to intracellular bacterial infection by targeting interferon-gamma. *Nat Immunol*. 2011;12:861–869.
- Liu B, Wu X, Wang C, et al. MiR-26a enhances metastasis potential of lung cancer cells via AKT pathway by targeting PTEN. *Biochim Biophys Acta*. 2012;1822:1692–1704.
- Acuff HB, Carter KJ, Fingleton B, Gordon DL, Matrisian LM. Matrix metalloproteinase-9 from bone marrow-derived cells contributes to survival but not growth of tumor cells in the lung microenvironment. *Cancer Res*. 2006;66:259–266.
- Kwon Y, Kim Y, Eom S, et al. MicroRNA-26a/-26b-COX-2-MIP-2 loop regulates allergic inflammation and allergic inflammation-promoted enhanced tumorigenic and metastatic potential of cancer cells. *J Biol Chem*. 2015;290:14245–14266.