

CRYSTALLOGRAPHIC
COMMUNICATIONS

ISSN 2056-9890

Crystal structure of *catena*-poly[[*(N,N*-diethyl-3-mesitylsulfonyl-1*H*-1,2,4-triazole-1-carboxamide- κ N¹)silver(I)]- μ -nitrate- κ^3 O,O':O]

Hyunjin Park,^a Eunjin Kwon,^a Il Yoon^{b*} and Jineun Kim^{a*}

Received 21 September 2016

Accepted 18 October 2016

Edited by A. Van der Lee, Université de Montpellier II, France

Keywords: crystal structure; cafenstrole; coordination polymer; silver; triazole; herbicide.**CCDC reference:** 1510357**Supporting information:** this article has supporting information at journals.iucr.org/e

^aDepartment of Chemistry (BK21 plus) and Research Institute of Natural Sciences, Gyeongsang National University, Jinju 52828, Republic of Korea, and ^bPhotodynamic Therapy Research Institute, School of Nanoscience and Engineering, Inje University, 197 Injero, Gimhae, Gyeongnam 50834, Republic of Korea. *Correspondence e-mail: yoonil71@inje.ac.kr, jekim@gnu.ac.kr


The reaction of silver nitrate and cafenstrole (*N,N*-diethyl-3-mesitylsulfonyl-1*H*-1,2,4-triazole-1-carboxamide), a triazole herbicide, leads to the title coordination polymer, [Ag(NO₃)(C₁₆H₂₂N₄O₃S)]_n, whose asymmetric unit comprises one cafenstrole ligand molecule, one Ag^I atom and one nitrate ion. The Ag^I atom, with a distorted trigonal–pyramidal environment, is coordinated by one nitrogen atom of a triazole ring, two oxygen atoms of a nitrate ion and one oxygen atom of a neighboring nitrate ion. The coordination bonds between silver and oxygen atoms give rise to a one-dimensional (1D) coordination polymer structure along [001]. The dihedral angle between the planes of the triazole and benzene rings is 87.13 (11)°. In the crystal, the coordination polymer is stabilized by C—H...O hydrogen bonds and C—H... π interactions, resulting in a three-dimensional architecture.

1. Chemical context

Recently, we have reported the crystal structure of the ligand cafenstrole (**L**; Kang *et al.*, 2015). Cafenstrole is a triazole herbicide and has been used for rice cultivation as an inhibitor of the germination of grass weeds (Takahashi *et al.*, 2001). Triazole derivatives have been investigated intensively over the years for pharmaceutical and agricultural purposes (Kumar *et al.*, 2013; Zhang *et al.*, 2014). It is very likely that triazole–metal interactions play a major role in the biological actions of triazole-containing drugs and agricultural chemicals. 1,2,4-Triazole and its derivatives have gained great attention as ligands to transition metals (Haasnoot, 2000). To understand the interactions of triazoles with metals, further research on the structures of triazole–metal compounds is of great necessity. Thus, our attention will be focused on the diversity of the coordination geometries of 1,2,4-triazole complexes with transition metal ions. Herein, we report the reaction of silver nitrate and cafenstrole to produce the title compound, which is a 1D silver(I) coordination polymer.


2. Structural commentary

The asymmetric unit of the title compound is shown in Fig. 1. It contains one **L** ligand and one silver nitrate ion. Reaction between silver nitrate and **L** afforded a 1D coordination polymer, in which the Ag^I atom has a distorted trigonal–pyramidal environment with one nitrogen atom (N1) [Ag1—N1 = 2.250 (3) Å] and three oxygen atoms (O4, O5, O5ⁱ)



OPEN ACCESS

[Ag1—O4 = 2.708 (3), Ag1—O5 = 2.450 (3) and Ag1—O5ⁱ = 2.396 (3) Å; symmetry code: (i) $-x + 1, -y + 1, z - \frac{1}{2}$], as shown in Fig. 2.



Atom Ag1 lies almost in the plane constituted by atoms O5, N1, and O5ⁱ [deviation = 0.0436 (12) Å]. The Ag1, O5, N1, and O5ⁱ atoms form a slightly distorted triangular basal plane with bond angles O5—Ag1—O5ⁱ = 106.52 (5), O5—Ag1—N1 = 118.75 (11) and O5ⁱ—Ag1—N1 = 134.63 (11)°. The apex atom, O4, deviates considerably from the normal to the basal plane, as indicated by the O4—Ag1—N1 bond angle of 149.66 (10)°. Other bond angles are 48.93 (10) and 67.18 (10)° for O4—Ag1—O5 and O4—Ag1—O5ⁱ, respectively. One oxygen atom of the nitrate ion (O6) is not bound to the Ag^I ion, whereas the other two oxygen atoms of the nitrate ion (O4 and O5) are bound to the Ag^I ion. One of the bound O atoms (O5) links




Figure 1
The asymmetric unit of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are shown as small spheres of arbitrary radius.




Figure 2
The coordination environment of the Ag^I atom in the title compound. [Symmetry codes: (i) $-x + 1, -y + 1, z - \frac{1}{2}$; (ii) $-x + 1, -y + 1, z + \frac{1}{2}$]

neighbouring Ag^I ion ions, thus forming a 1D polymer along [001]. The triazole plane is rotated about the S1—C10 axis in the opposite direction in comparison with free cafenstrol (Kang *et al.*, 2015). Thus, the diethyl amino group is located above the phenyl ring in the title compound, while that of free cafenstrol is placed outside the phenyl ring.

3. Supramolecular features

The O5 atom is bound to both Ag1 and neighboring Ag1ⁱⁱ [symmetry code: (ii) $-x + 1, -y + 1, z + \frac{1}{2}$], where the neighbouring asymmetric unit is related to the asymmetric unit by 2₁ symmetry, resulting in a 1D chain along [001] (Fig. 3). C—H...O hydrogen bonds between the 1D chains (yellow dashed lines) lead to the formation of layers parallel to (100). The layers are packed in an *ABAB* pattern along [010] (Fig. 4). Weak intermolecular C—H... π interactions (black dashed lines) between the *A* and *B* layers generate a three-dimensional network structure (Fig. 4). Thus the structure of the Ag^I coordination polymer is stabilized by C13—H13B...O2 and




Figure 3
The packing of the title compound showing chains along [001].

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C1–C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C8-H8C\cdots Cg1^i$	0.98	2.66	3.614 (5)	166
$C13-H13B\cdots O2^{ii}$	0.99	2.58	3.395 (4)	140
$C16-H16B\cdots O2^{iii}$	0.98	2.52	3.412 (6)	152

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{3}{2}, z$; (ii) $x + 1, y, z$; (iii) $x + 1, y, z + 1$.

$C16-H16B\cdots O2$ hydrogen bonds and weak intermolecular $C8-H8C\cdots Cg1$ ($Cg1$ is the centroid of the C1–C6 ring) interactions (Fig. 4 and Table 1).

4. Database survey

The crystal structure of cafenstrole has been reported (Kang *et al.*, 2015). The crystal structure of a 1,2,3-thiadiazole compound containing a 1,2,4-triazole moiety, $C_{15}H_{14}FN_5O_2S_2$, has been determined by Min *et al.* (2014) whereas the structure of a similar triazole herbicide, methyl 2-(1-diethylcarbamoyl-1,2,4-triazole-3-yl)sulfonyl)acetate, has been reported by Ohkita *et al.* (2002). The structure of 5-[4-cyclopropyl-5-[(3-fluorobenzyl)sulfinyl]-4H-1,2,4-triazol-3-yl]-4-methyl-1,2,3-thiadiazole ($C_{15}H_{14}FN_5OS_2$), was determined by Min *et al.* (2015) and the crystal structure of 1-(mesityl-2-sulfonyl)-3-

Table 2

Experimental details.

Crystal data	$[Ag(NO_3)(C_{16}H_{22}N_4O_3S)]$
Chemical formula	520.31
M_r	Orthorhombic, $Pna2_1$
Crystal system, space group	173
Temperature (K)	9.0947 (2), 31.0133 (6), 7.1934 (1)
a, b, c (Å)	2028.95 (7)
V (Å ³)	4
Z	Mo $K\alpha$
Radiation type	1.14
μ (mm ⁻¹)	$0.48 \times 0.10 \times 0.02$
Crystal size (mm)	
Data collection	
Diffractometer	Bruker APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2014)
T_{min}, T_{max}	0.579, 0.746
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	17454, 4760, 4259
R_{int}	0.042
$(\sin \theta/\lambda)_{max}$ (Å ⁻¹)	0.667
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.030, 0.053, 0.98
No. of reflections	4760
No. of parameters	267
No. of restraints	1
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{max}, \Delta\rho_{min}$ (e Å ⁻³)	0.65, -0.39
Absolute structure	Flack x determined using 1577 quotients $[(I^+)-(I^-)]/[(I^+)+(I^-)]$ (Parsons <i>et al.</i> , 2013)
Absolute structure parameter	0.003 (14)

Computer programs: *APEX2* and *SAINT* (Bruker, 2014), *SHELXS97* and *SHELXTL* (Sheldrick, 2008), *SHELXL2014* (Sheldrick, 2015), *DIAMOND* (Brandenburg, 2010) and *publCIF* (Westrip, 2010).

nitro-1,2,4-triazole has been determined by Kuroda *et al.* (1982). The complex, $[Pr(C_7H_5O_3)_2(NO_3)(C_{12}H_8N_2)] \cdot 2C_{12}H_8N_2$, has a polymeric chain structure, where nitrate ions show similar coordination bonds compared to those in the title compound, but with Ag^I ions replaced by with Pr^{III} atoms (Wang *et al.*, 2012).

5. Synthesis and crystallization

The title compound was prepared from a mixed solution of the cafenstrole ligand (0.05 g, 0.14 mmol) in acetone (5 mL) and $Ag(NO_3)$ (0.06 g, 0.35 mmol) in methanol (5 mL). The ligand was purchased from the Dr Ehrenstorfer GmbH Company. Single crystals suitable for X-ray crystallography were obtained by slow evaporation of the solvent at room temperature after one week.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All H atoms were positioned geometrically and refined using a riding model with $d(C-H) = 0.98$ Å, $U_{iso}(H) = 1.5U_{eq}(C)$ for methyl group, $d(C-H) = 0.99$ Å, $U_{iso}(H) = 1.2U_{eq}(C)$ for Csp^3-H and $d(C-H) = 0.95$ Å, $U_{iso}(H) = 1.2U_{eq}(C)$ for aromatic $C-H$.




Figure 4

The packing diagram showing the three-dimensional network structure formed by $C-H\cdots O$ hydrogen bonds and $C-H\cdots \pi$ interactions (shown as yellow and black dashed lines, respectively).

Acknowledgements

This work was supported by the National Research Foundation (NRF) of Korea, Grant funded by the Ministry of Education, Science and Technology (2014R1A1A4A01008346 and 2014R1A1A4A01009105).

References

- Brandenburg, K. (2010). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2014). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Haasnoot, J. G. (2000). *Coord. Chem. Rev.* **200–202**, 131–185.
- Kang, G., Kim, J., Park, H. & Kim, T. H. (2015). *Acta Cryst.* **E71**, o614.
- Kumar, R., Yar, M. S., Chaturvedi, S. & Srivastava, A. (2013). *Int. J. PharmTech Res.* **5**, 1844–1869.
- Kuroda, R., Sanderson, M. R., Neidle, S. & Reese, C. B. (1982). *J. Chem. Soc. Perkin Trans. 2*, pp. 617–620.
- Min, L.-J., Tan, C.-X., Weng, J.-Q. & Liu, X.-H. (2014). *Phosphorus Sulfur Silicon*, **189**, 379–386.
- Min, L.-J., Yang, M.-Y., Mu, J.-X., Sun, Z.-H., Tan, C.-X., Weng, J.-Q., Liu, X.-H. & Zhang, Y.-G. (2015). *Phosphorus Sulfur Silicon*, **190**, 1884–1892.
- Ohkata, K., Yano, T., Kojima, S., Hiraga, Y., Yoshii, T. & Hori, M. (2002). *Bull. Chem. Soc. Jpn*, **75**, 567–574.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Takahashi, H., Ohki, A., Kanzaki, M., Tanaka, A., Sato, Y., Matthes, B., Böger, P. & Wakabayashi, K. (2001). *J. Biosci.* **56**, 781–786.
- Wang, P., Xu, D. & Wang, X. (2012). *Acta Cryst.* **E68**, m1148.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zhang, H.-Z., Damu, G. L. V., Cai, G.-X. & Zhou, C.-H. (2014). *Curr. Org. Chem.* **18**, 359–406.

supporting information

Acta Cryst. (2016). E72, 1610-1613 [https://doi.org/10.1107/S2056989016016662]

Crystal structure of *catena*-poly[[*(N,N*-diethyl-3-mesitylsulfonyl-1*H*-1,2,4-triazole-1-carboxamide- κ N¹)silver(I)]- μ -nitrate- κ^3 O,*O'*:O]

Hyunjin Park, Eunjin Kwon, Il Yoon and Jineun Kim

Computing details

Data collection: *APEX2* (Bruker, 2014); cell refinement: *SAINT* (Bruker, 2014); data reduction: *SAINT* (Bruker, 2014); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *DIAMOND* (Brandenburg, 2010); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *publCIF* (Westrip, 2010).

catena-Poly[[*(N,N*-diethyl-3-mesitylsulfonyl-1*H*-1,2,4-triazole-1-carboxamide- κ N¹)silver(I)]- μ -nitrate- κ^3 O,*O'*:O]

Crystal data

[Ag(C₁₆H₂₂N₄O₃S)(NO₃)]

M_r = 520.31

Orthorhombic, *Pna*2₁

a = 9.0947 (2) Å

b = 31.0133 (6) Å

c = 7.1934 (1) Å

V = 2028.95 (7) Å³

Z = 4

F(000) = 1056

D_x = 1.703 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 7087 reflections

θ = 2.3–27.1°

μ = 1.14 mm⁻¹

T = 173 K

Plate, colourless

0.48 × 0.10 × 0.02 mm

Data collection

Bruker APEXII CCD

diffractometer

φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2014)

*T*_{min} = 0.579, *T*_{max} = 0.746

17454 measured reflections

4760 independent reflections

4259 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.042

θ_{max} = 28.3°, θ_{min} = 2.3°

h = -12→11

k = -39→41

l = -9→9

Refinement

Refinement on *F*²

Least-squares matrix: full

R[[*F*² > 2σ(*F*²)] = 0.030

wR(*F*²) = 0.053

S = 0.98

4760 reflections

267 parameters

1 restraint

Hydrogen site location: inferred from

neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.0179*P*)²]

where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} = 0.002

Δρ_{max} = 0.65 e Å⁻³

Δρ_{min} = -0.39 e Å⁻³

Absolute structure: Flack *x* determined using
1577 quotients [(*I*⁺)-(*I*)]/[(*I*⁺)+(*I*)] (Parsons et
al., 2013)

Absolute structure parameter: 0.003 (14)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.62392 (3)	0.52477 (2)	0.76161 (6)	0.03314 (9)
S1	0.73402 (9)	0.64406 (3)	0.83512 (13)	0.01964 (19)
O1	0.6829 (3)	0.66757 (8)	0.9943 (4)	0.0300 (7)
O2	0.6265 (2)	0.62494 (9)	0.7166 (4)	0.0309 (8)
O3	1.1685 (2)	0.53847 (7)	1.2570 (5)	0.0290 (5)
O4	0.3336 (4)	0.51667 (11)	0.6854 (5)	0.0486 (9)
O5	0.4227 (3)	0.49348 (10)	0.9434 (4)	0.0382 (7)
O6	0.1895 (3)	0.48605 (10)	0.8835 (5)	0.0424 (8)
N1	0.8072 (3)	0.55871 (9)	0.9126 (4)	0.0206 (7)
N2	0.9749 (3)	0.60970 (9)	0.9892 (4)	0.0187 (6)
N3	1.0242 (3)	0.56942 (9)	1.0338 (4)	0.0183 (6)
N4	1.2736 (3)	0.58767 (9)	1.0626 (4)	0.0206 (7)
N5	0.3120 (4)	0.49853 (10)	0.8351 (5)	0.0266 (8)
C1	0.8614 (3)	0.67468 (11)	0.7068 (5)	0.0205 (9)
C2	0.9148 (3)	0.71461 (10)	0.7730 (7)	0.0218 (7)
C3	1.0110 (4)	0.73701 (13)	0.6569 (6)	0.0286 (9)
H3	1.0465	0.7642	0.6973	0.034*
C4	1.0571 (4)	0.72185 (14)	0.4873 (6)	0.0300 (9)
C5	1.0073 (4)	0.68155 (14)	0.4304 (6)	0.0303 (10)
H5	1.0412	0.6703	0.3152	0.036*
C6	0.9102 (4)	0.65722 (12)	0.5352 (5)	0.0232 (8)
C7	0.8791 (4)	0.73432 (14)	0.9579 (6)	0.0353 (10)
H7A	0.9450	0.7587	0.9811	0.053*
H7B	0.8919	0.7127	1.0559	0.053*
H7C	0.7769	0.7444	0.9576	0.053*
C8	1.1603 (5)	0.74778 (18)	0.3669 (7)	0.0526 (14)
H8A	1.1032	0.7677	0.2896	0.079*
H8B	1.2166	0.7283	0.2868	0.079*
H8C	1.2279	0.7642	0.4459	0.079*
C9	0.8679 (5)	0.61345 (14)	0.4610 (6)	0.0391 (11)
H9A	0.7697	0.6150	0.4054	0.059*
H9B	0.8673	0.5925	0.5630	0.059*
H9C	0.9391	0.6044	0.3665	0.059*
C10	0.8450 (3)	0.60100 (11)	0.9190 (5)	0.0177 (8)
C11	0.9224 (4)	0.53946 (12)	0.9906 (5)	0.0216 (8)
H11	0.9315	0.5094	1.0124	0.026*
C12	1.1648 (4)	0.56330 (12)	1.1294 (5)	0.0193 (8)
C13	1.2803 (4)	0.60769 (12)	0.8765 (6)	0.0254 (9)
H13A	1.1939	0.5983	0.8031	0.030*

H13B	1.3697	0.5974	0.8116	0.030*
C14	1.2828 (4)	0.65612 (13)	0.8847 (7)	0.0365 (11)
H14A	1.1910	0.6666	0.9400	0.055*
H14B	1.2929	0.6678	0.7587	0.055*
H14C	1.3661	0.6656	0.9608	0.055*
C15	1.4092 (4)	0.58914 (14)	1.1768 (6)	0.0319 (10)
H15A	1.4890	0.6026	1.1035	0.038*
H15B	1.4398	0.5593	1.2073	0.038*
C16	1.3886 (5)	0.6139 (2)	1.3531 (8)	0.0642 (17)
H16A	1.3571	0.6433	1.3238	0.096*
H16B	1.4816	0.6148	1.4217	0.096*
H16C	1.3134	0.5998	1.4294	0.096*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.01833 (12)	0.03174 (15)	0.04934 (19)	−0.00517 (11)	−0.00399 (18)	−0.01117 (19)
S1	0.0136 (4)	0.0221 (4)	0.0232 (5)	0.0013 (3)	0.0017 (4)	0.0027 (4)
O1	0.0276 (14)	0.0292 (15)	0.0333 (18)	0.0050 (11)	0.0131 (13)	0.0002 (13)
O2	0.0178 (11)	0.0376 (15)	0.037 (2)	−0.0045 (10)	−0.0073 (12)	0.0054 (13)
O3	0.0248 (12)	0.0308 (13)	0.0315 (14)	−0.0015 (9)	−0.0036 (18)	0.0114 (18)
O4	0.064 (2)	0.044 (2)	0.037 (2)	−0.0106 (17)	0.0084 (16)	0.0004 (17)
O5	0.0227 (13)	0.057 (2)	0.0346 (19)	0.0013 (14)	−0.0001 (14)	−0.0022 (16)
O6	0.0200 (14)	0.053 (2)	0.054 (2)	−0.0094 (13)	0.0092 (14)	−0.0063 (16)
N1	0.0154 (14)	0.0192 (16)	0.0272 (19)	−0.0024 (12)	0.0000 (13)	−0.0001 (14)
N2	0.0189 (14)	0.0181 (15)	0.0192 (17)	0.0011 (11)	−0.0022 (13)	0.0017 (13)
N3	0.0164 (14)	0.0183 (16)	0.0203 (17)	−0.0009 (11)	−0.0002 (13)	0.0040 (14)
N4	0.0152 (14)	0.0212 (16)	0.0253 (19)	−0.0034 (12)	−0.0062 (13)	0.0060 (14)
N5	0.0290 (18)	0.0226 (18)	0.028 (2)	−0.0009 (14)	0.0059 (16)	−0.0059 (16)
C1	0.0175 (16)	0.0221 (18)	0.022 (2)	0.0025 (13)	0.0002 (15)	0.0049 (14)
C2	0.0220 (15)	0.0231 (16)	0.0204 (19)	0.0025 (12)	−0.006 (2)	0.003 (2)
C3	0.029 (2)	0.028 (2)	0.029 (2)	−0.0089 (17)	−0.0079 (18)	0.0088 (19)
C4	0.0210 (19)	0.045 (3)	0.024 (2)	−0.0072 (17)	−0.0035 (18)	0.013 (2)
C5	0.026 (2)	0.045 (3)	0.019 (2)	0.0022 (18)	0.0054 (18)	0.0061 (19)
C6	0.0224 (18)	0.026 (2)	0.021 (2)	0.0042 (15)	0.0016 (16)	0.0023 (17)
C7	0.044 (2)	0.025 (2)	0.037 (3)	−0.0028 (18)	−0.001 (2)	−0.0045 (19)
C8	0.043 (3)	0.078 (4)	0.037 (3)	−0.026 (3)	0.002 (2)	0.019 (3)
C9	0.059 (3)	0.031 (2)	0.027 (3)	−0.004 (2)	0.008 (2)	−0.008 (2)
C10	0.0126 (16)	0.0185 (18)	0.022 (2)	0.0003 (13)	0.0030 (15)	0.0006 (16)
C11	0.0222 (18)	0.0211 (18)	0.021 (2)	−0.0047 (14)	0.0032 (17)	0.0029 (16)
C12	0.0166 (17)	0.0185 (19)	0.023 (2)	0.0010 (13)	0.0000 (16)	0.0005 (16)
C13	0.0181 (17)	0.031 (2)	0.027 (2)	−0.0019 (15)	0.0039 (16)	0.0053 (17)
C14	0.030 (2)	0.027 (2)	0.052 (3)	−0.0036 (17)	0.000 (2)	0.016 (2)
C15	0.0199 (19)	0.033 (2)	0.042 (3)	−0.0039 (17)	−0.0091 (18)	0.007 (2)
C16	0.045 (3)	0.095 (5)	0.053 (3)	−0.001 (3)	−0.026 (3)	−0.025 (3)

Geometric parameters (Å, °)

Ag1—N1	2.250 (3)	C4—C5	1.391 (6)
Ag1—O5 ⁱ	2.396 (3)	C4—C8	1.509 (5)
Ag1—O5	2.450 (3)	C5—C6	1.385 (5)
S1—O2	1.426 (3)	C5—H5	0.9500
S1—O1	1.435 (3)	C6—C9	1.509 (5)
S1—C1	1.760 (4)	C7—H7A	0.9800
S1—C10	1.779 (3)	C7—H7B	0.9800
O3—C12	1.198 (5)	C7—H7C	0.9800
O4—N5	1.231 (4)	C8—H8A	0.9800
O5—N5	1.282 (4)	C8—H8B	0.9800
O5—Ag1 ⁱⁱ	2.396 (3)	C8—H8C	0.9800
O6—N5	1.230 (4)	C9—H9A	0.9800
N1—C11	1.330 (5)	C9—H9B	0.9800
N1—C10	1.357 (4)	C9—H9C	0.9800
N2—C10	1.313 (4)	C11—H11	0.9500
N2—N3	1.365 (4)	C13—C14	1.503 (5)
N3—C11	1.348 (4)	C13—H13A	0.9900
N3—C12	1.465 (4)	C13—H13B	0.9900
N4—C12	1.334 (4)	C14—H14A	0.9800
N4—C13	1.477 (5)	C14—H14B	0.9800
N4—C15	1.483 (5)	C14—H14C	0.9800
C1—C2	1.413 (5)	C15—C16	1.494 (6)
C1—C6	1.419 (5)	C15—H15A	0.9900
C2—C3	1.395 (5)	C15—H15B	0.9900
C2—C7	1.499 (7)	C16—H16A	0.9800
C3—C4	1.373 (6)	C16—H16B	0.9800
C3—H3	0.9500	C16—H16C	0.9800
N1—Ag1—O5 ⁱ	134.63 (11)	H7A—C7—H7C	109.5
N1—Ag1—O5	118.75 (11)	H7B—C7—H7C	109.5
O5 ⁱ —Ag1—O5	106.52 (5)	C4—C8—H8A	109.5
O2—S1—O1	117.78 (16)	C4—C8—H8B	109.5
O2—S1—C1	111.23 (17)	H8A—C8—H8B	109.5
O1—S1—C1	110.96 (16)	C4—C8—H8C	109.5
O2—S1—C10	106.22 (17)	H8A—C8—H8C	109.5
O1—S1—C10	107.14 (17)	H8B—C8—H8C	109.5
C1—S1—C10	102.10 (16)	C6—C9—H9A	109.5
N5—O5—Ag1 ⁱⁱ	118.1 (2)	C6—C9—H9B	109.5
N5—O5—Ag1	102.3 (2)	H9A—C9—H9B	109.5
Ag1 ⁱⁱ —O5—Ag1	137.37 (12)	C6—C9—H9C	109.5
C11—N1—C10	102.7 (3)	H9A—C9—H9C	109.5
C11—N1—Ag1	125.2 (2)	H9B—C9—H9C	109.5
C10—N1—Ag1	131.0 (2)	N2—C10—N1	116.1 (3)
C10—N2—N3	101.4 (3)	N2—C10—S1	119.1 (3)
C11—N3—N2	110.5 (3)	N1—C10—S1	124.8 (3)
C11—N3—C12	128.2 (3)	N1—C11—N3	109.2 (3)

N2—N3—C12	121.0 (3)	N1—C11—H11	125.4
C12—N4—C13	126.5 (3)	N3—C11—H11	125.4
C12—N4—C15	115.7 (3)	O3—C12—N4	128.3 (3)
C13—N4—C15	117.1 (3)	O3—C12—N3	117.9 (3)
O6—N5—O4	122.4 (4)	N4—C12—N3	113.9 (3)
O6—N5—O5	120.0 (4)	N4—C13—C14	112.6 (3)
O4—N5—O5	117.6 (4)	N4—C13—H13A	109.1
C2—C1—C6	121.3 (3)	C14—C13—H13A	109.1
C2—C1—S1	121.5 (3)	N4—C13—H13B	109.1
C6—C1—S1	117.2 (3)	C14—C13—H13B	109.1
C3—C2—C1	116.8 (4)	H13A—C13—H13B	107.8
C3—C2—C7	117.6 (3)	C13—C14—H14A	109.5
C1—C2—C7	125.6 (3)	C13—C14—H14B	109.5
C4—C3—C2	123.6 (4)	H14A—C14—H14B	109.5
C4—C3—H3	118.2	C13—C14—H14C	109.5
C2—C3—H3	118.2	H14A—C14—H14C	109.5
C3—C4—C5	118.0 (4)	H14B—C14—H14C	109.5
C3—C4—C8	121.2 (4)	N4—C15—C16	112.4 (4)
C5—C4—C8	120.8 (4)	N4—C15—H15A	109.1
C6—C5—C4	122.5 (4)	C16—C15—H15A	109.1
C6—C5—H5	118.8	N4—C15—H15B	109.1
C4—C5—H5	118.8	C16—C15—H15B	109.1
C5—C6—C1	117.7 (4)	H15A—C15—H15B	107.9
C5—C6—C9	117.4 (4)	C15—C16—H16A	109.5
C1—C6—C9	124.8 (3)	C15—C16—H16B	109.5
C2—C7—H7A	109.5	H16A—C16—H16B	109.5
C2—C7—H7B	109.5	C15—C16—H16C	109.5
H7A—C7—H7B	109.5	H16A—C16—H16C	109.5
C2—C7—H7C	109.5	H16B—C16—H16C	109.5
C10—N2—N3—C11	−0.7 (4)	N3—N2—C10—N1	−0.7 (4)
C10—N2—N3—C12	−176.0 (3)	N3—N2—C10—S1	−179.2 (2)
Ag1 ⁱⁱ —O5—N5—O6	−14.1 (4)	C11—N1—C10—N2	1.8 (4)
Ag1—O5—N5—O6	179.9 (3)	Ag1—N1—C10—N2	−166.7 (3)
Ag1 ⁱⁱ —O5—N5—O4	164.7 (3)	C11—N1—C10—S1	−179.8 (3)
Ag1—O5—N5—O4	−1.3 (4)	Ag1—N1—C10—S1	11.7 (5)
O2—S1—C1—C2	140.4 (3)	O2—S1—C10—N2	161.7 (3)
O1—S1—C1—C2	7.2 (3)	O1—S1—C10—N2	−71.6 (3)
C10—S1—C1—C2	−106.7 (3)	C1—S1—C10—N2	45.1 (3)
O2—S1—C1—C6	−40.6 (3)	O2—S1—C10—N1	−16.7 (4)
O1—S1—C1—C6	−173.8 (3)	O1—S1—C10—N1	110.0 (3)
C10—S1—C1—C6	72.3 (3)	C1—S1—C10—N1	−133.3 (3)
C6—C1—C2—C3	3.5 (5)	C10—N1—C11—N3	−2.1 (4)
S1—C1—C2—C3	−177.6 (3)	Ag1—N1—C11—N3	167.3 (2)
C6—C1—C2—C7	−175.3 (3)	N2—N3—C11—N1	1.9 (4)
S1—C1—C2—C7	3.6 (5)	C12—N3—C11—N1	176.8 (3)
C1—C2—C3—C4	−1.4 (5)	C13—N4—C12—O3	160.4 (4)
C7—C2—C3—C4	177.5 (4)	C15—N4—C12—O3	−10.0 (6)

C2—C3—C4—C5	−1.3 (6)	C13—N4—C12—N3	−21.2 (5)
C2—C3—C4—C8	179.4 (4)	C15—N4—C12—N3	168.4 (3)
C3—C4—C5—C6	2.1 (6)	C11—N3—C12—O3	−39.4 (5)
C8—C4—C5—C6	−178.6 (4)	N2—N3—C12—O3	135.0 (4)
C4—C5—C6—C1	−0.1 (6)	C11—N3—C12—N4	142.0 (4)
C4—C5—C6—C9	−178.3 (4)	N2—N3—C12—N4	−43.6 (4)
C2—C1—C6—C5	−2.8 (5)	C12—N4—C13—C14	116.8 (4)
S1—C1—C6—C5	178.2 (3)	C15—N4—C13—C14	−73.0 (4)
C2—C1—C6—C9	175.2 (4)	C12—N4—C15—C16	−70.8 (5)
S1—C1—C6—C9	−3.8 (5)	C13—N4—C15—C16	117.9 (4)

Symmetry codes: (i) $-x+1, -y+1, z-1/2$; (ii) $-x+1, -y+1, z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of the C1—C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C8—H8C \cdots Cg1 ⁱⁱⁱ	0.98	2.66	3.614 (5)	166
C13—H13B \cdots O2 ^{iv}	0.99	2.58	3.395 (4)	140
C16—H16B \cdots O2 ^v	0.98	2.52	3.412 (6)	152

Symmetry codes: (iii) $x+1/2, -y+3/2, z$; (iv) $x+1, y, z$; (v) $x+1, y, z+1$.