

A Contribution to the Knowledge of the Tenthredinidae (Symphyta, Hymenoptera) Fauna of Turkey

Part I: The Subfamily Tenthredininae*

Önder ÇALMAŞUR, Hikmet ÖZBEK
Atatürk University, Faculty of Agriculture, Department of Plant Protection,
25240 Erzurum - TURKEY

Received: 07.02.2003

Abstract: The subfamily Tenthredininae in the family Tenthredinidae was treated in this part of the study regarding the sawfly (Symphyta, Hymenoptera) fauna of Turkey. The materials were collected from various localities around the country, though examples from eastern Turkey are predominant. After examining more than 2500 specimens, 57 species in 8 genera were recorded. Eleven species were new for Turkish fauna; of these 3 species were recorded for the first time as Asian fauna. Furthermore, 3 species were endemic for Turkey. The distribution and new areas as well as the host plants of some species around Turkey and the world were given. For each species its chorotype was reported.

Key Words: Hymenoptera, Tenthredinidae, Tenthredininae, Fauna, Turkey

Türkiye'nin Tenthredinidae (Symphyta, Hymenoptera) Faunasına Katkılar Bölüm:I Tenthredininae Altfamilyası

Özet: Türkiye'nin testerelileri (Symphyta, Hymenoptera) faunasının tespitine yönelik çalışmaların bu bölümünde; Tenthredininae (Tenthredinidae) altfamilyası ele alınmış; incelenen 2500'den fazla örnek sonucu, sekiz cinse bağlı toplam 57 tür saptanmıştır. Türkiye faunası için yeni oldukları saptanan 11 türden üçünün Asya faunası için de yeni kayıt oldukları belirlenmiştir. Üç türün Türkiye için endemik oldukları saptanmıştır. Tespit edilen türlerin hemen tamamı için yeni yayılma alanları belirlenmiş, birçoğunun konukçuları bulunmuştur. Türkiye ve dünyadaki dağılışları chorotype'leri ile birlikte verilmiştir.

Anahtar Sözcükler: Hymenoptera, Tenthredinidae, Tenthredininae, Fauna, Türkiye

Introduction

The topographic and climatic structures of Turkey give it a rich and diverse fauna and flora. Turkey's location at the junction of the Euro-Siberian, Mediterranean and Turanian zoogeographic regions, and the fact that it is a crossroad between the continents of Europe, Asia and Africa, only adds to this richness.

The family Tenthredinidae (Symphyta, Hymenoptera) are called "Typical Sawflies"; it is a large family, including more than seven-eighths of all members of the suborder Symphyta (Chalastogastra) (Comstock, 1964). The adults are often brightly colored, and are usually found on

flowers and foliage; they are small to medium size, being 2.5-15.0 mm in length. The larvae are euriciform, mostly free-living, and most of them are external feeders on foliage, though some of them are extremely destructive pests. A few species are gall makers, and a few are leaf miners (Benson, 1952).

Tenthredinidae is best represented in the north temperate region; species richness tends to decrease from north to south. There are more than 6000 species in 360 genera in temperate regions of the Northern Hemisphere (Liston, 1995; Goulet and Huber, 1993; Lacourt, 1999; Taeger and Blank, 1998). More than 900 species occur in Europe (Liston, 1995).

* This study is part of the requirements for the doctoral degree submitted to Atatürk University on 01.11.2002.

There is considerable disagreement over the composition of the various subordinate groups. Fitton et al. (1978) adopted the system proposed by Benson (1952, 1958). Gauld and Bolton (1988) followed the classification of Smith (1979) and recognized 7 subfamilies. More recently, Lacourt (1999) divided the family Tenthredinidae into 14 subfamilies in his catalog.

Although various contributions have been conducted on their taxonomy, biology and distribution as well as the damage caused to agricultural and ornamental plants by sawflies in the Palearctic region (Enslin, 1918; Zhelokachev, 1928; Benson, 1950, 1951, 1952, 1958, 1967, 1968; Kontuniemi, 1965; Zombori, 1972, 1973, 1974a, 1974b, 1977, 1978, 1979, 1981, 1982a, 1982b, 1984, 1985; Moczar and Zombori, 1973; Chevin, 1975, 1984; Muche, 1975; Vikberg, 1975; Smith, 1976, 1981, 1982, 1986; Shaw, 1978; Vassilev, 1978; Altenhofer, 1980; Cingovski, 1980, 1982; Schedl, 1981, 1991; Scobiola-Palade, 1981; Liston, 1981, 1983a, 1983b, 1983c; Meitzner and Taeger, 1982; Viitasaari and Vikberg, 1985; Magis, 2002), the Turkish fauna is comparatively poorly studied despite its zoogeographic interest. The first comprehensive study on the sawfly fauna of Turkey was conducted by Guichard and Harvey (1967). Benson (1966, 1968) made a faunistical and systematical study on the Symphyta of Turkey, and described new taxa and constructed keys to some genera and species occurring in Turkey. Later, Wolf (1968) and Chevin and Chenon (1982) recorded some sawfly species from Turkey. Since Chevin and Chenon (1982) no additional faunal study has been conducted, although some studies have appeared dealing with different sawfly pests on various cultivated, forest, and ornamental plants (Alkan, 1948; Özeren, 1952, 1970; Öge, 1958; Birkardeşler, 1964, 1971; Kiroğlu and Baysal, 1971; Baş, 1973; Özbek, 1986; Erden, 1988; Güçlü and Özbeş, 1999). The objectives of this study are to contribute to the knowledge of the typical sawfly fauna of Turkey. In this first article, the subfamily Tenthredininae is described.

Materials and Methods

The present paper is based on more than 2500 specimens collected during 1996-2002 in various localities of Turkey by the authors. In addition, material accumulated since the 1970s in the Entomology Museum

Erzurum, Turkey (EMET), was evaluated and included in this work. Although the specimens have been collected in different parts of the country, the majority of the sawfly samples were collected from eastern Turkey. The habitats were mainly alpine meadows, mountain tundra, roadsides, and uncultivated areas between cultivated lands, orchards and open areas in forests. Besides the adults collected by insect net, immature stages of sawflies were also collected and reared under laboratory conditions to obtain adults. The materials mentioned in this study were deposited in EMET, Turkey. For determination of the material, Benson (1952) and Medvedev (1994) were used, and for taxa synonyms Lacourt (1999) was followed. Undetermined specimens were identified and others reviewed by Dr. David R. Smith and Dr. Lajos Zombori.

Genera and species were treated in alphabetical order in the subfamilies. All records were arranged in the following way: "province (alphabetically ordered): town, local place, geographic name and altitude if available, date of collecting, number of individuals as male and female" (except the widespread species occurring in Erzurum province). The name of the province from which a particular species was recorded for the first time was given in bold type. Species new for Turkish fauna were specified.

According to their distribution patterns, the species were classified into major chorotypes, in accordance with guidelines proposed by Vigna Taglianti et al. (1999). This chorotype classification is based only on distribution patterns, without paleogeographical or ecological assumptions.

Results

In the subfamily Tenthredininae 57 species in 8 genera were recorded.

Cuneala Zirngiebl, 1956

Synonym: *Parastasis* Kirby, 1881; *Labidia* Provancher, 1886; *Rethrax* Cameron, 1899; *Fethali* Cameron, 1902; *Clydostomus* Konow, 1908; *Tenthredella* Rohwer, 1910; *Zamacrophya* Rohwer, 1912.

Cuneala amasiensis (Kriechbaumer, 1869)

Material Examined: Denizli: Pamukkale, 5.V.1994, 1♀. İzmir: Efes, 5.V.1998, 2♀.

Distributed in Turkey: Amasya, Ankara, Aydın, Bursa and Çorum (Benson, 1968).

General Distribution and Chorotype: Turkey (Benson, 1968; Lacourt, 1999). It is endemic for Turkey.

Host Plant: Unknown

Cuneala confinis (Konow, 1886)

Synonym: *Allantus koehleri* sensu Puls, 1870, nec Klug, 1817; *A. atratus* André, 1883; *A. confinis* Konow, 1886; *A. parviceps* Konow, 1898.

Material Examined: Artvin: Borçka, Karagöl, 1050 m, 8.VII.1994, 2♂, 1♀; Genye Dağı, 1600-1800 m, 16.VIII.1990, 27♀. Erzurum: Aşkale, Kopdağı, 2000 m, 16.VI.2000, 1♂, 1♀. Rize: Çamlıhemşin, Ayder, 1550 m, 30.VII.2000, 19♀; İkizdere, Çamlık, 1600 m, 29.VII.2000, 1♂, 3♀, Ovit, 2500 m, 22.VII.2002, 4♀, 29.VII.2000, 3♀. Trabzon: Tonya, Erikbeli, 11.VII.1994, 1♀; Sümelamanastırı, 2.VII.2002, 2♂.

Distributed in Turkey: Bolu, Çankırı, Gümüşhane, Kastamonu, Kayseri, Rize and Trabzon (Benson, 1968).

General Distribution and Chorotype: Turkey, Transcaucasia, Iran (Benson, 1968; Lacourt, 1999); Turano-Mediterranean.

Host Plant: *Geranium psilostemum* (Lacourt, 1999).

Cuneala koehleri Klug, 1817

Material Examined: Artvin: Borçka, Karagöl, 1050 m, 8.VIII.1994, 7♀; Genye Dağı, 1600 m, 16.VIII.1990, 2♀. Erzurum: Şenkaya, Ormanlı, 1800 m, 20.V.1994, 1♀. Rize: Çamlıhemşin, 500-1200 m, 30.VII.2000, 2♀; İkizdere, Çamlık, 1600 m, 29.VII.2000, 2♀, Ovit, 1600-2400 m, 29.VII.2000, 3♀.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Europe, Yugoslavia, Russia (Liston, 1995; Lacourt, 1999); Sibero-European.

Remarks: Present data show that its easternmost record is eastern Turkey.

Host Plant: *Granium silvaticum*, *Sanguineum* spp., *Robertianum* spp., *Cardamine* spp. (Liston, 1995; Taeger and Blank, 1998; Lacourt, 1999).

Elinora Benson, 1946

Synonym: *Cuneala* Zirngiebl, 1956

Elinora asiatica Enslin, 1910

Material Examined: Bayburt: Çalidere, 1650 m, 16.VI.2000, 1♀. Erzurum: University field, 1850 m, 26.V.1994, 1♀, 31.V.2001, 1♂, 4♀, 1-11.VI.1971, 1♂, 9♀, 3.VI.2000, 5♀, 12.VI.1994, 1♀; Dumlu, 1850 m, 25.VI.2002, 1♀; Pasinler, 1700 m, 10.VI.1983, 1♀, 11.VI.1981, 1♂, 12.VI.1996, 1♂, 3♀, 2.VII.1983, 2♀; Köprüköy, Örentaş, 1900 m, 12.VI.2001, 1♀; Narman, Savaşcılar, 2300 m, 13.VI.1999, 1♀; Oltu, Başaklı, 1850 m, 20.V.1972, 1♂; Şenkaya, Ormanlı, 1800 m, 25.V.1994, 1♂, 1♀, Turnalı, 1750 m, 25.V.1994, 2♀; Uzungere, Şelale, 1000 m, 9.VI.1996, 1♀. İğdır: 850 m, 25.V.1971, 1♀. Isparta: S.Karaağaç, 1.V.1994, 1♀. Kars: Sarıkamış, Karakurt, TCK Çeşmesi, 1600 m, 2.VI.1999, 1♂, 1500 m, 4.VI.2000, 1♂, 1♀. Muğla: 27.VII.1983, 1♂.

Distributed in Turkey: Benson (1968) and Lacourt (1999) indicated the presence of this species in Turkey, but no locality was noted.

General Distribution and Chorotype: Present knowledge shows that this species is endemic for Turkey.

Host Plant: Adults were collected from pastures.

Elinora baetica Spinola, 1843

Synonym: *Allantus fulviventris* Mocsary, 1880; *A. dusmeti* Konow, 1894; *A. antigae* var. *atrosutellatus* Zirngiebl, 1937.

Material Examined: Erzurum: University field, 1850 m, 7.V.1970, 1♀, 25.V.1983, 1♀, 1.VI.2000, 1♂, 2♀, 8.VI.1994, 2♀, 11.VI.1971, 1♀, 30.VI.1992, 1♀; Dutçu, 1950 m, 20.VI.1994, 1♀, Nenehatun, 2100 m, 27.VI.1999, 1♀; Aşkale, Küçükgeçit, 1750 m, 15.VI.2002, 1♀; Köprüköy, Örentaş, 1900 m, 21.VI.2001, 1♂, 3♀. Kars: Sarıkamış, Karakurt, 1550 m, 2.VI.1999, 1♂, 4.VI.2000, 5♂, 1♀.

Distributed in Turkey: New record for Turkish fauna and Asia.

General Distribution and Chorotype: Spain and Portugal (Liston, 1995; Lacourt, 1999); Turano-European.

Host Plant: Larvae were collected on *Sinapis arvensis*. Lacourt (1999) gave the same plant as host.

Remarks: It is a new record for Asia as well as Turkey.

Elinora ornata André, 1881

Synonym: *Allantus discolor* Konow, 1890; *A. ornatus* André, 1881; *A. mocsaryi* Enslin, 1910; *Tenthredo baja* Enslin, 1912; *T. camilla* Enslin, 1912.

Material Examined: **Kars:** Sarıkamış, Karakurt, 1550 m, 23.V.2001, 3♂♀, 27.V.2000, 1♂, 1♀, 2.VI.1999, 1♂, 4:VI.2000, 5♂♂, 10♀♀, 16.VI.2001, 1♀.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Transcaucasia (Benson, 1968; Lacourt, 1999); Turano-Anatolian.

Host Plant: Unknown.

Remarks: Its westernmost distribution area is west of Kars province.

***Macrophya* Dahlbom, 1835**

Synonym: *Zalagium* Rohwer, 1912; *Beldonea* Enderlein, 1919, nec Cameron, 1899; *Paramacrophya* Forsius, 1918.

***Macrophya albicincta* (Schrank, 1776)**

Synonym: *M. leucopoda* Palma, 1861; *M. melasoma* Rudow, 1871; *Tenthredo albipes* Geoffroy, 1785; *T. fera* Fallen, 1808, nec Scopoli, 1786; *T. (Allantus) lugubris* Audinet-Serville, 1823; *T. (A.) luctuosa* Audinet-Serville, 1823 nec Schrank, 1802.

Material Examined: **Artvin:** Borçka, Karagöl, 1050 m, 8.VII.1994, 1♂.

Distributed in Turkey: Bolu, Giresun, Bursa, Rize, Samsun, Trabzon and Zonguldak (Benson, 1968; Chevin and Chenon, 1982).

General Distribution and Chorotype: Europe, Turkey, North Iran, Russia, Caucasus, Transcaucasia, Siberia, Baltic Coasts, Algeria (Benson, 1968; Cingovski, 1982; Liston, 1995; Lacourt, 1999; Medvedev, 1994); Centralasiatic-Mediterranean.

Host Plant: *Sambucus ebulus*, *S. nigra*, *S. racemosa*, *Valeriana officinalis*, *Viburnum opalus* (Liston, 1995; Taeger and Blank, 1998; Lacourt, 1999).

***Macrophya annulata* Geoffroy, 1785**

Synonym: *Tenthredo ligata* Müler, 1776; *T. similis* Spinola, 1808; *T. (Allantus) neglecta* Klug, 1817; *T. (A.) abietis* Audinet-Serville, 1823, nec Linne, 1758.

Material Examined: **Gümüşhane:** Vaukgeçidi, 1850 m, 21.VI.2001, 3♂♂, 1♀. **Rize:** Çamlıhemşin, Ayder, 1200-1550 m, 30.VII.2000, 2♂♂; İkizdere, Çamlık, 1600-2400 m, 29.VII.2000, 3♂♂.

Distributed in Turkey: Ankara and Samsun (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Iran, Russia, Siberia, Crimea, Caucasus (Benson, 1968; Liston, 1995; Lacourt, 1999); West Palearctic.

Host Plant: *Potentilla reptans*, *Euphorbia*, *Origanum vulgare*, *Rosa* spp., *Rubus* spp., *Sambucus* spp. (Liston, 1995; Taeger and Blank, 1998; Lacourt, 1999).

***Macrophya blanda* Fabricius, 1775**

Synonym: *Tenthredo solitaria* Schrank, 1781, nec Scopoli, 1763; *T. ligustrina* Geoffroy, 1785; *T. cylindrica* Fabricius, 1804; *T. (Allantus) lacrymosa* Audinet-Serville, 1823; *T. cognata* Fallen, 1829; *T. nyctea* Fischer de Waldheim, 1843.

Material Examined: **Adiyaman:** Nemrut Mountain, 1950 m, 29.V.1999, 1♀. **Artvin:** Ardanuç, Akarsu, 900 m, 7.VII.1994, 1♀, Ferhatlı, 450 m, 13.V.2000, 1♀; İşhan, 500 m, 16.VI.2001, 1♂; Şavşat, Karagöl, 1650 m, 8.VII.1998, 1♀; Yusufeli, 500 m, 7.VII.1992, 1♀, Ormanlı, 300 m, 12.V.2000, 1♀, Sarıgöl, 700 m, 5.VII.1994, 3♂♂, 5♀♀, Altıparmak, 1100 m, 5.VII.1994, 1♀. **Bayburt:** Kopdağı, 2100 m, 16.VI.2000, 1♀. **Erzincan:** Avcılar, 1300 m, 26.V.2001, 1♀; Dumanlı, 1300 m, 25.V.2001, 1♀. **Erzurum:** It is common and abundant in Erzurum province. The materials were collected from the end of May to the end of July from University field, Dutçu, Köşk, Çat, Hınıs, İlıca (Atılıkonak), İspir (Madenköprübaşı), Köprüköy (Örentaş), Narman (Kireçlidağı), Oltu (Başaklı, Çamlıbel, Sütkans, Sarısaz, Tutmaç), Olur, Pasınler (Çalıyazı, Pelitli), Pazaryolu (Karataş, Toprakkale), Şenkaya (Turnalı), Tortum (Aksukapı, Çataldere, Doruklu) and Uzundere (Dikyar) at altitudes of 1300-2400 m, 18♂♂, 61♀♀. **Gümüşhane:** Torul, Zigana, 1900 m, 14.VI.1990, 1♀. **Kars:** Sarıkamış, Karakurt, Aras Vadisi, 1500 m, 4.VI.2000, 1♀, 18.VII.2002, 1♀. **Hatay:** Dörtyol, Topakbaş Yaylası, 1200 m, 25.V.2002, 1♀. **Rize:** Çamlıhemşin, Ayder, 1200-1550 m, 30.VII.2000, 1♀; İkizdere, Ovit Dağı, 2400 m, 29.VII.2000, 1♀. **Trabzon:** Hamsiköy, 1600 m, 24.VI.1998, 1♀.

Distributed in Turkey: Amasya, Ankara, Artvin, Erzurum, Giresun, Konya, Mersin, Samsun, Trabzon, and Uşak (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Moldova, Transcaucasia, Caucasus, Central Asia, Siberia, Crimea, Iran (Benson, 1968; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Sibero-European.

Host Plant: Adults were collected on flowers of *Fragaria* sp. and *Rubus* sp. Liston (1995) indicated that adults of some sawfly species show a strong association with the flowers of the larval host.

***Macrophya crassula* (Klug, 1817)**

Synonym: *M. klugii* Snellen van Vollenhoven, 1869; *M. cora* Kirby, 1886; *Tenthredo (Allantus) maculosa* Audinet-Serville, 1823.

Material Examined: Artvin: Şavşat, Karagöl, 1650 m, 8.VII.1998, 1♂, 3♀, Yusufeli, Altıparmak, 1100 m, 5.VII.1994, 2♂♂.

Distributed in Turkey: Artvin, Bursa, Mersin, Samsun, Sinop and Trabzon (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia (Benson, 1968; Cingovski, 1982; Medvedev, 1994; Lacourt, 1999).

Host Plant: *Sambucus ebulus* (Liston, 1995; Taeger and Blank, 1998).

***Macrophya cyrus* Benson, 1954**

Material Examined: Bayburt: 25 km S., 1750 m, 21.VI.2001, 1♀; Kop Dağı, 2100 m, 16.VI.2000, 2♀♀, Maden, 1650 m, 16.VI.2000, 3♂♂, 7♀♀. Bingöl: Çirişli Geçidi, 2000 m, 11.VII.2002, 2♀♀, Çobantaş, 1400 m, 2.VI.2001, 5♂♂, 5♀♀. Bitlis: 28.V.1989, 1♀. Elazığ: Keban, 900 m, 16.V.2002, 1♂, 1♀. Erzincan: Avcılar, 1300 m, 26.V.2001, 1♂, 1♂, Dumanlı, 1200 m, 25.V.2001, 2♂♂, 3♀♀, Bahçeli, 1350 m, 26.V.2001, 4♂♂, 10♀♀, Gülkaynak, 1200 m, 25.V.2001, 1♂, 1♀, Sansa Boğazı, 1350 m, 25.V.2001, 1♀, Üzümlü, 1200 m, 25.V.2001, 2♂♂, 5♀♀. Erzurum: It is common and abundant in Erzurum province. The materials were collected from the end of May to the end of July from Akdağ, Dutçu, Güngörmez, Karagöbek, Köşk, Nenehatun, Umudum, Yeşildere, Aşkale, Çat (Çimisi), Hınıs, İlıca (Atılıkonak, Gülyurt Geçidi), İspir (Madenköprübaşı), Köprüköy (Örentaş), Narman (Kireçli), Oltu (Başaklı), Sarısaç, Sütkans, Tutmaç), Pasınler (Çalıyazı, Pelitli), Pazaryolu (Karataş), Şenkaya (Soğanlı Dağı), Tortum (Çataldere, Pehlivanlı, Şenyurt) and Uzundere (Yedigöller) at altitudes of 1100-2400 m, 19♂♂, 94♀♀. Gümüşhane: Vauk Geçidi, 1850 m, 21.VI.2001, 2♀♀. Kars: Sarıkamış, İsisu, 13.VII.1991, 3♂♂, 2♀♀, Karaurgan, 2000 m, 24.VI.1999, 2♂♂, Karakurt, 1500 m, 11.VI.1998, 1♀, 4.VI.2000, 1♂. Konya: Güneysünir, Güragaç, 1000 m, 2.VI.2002, 1♀.

Distributed in Turkey: Afyon, Amasya, Ankara, Niğde, Gümüşhane, Erzurum and Konya (Benson, 1968; Chevin and Chenon, 1982).

General Distribution and Chorotype: Turkey, Armenia, Iran (Benson, 1968; Lacourt, 1999); Irano-Anatolian.

Host Plant: Unknown.

Remarks: Present data show that its westernmost distribution point is Afyon.

***Macrophya diversipes* (Schrink, 1782)**

Synonym: *M. flavipes* Tischbein, 1852; *M. haematopus* var. *maculativentris* Costa, 1866; *M. eximia* Mocsary, 1877; *M. caucasica* André, 1881; *M. rubripes* André, 1881; *M. saundersi* Kirby, 1886; *M. sanguinipes* Mocsary, 1891; *M. dalmatina* Gasparini, 1891; *Tenthredo haematopus* Villers, 1789; *T. ochreata* Panzer, 1804, *T. orallipes* Eversmann, 1847; *Tenthredo halensis* Aichinger, 1870.

Material Examined: Artvin: Şavşat, Çayağzı, 500 m, 27.V.1997, 1♂; Yusufeli, Altıparmak, 12.VI.1997, 1♀, Demirkent, 450 m, 6.VII.1994, 1♀. Bayburt: Çalıdere, 1850 m, 16.VI.2000, 6♂♂, 3♀♀; Maden, 1650 m, 16.VI.2000, 10♂♂, 9♀♀, 1700 m, 21.VI.2001, 1♂; Kopdağı, 2100-2300 m, 16.VI.2000, 13♂♂, 14♀♀, 21.VI.2001, 4♀♀. Bingöl: Çirişligeçidi, 2000 m, 11.VII.2002, 2♂♂. Erzincan: Bahçeli, 1300 m, 26.V.2001, 2♀♀; Cevizli, 1300 m, 4.VI.1990, 1♀; Dumanlı, 1200 m, 25.V.2001, 1♀; Gulkaymak, 1200 m, 25.V.2001, 1♂, 1♀; Sansa Boğazı, 1350 m, 25.V.2001, 2♂♂, 5♀♀; Üzümlü, 1200 m, 25.V.2001, 2♀♀. Erzurum: It is common and abundant in Erzurum province. The materials were collected from the end of May to the middle of August from Aribahçe, Dutçu, Güngörmez, Karagöbek, Kargapazarı, Köşk, Nenehatun, Palandöken, University field, Yeşildere, Aşkale (Mezra), Çat (Çimisi), Hınıs, Horasan (Karaçuha), İlıca (Atılıkonak, Eğerti, Gülyurt Bakımevi, Rizekent), İspir (Çayırozü, Fısırik Kalesi, Madenköprübaşı), Köprüköy (Örentaş), Narman (Kireçli), Oltu (Başaklı, Çamlıbel, Sarısaç, Sütkans, Tutmaç), Olur (Salaçur, Süngübayı), Pasınler (Çalıyazı, Pelitli), Pazaryolu (Karataş, Toprakkale), Şenkaya (Soğanlı Dağı, Timurkişla, Turnalı), Tortum (Aksuyayla, Aktaş, Çataldere, Pehlivanlı, Şenyurt, Yukarisivri), and Uzundere (Dikyar, Şelale) at altitudes of 1000-2400 m, 129♂♂, 188♀♀. Gümüşhane: Torul, Zigana, 1350 m, 24.VI.1998, 1♀; Vauk Geçidi, 1850 m, 21.VI.2001, 2♂♂, 2♀♀. İğdır: 850 m, 25.VI.1971, 1♀. Kars: Sarıkamış,

Karakurt, 1500 m, 27.V.2000, 1♂, 4.VI.2000, 2♀, Şeytangeçmez, 1500 m, 2.VI.1999, 4♀, 11.VI.1996, 1♀. Rize: Çamlıhemşin, Ayder, 1550 m, 30.VII.2000, 1♀; İkizdere, Ovit, 2400 m, 29.VII.2000, 2♀.

Distributed in Turkey: Amasya, Ankara, Artvin, Çorum, Erzurum, Gümüşhane, Muğla and Mersin (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Caucasus, Kazakhstan, Urals, Iran (Benson, 1968; Medvedev, 1994; Lacourt, 1999); Asiatic-European.

Host Plant: Larvae were observed feeding on *Fragaria* sp. and adults were collected from the flowers of *Fragaria* sp. and *Rubus* sp.

***Macrophya duodecimpunctata* Linne, 1758**

Synonym: *M. novemguttata* Costa, 1859; *M. luridicarpa* Costa, 1894; *Tenthredo signata* Scopoli, 1763; *T. fera* Scopoli, 1786; *T. melanoleuca* Gmelin, 1789; *T. albamacula* Audinet-Serville, 1823; *T. idriensis* Lepeletier, 1823; *T. dolens* Eversmann, 1847; *T. curvipes* Gimmerthal, 1847; *Pachyprotasis dolens* André, 1881.

Material Examined: Bayburt: Maden, 1650 m, 16.VI.2000, 1♀. Erzurum: İlica, Atlikonak, 2000-2400 m, 29.VI.1999, 6♀, 29.VI.2001, 1♂, 1♀; Hınıs, 1800 m, 14.VI.2002, 1♀; Horasan, Aras Vadisi, 1600 m, 23.V.2001, 2♀; Oltu, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 1♀.

Distributed in Turkey: Samsun (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Caucasus, Siberia, Japan, North Africa (Benson, 1968; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Palearctic.

Host Plant: *Gramineae*, *Carex* sp., *Cyperaceae* (Liston, 1995; Taeger and Blank, 1998).

***Macrophya erythrocnema* A.Costa, 1859**

Synonym: *Tenthredo (Macrophya) femoralis* Kawall, 1864, nec Stephens, 1835

Material Examined: Bayburt: Kopdağı, 2100-2300 m, 16.VI.2000, 1♂, 2♀. Erzurum: Köprüköy, Örentaş, 1900-2100 m, 12.VI.2001, 1♀; Oltu, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 2♂, 2♀; Şenkaya, Turnalı, 1750 m, 25.VII.1996, 1♀; Tortum, Çataldere, 1950-2500 m, 12.VII.2001, 1♀. Rize: İkizdere, Çamlık, 1650 m, 29.VII.2000, 1♀, Ovit Dağı, 2300 m, 29.VII.2000, 1♀;

Çamlıhemşin, Ayder, 1200-1550 m, 30.VII.2000, 1♀. Trabzon: Hamsiköy, 1600 m, 24.VI.1998, 1♀.

Distributed in Turkey: Artvin and Rize (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Transcaucasia, Israel (Benson, 1968; Medvedev, 1994; Lacourt, 1999); Asiatic-European.

Host Plant: Unknown.

***Macrophya militaris* Klug, 1817**

Synonym: *M. lepeletieri* Costa, 1859; *Tenthredo analis* Spinola, 1808, nec Fabricius, 1793; *T. (Allantus) schaefferi* Audinet-Serville, 1823, nec Klug, 1817.

Material Examined: Artvin: Ormanlı, 300 m, 12.V.2000, 1♀. Erzurum: Oltu, Sarısaz, 1450 m, 21.VI.2000, 3♀.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Russia, Caucasus, Balkans, Central and Southern Europe (Medvedev, 1994; Liston, 1995; Lacourt, 1999); Central Asiatic-European.

Host Plant: *Rubus idaeus*, *Origanum vulgare*, (Medvedev, 1994; Liston, 1995; Taeger and Blank, 1998; Lacourt, 1999).

***Macrophya montana* Scopoli, 1753**

Material Examined: Bayburt: Kopdağı, 2300 m, 16.VI.2000, 2♂, Maden, 1650 m, 16.VI.2000, 2♀. Erzurum: İspir, Madenköprübaşı, 1400 m, 18.VI.1994, 2♀; Şenkaya, Turnalı, 1750 m, 6.VII.1990, 1♂; Tortum, Aşağımeydanlar, 4.VII.1991, 1♂, Pehlivanlı, 1400 m, 12.VII.1992, 2♂, 1♀; Uzunkavak, 1800 m, 12.VII.2001, 1♀.

Distributed in Turkey: Afyon, Amasya, Isparta, Rize, Samsun and Sivas (Benson, 1968; Chevin and Chenon, 1982).

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia, Crimea, Turkistan, Urals, Iran, North Africa (Benson, 1968; Chevin and Chenon, 1982; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Palearctic.

Host Plant: *Rubus caesius* (Liston, 1995; Taeger and Blank, 1998).

***Macrophya postica* Brullé, 1832**

Synonym: *M. ratzeburgii* Tischbein, 1852; *M. histrionica* Snellen, 1878.

Material Examined: Adana: Ceyhan, 22.V.1992, 2♀; Yumurtalık, 22.V.1992, 4♀. Artvin: Kalburlu, 300 m, 13.V.2000, 1♂, 1♀; Ardanuç, Akarsu, 900 m, 7.VII.1994, 1♂, 1♀, Cehennem Deresi, 600 m, 13.V.2000, 2♂, Ferhatlı, 550 m, 13.V.2000, 7♂, 7♀, 7.VII.1994, 1♂, Zeytinlik, 22.V.2002, 500 m, 1♂; Borçka, Karagöl, 1050 m, 8.VII.1994, 1♂; İshان, 600 m, 16.VI.2001, 1♂; Yusufeli, Altıparmak, 1100 m, 12.VI.1997, 1♀, 5.VII.1994, 1♀, Civnar, 15.VII.1992, 1♀, Demirkent, 450 m, 6.VII.1994, 1♀, Sarigöl, 700 m, 5.VII.1994, 8♀, Şavşat, Bahçeli, 500 m, 22.V.1997, 1♂, Oruçlu, 21.V.1997, 1♂; Bayburt: Maden, 1650 m, 16.VI.2000, 1♂; Bingöl: 1150 m, 30.V.2002, 3♀, 7♂. Diyarbakır: 700 m, 1.VI.2002, 1♀. Erzincan: Üzümlü, 1300 m, 16.VI.1994, 1♀. Erzurum: Abdurrahmangazi, 2200 m, 14.V.2000, 2♀; Horasan, 1700 m, 4.VI.2000, 1♀; İlica, Atlıkonak, 2000 m, 29.VI.1999, 1♀; İspir, Madenköprübaşı, 1300 m, 16.VI.1994, 1♂, 1♀; Oltu, 1400 m, 11.VII.1991, 2♂, 1♀, Kaleboğazı, 1450 m, 28.VII.1990, 2♂, Sarısaz, 1450 m, 21.VI.2000, 3♀, Sütkans, 1500 m, 25.VI.1996, 1♀, Taşlıköy, 12.VI.1991, 2♂; Olur, Coşkunlar, 12.VII.1991, 8♂; Şenkaya, İçmesu, 1800 m, 24.VI.1999, 1♀, Timurkişla, 24.VI.1999, 1♀, Turnalı, 1750 m, 12.VI.1992, 1♀; Tortum, Aksukapı, 25.VI.2002, 1400 m, 2♂, Çakılı, 1.VII.1991, 1♂, Pehlivanalı, 1400 m, 12.VII.1992, 1♀; Uzungere, 4.VI.1989, 2♀, Dikyar, 1100 m, 31.VII.1993, 1♂, Çamlıyamaç, 1200 m, 21.VI.1998, 3♂, 2♀, Şelale, 1000 m, 9.VI.1996, 1♀. Gümüşhane: 15.VI.1992, 1♂, 1♀. İzmir: Efes, 5.V.1998, 2♀. Kars: Sarıkamış, Karakurt, Aras Vadisi, 1500 m, 27.V.2000, 1♂, 11.VI.1998, 3♀, 23.VI.2000, 4♂, 5♀, 18.VII.2002, 1♂, Şeytangeçmez, 1500 m, 2.VI.1999, 1♂, TCK Çeşmesi, 1700 m, 4.VI.2000, 1♂, 3♀. Ordu: 28.VI.1991, 5♀. Osmaniye: Zorkun Yaylası, 1500 m, 24.VI.1990, 1♀; Kadirli, Karatepe, 500 m, 14.V.2002, 2♀. Rize: Çamlıhemşin, Ayder, 1550 m, 30.VII.2000, 1♀.

Distributed in Turkey: Afyon, Amasya, Ankara, Antalya, Artvin, Aydın, Balıkesir, Bolu, Gümüşhane, Erzurum, Mersin, Muğla, Samsun and Sinop (Benson, 1968; Özeren, 1970; Chevin and Chenon, 1982).

General Distribution and Chorotype: Russia, Crimea, Caucasus, Turkey, Balkans, Hungary, Italy (Medvedev, 1994; Liston, 1995; Lacourt, 1999); Asiatic-European.

Host Plant: Adults were collected from flowers of *Fragaria* sp. and *Rubus* sp.

Macrophya rufipes Linne, 1758

Synonym: *Tenthredo pavida* Fabricius, 1775; *T. dumetorum* Geoffroy, 1785; *T. strigosa* Fabricius, 1798; *T. (Allantus) citreipes* Audinet-Serville, 1823.

Material Examined: Artvin: Ardanuç, Ferhatlı, 450 m, 13.V.2000, 1♂, İshان, 600 m, 16.VI.2001, 1♂, 2♀. Bayburt: Çalidere, 1850 m, 16.VI.2000, 2♀, Kopdağı, 2100 m, 16.VI.2000, 1♀. Bingöl: Çobantaşı, 1400 m, 2.VI.2001, 1♂. Erzurum: Oltu, 1400 m, 25.VI.1971, 1♂, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 1♂, 1♀; Tortum, Çataldere, 1950-2500 m, 12.VII.2001, 1♀, Şenyurt, 1500 m, 27.VI.1996, 1♀, Yukarımeydanlar, 3.VII.1991, 1♀. Gümüşhane: Vaukgeçidi, 1850 m, 21.VI.2001, 1♂. Kars: Sarıkamış, Karakurt, TCK Çeşmesi, 1500 m, 4.VI.2000, 2♀.

Distributed in Turkey: Balıkesir (Chevin and Chenon, 1982).

General Distribution and Chorotype: Europe, Turkey, Russia, Urals, Crimea, Caucasus, Central Asia (Medvedev, 1994; Liston, 1995; Lacourt, 1999); Asiatic-European.

Host Plant: *Agrimonia eupatoria*, *Vitis vinifera* (Liston, 1995; Taeger and Blank, 1998).

Macrophya superba Tischbein, 1852

Synonym: *M. flavipennis* Kriechbaumer, 1891; *Tenthredo erythropus* Brullé, 1832.

Material Examined: Artvin: Ardanuç, Ferhatlı, 450 m, 13.V.2000, 1♀. Bayburt: 25 km SE, 21.VI.2001, 1700 m, 1♀, Çalidere, 1850 m, 16.VI.2000, 4♂, 4♀; Kopdağı, 2100-2300 m, 16.VI.2000, 2♂, 2♀, 21.VI.2001, 1♀, Maden, 1650 m, 16.VI.2000, 13♂, 5♀, 1700 m, 21.VI.2001, 2♀. Erzincan: 1350 m, Avcılar, 6.VI.1999, 1♀, Bahçeli, 1200 m, 26.V.2001, 1♂. Erzurum: It is common and abundant in Erzurum province. The materials were collected from the middle of May to the end of July in a University field, Yeşildere, Çat (Çimis), İlica (Atlıkonak), İspir (Madenköprübaşı), Oltu (Başaklı, Çamlıbel, Sarısaz, Subatık, Sütkans, Tutmaç), Pasinler, Pazaryolu (Karataş), Şenkaya (İçmesu, Turnalı), Tortum (Aksuyayla, Çakılı, Çataldere, Doruklu, and Kaledibi, Pehlivanalı) at altitudes of 1300-2300 m, 38♂, 49♀. Gümüşhane: Torul, Zigana, 1600 m, 14.VI.1990, 1♂; Vauk Geçidi, 1850 m, 24.VI.2001, 6♂, 6♀. Kars: Sarıkamış, Karaurgan, 2000 m, 24.VI.1999, 1♀.

Distributed in Turkey: Afyon, Amasya, Ankara, Artvin, Balıkesir, Çankırı, Erzurum, Gümüşhane, İzmit, Manisa and Niğde (Benson, 1968; Chevin and Chenon, 1982).

General Distribution and Chorotype: Italy, Balkans, Turkey, Caucasus (Benson, 1968; Liston, 1995; Lacourt, 1999); Turano-European.

Host Plant: Adults were collected from *Fragaria* sp., *Rubus* sp. and *Euphorbia* spp.

***Macrophya teutonia* Panzer, 1799**

Synonym: *M. marginata* Mocsary, 1881; *Tenthredo aureatensis* Schrank, 1802.

Material Examined: **Erzincan**: Avcılar, 1300 m, 26.V.2001, 1♀. **Erzurum**: University field, 1850 m, 3.VI.2000, 1♀, Palandöken, 2400 m, 12.VI.1994, 1♀, 1.VII.1996, 1♀; Çat, Çıraklı Geçidi, 2000-2100 m, 12.VI.2001, 1♀; Oltu, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 2♂. **Kars**: Sarıkamış, Karakurt, TCK Çeşmesi, 1500 m, 12.VI.2002. 1♀.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Europe, Russia, Transcaucasia (Liston, 1995; Lacourt, 1999); Sibero-European.

Host Plant: Several adults were collected on the flowers of *Euphorbia* sp.; *Euphorbia cyparissias* (Liston, 1995; Taeger and Blank, 1998).

***Pachyprotasis* Hartig, 1837**

Synonym: *Lithracia* Cameron, 1902

***Pachyprotasis rapae* Linne, 1767**

Synonym: *Tenthredo scripta* Gmelin, 1790; *T. luctuosa* Schrank, 1802; *T. floricola* Gravenhorst, 1807; *T. nigrofasciata* Eschscholtz, 1822; *Macrophya (Pachyprotasis) omega* Norton, 1867; *Rhogogaster sayi* Rohwer, 1908.

Material Examined: **Artvin**: Şavşat, 1650 m, 8.VII.1998, 1♀. **Erzurum**: Şenkaya, Turnalı, 1750 m, 1.VII.1990, 1♀. **Rize**: Çamlıhemşin, Ayder, 1200-1550 m, 30.VII.2000, 2♀. **Trabzon**: Hamsiköy, 1700 m, 20.VI.1999, 1♀.

Distributed in Turkey: Giresun and Trabzon (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia, Pacific Coastal Region, Siberia, Mongolia, India, China, Japan, North America, (Benson, 1968; Liston, 1995; Lacourt, 1999); Holarctic.

Host Plant: *Solanum tuberosum*, *Pedicularis palustris*, *Angelica sylvestris*, *Veronica beccabunga*, *Betonica*

officinalis, *Scrophularia* sp., *Solidago* sp., *Verbascum* sp., *Corylus avellana*, *Origanum* sp., *Fraxinus excelsior*, *Tussilago farfara*, *Symphoricarpos albus*, *Atropa* sp., *Salix caprea*, *Sarothamnus* sp., *Senecio* spp., *Polygonum* sp., *Aspidium* sp., *Epilobium* spp., *Hypericum* spp., *Galeopsis* spp., *Mentha* sp., *Polystichum* spp., *Plantago* spp., *Quercus* spp., *Stachys* spp. (Liston, 1995; Taeger and Blank, 1998).

***Rhogogaster* Konow, 1884**

Synonym: *Rhogogastera* Konow, 1885

***Rhogogaster chlorosoma* Benson, 1943**

Synonym: *R. scalaris* Klug, 1817.

Material Examined: **Artvin**: Şavşat, Karagöl, 1650 m, 8.VII.1998, 1♀. **Bayburt**: Çalıdere, 1850 m, 16.VI.2000, 3♀, Kopdağı 2100 m, 16.VI.2000, 2♂. **Bingöl**: Çıraklı Geçidi, 2000 m, 11.VII.2002, 1♀. **Erzurum**: Aribahçe, 2400 m, 20.VII.2001, 1♀, Dutçu, Komyolu, 2200 m, 18.VII.1999, 1♀; İlica, Atlikonak, 2000 m, 29.VI.1999, 1♂; İspir, Madenköprübaşı, 1400 m, 7.VII.1996, 1♀; Köprüköy, Örenteş, 1900 m, 12.VI.2001, 1♂; Oltu, Çamlıbel, 1750 m, 2.VII.1997, 1♀, 14.VII.1996, 1♀, 22.VII.2001, 1♀, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 1♀; Pasinler, Çalıyazı, 2400 m, 10.VII.1997, 1♀, Pelitli, 2000 m, 14.VII.1998, 1♀; Tortum, 1500 m, 4.VII.1994, 1♂, Kireçli Dağı 2200 m, 2.VII.2000, 1♀, Uzunkavak, 1800 m, 12.VII.2001, 2♀. **Gümüşhane**: Vauk Geçidi, 1850 m, 21.VI.2001, 2♀. **Rize**: Çamlıhemşin, Ayder, 1550 m, 30.VII.2000, 3♀.

Distributed in Turkey: Ankara, Erzurum, Giresun, İstanbul, Konya and Tokat (Benson, 1968).

General Distribution and Chorotype: Turkey, Caucasus, Mongolia, Siberia (Benson, 1968; Medvedev, 1994; Lacourt, 1999); Sibero-European.

Host Plant: Larvae feed on *Salix* spp., adults were collected on the same plants. It can be considered a minor pest of *Salix* spp. Taeger and Blank (1998) and Lacourt (1999) listed the following plant species as hosts: *Pteridium aquilinum*, *Alnus glutinosa*, *Circaeaa*, *Prunus* spp., *Ranunculus* spp., *Rosa* spp., *Salix alba*, *S. purpurea*, *Stellaria* spp., *Filipendula ulmaria*, *Populus tremula*, *Padus* spp., *Betula* spp., *Corylus avellana* and *Sorbus* spp.

***Sciapteryx* Stephens, 1835**

Synonym: *Eniscia* Thomson, 1870; *Sciapterina* Ushinskij, 1940.

Sciapteryx circassica Dovnar-Zapolski, 1930

Material Examined: Erzurum: Şenkaya, Ormanlı, 1800 m, 20.V.1994, 4♀.

Distributed in Turkey: Gümüşhane and Trabzon (Benson, 1968).

General Distribution and Chorotype: Russia, Turkey, Transcaucasia (Benson, 1968; Liston, 1995; Zhelochovtsev and Zinovjev, 1996; Lacourt, 1999); Sibero-European.

Host Plant: Unknown.

Sciapteryx laeta Konow, 1891

Synonym: *S. laeta antennata* Muche, 1973

Material Examined: Erzurum: Pazaryolu, Karataş, 2000 m, 5.VII.2000, 3♂, 4♀.

Distributed in Turkey: Gümüşhane (Benson, 1968).

General Distribution and Chorotype: Turkey, Transcaucasia, Georgia (Benson, 1968; Zhelochovtsev and Zinovjev, 1996; Lacourt, 1999); Turanian.

Host Plant: *Ranunculus* spp. (Lacourt, 1999).

Remarks: Present data show that its westernmost record is Erzurum and Gümüşhane.

3.7. *Tenthredo* Linne, 1758

Synonym: *Parastatis* Kirby, 1881; *Labidia* Provancher, 1886; *Rethrax* Cameron, 1899; *Fethalia* Cameron, 1902; *Clydostomus* Konow, 1908; *Tenthredina* Rohwer, 1910; *Zamacrophya* Rohwer, 1912.

Tenthredo acerrima Benson, 1952

Synonym: *T. nitidior* (Konow, 1888); *T. marginella* Panzer, 1804, nec Fabricius, 1793; *T. aegra* Enslin, 1912; *T. sulphuripes* (Kriechbaumer, 1869); *Allantus arcuatus* var. *atricereus* Zirngiebl, 1937.

Material Examined: Ardahan: Posof, Yurtbekler, 1600 m, 27.VII.1991, 1♂. Artvin: Şavşat, Karagöl, 1650 m, 8.VII.1998, 2♂. Erzurum: Pazaryolu, Karataş, 2200 m, 5.VII.2000, 1♂. Rize: Çamlıhemşin, Ayder, 1200-1550 m, 30.VII.2000, 1♂, 1♀, İkizdere, Çamlık, 1600 m, 29.VII.2000, 4♂, 1♀, Ovit Dağı 1600-2400 m, 29.VII.2000, 8♂, 7♀.

Distributed in Turkey: Artvin, Gümüşhane and Trabzon, (Benson, 1968).

General Distribution and Chorotype: Europe, Russia, Turkey, Caucasus, Transcaucasia (Benson, 1968;

Cingovski, 1982; Liston, 1995; Lacourt, 1999); Sibero-European.

Host Plant: *Lotus corniculatus* (Liston, 1995, Lacourt, 1999, Taeger and Blank, 1998).

Tenthredo albiventris Mocsary, 1880

Synonym: *T. trivittata* André, 1881; *T. lederi* Konow, 1890; *T. (Allantus) albiventris hostia* Enslin, 1912; *T. pallidula* Benson, 1930; *T. kerketa* Dovnar-Zapolskij, 1930.

Material Examined: Gümüşhane: Vauk Geçidi, 1850 m, 21.VI.2001, 2♂, 1♀, 24.VI.2001, 16♂, 23♀.

Distributed in Turkey: Lacourt (1999) indicated that it occurred in Turkey, but no locality was noted.

General Distribution and Chorotype: Turkey, Transcaucasia (Lacourt, 1999); Turanian.

Host Plant: Adults were collected from flowers of *Conium* sp.

Tenthredo albopicta Puls, 1870

Synonym: *T. nigroclipeata* Mocsary, 1909

Material Examined: Artvin: Borçka, Karagöl, 1650 m, 8.VII.1994, 1♀. Trabzon: Sümelamanastırı, 2.VII.1996, 1♀.

Distributed in Turkey: Trabzon (Benson, 1968).

General Distribution and Chorotype: Turkey, Transcaucasia (Benson, 1968; Lacourt, 1999); Turanian.

Host Plant: Unknown.

Remarks: Present knowledge shows that Trabzon is the westernmost point of the distribution area of *T. albopicta* around the world.

Tenthredo arcuata Forster, 1771

Synonym: *T. labiata* Geoffroy, 1785; *T. virgata* Geoffroy, 1785; *T. viminalis* Schrank, 1796; *T. marginella* Panzer, 1804, nec Fabricius, 1793, p.p.; *Allantus melanotus* Rudow, 1872; *A. dispar* Rudow, 1872, nec Klug, 1817; *A. cipealis* Konow, 1888.

Material Examined: Artvin: Genye Dağı 1800 m, 16.VIII.1990, 2♀. Erzurum: Şenkaya, Turnalı, 1750 m, 25.VII.1996, 2♂, 6♀, 6.VIII.1990, 6♀; Tortum, Dorukluyaylaşı, 4.VII.1991, 2♀; Uzundere, Dikyar, 1200 m, 2.VII.1994, 3♀. Rize: Çamlıhemşin, Ayder, 1550 m, 30.VII.2000, 1♂, 2♀; İkizdere, Ovit dağı, 1600-2500 m, 22.VII.2002, 3♂, 6♀, 29.VII.2000, 9♂, 18♀.

Distributed in Turkey: Medvedev (1994) indicated that it was present in Turkey but no locality was noted.

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia, Caucasus, Pacific Coastal Region, Central Asia, Siberia, Mongolia, Japan (Medvedev, 1994; Liston, 1995; Lacourt, 1999); Asiatic-European.

Host Plant: Adults were collected from the flowers of *Trifolium pratense*. *T. repens* was given as host plant (Liston, 1995; Taeger and Blank, 1998; Lacourt, 1999).

***Tenthredo caligator* Eversmann, 1847**

Synonym: *T. nigritarsis* Puls, 1870; *T. morawitzi* Jakovlev, 1888.

Material Examined: Erzurum: 2400 m, 11.VII.1996, 1♀; İlçe, Rizekent, 2200 m, 30.VII.1998, 1♀; İspir, Çayırozungü, 4.VII.1997, 1♀; Oltu, Sütkans, 1500 m, 25.VI.1996, 2♀, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 5♂, 4♀; Pasinler, Çalıyazı, 2400 m, 10.VII.1997, 3♂, 4♀, 11.VII.1997, 2♀, Pelitli, 2000 m, 14.VII.1998, 2♀; Pazaryolu, Toprakkale, 2200 m, 30.VII.1998, 1♀; Şenkaya, Turnalı, 1750 m, 1.VII.1990, 1♂, 6.VII.1990, 2♀, 25.VII.1990, 1♂, 1♀, 25.VI.1991, 1♂, 10.VII.1999, 1♀; Tortum, Aşağımeydanlar, 4.VII.1991, 1♂, Kireçli Dağı 2200 m, 25.VI.1996, 3♂, 1♀, 2.VII.2000, 1♂. Rize: İkizdere, Çamlık, 1600 m, 29.VII.2000, 2♂, 4♀, Ovit Dağı 1600-2400 m, 29.VII.2000, 6♂, 4♀; Trabzon: Hamsiköy, 1600 m, 24.VI.1998, 2♂, 3♀.

Distributed in Turkey: Trabzon (Benson, 1968).

General Distribution and Chorotype: Turkey, Transcaucasia (Benson, 1968; Lacourt, 1999); Turanian.

Host Plant: Adults were collected on flowers adjacent to forest areas.

***Tenthredo caucasica* Eversmann, 1847**

Synonym: *Allantus moestus* Mocsary, 1883

Material Examined: Erzurum: İspir, Madenköprübaşı, 1400 m, 18.VI.1994, 1♂; Oltu, Sütkans, 1500 m, 18.VI.1996, 1♂; Pazaryolu, Karataş, 2000-2200 m, 5.VII.2000, 1♀; Şenkaya, Turnalı, 1750 m, 6.VII.1990, 1♂, 25.V.1994, 1♀, 25.VII.1996, 1♂, 6.VIII.1990, 1♂, 1♀; Tortum, Çataldere, 1900-2500 m, 12.VII.2001, 3♀, Doruklu, 4.VII.1991, 1♀, Uzunkavak, 1800 m, 12.VII.2001, 1♂, 4♀. Rize: İkizdere, Çamlık, 1600 m, 29.VII.2000, 2♂, 11♀, Ovit Dağı 1600-2400 m, 29.VII.2000, 7♀; Çamlıhemşin, Ayder, 1200-1550 m, 30.VII.2000, 1♀. Trabzon: 10.IX.1992, 1♂.

Distributed in Turkey: Erzurum, Gümüşhane, Isparta and Trabzon (Benson, 1968; Chevin and Chenon, 1982).

General Distribution and Chorotype: Europe, Turkey, Transcaucasia (Benson, 1968; Cingovski, 1980; Medvedev, 1994; Lacourt, 1999); Turano-European.

Host Plant: Adults were collected from the flowers of *Euphorbia* spp.

***Tenthredo colon* Klug, 1817**

Synonym: *T. obscura* Panzer, 1805, nec Gmelin, 1790; *T. rudowi* André, 1881; *T. nigricollis* Kirby, 1882; *T. jocosa* Provancher, 1882; *T. punctulata* Konow, 1887; *T. livida* Linne sensu Ross, 1951; *Tenthredella colon* var. *nigriventris* Enslin, 1912.

Material Examined: Artvin: Yusufeli, Sarıgöl, 700 m, 5.VII.1994, 1♂.

Distributed in Turkey: Kütahya and Trabzon (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia, Caucasus, North Asia, Central Asia, Mongolia, Siberia, Korea, Japan, North America, (Benson, 1968; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Holarctic.

Host Plant: *Epilobium* sp., *Fuchsia* spp., *Circeae lutetiana*, *Pteridium aquilinum*, *Aconitum* spp., *Dasiphora* sp., *Geranium* spp., *Potentilla* spp., *Ranunculus borealis*, *Sanguisorba* spp., *Salix* spp. (Liston, 1995; Taeger and Blank, 1998; Lacourt, 1999).

***Tenthredo costata* Klug, 1817**

Synonym: *T. sareptana* Eversmann, 1847; *Allantus subcostatus* Jakovlev, 1888; *A. graecus* Konow, 1888; *A. parnasius* Konow, 1888; *A. violaceipennis* Costa, 1890.

Material Examined: Artvin: Ardanuç, Akarsu, 600 m, 7.VII.1994, 1♂, 1♀, Cehennemderesi, 500 m, 13.V.2000, 1♂. Bayburt: Maden, 1650 m, 16.VI.2000, 5♂, 2♀, Kopdağı, 2100-2300 m, 16.VI.2000, 3♂, 2♀, 21.VI.2001, 1♂. Bingöl: Çirişli Geçidi, 2000 m, 11.VII.2002, 1♀; Çobantaşı, 1400 m, 2.VI.2001, 1♀. Bitlis: Adilcevaz, Yıldız, 15.VII.1999, 1♂, 1♀. Elazığ: Keban, 900 m, 16.V.2002, 3♂, 2♀. Erzincan: 1350 m, 5.VI.1992, 1♀, Dumanlı, 1200 m, 25.V.2001, 1♀; Avcılar, 1300 m, 26.V.2001, 5♂, 5♀; Üzümlü, 1300 m, 16.VI.1994, 1♀, 25.V.2001, 2♂, 5♀; Kemah, 1200 m, 25.V.2001, 1♀; Bahçeli, 1350 m, 26.V.2001, 3♀; Sansa Boğazı, 1350 m, 25.V.2001, 3♂, 6♀. Erzurum:

It is common and abundant in Erzurum province. The materials were collected from the middle of May to the middle of August in Abdurrahman Gazi, Dumlu, Köşk, Nenehatun, Palandöken, University field, Yeşildere, Aşkale (Kopdağı), Horasan, İlica (Sorkunlu), İspir (Madenköprübaşı), Olur (Salaçur, Sungübeyir), Oltu (Başaklı, Çamlıbel, Sarısaç, Subatık, Sütkans, Tutmaç), Pasinler (Pelitli), Pazaryolu, Şenkaya (İçmesu, Turnalı), Tortum (Aşağımeydanlar, Çataldere, Doruklu, Pehlivanlı, Tortumkale), and Uzundere (Çamlıyamaç, Dikyar, Şelale) at altitudes of 1300-2500 m, 91♂, 98♀. Gümüşhane: Vauk Geçidi, 1850 m, 24.VI.2001, 3♂, 7♀. Konya: Güneysünir, Güragaç, 1050 m, 2.VI.2002, 3♂, 6♀. Kars: Sarıkamış, İsisu, 3.VII.1991, 1♂, 1♀, 7.VII.1991, 1♂, Karakurt, 1550 m, 2.VI.1999, 1♂, 3♀, 1700 m, 4.VI.2000, 1♂, 1♀, 23.VI.2000, 1♀.

Distributed in Turkey: Amasya, Ankara, Artvin, Erzurum, Giresun, Niğde and Sivas (Benson, 1968).

General Distribution and Chorotype: Turkey, Russia, Caucasus, Transcaucasia, Iran, Austria, Hungary, Italy, Balkans (Benson, 1968; Cingovski, 1980; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Turano-European.

Host Plant: Adults were collected from flowers of *Euphorbia* spp. Lacourt (1999) gave *Chondrilla junce* and *C. ramosissima* as larval host plants.

Tenthredo discophora Konow, 1896

Synonym: *T. discophora* var. *pullata* Konow, 1898; *T. bigemmis* Konow, 1903.

Material Examined: Gümüşhane: Torul, Zigana Dağı 1900 m, 14.VI.1990, 2♀.

Distributed in Turkey: Bolu, Gümüşhane, Rize and Trabzon (Benson, 1968).

General Distribution and Chorotype: Turkey, Transcaucasia (Benson, 1968; Lacourt, 1999); Turanian.

Host Plant: Unknown.

Tenthredo excellens Konow, 1886

Synonym: *T. annulata* Klug, 1817, nec Geoffroy, 1785; *T. persa* var. *caja* Enslin, 1912; *Allantus persa* Konow, 1888; *A. persa* var. *mandibularis* Enslin, 1910.

Material Examined: Bayburt: Maden, 1650 m, 16.VI.2000, 4♂, 3♀. Erzincan: 1250 m, 25.VI.1992, 1♀, 14.VI.1994, 1♀, 23.VI.1994, 1♀; Avcılar, 1300 m, 26.V.2001, 2♀, Bahçeli, 1350 m, 26.V.2001, 1♀, Sansa Boğazı, 1350 m, 25.V.2001, 1♀, Cevizli, 1300 m,

4.VI.1990, 1♂. Erzurum: It is common and abundant in Erzurum province. The materials were collected from the middle of May to the beginning of August in Gündömez, Köşk, Palandöken, University field, Aşkale (Kopdağı), Horasan (Karaçuha), İlica (Atlikonak), İspir (Madenköprübaşı, Gölyurt geçidi), Köprüköy (Örenteş), Olur (Sungübeyir), Oltu (Başaklı, Çamlıbel, Sarısaç, Subatık, Sütkans, Tutmaç), Pasinler, Şenkaya (Turnalı), Tortum (Aksuyayla, Çakılı, Doruklu, Pehlivanlı, Uzunkavak), and Uzundere (Dikyar, Şelale) at altitudes of 1000-2500 m, 28♂, 74♀. Kars: Digor, Kocaköy, 22.VIII.1997, 1♂, Sarıkamış, Karakurt, Aras Vadisi, 1600 m, 23.V.2001, 1♀, 25.VIII.1997, 1♂. Sivas: University field, 1350 m, 7.VI.1999, 1♂, 1♀.

Distributed in Turkey: Amasya, Ankara, Balıkesir, Erzurum, Kayseri, Mersin and Sivas (Benson, 1968; Chevin and Chenon, 1982).

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia, Caucasus, Central Asia, Iran (Benson, 1968; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Asiatic-European.

Host Plant: Unknown.

Tenthredo flavipennis (Brullé, 1832)

Synonym: *Allantus frivaldszkyi* Mocsary, 1879; *A. laetus* Konow, 1891.

Material Examined: Adiyaman: Nemrut Mountain, 1950 m, 29.V.1999, 1♀. Bayburt: Maden, 1650 m, 16.VI.2000, 1♀; Kopdağı 2100-2300 m, 16.VI.2000, 2♀, Çalidere, 1850 m, 16.VI.2000, 1♀. Erzincan: Avcılar, 1300 m, 26.V.2001, 1♂; Cevizli, 1300 m, 4.VI.1990, 1♂, 1♂, Sansa Boğazı, 1350 m, 25.V.2001, 1♂. Erzurum: Köşk, 1850 m, 20.VI.1996, 2♂, 1♀, Yeşildere, 1850 m, 25.VI.2002, 1♂; İlica, Atlikonak, 2000-2400 m, 8.VI.2001, 3♂, 8♀, 29.VI.1999, 1♂, 1♀; Köprüköy, Örenteş, 1900-2100 m, 12.VI.2001, 2♂, 2♀; Tortum, Aktaş, 1700 m, 25.V.1994, 1♀. Kars: Karakurt, Aras Vadisi, TCK Çeşmesi, 1650 m, 23.V.2001, 1♀.

Distributed in Turkey: Erzurum and Gümüşhane (Benson, 1968).

General Distribution and Chorotype: Balkans, Turkey, Transcaucasia (Benson, 1968; Liston, 1995; Lacourt, 1999); Central Asiatic-European.

Host Plant: Some adults collected from *Euphorbia* spp. It was observed that 1 female was feeding on the body fluid of a cantarid beetle.

Tenthredo frauendorfii Giraud, 1857

Synonym: *Allantus helveticus* Konow, 1898; *A. merceti* Konow; *A. montanus* Enslin (Benson, 1968); *Paratenthredo dilleri* Taeger, 1991.

Material Examined: Artvin: Şavşat, Bahçeli, 500 m, 22.V.1997, 1♀, Karagöl, 1650 m, 22.V.1997, 1♀. Erzurum: Oltu, Tutmaç-Başaklı, 2000 m, 2.VII.2000, 1♀. Gümüşhane: Vaukçeşidi, 1850 m, 21.VI.2001, 3♂, 3♀, 24.VI.2001, 7♂, 5♀. Rize: İkizdere, Çamlık, 1600 m, 29.VII.2000, 1♀.

Distributed in Turkey: Gümüşhane and Samsun (Benson, 1968).

General Distribution and Chorotype: Turkey, Iran; Switzerland, France, Italy, Yugoslavia (Liston, 1995); Central-Asiatic-European.

Host Plant: Some adults were collected from *Conium* spp. Lacourt (1999) gave *Paeonia officinalis* as a larval host of *T. frauendorfii*.

Tenthredo hyrcana Benson, 1968

Material Examined: Artvin: Borçka, Karagöl, 1050 m, 8.VII.1994, 1♂. Erzincan: Sansa, 1350 m, 25.V.2000, 1♀. Erzurum: İspir, 1500 m, 17.VII.1992, 1♂; Şenkaya, Turnalı, 1750 m, 25.V.1994, 1♂; Tortum, Kireçlidağı, 2200 m, 2.VII.2000, 1♂.

Distributed in Turkey: Amasya, Ankara and Erzurum (Benson, 1968).

General Distribution and Chorotype: Armenia, Turkey (Benson, 1968); Turanian.

Host Plant: Unknown.

Tenthredo litorata (Mocsary, 1886)

Synonym: *T. reitteri* var. *egregia* Enslin, 1914; *Allantus reitteri* Konow, 1886.

Material Examined: Artvin: Borçka, Karagöl, 1650 m, 8.VII.1994, 2♂. Erzurum: Şenkaya, Ormanlı, 1800 m, 20.V.1994, 1♂, Turnalı, 1750 m, 25.VI.1991, 1♂. Rize: İkizdere, Ovit Dağı 2400 m, 29.VII.2000, 1♂.

Distributed in Turkey: Çankırı, Gümüşhane, Kütahya and Trabzon (Benson, 1968).

General Distribution and Chorotype: Turkey, Transcaucasia (Benson, 1968; Lacourt, 1999); Turanian.

Host Plant: Unknown.

Tenthredo livida Linne, 1758

Synonym: *T. dubia* Ström, 1768; *T. lurida* Müler, 1775; *T. pellucida* Müler, 1776; *T. bipunctata* Müler, 1776; *T. annularis* Schrank, 1776; *T. albicornis* Geoffroy, 1785; *T. maura* Fabricius, 1793; *T. carpini* Panzer, 1799; *Allantus aterrimus* Stephens, 1835.

Material Examined: Bayburt: Kopdağı, 2100 m, 16.VI.2000, 1♀. Erzurum: Köşk, 1850 m, 1.VI.1995, 2♀; Tortum, Kireçlidağı, 2200 m, 2.VII.2000, 1♀, Uzunkavak, 1800 m, 12.VII.2001, 1♀. Rize: İkizdere, Çamlık, 1600 m, 29.VII.2000, 5♂, 6♀, Ovit Dağı 1600-2500 m, 22.VII.2002, 1♀, 29.VII.2000, 1♂, 13♀.

Distributed in Turkey: Zonguldak (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Transcaucasia, Siberia, Japan (Benson, 1968; Liston, 1995; Lacourt, 1999); Asiatic-European.

Host Plant: *Epilobium* spp., *Rosa* spp., *Sorbus aucuparia*, *Salix* spp., *Corylus avellana*, *Corpinus betulus*, *Plantago* spp., *Fraxinus excelsior*, *Viburnum album*, *Lonicera* spp., *Symporicarpus alba*, *Arctium* spp., *Lamium album*, *Melissa officinalis*, *Pteridium aquilinum*, *Athyrium filix-femina* (Taeger and Blank, 1998; Lacourt, 1999).

Tenthredo luteipennis Eversmann, 1847

Synonym: *T. fallax* Mocsary, 1880 nec Smith, 1878, nec Audinet-Serville, 1823; *T. mocsaryi* André, 1882; *T. menetriesi* Kirby, 1882; *Tenthredella luteipennis* var. *obscuraria* Enslin, 1913.

Material Examined: Erzurum: Oltu, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 1♂; Şenkaya, Turnalı, 1750 m, 25.VII.1996, 2♀, 6.VIII.1990, 1♀; Tortum, Uzunkavak, 1800 m, 12.VII.2001, 1♂, 1♀. Rize: İkizdere, Çamlık, 1600 m, 29.VII.2000, 2♀, Ovit Dağı 1600-2400 m, 29.VII.2000, 1♀.

Distributed in Turkey: Trabzon (Benson, 1968).

General Distribution and Chorotype: Turkey, Caucasus, Pamirs (Benson, 1968; Lacourt, 1999); Turanian.

Host Plant: Unknown.

Tenthredo maculata Geoffroy, 1785

Synonym: *T. unifasciata* Geoffroy, 1785; *T. zonta* Christ, 1791; *T. equestris* Panzer, 1809; *T. (Allantus) latizona* Audinet-Serville, 1823; *T. Andréi* Mocsary, 1883; *Tenthredella maculata* var. *atricollaris* Enslin, 1912.

Material Examined: Erzurum: Oltu, Sütkans, 1500 m, 25.VI.1996, 1♀. Gümüşhane: Torul, Zigana Dağı 1900 m, 14.VI.1990, 1♀. Trabzon: Hamsiköy, 1600 m, 24.VI.1998, 1♂, 4♀.

Distributed in Turkey: New record for Turkish fauna. It is also new for the Asian continent.

General Distribution and Chorotype: Europe (Liston, 1995; Lacourt, 1999); Turano-European.

Host Plant: *Brachypodium* spp., *Dactylis* spp. (Liston, 1995).

Tenthredo mesomelas Linne, 1758

Synonym: *T. viridis* Klug, 1817, nec Linne, 1758; *T. bernardi* Konow, 1888; *T. mesomelaena* Konow, 1905; *T. m. montana* Pasteels, 1946; *Tenthredella mesomelas* var. *mesopleuralis* Enslin, 1912; *T. m.* var. *atramentaria* Enslin, 1912.

Material Examined: Erzurum: Şenkaya, Turnalı, 1750 m, 5.IX.1990, 1♀; Oltu, Sütkans, 1500 m, 25.VI.1996, 1♀, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 1♂; Tortum, Kireçlıdağı, 2200 m, 2.VII.2000, 2♀, Uzunkavak, 1800 m, 12.VII.2001, 1♀. Gümüşhane: Torul, Zigana Dağı 1900 m, 14.VI.1990, 1♂, 1♀; Vaukgeçidi, 1850 m, 24.VI.2001, 1♂. Rize: İkizdere, Ovit Dağı 2500 m, 22.VII.2002, 1♂, 1♀. Trabzon: Sumelamanastırı, 2.VII.1996, 1♂.

Distributed in Turkey: Bolu, Gümüşhane, Trabzon and Zonguldak (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia, Siberia, Pacific Coastal Region, Mongolia, China, Japan, Tunisia (Benson, 1968; Cingovski, 1980; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Palearctic.

Host Plant: Larvae were collected on *Arctium* sp. and *Rumex* sp. Liston (1995) and Lacourt (1999) listed the following plant species as host plants: *Polygonum persicaria*, *Arctium lappa*, *Heracleum* spp., *Ranunculus* spp., *Epilobium* spp., *Rumex* spp., *Salix* spp., *Veronica* sp., *Tussilago* spp., *Petasites* sp., *Senecio* sp., *Solidago* sp., *Stachys* spp.

Tenthredo notha Klug, 1817

Synonym: *Allantus perkinsi* Morice, 1919; *A. arcuatus* var. *niger* Zirngiebl, 1937.

Material Examined: Erzurum: Güngörmez, 2400 m,

5.VIII.1990, 1♀; Pasinler, Çalıyazı, 2400 m, 10.VII.1997, 1♀.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Europe, Russia, North Africa, Transcaucasia, Iran (Liston, 1995; Lacourt, 1999); Palearctic.

Host Plant: *Trifolium repens*, *T. pratense*, *Vicia cracca* (Liston, 1995; Lacourt, 1999).

Tenthredo omissa Förster, 1844

Synonym: *T. vienensis* Schrank, sensu Panzer, 1799; *T. marginella* Fabricius sensu Thomson, 1871; *T. omissa* var. *melanoceraea* Enslin, 1912.

Material Examined: Erzurum: Konaklı, 2000-2400 m, 22.VII.2000, 2♂, 2♀, Palandöken, 2400 m, 1.VII.1996, 1♀; İlica, Sorkunlu, 1900 m, 10.VIII.1999, 1♀; Oltu, Tutmaç-Başaklı, 1700-2000 m, 31.VII.2001, 1♀.

Distributed in Turkey: Bolu (Wolf, 1968).

General Distribution and Chorotype: Turkey, East Europe, East Siberia, Transcaucasia (Wolf, 1968; Medvedev, 1994; Lacourt, 1999); Sibero-European.

Host Plant: *Plantago media*, *P. lanceolata* (Lacourt, 1999).

Tenthredo propinqua Klug, 1817

Material Examined: Artvin: Borçka, Karagöl, 1600 m, 8.VII.1994, 1♀; Genye Dağı 1600 m, 16.VIII.1990, 1♀; Yusufeli, Kınalıçam, 400 m, 21.V.2002, 1♀. Erzurum: Aşkale, Kopdağı, 2000 m, 16.VI.2000, 2♂; İspir, 17.VII.1992, 1♀, Fisırıkkalesi, 2300 m, 12.VII.2001, 1♀; Olur, Salaçur, 2400 m, 10.VII.2001, 1♀; Oltu, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 3♀; Pazaryolu, 2200 m, 5.VII.2000, 1♀; Şenkaya, Turnalı, 1750 m, 1.VII.1990, 1♂, 6.VII.1990, 1♀; Tortum, Çataldere, 1950-2500 m, 12.VII.2001, 2♂, 2♀; Uzundere, Dikyar, 1100 m, 24.V.1994, 1♀. Rize: İkizdere, Ovit Dağı 1600-2400 m, 29.VII.2000, 1♀.

Distributed in Turkey: Kastamonu-Çankırı border (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Ukraine, Transcaucasia (Benson, 1968; Zhelochovtsev and Zinovjev, 1996; Lacourt, 1999); Sibero-European.

Host Plant: Unknown.

Tenthredo scrophulariae Linne, 1758

Material Examined: Artvin: Genye Dağı 1600 m, 16.VIII.1990, 1♀; Bayburt: Kopdağı 2300 m, 25.VII.2001, 1♀. Erzincan: Çatalarmut, 9.VIII.1990, 1♀. Erzurum: Arıbahçe, 2400 m, 20.VII.2001, 2♂, 3♀; Güngörmez, 2400 m, 5.VIII.1990, 7♀, 2500 m, 28.VII.1998, 1♀; Kargapazarı Dağları, Radar yolu, 2200 m, 11.VIII.1990, 1♀; Dutçu, 2200 m, 18.VII.1999, 3♀; Kö_k, 1850 m, 20.VI.1996, 1♂; İspir, Madenköprübaşı, 1400 m, 7.VII.1996, 1♂; Şenkaya, 2400 m, 9.VII.1992, 2♂, Turnalı, 1750 m, 1.VII.1990, 2♀, 25.VI.1991, 1♂, 2♀. Rize: Çamlıhemşin, Ayder, 1200-1550 m 30.VII.2000, 1♀; İkizdere, Çamlık, 1600 m, 29.VII.2000, 1♂, Ovit Dağı 1600-2400 m, 29.VII.2000, 1♀.

Distributed in Turkey: Sivas and Tokat (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia, Caucasus (Benson, 1968; Liston, 1995; Lacourt, 1999); Sibero-European.

Host Plant: *Scrophularia* spp., *Verbascum* spp. (Liston, 1995, Lacourt, 1999).

Tenthredo thompsoni Curtis, 1839

Synonym: *T. marginella* var. *basanalis* Enslin, 1912; *T. marginella* Fabricius sensu Chevin, 1983; *Allantus semifasciatus* Rudow, 1872.

Material Examined: Erzincan: Cevizli, 1300 m, 10.VIII.1990, 1♂, 1.VIII.1991, 1♂, 1♀. Erzurum: Konaklı, 2400 m, 22.VII.2000, 3♂, 2♀, Arıbahçe, 2400 m, 20.VII.2001, 1♂; İlica, Sorkunlu, 1900 m, 10.VIII.1999, 1♂; İspir, 30.VII.1990, 2♂, Madenköprübaşı, 1400 m, 18.VI.1994, 1♂; Oltu, İnanmış, 1700 m, 26.VII.2000, 1♂, Kaleboğazı, 1400 m, 28.VII.1990, 1♂, Tutmaç, 2000 m, 3.VII.2001, 1♂; Pasinler, Çalıyazı, 2100 m, 10.VII.1997, 1♂; Şenkaya, Turnalı, 1750 m, 25.VII.1996, 1♂; Tortum, Aşağımeydanlar, 7.VIII.1996, 1♂, Pehlivانlı, 1400 m, 13.VIII.1991, 1♂, 1♀.

Distributed in Turkey: Bursa, Gümüşhane and Trabzon (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Caucasus, Kazakhstan (Medvedev, 1994; Liston, 1995; Lacourt, 1999); Asiatic-European.

Host Plant: *Pimpinella major* (Liston, 1995).

Tenthredo vespa Retzius, 1783

Synonym: *T. multifasciata* Geoffroy, 1785; *T. sexannulata* Schrank, 1802; *T. tricincta* Fabricius, 1804, nec Christ, 1791; *T. quadricincta* Fallen, 1808; *T. vespiformis* Audinet-Serville, 1823; *Allantus multicinctus* Rudow, 1872; *A. nigrilabris* Frivaldszky, 1877.

Material Examined: Erzurum: İlica, Rizekent, 2200 m, 30.VII.1998, 2♀; Pazaryolu, Toprakkale, 2200 m, 30.VII.1998, 1♂, 1♀; Uzundere, Dikyar, 1100 m, 2.VII.1994, 1♀. Rize: İkizdere, Çamlık, 1600 m, 29.VI.2000, 2♀.

Distributed in Turkey: Trabzon (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Transcaucasia, Siberia, Caucasus, China, Mongolia (Benson, 1968; Medvedev, 1994; Lacourt, 1999); Asiatic-European.

Host Plant: *Lonicera caprifolium*, *Syringa vulgaris*, *Viburnum opulus*, *Fraxinus excelsior*, *Jasminum officinale*, *Rosa*, *Spiraea*, *Acer platanoides*, *Ligustrum vulgare*, *Symporicarpos albus*, *S. alba* (Liston, 1995; Taeger and Blank, 1998; Lacourt, 1999).

Tenthredo vespiformis Schrank, 1781

Material Examined: Erzincan: Sansa Boğazı, 1350 m, 25.V.2001, 1♀. Erzurum: Göngörmez, 2400 m, 5.VIII.1990, 1♀; Horasan, Aras Vadisi, 1600 m, 23.V.2001, 1♀; Köprüköy, Örentaş, 1900-2100 m, 12.VI.2001, 1♀; Oltu, Sütkans, 1500 m, 18.VI.1996, 1♀, 25.VI.1996, 1♀; Pasinler, Çalıyazı, 2400 m, 10.VII.1997, 1♀; Pelitli, 2000 m, 14.VII.1998, 1♀; Uzundere, Dikyar, 1100 m, 24.V.1994, 1♀. Kars: Posof, 30.V.1973, 1♀; Sarıkamış, Karakurt, TCK Çeşmesi, 1500 m, 4.VI.2000, 4♀.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Europe, Russia, Ukraine (Medvedev, 1994; Liston, 1995; Zhelochovtsev and Zinovjev, 1996); Sibero-European.

Host Plant: Unknown.

Tenthredo zonula Klug, 1817

Synonym: *T. fasciata* Scopoli, 1763 nec Linne, 1758; *T. bifasciata* Geoffroy, 1785; *T. bicinctaflava* Christ, 1791; *Allantus similis* Mocsary, 1880; *A. nazareensis* André; *A. serena* Konow; *A. calcaratus* André, 1881, *A. scutellaris* Konow, 1898.

Material Examined: Adiyaman: Nemrut Mountain, 1950 m, 29.V.1999, 2♀. Artvin: Ardanuç, Ferhatlı, 450 m, 13.V.2000, 1♂; Yusufeli, Sarıgöl, 700 m, 5.VII.1994, 1♀; Şavşat, Karagöl, 1650 m, 8.VII.1998, 1♂, 6♀. Bayburt: 25 km S., 1750 m, 21.VI.2001, 3♂, 3♀; Maden, 1650 m, 16.VI.2000, 5♂, 7♀, Çalidere, 1850 m, 16.VI.2000, 9♂, 4♀, Kopdağı 2100-2300 m, 28.VI.1990, 1♀, 16.VI.2000, 3♂, 11♀, 21.VI.2001, 3♂, 4♀. Bingöl: Çıraklı Geçidi, 2000 m, 3.VI.2001, 2♂, 2♀, 11.VII.2002, 1♂, 3♀. Diyarbakır: University field, 700 m, 1.VI.2002, 3♀. Erzincan: Bahçeli, 1350 m, 26.V.2001, 2♂; Cevizli, 1350 m, 4.VI.1990, 1♂, 2♀. Erzurum: It is common and abundant in Erzurum province. The materials were collected from the middle of May to the end of July in Akdağ, Dumlu, Gökçeyamaç, Palandöken, Yeşildere, Aşkale (Aşağıkop, Mezra, Tepebaşı) Horasan, İlica (Atlıkonak, Gülyurt Geçidi), İspir (Fısırik Kalesi, Madenköprübaşı), Köprüköy (Örentaş), Olur (Süngübayı), Oltu (Başaklı, Çamlıbel, Sütkans, Tutmaç), Pasinler (Çalıyazı), Pazaryolu (Karataş), Şenkaya (Ormanlı, Turnalı), Tortum (Aktaş, Kireçli Dağı Uzunkavak), and Uzundere (Dikyar) at altitudes of 1100-2400 m, 87♂, 184♀. Gümüşhane: 25 km SE, 1700 m, 19.VI.2001, 1♀, Vauk Geçidi, 1850 m, 21.VI.2001, 3♂, 2♀, 24.VI.2001, 1♀. Hatay: Bademli, 6 km W. to Hassa, 850 m, 13.V.2002, 3♂, 1♀. Kars: Sarıkamış, Karakurt, Şeytangeçmez, 1500 m, 2.VI.1999, 1♂, 2♀, Karakurt, 1550 m, 2.VI.1999, 1♀, Aras Vadisi, Karakurt, 1550 m, 23.V.2001, 7♂, 7♀, 4.VI.2001, 1♂, 5♀. Konya: University Campus, 9.VII.1997, 1♀. Osmaniye: Bahçe, 300 m, 13.V.2002, 1♀. Şanlıurfa: K.Köprü, 30.IV.1990, 2♂, 1♀.

Distributed in Turkey: Afyon, Amasya, Ankara, Antalya, Artvin, Aydın, Bursa, Çorum, Erzincan, Erzurum, Isparta, İstanbul, İzmir, Konya, Muğla and Niğde (Benson, 1968; Chevin and Chenon, 1982).

General Distribution and Chorotype: Europe, Turkey, Russia, Ukraine, Moldova, Transcaucasia, Caucasus, Central Asia, Armenia, Iran, Israel, Syria, Lebanon (Benson, 1968; Medvedev, 1994; Liston, 1995; Zhelochovtsev and Zinovjev, 1996; Lacourt, 1999); Asiatic-European.

Host Plant: *Hypericum perforatum* (Liston, 1995; Lacourt, 1999); Asiatic-European.

Tenthredopsis Costa, 1859

Synonym: *Ebolia Costa, 1859; Thomsonia Konow, 1884; Eutenthredopsis Enslin, 1913.*

Tenthredopsis albonotata (Brullé, 1832)

Synonym: *Tenthredo propinqua* Mocsary, 1880, nec Klug, 1817.

Material Examined: Diyarbakır: Silvan, 17.IV.1995, 5♀. Elazığ Keban, 900 m, 16.V.2002, 1♀. Hatay: Antakya, Yayladağı, 25 km S., 28.IV.2002, 1♂.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Europe, East Mediterranean, Lebanon, Syria, Israel (Benson, 1968; Lacourt, 1999); Asiatic-European.

Host Plant: Unknown

Tenthredopsis festiva Konow, 1890

Material Examined: Bayburt: Çalidere, 1850 m, 16.VI.2000, 1♂. Erzurum: İlica, Atlıkonak, 2000-2400 m, 11.VI.2000, 1♀, 29.VI.1999, 1♀, 29.VI.2001, 1♀; Aras Vadisi, 1700 m, 23.V.2001, 1♂. Gümüşhane: Vauk Geçidi, 1850 m, 21.VI.2001, 1♀, 24.VI.2001, 1♂, 1♀. Kars: Aras Vadisi, TCK Çeşmesi, 1500-1600 m, 23.V.2001, 2♀.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Transcaucasia (Benson, 1968; Zhelochovtsev and Zinovjev, 1996; Lacourt, 1999); Turanian.

Its westernmost point of distribution is Turkey.

Host Plant: Adults were collected from flowers of *Corium maculatum*.

Tenthredopsis harveyi Benson, 1968

Material Examined: Erzurum: Aşkale, Pyrnakapan, 1800 m, 23.V.2000, 1♀.

Distributed in Turkey: Bolu (Benson, 1968).

General Distribution and Chorotype: Turkey (Benson, 1968; Lacourt, 1999); Endemic to Turkey.

Host Plant: Unknown.

Remarks: Benson (1968) described this species from a female (Holotype) from Bolu. This is the second specimen collected in Erzurum since the time of description. The male remains unknown. During this study after such a dense collection of the material in Erzurum province, finding only 1 specimen shows that this species is very rare.

Tenthredopsis nassata Linne, 1767

Synonym: *T. albomaculatus* Cameron, 1881; *T. dorsivittatus* Cameron, 1881; *T. lividiventris* Cameron, 1881; *Tenthredo alneti* Schrank, 1781; *T. palmata* Geoffroy, 1785; *T. perlata* Geoffroy, 1785; *T. apicaris* Geoffroy, 1785; *T. subulata* Gmelin, 1790; *T. melanorrhoea* Gmelin, 1790; *T. tiliae* Panzer, 1805; *T. korlevici* Konow, 1887; *T. arroganus* Konow, 1890; *T. fenestrata* Konow, 1890; *T. nassata* var. *rufata* Konow, 1890; *T. konowi* Strobl, 1895, nec Lethierry, 1886; *T. nigroscutellata* Konow, 1904; *T. virgineus* Muche, 1965; *Thomsonia friesei* Konow, 1884.

Material Examined: Artvin: Borçka, Karagöl, 1650 m, 8.VII.1994, 1♂. Erzurum: Oltu, Sütkans, 1500 m, 25.VI.1996, 1♀.

Distributed in Turkey: Amasya, Ankara, Çankırı and Trabzon (Benson, 1968).

General Distribution and Chorotype: Europe, Turkey, Russia, Siberia, China, Mongolia, Japan (Benson, 1968; Cingovski, 1980; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Asiatic-European.

Host Plant: *Dactylis glomerata*, *Deschampsia cespitosa*, *D. calmagrostis*, *Flexuosa* spp., *Holcus* spp., *Lolium perenne*, *Agropyron* spp., *Carex* spp., *Anthriscus silvestris*, *Artemisia* spp. (Liston, 1995; Lacourt, 1999).

Tenthredopsis nigella Konow, 1891

Material Examined: Erzurum: Pasinler, Çalıyazı, 2400 m, 10.VII.1997, 1♂.

Distributed in Turkey: Benson (1968) and Lacourt (1999) indicated the presence of this species in Turkey but no locality was given.

General Distribution and Chorotype: Turkey, Transcaucasia (Benson, 1968; Zhelochovtsev and Zinovjev, 1996; Lacourt, 1999); Turanian.

Host Plant: Unknown.

Tenthredopsis sordida Klug, 1817

Material Examined: Erzurum: Dutçu, 2000 m, 26.VI.1997, 1♂; Şenkaya, Hoş, 1600 m, 12.VI.1996, 1♂.

References

- Alkan, B. 1948. Meyva destereli arıları (Hoplocampa'lar). Mahs. Hek. 1(5): 13-16.
Altenhofer, von E. 1980. Zur biologie der in baumblättern minierenden Blattwespen (Hym., Tenthred.) Z. Ang. Ent. 89: 122-134.

Distributed in Turkey: New record for Turkish fauna.

General Distribution and Chorotype: Austria, Belgium, Switzerland, Germany, Denmark, France, Hungary, Italy, Luxembourg, Netherlands, Romania, Albania, Bulgaria (Medvedev, 1994; Liston, 1995; Lacourt, 1999); Turano-European.

Host Plant: *Arrhenatherum elatius*, *Lolium perene*, *Carex* spp., *Calamagrostis* sp., *Dactylis glomerata* (Medvedev, 1994; Taeger and Blank, 1998; Lacourt, 1999).

Remarks: Previous data relevant to the distribution of *T. sordida* showed that it has been known only from Europe so far. Our study revealed that the distribution area of this species extends to Asia, making this the first record for Asian fauna as well.

Tenthredopsis tessellata Klug, 1817

Synonym: *T. tessellata* var. *alboplagiata* Konow, 1890; *Tenthredo ischiadica* Eversmann, 1847; *Perineura cylindrica* Rudow, 1871.

Material Examined: Erzurum: Oltu, Tutmaç-Başaklı, 2000 m, 1.VII.2000, 2♀.

Distributed in Turkey: Bursa, Erzurum and Samsun (Benson, 1968).

General Distribution and Chorotype: Europe, East Mediterranean, Turkey, Russia, Caucasus, Mongolia, Siberia, Iran (Benson, 1968; Cingovski, 1980; Medvedev, 1994; Liston, 1995; Lacourt, 1999); Asiatic-European.

Host Plant: *Deschampsia*, *Dactylis*, *Aira*, *Lolium* (Liston, 1995; Lacourt, 1999).

Acknowledgments

The authors are most grateful to Dr. David R. Smith and Dr. Lajos Zombori for their in valuable help in determining the material and providing literature.

Baş, R. 1973. Türkiye'de orman ağaçlarında zarar yapan zar kanatlılar (Hymenoptera) üzerinde araştırmalar. Orm. Bak. Orm. Gn. Md. Yay. No: 570, Seri No: 23; 169 pp.

- Benson, R.B. 1950. An introduction to the natural history of British sawflies (Hymenoptera, Symphyta). Trans. Soc. Br. Ent., Bournemouth 10: 45-142.
- Benson, R.B. 1951. Hymenoptera Symphyta. Handbooks for the Identification of British Insects 6(2a): 1-50.
- Benson, R.B. 1952. Hymenoptera (Symphyta). Handbooks for the Identification of British Insects, 6(2b): 51-138.
- Benson, R.B. 1958. Hymenoptera (Symphyta). Handbooks for the Identification of British Insects 6(2c): 139-252.
- Benson, R.B. 1966. A new genus of the Lycaotini (Blennocampinae) in Turkey (Hymenoptera, Tenthredinidae). Proc. R. Ent. Soc. Lond, (B) 35: 75-77.
- Benson, R.B. 1967. A revision of Athaliini (Hymenoptera: Tenthredinidae). Bull. British Mus. Nat. Hist. Ent., 11: 7.
- Benson, R.B. 1968. Hymenoptera from Turkey, Symphyta. Bulletin of the British Museum (N.H.) Entomology 22(4): 4-207.
- Birkardeşler, H. 1964. Kocaeli ve Sakarya Bölgesinde *Hoplocampa* türlerinin (Hymenoptera-Tenthredinidae) kısa biyolojisi ve mücadeleşi üzerinde ilk çalışmalar. M.Sc. thesis, Ankara Ü. Zir. Fak. Ent. Kürs.
- Birkardeşler, H. 1971. Marmara Bölgesinde Kiraz sülügü (*Caloria limacina* Retz.)'nın yayılışı, biyolojisi ve mücadelesi üzerinde araştırmalar. Bitki Koruma Bülteni. 11(3): 181-201.
- Chevin, H. 1975. Remarques taxinomiques et biologiques sur les Macrophyia (Hym., Tenthredinidae) se développant sur Sambucus (Caprifoliaceae). Ann. Soc. Ent. France, (N.S.) 11(2): 253-260.
- Chevin, H. 1984. Contribution a la faune de l'Iran 26. Hymenopteres Symphytes. Nouv. Rev. Ent. (N.S.), T.I. 4: 347-351.
- Chevin, H. and Chenon, R.D. 1982. Contribution a la faune des Hymenopteres Symphytes de Turquie. Bull. Ent. France 87: 44-47.
- Cingovski, J. 1980. Contribution to the knowledge on the fauna of sawflies on the mountain Galicica (Hymenoptera, Symphyta). Fragmenta Balcanica 10(20): 175-189.
- Cingovski, J. 1982. Symphyta aus der sammlung des Instituts für pflanzenschutz in Belgrad (Hym., Symphyta). Fragmenta Balcanica 9(11): 97-110.
- Comstock, C.H. 1964. An Introduction to Entomology. Comstock Publishing Comp., Binghamton, N.Y. 1064 pp.
- Enslin, E. 1918. Die Tenthredinoidea Mittel Europas. Deutsche Ent. Zeitschr. 790 pp.
- Erden, F. 1988. Erzincan Bölgesi yumuşak çekirdekli meye ağaçlarının böcek kökenli zararlıları, tanımları ve önemlilerinin zararlılık durumları üzerinde araştırmalar. T.C. Tar. Or. ve Köyişleri Bak. Kor. ve Kont. Gen. Müd. Yayın Dairesi Bşk., Yayın No: 4, Ankara.
- Fitton, M.G., de V. Graham, M.W.R., Boucek, Z.R.J., Fergusson, N.D.M., Huddleston, T., Quinlan, J. and Richards, O.W. 1978. A Check List of British Insects. Handbooks for the Identification of British Insects. 11(4): 1-159.
- Gauld, I. and Bolton, B. 1988. The Hymenoptera. Oxford University Press. Oxford.
- Goulet, H. and Huber, J.T. 1993. Hymenoptera of The World: An Identification Guide to Families. Canada Communication Group-Publishing Ottawa., Canada.
- Guichard, K.M. and Harvey, D.H. 1967. Collecting in Turkey 1959, 1960 and 1962. Bull. Br. Mus. Nat. Hist. (Ent.). 19(4): 223-250.
- Güçlü, Ş. and Özbek, H. 1999. *Messa hortulana* (Klug) (Hymenoptera, Tenthredinidae), A new record and a new poplar pest for Turkey. Acta Ento. Bulg. 2,3,4: 72-75.
- Kiroğlu, H. and Baysal, Y. 1971. Amasya'da Armut testereli arısı (*Hoplocampa brevis* Klug) üzerinde araştırmalar. Zir. Müc. Ar. Yill. 38-39.
- Kontuniemi, T. 1965. Die letzte larvale Hautung bei den Sagewespen (Hym., Symphyta) als taxonomisches Kriterium. Ann. Entomol. Fenn. 31(2): 115-117.
- Lacourt, P.J. 1999. Repertoire des Tenthredinidae ouest-palearctiques (Hymenoptera, Symphyta). Memories de la SEF, NO: 3, Societe entomologique de France, Juillet, Paris.
- Liston, A.D. 1981. *Pristiphora* (Lygaeophora) *lanifica* (Zaddach, 1882) New to Britain (Hymenoptera: Tenthredinidae). Entomologist's Gazette 32: 181-184.
- Liston, A.D. 1983a. Sawflies (Hymenoptera) from the Orkney Islands. Entomologist's Gazette 34: 113-116.
- Liston, A.D. 1983b. Some sawflies from Greece (Hym., Symphyta). Entom. Gesellschaft Basel 33(3): 85-88.
- Liston, A.D. 1983c. Abnormal morphology and venation in Sawflies (Hym., Symphyta). Entomologist's Records 95: 208-210.
- Liston, A.D. 1995. Compendium of European Sawflies. Chalastos Forestry, Gottfrieding, Germany.
- Magis, N. 2002. Tenthredinidae recoltes en Bulgarie à l'aide du "piège Malaise", sous les auspices du Musée d'Histoire Naturelle de Leiden. Notes fauniques de Gembloux 46: 13-22.
- Medvedev, G.S. 1994. Keys to the Insects of the European Part of the USSR. E.J. Brill, Volume III. New York.
- Meitzner, V. and Taeger, A. 1982. Beitrag zur Blattwespenfauna Bulgariens (Hymenoptera, Symphyta). Entomologische Nachrichten und Berichte 26: 127-130.
- Moczár, L. and Zombori, L. 1973. Levéldarazs-alkatuk I-Tenthredinoide I. Fauna Hungariae, Budapest. 111: 128pp.
- Muche, W.H. 1975. Die Symphyten meiner Mittelasienreise 1975 (hymenoptera, Symphyta). Faun. Abh. Mus. Tierk. Dresden, 5(13): 289-295.
- Öge, Z. 1958. Pancar zararlılarından Pancar Yaprak Arısı (*Athalia spinarum* and *Atomaria linearis* Steph. Pancar. 82: 5-6.
- Özbek, H. 1986. Erzurum'da Türkiye için yeni bir Huş (*Betula verrucosa* Ehrh.) zararlısı, *Fenusia pusilla* Lep. (Tenthredinidae, Hymenoptera). Türk. Bitki Kor. Derg. 10(2): 115-123.

- Özeren, M. 1952. *Hoplocampa brevis* (Armut testereli arısı). Tomurcuk 1 (6): 21 pp.
- Özeren, H. 1970. Ankara Bölgesinde bazı meyve ve orman ağaçları ile süs bitkilerine zarar veren Hymenoptera türleri, yayılışı ve taksonomileri üzerinde ön araştırmalar. A.Ü.Zir. Fak. Yay.: 396, B.A. Inc. 246: 73pp.
- Schedl, W. 1981. Die pflanzenwespen der Insel Kreta (Insecta: Hymenoptera, Symphyta). Ber. Nat. Med. Ver. Innsbruck 68: 145-157.
- Schedl, W. 1991. Hymenoptera unterordnung Symphyta. Pflanzenwespen. Handbuch der Zoologie, Walter de Gruyter, Berlin & New York 4(31): 1-136.
- Scobiola-Palade, X.G. 1981. Hymenoptera Symphyta Tenthredinoidea. Fauna Republicii Socialiste Romania, Bucarest 9(9): 1-328.
- Shaw, M.R. 1978. Some sawfly (Hym., Symphyta) records from Greater Manchester county. Lancashire & Cheshire fauna committee No: 73: 13-14.
- Smith, D.R. 1976. World genera of the leaf-mining sawfly tribe Fenusini (Hymenoptera: Tenthredinidae). Ent. Scand. 7: 253-260.
- Smith, D.R. 1979. Nearctic Sawflies IV. Allantinae: adults and larvae (Hymenoptera, Tenthredinidae). Technical Bulletin of the United States Department of Agriculture 1595. 1-172.
- Smith, D.R. 1981. Studies on the leaf-mining sawflies of the tribe Fenusini in Asia (Hymenoptera: Tenthredinidae). Proc. Entomol. Soc. Wash. 83(4): 763-771.
- Smith, D.R. 1982. List of the Symphyta (Hymenoptera) of Israel, with description of four new species. Israel J. Ent. 16: 1-25.
- Smith, D.R. 1986. Review of the sawfly genus *Lagium* (Hymenoptera: Tenthredinidae). Proc. Ent. Soc. Wash. 88(1): 179-184.
- Taeger, A. and Blank, S.M. 1998. Pflanzenwespen Deutschlands (Hymenoptera, Symphyta). Goecke und Evers, Keltern.
- Vassilev, I. 1978. Hymenoptera, Symphyta. Fauna Bulgarica, Sofia 8: 1-179.
- Vigna Taglianti, A., Audisio, P.A., Biondi, M., Bologna, M.A., Capaneto, G.M., De Biase, A., Fattorini, S., Piatella, E., Sindaco, R., Venchi, A. and Zaporoli, M. 1999. A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. Biogeographia XX(1): 31-59.
- Viitasaari, M. and Vikberg, V. 1985. A checklist of sawflies (Hymenoptera, Symphyta) of Finland. Notulae Entomologicae 65: 1-17.
- Vikberg, V. 1975. Notes on some Nematine sawflies feeding on Larix (Hym., Tenthredinidae). Ann. Ent. Fenn. 41(1): 1-10.
- Wolf, F. 1968. Hymenopteres Symphytes de Turquie. Bull. Rech. Agron. Gembloux (N.S.) 3(3): 562-565.
- Zhelokochev, A. 1928. Über palaarktische Dolerinae, Zool. Anzeiger, 79: 105-112.
- Zhelochovtsev, A.N., Zinovjev, A.G. 1996. A list of the sawflies and horntails (Hymenoptera, Symphyta) of the fauna of Russia and adjacent territories. II. Entomologicheskoe Obozrenie 75(2): 357-378.
- Zombori, L. 1972. Symphyta (Hymenoptera) from Mongolia, II. Acta Z. Acad. Sci. Hungaricae 18(3-4): 435-448.
- Zombori, L. 1973. Symphyta (Hymenoptera) from Mongolia III. Acta Z. Acad. Sci. Hungaricae 19(3-4): 445-460.
- Zombori, L. 1974a. Data to sawfly fauna of Yugoslavia (Hymenoptera: Symphyta). Musei Macedonici Scientiarum Naturalium 9(18): 173-185.
- Zombori, L. 1974b. Sawflies from Korea (Hymenoptera: Symphyta). Acta Zoologica Acad. Sci. Hungaricae 20(3-4): 453-461.
- Zombori, L. 1977. Two new sawfly species from Armenia (Hymenoptera: Symphyta). Acta Zoologica Acad. Sci. Hungaricae 24(3-4): 451-454.
- Zombori, L. 1978. New sawfly species from Korea (Hymenoptera: Symphyta). Acta Zoologica Acad. Sci. Hungaricae 24(1-2): 253-268.
- Zombori, L. 1979. The Symphyta of the Dodero Collection 1: Description of six new taxa and notes synonymy (Hymenoptera). Frustula Ent. (N.S.) 1(14): 223-246.
- Zombori, L. 1981. The European genera of Selandriinae and Dolerinae (Hymenoptera: Symphta, Tenthredinidae). Acta Zoologica Acad. Sci. Hungaricae 27(3-4): 443-450.
- Zombori, L. 1982a. The European genera of Blennocampinae (Hymenoptera: Symphyta, Tenthredinidae). Acta Zoologica Acad. Sci. Hungaricae 28(1-2): 183-192.
- Zombori, L. 1982b. The European genera of Tenthredininae (Hymenoptera: Symphyta, Tenthredinidae). Acta Zoologica Acad. Sci. Hungaricae 28(3-4): 455-460.
- Zombori, L. 1984. The European genera of Nematinae (Hymenoptera: Symphyta, Tenthredinidae). Acta Zoologica Acad. Sci. Hungaricae 30(3-4): 545-550.
- Zombori, L. 1985. Sawflies from Nepal (Hymenoptera), I. Acta Zoologica Acad. Sci. Hungaricae 31(1-3): 291-293.