



Crystal structure of bis(4-acetyl-anilinium) tetrachloridocobaltate(II)

Manickam Thairiyaraja,^a Arumugam Elangovan,^b
Ramasamy Shanmugam,^b Kuthambalam Selvaraju^{a*} and
Subbiah Thamotharan^{c*}

^aPG & Research Department of Physics, Government Arts College, Ariyalur 621 713, India, ^bDepartment of Chemistry, Thiagarajar College, Madurai 625 009, India, and ^cBiomolecular Crystallography Laboratory, Department of Bioinformatics, School of Chemical and Biotechnology, SASTRA University, Thanjavur 613 401, India.

*Correspondence e-mail: selsphy@yahoo.com, thamu@scbt.sastra.edu

Received 9 November 2015; accepted 11 November 2015

Edited by M. Weil, Vienna University of Technology, Austria


The structure of the title salt, $(C_8H_{10}NO)_2[CoCl_4]$, is isotypic with the analogous cuprate(II) structure. The asymmetric unit contains one 4-acetylanilinium cation and one half of a tetrachloridocobaltate(II) anion for which the Co^{II} atom and two Cl^- ligands lie on a mirror plane. The $Co-Cl$ distances in the distorted tetrahedral anion range from 2.2519 (6) to 2.2954 (9) Å and the $Cl-Co-Cl$ angles range from 106.53 (2) to 110.81 (4)°. In the crystal, cations are self-assembled by intermolecular $N-H\cdots O$ hydrogen-bonding interactions, leading to a $C(8)$ chain motif with the chains running parallel to the b axis. $\pi-\pi$ stacking interactions between benzene rings, with a centroid-to-centroid distance of 3.709 Å, are also observed along this direction. The $CoCl_4^{2-}$ anions are sandwiched between the cationic chains and interact with each other through intermolecular $N-H\cdots Cl$ hydrogen-bonding interactions, forming a three-dimensional network structure.

Keywords: crystal structure; isotypism; cobalt(II); hydrogen bonding; $\pi-\pi$ stacking interactions.

CCDC reference: 967676

1. Related literature

For the structure of the isotypic tetrachloridocuprate(II) compound, see: Elangovan *et al.* (2007).



2. Experimental

2.1. Crystal data

$(C_8H_{10}NO)_2[CoCl_4]$
 $M_r = 473.07$
Orthorhombic, $Cmce$
 $a = 19.4605$ (6) Å
 $b = 15.5108$ (6) Å
 $c = 13.7374$ (5) Å

$V = 4146.6$ (3) Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 1.36$ mm⁻¹
 $T = 293$ K
 $0.3 \times 0.2 \times 0.2$ mm

2.2. Data collection

Bruker SMART APEX CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2004)
 $T_{min} = 0.687$, $T_{max} = 0.773$

24499 measured reflections
3329 independent reflections
2439 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.035$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.123$
 $S = 1.05$
3329 reflections
131 parameters
3 restraints

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{max} = 1.35$ e Å⁻³
 $\Delta\rho_{min} = -0.66$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N41-H41A\cdots O11^i$	0.91 (2)	1.88 (2)	2.781 (3)	174 (3)
$N41-H41B\cdots Cl2^{ii}$	0.92 (2)	2.31 (2)	3.211 (2)	168 (3)
$N41-H41C\cdots Cl3^{iii}$	0.88 (2)	2.48 (2)	3.309 (3)	157 (3)

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, z$; (ii) $x + \frac{1}{2}, y, -z + \frac{3}{2}$; (iii) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z + 2$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; method used to solve structure: coordinates taken from an isotypic structure; program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Acknowledgements

ST is very grateful to the management of SASTRA University for infrastructural and financial support (Professor TRR fund).

Supporting information for this paper is available from the IUCr electronic archives (Reference: WM5237).

References

- Bruker (2004). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Elangovan, A., Thamaraichelvan, A., Ramu, A., Athimoolam, S. & Natarajan, S. (2007). *Acta Cryst. E* **63**, m224–m226.
- Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2015). E71, m221–m222 [https://doi.org/10.1107/S2056989015021404]

Crystal structure of bis(4-acetylanilinium) tetrachloridocobaltate(II)


Manickam Thairiyaraja, Arumugam Elangovan, Ramasamy Shanmugam, Kuthambalam Selvaraju and Subbiah Thamocharan

S1. Synthesis and crystallization

A solution of 4-aminoacetophenone (20 mmol) in 2 ml of HCl and deionized water (10 ml) was added to a 10 ml solution of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (10 mmol). The resulting solution was concentrated and kept unperturbed at ambient temperature for crystallization. Dark green block-shaped crystals were obtained after 7 days.

S2. Refinement

Since the title complex is isotypic with its tetrachloridocuprate counterpart, it was refined with the coordinates of the latter (Elangovan *et al.*, 2007) as starting parameters. The amino H atoms were located from a difference Fourier map and refined with a distance restraint of $\text{N—H} = 0.89(2) \text{ \AA}$. The methyl H atoms were constrained to an ideal geometry ($\text{C—H} = 0.96 \text{ \AA}$) with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$, but were allowed to rotate freely about the C—C bond. The remaining H atoms were positioned in geometrically calculated positions and refined using a riding model with $\text{C—H} = 0.93 \text{ \AA}$ and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. At this stage, the maximum residual electron density of 1.35 e \AA^{-3} indicated the presence of a possible atom at Wyckoff position $4a$ at a distance of 2.81 \AA near atom H5. This peak was assumed to be the O atom of a water molecule and was refined with isotropic displacement parameters. However, the resultant model had slightly higher reliability factors and a very high isotropic atomic displacement parameter for this O atom. As a consequence, this water O atom was not included in the final model.

**Figure 1**

The molecular components in the structure of the title salt. Displacement ellipsoids are drawn at the 50% probability level. [Symmetry code: (i) $-x, y, z$.]




Figure 2

The crystal packing of the title salt viewed along the *c* axis. Hydrogen bonds are shown as dashed lines; H atoms bound to C were omitted for clarity.

Bis(4-acetylanilinium) tetrachloridocobaltate(II)

Crystal data

(C₈H₁₀NO)₂[CoCl₄]

M_r = 473.07

Orthorhombic, *Cmce*

a = 19.4605 (6) Å

b = 15.5108 (6) Å

c = 13.7374 (5) Å

V = 4146.6 (3) Å³

Z = 8

F(000) = 1928

D_x = 1.516 Mg m⁻³

Mo *Kα* radiation, λ = 0.71073 Å

Cell parameters from 14227 reflections

θ = 2.0–30.0°

μ = 1.36 mm⁻¹

T = 293 K

Block, green

0.3 × 0.2 × 0.2 mm

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

ω and φ scan

Absorption correction: multi-scan
(*SADABS*; Bruker, 2004)

T_{min} = 0.687, *T_{max}* = 0.773

24499 measured reflections

3329 independent reflections

2439 reflections with *I* > 2σ(*I*)

R_{int} = 0.035

θ_{max} = 30.9°, θ_{min} = 2.2°

h = −27→28

k = −18→22

l = −19→18

Refinement

Refinement on *F*²

Least-squares matrix: full

R [*F*² > 2σ(*F*²)] = 0.039

wR(*F*²) = 0.123

S = 1.05

3329 reflections

131 parameters

3 restraints

Primary atom site location: isomorphous
structure methods

Hydrogen site location: mixed
H atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0558P)^2 + 6.0921P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 1.35 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.66 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O11	0.19113 (10)	0.60656 (11)	0.87507 (15)	0.0548 (5)
N41	0.37078 (11)	0.26577 (13)	0.90637 (18)	0.0421 (5)
H41A	0.3519 (18)	0.2142 (15)	0.891 (3)	0.082 (12)*
H41B	0.4081 (14)	0.276 (2)	0.867 (2)	0.068 (10)*
H41C	0.3860 (16)	0.268 (2)	0.9661 (14)	0.065 (10)*
C1	0.22194 (11)	0.46100 (12)	0.87145 (14)	0.0299 (4)
C2	0.20229 (11)	0.37474 (14)	0.86968 (16)	0.0353 (4)
H2	0.1563	0.3603	0.8613	0.042*
C3	0.25125 (12)	0.30994 (13)	0.88036 (16)	0.0366 (5)
H3	0.2384	0.2522	0.8793	0.044*
C4	0.31896 (11)	0.33283 (13)	0.89257 (15)	0.0322 (4)
C5	0.33997 (11)	0.41822 (14)	0.89347 (17)	0.0371 (5)
H5	0.3861	0.4324	0.9011	0.045*
C6	0.29092 (11)	0.48178 (13)	0.88278 (16)	0.0358 (4)
H6	0.3042	0.5394	0.8832	0.043*
C11	0.17101 (12)	0.53274 (14)	0.86411 (15)	0.0352 (4)
C12	0.09695 (12)	0.51393 (16)	0.8452 (2)	0.0483 (6)
H12A	0.0727	0.5670	0.8343	0.072*
H12B	0.0776	0.4848	0.9004	0.072*
H12C	0.0929	0.4779	0.7886	0.072*
Co1	0.0000	0.25104 (3)	0.86031 (3)	0.03357 (13)
Cl1	0.0000	0.33842 (5)	0.99225 (6)	0.04207 (19)
Cl2	0.0000	0.33034 (6)	0.71924 (6)	0.0462 (2)
Cl3	0.09841 (3)	0.17486 (5)	0.86639 (6)	0.0601 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O11	0.0497 (10)	0.0283 (8)	0.0863 (14)	0.0022 (7)	−0.0056 (9)	−0.0036 (8)
N41	0.0326 (10)	0.0337 (10)	0.0599 (14)	0.0014 (8)	−0.0043 (9)	0.0067 (9)
C1	0.0330 (9)	0.0256 (8)	0.0311 (10)	−0.0005 (7)	0.0005 (8)	−0.0006 (8)
C2	0.0280 (9)	0.0304 (9)	0.0476 (12)	−0.0044 (8)	−0.0006 (8)	0.0006 (8)
C3	0.0337 (10)	0.0261 (9)	0.0499 (12)	−0.0046 (8)	−0.0011 (9)	0.0022 (8)
C4	0.0311 (10)	0.0297 (9)	0.0358 (10)	0.0002 (8)	−0.0012 (8)	0.0030 (8)

C5	0.0306 (10)	0.0329 (10)	0.0479 (12)	−0.0054 (8)	−0.0054 (9)	0.0005 (9)
C6	0.0371 (11)	0.0270 (9)	0.0432 (12)	−0.0060 (8)	−0.0026 (9)	−0.0009 (8)
C11	0.0387 (11)	0.0310 (9)	0.0360 (11)	0.0017 (8)	0.0020 (9)	−0.0008 (8)
C12	0.0355 (12)	0.0417 (12)	0.0676 (17)	0.0053 (10)	0.0028 (11)	−0.0038 (11)
Co1	0.0253 (2)	0.0355 (2)	0.0399 (2)	0.000	0.000	−0.00463 (16)
Cl1	0.0469 (4)	0.0377 (4)	0.0416 (4)	0.000	0.000	−0.0080 (3)
Cl2	0.0456 (4)	0.0525 (5)	0.0405 (4)	0.000	0.000	0.0007 (3)
Cl3	0.0399 (3)	0.0581 (4)	0.0824 (5)	0.0193 (3)	−0.0109 (3)	−0.0218 (3)

Geometric parameters (Å, °)

O11—C11	1.220 (3)	C4—C5	1.386 (3)
N41—C4	1.461 (3)	C5—C6	1.380 (3)
N41—H41A	0.904 (18)	C5—H5	0.9300
N41—H41B	0.921 (18)	C6—H6	0.9300
N41—H41C	0.872 (17)	C11—C12	1.493 (3)
C1—C6	1.389 (3)	C12—H12A	0.9600
C1—C2	1.392 (3)	C12—H12B	0.9600
C1—C11	1.494 (3)	C12—H12C	0.9600
C2—C3	1.393 (3)	Co1—Cl3 ⁱ	2.2519 (6)
C2—H2	0.9300	Co1—Cl3	2.2519 (6)
C3—C4	1.375 (3)	Co1—Cl1	2.2631 (9)
C3—H3	0.9300	Co1—Cl2	2.2954 (9)
C4—N41—H41A	109 (2)	C4—C5—H5	120.7
C4—N41—H41B	110 (2)	C5—C6—C1	120.96 (19)
H41A—N41—H41B	110 (3)	C5—C6—H6	119.5
C4—N41—H41C	109 (2)	C1—C6—H6	119.5
H41A—N41—H41C	112 (3)	O11—C11—C12	121.0 (2)
H41B—N41—H41C	106 (3)	O11—C11—C1	118.6 (2)
C6—C1—C2	119.37 (19)	C12—C11—C1	120.44 (19)
C6—C1—C11	118.42 (18)	C11—C12—H12A	109.5
C2—C1—C11	122.2 (2)	C11—C12—H12B	109.5
C1—C2—C3	120.3 (2)	H12A—C12—H12B	109.5
C1—C2—H2	119.9	C11—C12—H12C	109.5
C3—C2—H2	119.9	H12A—C12—H12C	109.5
C4—C3—C2	118.82 (19)	H12B—C12—H12C	109.5
C4—C3—H3	120.6	Cl3 ⁱ —Co1—Cl3	116.52 (4)
C2—C3—H3	120.6	Cl3 ⁱ —Co1—Cl1	106.53 (2)
C3—C4—C5	122.04 (19)	Cl3—Co1—Cl1	106.53 (2)
C3—C4—N41	119.57 (19)	Cl3 ⁱ —Co1—Cl2	108.21 (3)
C5—C4—N41	118.39 (19)	Cl3—Co1—Cl2	108.21 (3)
C6—C5—C4	118.5 (2)	Cl1—Co1—Cl2	110.81 (4)
C6—C5—H5	120.7		
C6—C1—C2—C3	0.8 (3)	C4—C5—C6—C1	0.1 (3)
C11—C1—C2—C3	−177.6 (2)	C2—C1—C6—C5	−0.8 (3)
C1—C2—C3—C4	0.0 (3)	C11—C1—C6—C5	177.7 (2)

C2—C3—C4—C5	−0.7 (3)	C6—C1—C11—O11	−5.0 (3)
C2—C3—C4—N41	178.4 (2)	C2—C1—C11—O11	173.4 (2)
C3—C4—C5—C6	0.7 (3)	C6—C1—C11—C12	175.8 (2)
N41—C4—C5—C6	−178.5 (2)	C2—C1—C11—C12	−5.8 (3)

Symmetry code: (i) $-x, y, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N41—H41A \cdots O11 ⁱⁱ	0.91 (2)	1.88 (2)	2.781 (3)	174 (3)
N41—H41B \cdots C12 ⁱⁱⁱ	0.92 (2)	2.31 (2)	3.211 (2)	168 (3)
N41—H41C \cdots C13 ^{iv}	0.88 (2)	2.48 (2)	3.309 (3)	157 (3)

Symmetry codes: (ii) $-x+1/2, y-1/2, z$; (iii) $x+1/2, y, -z+3/2$; (iv) $-x+1/2, -y+1/2, -z+2$.