

CRYSTALLOGRAPHIC
COMMUNICATIONS

ISSN 2056-9890

Crystal structure and Hirshfeld-surface analysis of (benzenecarbothioamide- κ S)bromidobis(triphenylphosphane- κ P)silver(I)

Wattana Ruangwut, Saowanit Saithong and Chaveng Pakawatchai*

Department of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90112, Thailand. *Correspondence e-mail: chavengp@gmail.com

Received 10 June 2016

Accepted 13 June 2016

Edited by W. T. A. Harrison, University of Aberdeen, Scotland

Keywords: crystal structure; Ag^I complex; benzenecarbothioamide; triphenylphosphane; hydrogen bonding; Hirshfeld surface; supra-molecular interaction.



CCDC reference: 1484796

Supporting information: this article has supporting information at journals.iucr.org/e

The title complex, [AgBr(C₇H₇NS)(C₁₈H₁₅P)₂], was obtained from the reaction of silver(I) bromide with benzenecarbothioamide (C₇H₇NS) and triphenylphosphane (C₁₈H₁₅P) in the mixed solvent of acetonitrile and ethanol. The mononuclear complex exhibits a distorted tetrahedral coordination geometry about the metal atom, arising from one S atom of a benzenecarbothioamide ligand, two P atoms of two triphenylphosphane molecules and one bromide ion. An intramolecular N—H...Br hydrogen bond is observed and in the crystal structure, inversion dimers linked by pairs of N—H...Br and C—H...Br hydrogen bonds are observed. In addition, C—H... π interactions occur, leading to [101] chains. Hirshfeld-surface analyses are presented and discussed.

1. Chemical context

Mixed-ligand complexes of Ag^I-containing phosphorus and sulfur donor ligands have been studied and published extensively in recent years (Dennehy *et al.*, 2007; Ruangwut & Pakawatchai, 2014) because of their potential ability to inhibit bacteria (Isab *et al.*, 2010; Nawaz *et al.*, 2011). Triphenylphosphane and thione ligands, which contain P and S donor atoms, respectively, are capable of forming mixed-ligand silver(I) complexes as mononuclear (Aslanidis *et al.*, 1997) and dinuclear models (Cox *et al.*, 2000). In this paper, we report the synthesis and structure of the mixed-ligand complex of silver(I) bromide with triphenylphosphane and benzenecarbothioamide ligands.



OPEN ACCESS




Figure 1
The molecular structure of the title compound, showing 30% probability displacement ellipsoids.

2. Structural commentary

The monomeric complex of the title compound crystallizes in the monoclinic crystal system, space group $P2_1/n$, and is shown in Fig. 1. The silver ion is four-coordinated exhibiting a distorted tetrahedral environment. This deviation can be explained by $P1-Ag1-P2$ angle which has the highest value of $121.60(2)^\circ$ due to the steric hindrance and the repulsion between two bulky triphenylphosphane molecules. The range of angles around the Ag atom of $97.338(18)$ – $121.60(2)^\circ$ is similar to that observed in the analogous mononuclear silver(I) complex $[AgCl(C_7H_7NS)(C_{18}H_{15}P)_2]$ previously synthesized by us (Ruangwut & Pakawatchai, 2014), in which the angles about the metal ion are $97.298(16)$ – $120.053(16)^\circ$. The Ag–S bond length of $2.6015(8)$ Å is slightly longer than




Figure 2
An inversion dimer in the crystal of the title compound linked by two pairs of $N-H\cdots Br$ interactions, forming $R_4^2(8)$ loops, and pairs of $C-H\cdots Br$ interactions, forming $R_2^2(14)$ loops.

Table 1

Hydrogen-bond geometry (Å, °).

$Cg3$ is the centroid of the $C13-C18$ ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N1-H1A\cdots Br1^i$	0.85 (1)	2.54 (1)	3.357 (3)	161 (3)
$N1-H1B\cdots Br1$	0.85 (1)	2.58 (1)	3.413 (3)	166 (3)
$C17-H17\cdots Br1^i$	0.93	2.91	3.789 (3)	158

in $[AgCl(C_7H_7NS)(C_{18}H_{15}P)_2]$, $2.5580(5)$ Å. The Ag–P bond lengths of $2.4682(7)$ and $2.4671(6)$ Å for $Ag1-P1$ and $Ag1-P2$, respectively, are similar to those of the Ag–P bond lengths in $[AgCl(C_7H_7NS)(C_{18}H_{15}P)_2]$ [$2.4529(5)$ and $2.4578(5)$ Å], and similar to the Ag–P distances of analogous tetrahedrally coordinated Ag^I complexes such as $[Ag(NO_3)(C_2H_3N_3S)(C_{18}H_{15}P)_2]$ [$2.4485(6)$ and $2.4493(6)$ Å; Wattananajana *et al.*, 2014], $[Ag(Htsa)(PPh_3)_3]$ [$2.574(7)$ – $2.611(6)$ Å; Nomiya *et al.*, 1998] and $[Ag(PPh_3)_2(bzoxth)]\cdot 2NO_3$ [$2.480(1)$ and $2.514(2)$ Å; McFarlane *et al.*, 1998]. An intramolecular hydrogen bond $N1-H1B\cdots Br1$ [$3.413(3)$ Å; Table 1] is found between one of the H atoms from an amine group of the benzenecarbothioamide molecule and the bromide ion, as depicted in Fig. 2, which also shows the intermolecular dimeric hydrogen bonds.

3. Supramolecular features

In the crystal, the dimeric intermolecular interactions are generated through a crystallographic inversion center by




Figure 3
The supramolecular $C-H\cdots\pi$ chain in the title compound.




Figure 4
 d_{norm} mapped on the Hirshfeld surface for visualizing the intermolecular interactions of the title compound. Dotted red lines represent hydrogen bonds.

linking through the $\text{N1-H1A}\cdots\text{Br1}^i$ [3.357 (3) Å] and $\text{C17-H17}\cdots\text{Br1}^i$ [3.789 (3) Å] [symmetry code: (i) $1-x, 1-y, 1-z$] hydrogen bonds between a pair of adjacent complex molecules; these are similar to the those in the above-mentioned complex $[\text{AgCl}(\text{C}_7\text{H}_7\text{NS})(\text{C}_{18}\text{H}_{15}\text{P})_2]$ (Ruangwut & Pakawatchai, 2014). There are two cyclic patterns of $R_4^2(8)$ loops formed by two pairs of $\text{N1-H1A}\cdots\text{Br1}$ and $\text{N1-H1B}\cdots\text{Br1}$ interactions and of $R_2^2(14)$ loops forming by a pair of $\text{C17-H17}\cdots\text{Br1}$ interactions, as illustrated in Fig. 2. In addition, supramolecular $\text{C-H}\cdots\pi$ chains (Fig. 3) are formed between the Csp^2 atoms of the phenyl rings and the centroids of another phenyl ring [$\text{C22-H22}\cdots\text{Cg3} = 3.782$ (3) Å].

4. Hirshfeld surface analysis

For the title complex, the Hirshfeld-surfaces analysis (McKinnon *et al.*, 2004; Spackman & Jayatilaka, 2009) was generated by *Crystal Explorer 3.1* (Wolff *et al.*, 2012) and mapped over d_{norm} , d_e and d_i fingerprint plot (Spackman & McKinnon 2002; McKinnon *et al.*, 2007). The contact distances to the closest atom inside (d_i) and outside (d_e) of the Hirshfeld surface analyse the intermolecular interactions *via* the mapping of d_{norm} , as depicted in Fig. 4. The interactions are shown on the Hirshfeld surfaces with short contacts indicated in red. The corresponding fingerprint plots (Fig. 5*a-d*) for Hirshfeld surfaces of the complex are shown with characteristic pseudo-symmetry wings in the upper left and lower right sides of the d_e and d_i diagonal axes that represent the overall 2D fingerprint plot and those delineated into $\text{H}\cdots\text{H}$, $\text{H}\cdots\text{Br}/\text{Br}\cdots\text{H}$, and $\text{C}\cdots\text{H}/\text{H}\cdots\text{C}$ contacts are shown in Fig. 5*a-d*, respectively. The fingerprint plot of $\text{H}\cdots\text{H}$ contacts represented by the largest contribution within the Hirshfeld surfaces (60.8%) are shown as one distinct pattern with a minimum value of $d_e + d_i \sim 2.6$ Å. The reciprocal $\text{H}\cdots\text{Br}/\text{Br}\cdots\text{H}$ contacts consist of 5.4% of the total Hirshfeld surface with $d_e + d_i \sim 3.3$ Å, exhibited by two symmetrical narrow pointed wings indicating the intermolecular hydrogen-bond interactions $\text{N1-H1A}\cdots\text{Br1}$ and $\text{C17-H17}\cdots\text{Br1}$ in the crystal packing. The presence of $\text{C-H}\cdots\pi$ interactions on the

fingerprint plot, which contribute 29.7% of overall Hirshfeld surface, are indicated by $d_e + d_i \sim 3.0$ Å.

5. Synthesis and crystallization

Silver(I) bromide (0.10 g, 0.5 mmol) was dissolved in the mixed solvent of 15 ml of acetonitrile and 15 ml of ethanol and then triphenylphosphane (0.27 g, 1 mmol) was added. The mixture was refluxed for 2 h at 343 K and a white precipitate




Figure 5
 Two-dimensional fingerprint plots of the title complex showing the percentage contributions of individual types of interactions: (a) all intermolecular interactions, (b) $\text{H}\cdots\text{H}$ contacts, (c) $\text{H}\cdots\text{Br}/\text{Br}\cdots\text{H}$ contacts and (d) $\text{C}\cdots\text{H}/\text{H}\cdots\text{C}$ contacts. d_e and d_i represent the distances from the surface to nearest external and internal atoms and the blue-cyan color represents increasing numbers of surface contributors at individual d_e/d_i points

Table 2
Experimental details.

Crystal data	
Chemical formula	[AgBr(C ₇ H ₇ NS)(C ₁₈ H ₁₅ P) ₂]
<i>M_r</i>	849.51
Crystal system, space group	Monoclinic, <i>P</i> 2 ₁ / <i>n</i>
Temperature (K)	293
<i>a</i> , <i>b</i> , <i>c</i> (Å)	14.4354 (5), 14.1925 (5), 19.1682 (6)
β (°)	98.786 (1)
<i>V</i> (Å ³)	3881.0 (2)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ^{−1})	1.72
Crystal size (mm)	0.23 × 0.13 × 0.08
Data collection	
Diffraction	Broker <i>APEX</i> CCD area-detector
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2003)
<i>T</i> _{min} , <i>T</i> _{max}	0.885, 1.000
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	52150, 9261, 6704
<i>R</i> _{int}	0.048
(sin θ /λ) _{max} (Å ^{−1})	0.658
Refinement	
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.036, 0.085, 1.02
No. of reflections	9261
No. of parameters	450
No. of restraints	2
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
Δρ _{max} , Δρ _{min} (e Å ^{−3})	0.47, −0.24

Computer programs: *SMART* and *SAINT* (Bruker, 2003), *SHELXT2014* (Sheldrick, 2015a), *SHELXL2014* (Sheldrick, 2015b), *Mercury* (Macrae *et al.*, 2008), *WinGX* (Farrugia, 2012) and *publCIF* (Westrip, 2010).

was formed. After that, benzenecarbothioamide (0.13 g, 1 mmol) was added and continually refluxed for 5 h. At that time, the white precipitate dissolved. The clear yellow solution was filtered and left to evaporate at room temperature. After a day, pale-yellow blocks of the title compound were filtered off and dried *in vacuo*. Calculated for C₄₃H₃₇AgBrNP₂S: C 61.07, H 4.37, N 1.65 and S 3.78%. Found: C 60.50, H 4.21, N 1.43 and S 3.70%.

6. Refinement

Crystal data and details of structure determination are summarized in Table 2. All H atoms on carbon atoms were positioned geometrically and refined using a riding-model

approximation with C—H = 0.93 Å with *U*_{iso}(H) = 1.2 *U*_{eq}(C). N-bound H atoms were found from difference maps and refined isotropically with distance restraint N—H = 0.85–0.86 Å.

Acknowledgements

We are grateful for financial support from the Center of Excellence for Innovation in Chemistry (PERCH-CIC), Office of the Higher Education Commission, Ministry of Education, the Department of Chemistry and the Graduate school, Prince of Songkla University.

References

- Aslanidis, P., Karagiannidis, P., Akrivos, P. D., Krebs, B. & Läge, M. (1997). *Inorg. Chim. Acta*, **254**, 277–284.
- Bruker (2003). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cox, P. J., Aslanidis, P., Karagiannidis, P. & Hadjikakou, S. (2000). *Inorg. Chim. Acta*, **310**, 268–272.
- Dennehy, M., Quinzani, O. V. & Jennings, M. (2007). *J. Mol. Struct.* **841**, 110–117.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Isab, A. A., Nawaz, S., Saleem, M., Altaf, M., Monim-ul-Mehboob, M., Ahmad, S. & Evans, H. S. (2010). *Polyhedron*, **29**, 1251–1256.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- McFarlane, W., Akrivos, P. D., Aslanidis, P., Karagiannidis, P., Hatzisymeon, C., Numan, M. & Kokkou, S. (1998). *Inorg. Chim. Acta*, **281**, 121–125.
- McKinnon, J. J., Jayatilaka, D. & Spackman, M. A. (2007). *Chem. Commun.* pp. 3814–3816.
- McKinnon, J. J., Spackman, M. A. & Mitchell, A. S. (2004). *Acta Cryst.* **B60**, 627–668.
- Nawaz, S., Isab, A. A., Merz, K., Vasylyeva, V., Metzler-Nolte, N., Saleem, M. & Ahmad, S. (2011). *Polyhedron*, **30**, 1502–1506.
- Nomiya, K., Kasuga, N. C., Takamori, I. & Tsuda, K. (1998). *Polyhedron*, **17**, 3519–3530.
- Ruangwut, W. & Pakawatchai, C. (2014). *Acta Cryst.* **E70**, m310–m311.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Spackman, M. A. & Jayatilaka, D. (2009). *CrystEngComm*, **11**, 19–32.
- Spackman, M. A. & McKinnon, J. J. (2002). *CrystEngComm*, **4**, 378–392.
- Wattananajana, Y., Palamae, S., Ratthiwan, J. & Nimthong, R. (2014). *Acta Cryst.* **E70**, m61–m62.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Wolff, S. K., Grimwood, D. J., McKinnon, J. J., Turner, M. J., Jayatilaka, D. & Spackman, M. A. (2012). *Crystal Explorer*. The University of Western Australia.

supporting information

Acta Cryst. (2016). E72, 984–987 [https://doi.org/10.1107/S2056989016009518]

Crystal structure and Hirshfeld-surface analysis of (benzenecarbothioamide- κ S)bromidobis(triphenylphosphane- κ P)silver(I)

Wattana Ruangwut, Saowanit Saithong and Chaveng Pakawatchai

Computing details

Data collection: *SMART* (Bruker, 2003); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT* (Bruker, 2003); program(s) used to solve structure: *SHELXT2014* (Sheldrick, 2015); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *publCIF* (Westrip, 2010).

(Benzenecarbothioamide- κ S)bromidobis(triphenylphosphane- κ P)silver(I)

Crystal data

[AgBr(C₇H₇NS)(C₁₈H₁₅P)₂]

$M_r = 849.51$

Monoclinic, $P2_1/n$

$a = 14.4354$ (5) Å

$b = 14.1925$ (5) Å

$c = 19.1682$ (6) Å

$\beta = 98.786$ (1)°

$V = 3881.0$ (2) Å³

$Z = 4$

$F(000) = 1720$

$D_x = 1.454$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7218 reflections

$\theta = 2.4$ – 22.5°

$\mu = 1.72$ mm⁻¹

$T = 293$ K

Block, pale yellow

$0.23 \times 0.13 \times 0.08$ mm

Data collection

Bruker APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Frames, each covering 0.3° in ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2003)

$T_{\min} = 0.885$, $T_{\max} = 1.000$

52150 measured reflections

9261 independent reflections

6704 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.048$

$\theta_{\max} = 27.9^\circ$, $\theta_{\min} = 1.7^\circ$

$h = -18 \rightarrow 18$

$k = -18 \rightarrow 18$

$l = -25 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.085$

$S = 1.02$

9261 reflections

450 parameters

2 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0406P)^2 + 0.3053P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 0.47$ e Å⁻³

$\Delta\rho_{\min} = -0.24$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.46287 (2)	0.62099 (2)	0.69031 (2)	0.04396 (7)
Br1	0.58251 (2)	0.50868 (2)	0.63144 (2)	0.05659 (9)
P2	0.48182 (4)	0.57124 (5)	0.81500 (3)	0.03691 (15)
P1	0.52341 (5)	0.77367 (5)	0.65520 (3)	0.03869 (15)
S1	0.29135 (5)	0.58599 (6)	0.63246 (4)	0.0582 (2)
N1	0.36464 (19)	0.50017 (19)	0.53303 (13)	0.0570 (6)
C31	0.46028 (17)	0.44578 (18)	0.82667 (13)	0.0400 (6)
C13	0.52381 (17)	0.79219 (17)	0.56084 (13)	0.0400 (6)
C19	0.59722 (17)	0.58766 (17)	0.86800 (13)	0.0398 (6)
C25	0.40173 (17)	0.63235 (18)	0.86506 (13)	0.0418 (6)
C1	0.64521 (18)	0.78817 (18)	0.69490 (14)	0.0446 (6)
C7	0.46305 (18)	0.87882 (18)	0.68001 (13)	0.0431 (6)
C38	0.20344 (19)	0.5373 (2)	0.49980 (13)	0.0479 (6)
C8	0.5067 (2)	0.95988 (19)	0.70832 (14)	0.0496 (7)
H8	0.5717	0.9634	0.7165	0.059*
C14	0.5523 (2)	0.87597 (18)	0.53392 (15)	0.0505 (7)
H14	0.5707	0.9259	0.5643	0.061*
C37	0.29051 (18)	0.53868 (19)	0.55164 (13)	0.0438 (6)
C26	0.3653 (2)	0.5943 (2)	0.92112 (15)	0.0575 (7)
H26	0.3812	0.5330	0.9354	0.069*
C18	0.4949 (2)	0.7197 (2)	0.51477 (14)	0.0567 (8)
H18	0.4745	0.6633	0.5319	0.068*
C36	0.4922 (2)	0.39729 (19)	0.88842 (15)	0.0527 (7)
H36	0.5274	0.4286	0.9261	0.063*
C15	0.5538 (2)	0.8863 (2)	0.46293 (16)	0.0570 (8)
H15	0.5739	0.9427	0.4456	0.068*
C32	0.4108 (2)	0.39653 (19)	0.77074 (15)	0.0533 (7)
H32	0.3904	0.4272	0.7283	0.064*
C9	0.4535 (3)	1.0364 (2)	0.72469 (15)	0.0627 (8)
H9	0.4832	1.0904	0.7444	0.075*
C30	0.3763 (2)	0.7235 (2)	0.84460 (17)	0.0589 (8)
H30	0.3992	0.7501	0.8063	0.071*
C6	0.7174 (2)	0.7775 (2)	0.65652 (18)	0.0662 (8)
H6	0.7044	0.7668	0.6081	0.079*
C35	0.4722 (2)	0.3025 (2)	0.89437 (16)	0.0629 (8)
H35	0.4934	0.2708	0.9362	0.075*
C20	0.6114 (2)	0.6257 (2)	0.93455 (15)	0.0625 (8)
H20	0.5601	0.6456	0.9547	0.075*
C16	0.5258 (2)	0.8140 (2)	0.41780 (15)	0.0638 (8)

H16	0.5267	0.8212	0.3697	0.077*
C34	0.4216 (2)	0.2551 (2)	0.83930 (18)	0.0647 (8)
H34	0.4078	0.1916	0.8437	0.078*
C12	0.3662 (2)	0.8761 (2)	0.66785 (19)	0.0675 (9)
H12	0.3354	0.8221	0.6490	0.081*
C22	0.7760 (2)	0.6065 (2)	0.9435 (2)	0.0730 (10)
H22	0.8361	0.6122	0.9689	0.088*
C17	0.4962 (3)	0.7304 (2)	0.44316 (16)	0.0703 (9)
H17	0.4772	0.6812	0.4123	0.084*
C33	0.3916 (2)	0.3018 (2)	0.77778 (18)	0.0676 (9)
H33	0.3577	0.2695	0.7400	0.081*
C43	0.1355 (2)	0.6041 (3)	0.50190 (18)	0.0769 (11)
H43	0.1446	0.6509	0.5363	0.092*
C27	0.3056 (2)	0.6455 (3)	0.95658 (19)	0.0755 (10)
H27	0.2818	0.6189	0.9945	0.091*
C24	0.6743 (2)	0.5599 (2)	0.83923 (16)	0.0646 (8)
H24	0.6667	0.5344	0.7940	0.077*
C28	0.2820 (2)	0.7350 (3)	0.9356 (2)	0.0835 (11)
H28	0.2413	0.7693	0.9592	0.100*
C39	0.1878 (2)	0.4692 (2)	0.44791 (17)	0.0699 (9)
H39	0.2329	0.4232	0.4450	0.084*
C2	0.6671 (2)	0.8013 (2)	0.76686 (15)	0.0621 (8)
H2	0.6190	0.8066	0.7939	0.075*
C3	0.7590 (3)	0.8066 (3)	0.7994 (2)	0.0811 (11)
H3	0.7727	0.8166	0.8478	0.097*
C29	0.3173 (2)	0.7755 (3)	0.8803 (2)	0.0775 (10)
H29	0.3019	0.8372	0.8669	0.093*
C23	0.7633 (2)	0.5695 (3)	0.8772 (2)	0.0807 (11)
H23	0.8151	0.5505	0.8573	0.097*
C21	0.7004 (2)	0.6350 (3)	0.97230 (18)	0.0788 (10)
H21	0.7085	0.6608	1.0174	0.095*
C4	0.8294 (3)	0.7974 (3)	0.7605 (2)	0.0887 (12)
H4	0.8914	0.8012	0.7823	0.106*
C10	0.3588 (3)	1.0325 (3)	0.71206 (19)	0.0775 (11)
H10	0.3236	1.0839	0.7228	0.093*
C41	0.0393 (3)	0.5356 (3)	0.40405 (18)	0.0859 (12)
H41	−0.0162	0.5347	0.3722	0.103*
C11	0.3150 (3)	0.9533 (3)	0.6836 (2)	0.0874 (12)
H11	0.2499	0.9512	0.6746	0.105*
C5	0.8099 (2)	0.7825 (3)	0.6898 (2)	0.0874 (11)
H5	0.8586	0.7757	0.6636	0.105*
C40	0.1056 (3)	0.4688 (3)	0.4003 (2)	0.0889 (12)
H40	0.0957	0.4226	0.3656	0.107*
C42	0.0537 (3)	0.6035 (3)	0.4541 (2)	0.0978 (14)
H42	0.0085	0.6497	0.4562	0.117*
H1A	0.370 (2)	0.485 (2)	0.4908 (7)	0.064 (10)*
H1B	0.4142 (14)	0.498 (2)	0.5637 (13)	0.071 (10)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.05487 (13)	0.04402 (12)	0.03328 (10)	−0.00277 (9)	0.00765 (8)	0.00067 (8)
Br1	0.06087 (19)	0.06224 (19)	0.04602 (16)	0.01813 (14)	0.00608 (13)	−0.00806 (13)
P2	0.0384 (4)	0.0428 (4)	0.0294 (3)	−0.0038 (3)	0.0046 (3)	−0.0006 (3)
P1	0.0405 (4)	0.0387 (4)	0.0364 (3)	−0.0011 (3)	0.0043 (3)	0.0022 (3)
S1	0.0462 (4)	0.0899 (6)	0.0381 (4)	0.0004 (4)	0.0049 (3)	−0.0146 (4)
N1	0.0511 (16)	0.0765 (18)	0.0416 (15)	0.0049 (13)	0.0016 (12)	−0.0149 (13)
C31	0.0390 (14)	0.0432 (14)	0.0395 (14)	−0.0028 (11)	0.0117 (11)	−0.0001 (11)
C13	0.0413 (14)	0.0394 (14)	0.0392 (13)	0.0008 (11)	0.0056 (11)	0.0026 (11)
C19	0.0410 (14)	0.0418 (13)	0.0357 (13)	−0.0033 (11)	0.0032 (11)	0.0038 (11)
C25	0.0335 (13)	0.0524 (16)	0.0381 (13)	−0.0018 (11)	0.0009 (10)	−0.0091 (11)
C1	0.0417 (15)	0.0399 (14)	0.0508 (16)	−0.0016 (11)	0.0031 (12)	0.0019 (12)
C7	0.0475 (15)	0.0438 (15)	0.0387 (14)	0.0029 (12)	0.0090 (11)	0.0079 (11)
C38	0.0470 (16)	0.0564 (17)	0.0388 (15)	−0.0034 (13)	0.0017 (12)	−0.0017 (12)
C8	0.0599 (18)	0.0475 (16)	0.0412 (15)	0.0022 (14)	0.0076 (13)	0.0020 (12)
C14	0.0604 (18)	0.0443 (15)	0.0478 (16)	−0.0077 (13)	0.0111 (13)	−0.0001 (12)
C37	0.0480 (16)	0.0474 (15)	0.0358 (13)	−0.0048 (12)	0.0062 (11)	−0.0004 (11)
C26	0.0495 (17)	0.076 (2)	0.0486 (17)	0.0003 (15)	0.0137 (13)	−0.0011 (15)
C18	0.081 (2)	0.0434 (16)	0.0426 (16)	−0.0081 (14)	−0.0016 (14)	0.0039 (12)
C36	0.0664 (19)	0.0514 (17)	0.0409 (15)	−0.0057 (14)	0.0105 (13)	0.0014 (12)
C15	0.069 (2)	0.0506 (17)	0.0537 (18)	−0.0045 (14)	0.0185 (15)	0.0128 (14)
C32	0.0589 (18)	0.0466 (17)	0.0518 (17)	−0.0013 (13)	−0.0006 (14)	−0.0021 (13)
C9	0.099 (3)	0.0502 (18)	0.0401 (16)	0.0130 (17)	0.0152 (16)	0.0038 (13)
C30	0.0547 (18)	0.0578 (19)	0.0644 (19)	0.0051 (14)	0.0103 (15)	−0.0063 (15)
C6	0.0484 (18)	0.083 (2)	0.067 (2)	0.0052 (16)	0.0095 (15)	−0.0037 (17)
C35	0.086 (2)	0.0514 (18)	0.0558 (18)	0.0008 (16)	0.0264 (17)	0.0143 (15)
C20	0.0428 (16)	0.094 (2)	0.0492 (17)	−0.0026 (16)	0.0020 (13)	−0.0187 (16)
C16	0.088 (2)	0.067 (2)	0.0371 (15)	0.0064 (18)	0.0116 (15)	0.0091 (15)
C34	0.075 (2)	0.0433 (17)	0.080 (2)	−0.0069 (15)	0.0283 (18)	−0.0005 (16)
C12	0.0484 (18)	0.0524 (18)	0.102 (3)	0.0035 (15)	0.0115 (17)	0.0094 (17)
C22	0.0412 (18)	0.090 (3)	0.081 (3)	−0.0003 (17)	−0.0120 (16)	0.005 (2)
C17	0.112 (3)	0.0534 (19)	0.0414 (16)	−0.0049 (18)	−0.0030 (17)	−0.0042 (14)
C33	0.075 (2)	0.0524 (19)	0.072 (2)	−0.0125 (16)	0.0024 (17)	−0.0101 (16)
C43	0.074 (2)	0.093 (3)	0.057 (2)	0.022 (2)	−0.0118 (17)	−0.0258 (18)
C27	0.055 (2)	0.107 (3)	0.069 (2)	−0.001 (2)	0.0270 (17)	−0.015 (2)
C24	0.0506 (18)	0.085 (2)	0.0585 (19)	0.0097 (17)	0.0088 (15)	−0.0137 (17)
C28	0.057 (2)	0.107 (3)	0.088 (3)	0.007 (2)	0.019 (2)	−0.039 (2)
C39	0.068 (2)	0.071 (2)	0.066 (2)	0.0066 (17)	−0.0053 (16)	−0.0185 (17)
C2	0.0574 (19)	0.073 (2)	0.0520 (18)	−0.0008 (16)	−0.0031 (14)	0.0007 (15)
C3	0.076 (3)	0.085 (3)	0.072 (2)	−0.008 (2)	−0.022 (2)	−0.0021 (19)
C29	0.065 (2)	0.068 (2)	0.097 (3)	0.0170 (18)	0.006 (2)	−0.018 (2)
C23	0.0428 (18)	0.110 (3)	0.089 (3)	0.0174 (19)	0.0091 (17)	−0.007 (2)
C21	0.059 (2)	0.111 (3)	0.060 (2)	−0.005 (2)	−0.0114 (16)	−0.0200 (19)
C4	0.051 (2)	0.091 (3)	0.115 (4)	−0.007 (2)	−0.019 (2)	0.002 (2)
C10	0.101 (3)	0.065 (2)	0.076 (2)	0.036 (2)	0.040 (2)	0.0163 (18)
C41	0.064 (2)	0.128 (3)	0.057 (2)	0.006 (2)	−0.0167 (17)	−0.015 (2)

C11	0.058 (2)	0.082 (3)	0.128 (3)	0.022 (2)	0.031 (2)	0.020 (2)
C5	0.045 (2)	0.108 (3)	0.111 (3)	0.0058 (19)	0.017 (2)	0.005 (3)
C40	0.083 (3)	0.101 (3)	0.072 (2)	0.000 (2)	−0.020 (2)	−0.035 (2)
C42	0.082 (3)	0.130 (4)	0.071 (2)	0.043 (2)	−0.021 (2)	−0.027 (2)

Geometric parameters (Å, °)

Ag1—P2	2.4671 (6)	C30—H30	0.9300
Ag1—P1	2.4682 (7)	C6—C5	1.391 (5)
Ag1—S1	2.6015 (8)	C6—H6	0.9300
Ag1—Br1	2.7189 (3)	C35—C34	1.365 (4)
P2—C31	1.827 (3)	C35—H35	0.9300
P2—C19	1.829 (2)	C20—C21	1.381 (4)
P2—C25	1.830 (3)	C20—H20	0.9300
P1—C1	1.818 (3)	C16—C17	1.374 (4)
P1—C7	1.827 (3)	C16—H16	0.9300
P1—C13	1.828 (2)	C34—C33	1.364 (4)
S1—C37	1.687 (3)	C34—H34	0.9300
N1—C37	1.299 (4)	C12—C11	1.380 (5)
N1—H1A	0.852 (10)	C12—H12	0.9300
N1—H1B	0.854 (10)	C22—C21	1.358 (5)
C31—C32	1.383 (3)	C22—C23	1.361 (5)
C31—C36	1.386 (3)	C22—H22	0.9300
C13—C18	1.378 (4)	C17—H17	0.9300
C13—C14	1.384 (3)	C33—H33	0.9300
C19—C20	1.371 (4)	C43—C42	1.380 (5)
C19—C24	1.373 (4)	C43—H43	0.9300
C25—C26	1.377 (4)	C27—C28	1.361 (5)
C25—C30	1.385 (4)	C27—H27	0.9300
C1—C6	1.373 (4)	C24—C23	1.383 (4)
C1—C2	1.380 (4)	C24—H24	0.9300
C7—C8	1.382 (4)	C28—C29	1.372 (5)
C7—C12	1.383 (4)	C28—H28	0.9300
C38—C43	1.370 (4)	C39—C40	1.382 (4)
C38—C39	1.381 (4)	C39—H39	0.9300
C38—C37	1.479 (4)	C2—C3	1.380 (4)
C8—C9	1.393 (4)	C2—H2	0.9300
C8—H8	0.9300	C3—C4	1.355 (5)
C14—C15	1.372 (4)	C3—H3	0.9300
C14—H14	0.9300	C29—H29	0.9300
C26—C27	1.383 (4)	C23—H23	0.9300
C26—H26	0.9300	C21—H21	0.9300
C18—C17	1.384 (4)	C4—C5	1.357 (5)
C18—H18	0.9300	C4—H4	0.9300
C36—C35	1.385 (4)	C10—C11	1.363 (5)
C36—H36	0.9300	C10—H10	0.9300
C15—C16	1.363 (4)	C41—C42	1.353 (5)
C15—H15	0.9300	C41—C40	1.357 (5)

C32—C33	1.384 (4)	C41—H41	0.9300
C32—H32	0.9300	C11—H11	0.9300
C9—C10	1.352 (5)	C5—H5	0.9300
C9—H9	0.9300	C40—H40	0.9300
C30—C29	1.384 (4)	C42—H42	0.9300
P2—Ag1—P1	121.60 (2)	C34—C35—C36	120.6 (3)
P2—Ag1—S1	108.40 (2)	C34—C35—H35	119.7
P1—Ag1—S1	113.89 (3)	C36—C35—H35	119.7
P2—Ag1—Br1	104.528 (18)	C19—C20—C21	121.3 (3)
P1—Ag1—Br1	97.338 (18)	C19—C20—H20	119.3
S1—Ag1—Br1	109.550 (19)	C21—C20—H20	119.3
C31—P2—C19	102.34 (11)	C15—C16—C17	120.2 (3)
C31—P2—C25	105.36 (12)	C15—C16—H16	119.9
C19—P2—C25	104.08 (11)	C17—C16—H16	119.9
C31—P2—Ag1	113.69 (8)	C33—C34—C35	119.4 (3)
C19—P2—Ag1	117.29 (8)	C33—C34—H34	120.3
C25—P2—Ag1	112.76 (9)	C35—C34—H34	120.3
C1—P1—C7	105.53 (12)	C11—C12—C7	120.1 (3)
C1—P1—C13	104.41 (12)	C11—C12—H12	119.9
C7—P1—C13	102.34 (11)	C7—C12—H12	119.9
C1—P1—Ag1	110.20 (8)	C21—C22—C23	119.5 (3)
C7—P1—Ag1	116.31 (8)	C21—C22—H22	120.3
C13—P1—Ag1	116.78 (8)	C23—C22—H22	120.3
C37—S1—Ag1	109.90 (10)	C16—C17—C18	119.9 (3)
C37—N1—H1A	124 (2)	C16—C17—H17	120.1
C37—N1—H1B	118 (2)	C18—C17—H17	120.1
H1A—N1—H1B	117 (3)	C34—C33—C32	120.9 (3)
C32—C31—C36	118.4 (2)	C34—C33—H33	119.5
C32—C31—P2	118.3 (2)	C32—C33—H33	119.5
C36—C31—P2	123.2 (2)	C38—C43—C42	121.3 (3)
C18—C13—C14	118.7 (2)	C38—C43—H43	119.3
C18—C13—P1	118.69 (19)	C42—C43—H43	119.3
C14—C13—P1	122.6 (2)	C28—C27—C26	119.5 (3)
C20—C19—C24	118.1 (2)	C28—C27—H27	120.2
C20—C19—P2	124.1 (2)	C26—C27—H27	120.2
C24—C19—P2	117.8 (2)	C19—C24—C23	120.3 (3)
C26—C25—C30	118.1 (3)	C19—C24—H24	119.9
C26—C25—P2	124.8 (2)	C23—C24—H24	119.9
C30—C25—P2	117.1 (2)	C27—C28—C29	120.9 (3)
C6—C1—C2	118.3 (3)	C27—C28—H28	119.6
C6—C1—P1	121.9 (2)	C29—C28—H28	119.6
C2—C1—P1	119.5 (2)	C38—C39—C40	120.5 (3)
C8—C7—C12	118.5 (3)	C38—C39—H39	119.8
C8—C7—P1	125.1 (2)	C40—C39—H39	119.8
C12—C7—P1	116.3 (2)	C1—C2—C3	121.1 (3)
C43—C38—C39	117.8 (3)	C1—C2—H2	119.5
C43—C38—C37	120.7 (3)	C3—C2—H2	119.5

C39—C38—C37	121.5 (3)	C4—C3—C2	119.8 (4)
C7—C8—C9	120.2 (3)	C4—C3—H3	120.1
C7—C8—H8	119.9	C2—C3—H3	120.1
C9—C8—H8	119.9	C28—C29—C30	119.2 (3)
C15—C14—C13	120.8 (3)	C28—C29—H29	120.4
C15—C14—H14	119.6	C30—C29—H29	120.4
C13—C14—H14	119.6	C22—C23—C24	120.8 (3)
N1—C37—C38	117.5 (2)	C22—C23—H23	119.6
N1—C37—S1	121.9 (2)	C24—C23—H23	119.6
C38—C37—S1	120.6 (2)	C22—C21—C20	120.0 (3)
C25—C26—C27	121.2 (3)	C22—C21—H21	120.0
C25—C26—H26	119.4	C20—C21—H21	120.0
C27—C26—H26	119.4	C3—C4—C5	120.4 (3)
C13—C18—C17	120.3 (3)	C3—C4—H4	119.8
C13—C18—H18	119.9	C5—C4—H4	119.8
C17—C18—H18	119.9	C9—C10—C11	120.0 (3)
C35—C36—C31	120.4 (3)	C9—C10—H10	120.0
C35—C36—H36	119.8	C11—C10—H10	120.0
C31—C36—H36	119.8	C42—C41—C40	120.1 (3)
C16—C15—C14	120.0 (3)	C42—C41—H41	120.0
C16—C15—H15	120.0	C40—C41—H41	120.0
C14—C15—H15	120.0	C10—C11—C12	120.8 (3)
C31—C32—C33	120.2 (3)	C10—C11—H11	119.6
C31—C32—H32	119.9	C12—C11—H11	119.6
C33—C32—H32	119.9	C4—C5—C6	120.3 (4)
C10—C9—C8	120.3 (3)	C4—C5—H5	119.9
C10—C9—H9	119.8	C6—C5—H5	119.9
C8—C9—H9	119.8	C41—C40—C39	120.3 (3)
C29—C30—C25	121.0 (3)	C41—C40—H40	119.8
C29—C30—H30	119.5	C39—C40—H40	119.8
C25—C30—H30	119.5	C41—C42—C43	119.9 (3)
C1—C6—C5	120.2 (3)	C41—C42—H42	120.0
C1—C6—H6	119.9	C43—C42—H42	120.0
C5—C6—H6	119.9		
C19—P2—C31—C32	−145.9 (2)	P1—C13—C18—C17	−178.6 (2)
C25—P2—C31—C32	105.5 (2)	C32—C31—C36—C35	−2.1 (4)
Ag1—P2—C31—C32	−18.4 (2)	P2—C31—C36—C35	178.9 (2)
C19—P2—C31—C36	33.1 (2)	C13—C14—C15—C16	0.8 (5)
C25—P2—C31—C36	−75.4 (2)	C36—C31—C32—C33	2.1 (4)
Ag1—P2—C31—C36	160.6 (2)	P2—C31—C32—C33	−178.9 (2)
C1—P1—C13—C18	119.0 (2)	C7—C8—C9—C10	−1.0 (4)
C7—P1—C13—C18	−131.2 (2)	C26—C25—C30—C29	1.2 (4)
Ag1—P1—C13—C18	−3.0 (3)	P2—C25—C30—C29	−178.8 (2)
C1—P1—C13—C14	−60.7 (2)	C2—C1—C6—C5	−1.8 (5)
C7—P1—C13—C14	49.1 (2)	P1—C1—C6—C5	−175.4 (3)
Ag1—P1—C13—C14	177.32 (19)	C31—C36—C35—C34	0.7 (5)
C31—P2—C19—C20	−101.0 (3)	C24—C19—C20—C21	−0.8 (5)

C25—P2—C19—C20	8.6 (3)	P2—C19—C20—C21	179.1 (3)
Ag1—P2—C19—C20	133.9 (2)	C14—C15—C16—C17	−0.2 (5)
C31—P2—C19—C24	78.9 (2)	C36—C35—C34—C33	0.6 (5)
C25—P2—C19—C24	−171.6 (2)	C8—C7—C12—C11	0.3 (5)
Ag1—P2—C19—C24	−46.3 (3)	P1—C7—C12—C11	−178.6 (3)
C31—P2—C25—C26	24.4 (3)	C15—C16—C17—C18	0.0 (5)
C19—P2—C25—C26	−82.9 (2)	C13—C18—C17—C16	−0.5 (5)
Ag1—P2—C25—C26	148.9 (2)	C35—C34—C33—C32	−0.7 (5)
C31—P2—C25—C30	−155.6 (2)	C31—C32—C33—C34	−0.7 (5)
C19—P2—C25—C30	97.1 (2)	C39—C38—C43—C42	0.3 (6)
Ag1—P2—C25—C30	−31.0 (2)	C37—C38—C43—C42	−179.7 (3)
C7—P1—C1—C6	−129.4 (2)	C25—C26—C27—C28	0.3 (5)
C13—P1—C1—C6	−21.9 (3)	C20—C19—C24—C23	0.6 (5)
Ag1—P1—C1—C6	104.2 (2)	P2—C19—C24—C23	−179.2 (3)
C7—P1—C1—C2	57.0 (3)	C26—C27—C28—C29	−0.7 (6)
C13—P1—C1—C2	164.5 (2)	C43—C38—C39—C40	−0.5 (5)
Ag1—P1—C1—C2	−69.4 (2)	C37—C38—C39—C40	179.5 (3)
C1—P1—C7—C8	13.5 (3)	C6—C1—C2—C3	2.1 (5)
C13—P1—C7—C8	−95.4 (2)	P1—C1—C2—C3	175.9 (3)
Ag1—P1—C7—C8	136.0 (2)	C1—C2—C3—C4	−1.1 (5)
C1—P1—C7—C12	−167.7 (2)	C27—C28—C29—C30	1.3 (5)
C13—P1—C7—C12	83.3 (2)	C25—C30—C29—C28	−1.6 (5)
Ag1—P1—C7—C12	−45.2 (2)	C21—C22—C23—C24	−0.6 (6)
C12—C7—C8—C9	0.6 (4)	C19—C24—C23—C22	0.1 (6)
P1—C7—C8—C9	179.4 (2)	C23—C22—C21—C20	0.5 (6)
C18—C13—C14—C15	−1.3 (4)	C19—C20—C21—C22	0.2 (6)
P1—C13—C14—C15	178.4 (2)	C2—C3—C4—C5	−0.3 (6)
C43—C38—C37—N1	−154.4 (3)	C8—C9—C10—C11	0.4 (5)
C39—C38—C37—N1	25.6 (4)	C9—C10—C11—C12	0.6 (6)
C43—C38—C37—S1	27.4 (4)	C7—C12—C11—C10	−0.9 (6)
C39—C38—C37—S1	−152.6 (3)	C3—C4—C5—C6	0.6 (6)
Ag1—S1—C37—N1	19.3 (3)	C1—C6—C5—C4	0.5 (6)
Ag1—S1—C37—C38	−162.44 (19)	C42—C41—C40—C39	0.6 (7)
C30—C25—C26—C27	−0.5 (4)	C38—C39—C40—C41	0.1 (6)
P2—C25—C26—C27	179.5 (2)	C40—C41—C42—C43	−0.8 (7)
C14—C13—C18—C17	1.2 (4)	C38—C43—C42—C41	0.4 (7)

Hydrogen-bond geometry (Å, °)

Cg3 is the centroid of the C13—C18 ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots Br1 ⁱ	0.85 (1)	2.54 (1)	3.357 (3)	161 (3)
N1—H1B \cdots Br1	0.85 (1)	2.58 (1)	3.413 (3)	166 (3)
C17—H17 \cdots Br1 ⁱ	0.93	2.91	3.789 (3)	158
C22—H22 \cdots Cg3 ⁱⁱ	0.93	2.94	3.78 (3)	151

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x+1/2, -y+3/2, z+1/2$.