

## Crystal structure of bis(4-acetyl-anilinium) tetrachloridomercurate(II)

Manickam Thairiyaraja,<sup>a</sup> Arumugam Elangovan,<sup>b\*</sup> Ganesh Arivazhagan,<sup>c</sup> Kuthambalam Selvaraju<sup>a\*</sup> and Subbiah Thamotharan<sup>d\*</sup>

<sup>a</sup>PG & Research Department of Physics, Government Arts College, Ariyalur 621 713, India, <sup>b</sup>Department of Chemistry, Thiagarajar College, Madurai 625 009, India,

<sup>c</sup>Department of Physics, Thiagarajar College, Madurai 625 009, India, and

<sup>d</sup>Biomolecular Crystallography Laboratory, Department of Bioinformatics, School of Chemical and Biotechnology, SASTRA University, Thanjavur 613 401, India.

\*Correspondence e-mail: arumugam.elangovan@gmail.com, selsphy@yahoo.com, thamu@scbt.sastrau.edu

Received 16 November 2015; accepted 23 November 2015

Edited by M. Weil, Vienna University of Technology, Austria


The structure of the title salt,  $(C_8H_{10}NO)_2[HgCl_4]$ , is isotypic with that of the cuprate(II) and cobaltate(II) analogues. The asymmetric unit contains one 4-acetyl-anilinium cation and one half of a tetrachloridomercurate(II) anion (point group symmetry  $m$ ). The Hg–Cl distances are in the range 2.4308 (7)–2.5244 (11) Å and the Cl–Hg–Cl angles in the range of 104.66 (2)–122.94 (4)°, indicating a considerable distortion of the tetrahedral anion. In the crystal, cations are linked by an intermolecular N–H···O hydrogen-bonding interaction, leading to a  $C(8)$  chain motif with the chains extending parallel to the  $b$  axis. There is also a  $\pi$ – $\pi$  stacking interaction with a centroid-to-centroid distance of 3.735 (2) Å between neighbouring benzene rings along this direction. The anions lie between the chains and interact with the cations through intermolecular N–H···Cl hydrogen bonds, leading to the formation of a three-dimensional network structure.

**Keywords:** crystal structure; isotypism; mercury(II); hydrogen bonding.

**CCDC reference:** 1047866

### 1. Related literature

For the structures of the isotypic tetrachloridocuprate(II) and tetrachloridocobaltate(II) analogues, see: Elangovan *et al.* (2007) and Thairiyaraja *et al.* (2015), respectively.



### 2. Experimental

#### 2.1. Crystal data

$(C_8H_{10}NO)_2[HgCl_4]$	$V = 4208.1 (3) \text{ \AA}^3$
$M_r = 614.73$	$Z = 8$
Orthorhombic, $Cmce$	Mo $K\alpha$ radiation
$a = 19.9231 (6) \text{ \AA}$	$\mu = 7.84 \text{ mm}^{-1}$
$b = 15.3515 (6) \text{ \AA}$	$T = 293 \text{ K}$
$c = 13.7587 (5) \text{ \AA}$	$0.30 \times 0.25 \times 0.20 \text{ mm}$

#### 2.2. Data collection

Bruker SMART APEX CCD diffractometer	26289 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2004)	2988 independent reflections
$T_{\min} = 0.202$ , $T_{\max} = 0.303$	2152 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.031$

#### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.056$	$\Delta\rho_{\max} = 0.77 \text{ e \AA}^{-3}$
$S = 1.04$	$\Delta\rho_{\min} = -0.93 \text{ e \AA}^{-3}$
2988 reflections	
131 parameters	
3 restraints	

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N41–H41A···O1 <sup>i</sup>	0.90 (2)	1.92 (2)	2.792 (3)	162 (4)
N41–H41C···Cl2 <sup>ii</sup>	0.90 (2)	2.34 (2)	3.206 (3)	162 (3)
N41–H41B···Cl3 <sup>iii</sup>	0.89 (2)	2.49 (2)	3.326 (3)	158 (3)

Symmetry codes: (i)  $-x + \frac{1}{2}, y - \frac{1}{2}, z$ ; (ii)  $x + \frac{1}{2}, y, -z + \frac{3}{2}$ ; (iii)  $-x + \frac{1}{2}, -y + \frac{1}{2}, -z + 2$ .

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: coordinates taken from an isotypic compound; program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).

## Acknowledgements

ST is very grateful to the management of SASTRA University for infrastructural and financial support (Professor TRR fund).

Supporting information for this paper is available from the IUCr electronic archives (Reference: WM5243).

## References

- Bruker (2004). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Elangovan, A., Thamaraichelvan, A., Ramu, A., Athimoolam, S. & Natarajan, S. (2007). *Acta Cryst. E* **63**, m224–m226.
- Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Thairiyaraja, M., Elangovan, A., Shanmugam, R., Selvaraju, K. & Thamotharan, S. (2015). *Acta Cryst. E* **71**, m221–m222.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

# supporting information

*Acta Cryst.* (2015). E71, m236–m237 [https://doi.org/10.1107/S2056989015022355]

## Crystal structure of bis(4-acetylaniinium) tetrachloridomercurate(II)


**Manickam Thairiyaraja, Arumugam Elangovan, Ganesh Arivazhagan, Kuthambalam Selvaraju and Subbiah Thamotharan**

### S1. Synthesis and crystallization


A solution of 4-aminoacetophenone (20 mmol) in 2 ml of HCl and deionized water (10 ml) was added to a 10 ml solution of  $\text{HgCl}_2$  (10 mmol). The resulting solution was concentrated and kept unperturbed at ambient temperature for crystallization. Single crystals suitable for X-ray diffraction were obtained within a week.

### S2. Refinement

Since the title salt is isotopic with its tetrachloridocobaltate and tetrachloridocuprate analogues, it was refined with the coordinates of the tetrachloridocobaltate salt (Thairiyaraja *et al.*, 2015) as a starting model. The ammonium H atoms were located from a difference Fourier map and were refined with a distance restraint of N—H = 0.89 (2) Å. The methyl H atoms were constrained to an ideal geometry (C—H = 0.96 Å) with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ , but were allowed to rotate freely about the C—C bond. The remaining H atoms were positioned geometrically and refined using a riding model approximation with C—H = 0.93 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . At this stage, the maximum residual electron density of 0.77 e Å<sup>-3</sup> suggested the presence of a possible atom site at Wyckoff position 4a at a distance of 2.88 Å near atom H5. This electron density was assumed to be the O atom of a water molecule and was refined with isotropic displacement parameters. However, the resultant model had higher reliability factors and a very high isotropic atomic displacement parameter for this O atom. As a consequence, this water O atom was not included in the final model.

**Figure 1**

The molecular components of the title salt, showing displacement ellipsoids at the 50% probability level. [Symmetry code: (i)  $-x, y, z$ .]

**Figure 2**

The crystal packing of the title salt viewed along the  $a$  axis. Hydrogen bonds are shown as dashed lines; H atoms bound to C were omitted for clarity.

### Bis(4-acetylanilinium) tetrachloridomercurate(II)

#### *Crystal data*



$M_r = 614.73$

Orthorhombic,  $Cmce$

$a = 19.9231 (6)$  Å

$b = 15.3515 (6)$  Å

$c = 13.7587 (5)$  Å

$V = 4208.1 (3)$  Å<sup>3</sup>

$Z = 8$

$F(000) = 2352$

$D_x = 1.941 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 9224 reflections

$\theta = 2.2\text{--}29.3^\circ$

$\mu = 7.84 \text{ mm}^{-1}$

$T = 293$  K

Block, orange

$0.30 \times 0.25 \times 0.20$  mm

#### *Data collection*

Bruker SMART APEX CCD  
diffractometer

Radiation source: fine-focus sealed tube

$\omega$  and  $\varphi$  scan

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2004)

$T_{\min} = 0.202$ ,  $T_{\max} = 0.303$

26289 measured reflections

2988 independent reflections

2152 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 29.4^\circ$ ,  $\theta_{\min} = 2.2^\circ$

$h = -27 \rightarrow 27$

$k = -19 \rightarrow 21$

$l = -18 \rightarrow 19$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.023$$

$$wR(F^2) = 0.056$$

$$S = 1.04$$

2988 reflections

131 parameters

3 restraints

Primary atom site location: isomorphous  
structure methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent  
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0217P)^2 + 6.8872P]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.77 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.93 \text{ e \AA}^{-3}$$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O11	0.18874 (11)	0.60387 (14)	0.87716 (17)	0.0544 (6)
N41	0.37059 (13)	0.26610 (16)	0.9093 (2)	0.0430 (6)
H41A	0.359 (2)	0.2125 (16)	0.889 (3)	0.084 (14)*
H41B	0.3848 (19)	0.270 (2)	0.9703 (16)	0.072 (13)*
H41C	0.4057 (15)	0.276 (3)	0.870 (2)	0.067 (12)*
C1	0.22199 (12)	0.45798 (17)	0.87221 (17)	0.0297 (5)
C2	0.20483 (13)	0.37068 (17)	0.86957 (19)	0.0359 (6)
H2	0.1603	0.3546	0.8606	0.043*
C3	0.25367 (13)	0.30686 (17)	0.8803 (2)	0.0373 (6)
H3	0.2423	0.2481	0.8784	0.045*
C4	0.31897 (12)	0.33222 (17)	0.89352 (19)	0.0321 (5)
C5	0.33780 (13)	0.41877 (18)	0.8947 (2)	0.0374 (6)
H5	0.3825	0.4346	0.9029	0.045*
C6	0.28863 (13)	0.48110 (18)	0.88350 (19)	0.0349 (6)
H6	0.3005	0.5397	0.8835	0.042*
C11	0.17064 (14)	0.52843 (18)	0.86546 (19)	0.0341 (6)
C12	0.09920 (13)	0.5071 (2)	0.8458 (2)	0.0482 (8)
H12A	0.0742	0.5600	0.8372	0.072*
H12B	0.0811	0.4750	0.8996	0.072*
H12C	0.0962	0.4725	0.7878	0.072*
Hg1	0.0000	0.24841 (2)	0.85591 (2)	0.04734 (7)
Cl1	0.0000	0.34199 (7)	1.00252 (8)	0.0479 (2)
Cl2	0.0000	0.33901 (7)	0.70279 (8)	0.0455 (2)
Cl3	0.10720 (4)	0.17292 (6)	0.86099 (7)	0.0607 (2)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O11	0.0515 (12)	0.0266 (12)	0.0850 (17)	0.0019 (9)	-0.0071 (11)	-0.0038 (10)

N41	0.0335 (11)	0.0344 (16)	0.061 (2)	0.0014 (9)	-0.0018 (11)	0.0055 (12)
C1	0.0341 (12)	0.0274 (14)	0.0275 (14)	-0.0026 (9)	0.0008 (10)	-0.0008 (10)
C2	0.0309 (12)	0.0321 (15)	0.0448 (16)	-0.0053 (10)	-0.0016 (11)	-0.0005 (11)
C3	0.0361 (13)	0.0236 (14)	0.0523 (18)	-0.0041 (10)	-0.0008 (12)	0.0023 (11)
C4	0.0331 (12)	0.0269 (14)	0.0365 (14)	0.0009 (10)	-0.0009 (10)	0.0030 (11)
C5	0.0326 (12)	0.0351 (16)	0.0445 (16)	-0.0062 (11)	-0.0037 (11)	0.0013 (12)
C6	0.0387 (14)	0.0248 (14)	0.0410 (15)	-0.0054 (10)	-0.0025 (11)	-0.0022 (11)
C11	0.0387 (13)	0.0299 (15)	0.0336 (14)	0.0010 (10)	0.0017 (11)	-0.0014 (11)
C12	0.0359 (13)	0.0407 (17)	0.068 (2)	0.0042 (12)	0.0015 (14)	-0.0065 (15)
Hg1	0.03172 (8)	0.05938 (13)	0.05092 (11)	0.000	0.000	-0.00393 (9)
Cl1	0.0538 (5)	0.0448 (6)	0.0453 (6)	0.000	0.000	-0.0061 (5)
Cl2	0.0420 (5)	0.0490 (6)	0.0455 (6)	0.000	0.000	-0.0018 (5)
Cl3	0.0383 (4)	0.0555 (5)	0.0883 (6)	0.0145 (3)	-0.0098 (4)	-0.0202 (4)

Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )

O11—C11	1.224 (3)	C4—C5	1.381 (4)
N41—C4	1.461 (3)	C5—C6	1.378 (4)
N41—H41A	0.901 (19)	C5—H5	0.9300
N41—H41B	0.887 (19)	C6—H6	0.9300
N41—H41C	0.899 (19)	C11—C12	1.486 (4)
C1—C6	1.383 (4)	C12—H12A	0.9600
C1—C2	1.384 (4)	C12—H12B	0.9600
C1—C11	1.492 (4)	C12—H12C	0.9600
C2—C3	1.389 (4)	Hg1—Cl3 <sup>i</sup>	2.4308 (7)
C2—H2	0.9300	Hg1—Cl3	2.4308 (7)
C3—C4	1.370 (4)	Hg1—Cl1	2.4764 (11)
C3—H3	0.9300	Hg1—Cl2	2.5244 (11)
C4—N41—H41A	114 (3)	C4—C5—H5	120.9
C4—N41—H41B	109 (2)	C5—C6—C1	121.1 (2)
H41A—N41—H41B	116 (3)	C5—C6—H6	119.4
C4—N41—H41C	110 (3)	C1—C6—H6	119.4
H41A—N41—H41C	100 (3)	O11—C11—C12	121.0 (3)
H41B—N41—H41C	108 (3)	O11—C11—C1	118.4 (2)
C6—C1—C2	119.3 (2)	C12—C11—C1	120.6 (2)
C6—C1—C11	118.6 (2)	C11—C12—H12A	109.5
C2—C1—C11	122.1 (2)	C11—C12—H12B	109.5
C1—C2—C3	120.5 (2)	H12A—C12—H12B	109.5
C1—C2—H2	119.8	C11—C12—H12C	109.5
C3—C2—H2	119.8	H12A—C12—H12C	109.5
C4—C3—C2	118.6 (2)	H12B—C12—H12C	109.5
C4—C3—H3	120.7	Cl3 <sup>i</sup> —Hg1—Cl3	122.94 (4)
C2—C3—H3	120.7	Cl3 <sup>i</sup> —Hg1—Cl1	104.66 (2)
C3—C4—C5	122.2 (2)	Cl3—Hg1—Cl1	104.66 (2)
C3—C4—N41	119.4 (2)	Cl3 <sup>i</sup> —Hg1—Cl2	106.66 (3)
C5—C4—N41	118.4 (2)	Cl3—Hg1—Cl2	106.66 (3)
C6—C5—C4	118.3 (2)	Cl1—Hg1—Cl2	111.11 (4)

C6—C5—H5	120.9		
C6—C1—C2—C3	1.4 (4)	C4—C5—C6—C1	0.6 (4)
C11—C1—C2—C3	−177.2 (2)	C2—C1—C6—C5	−1.8 (4)
C1—C2—C3—C4	0.1 (4)	C11—C1—C6—C5	176.9 (2)
C2—C3—C4—C5	−1.3 (4)	C6—C1—C11—O11	−5.2 (4)
C2—C3—C4—N41	177.3 (3)	C2—C1—C11—O11	173.4 (3)
C3—C4—C5—C6	0.9 (4)	C6—C1—C11—C12	175.3 (2)
N41—C4—C5—C6	−177.6 (3)	C2—C1—C11—C12	−6.1 (4)

Symmetry code: (i)  $-x, y, z$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N41—H41A $\cdots$ O11 <sup>ii</sup>	0.90 (2)	1.92 (2)	2.792 (3)	162 (4)
N41—H41C $\cdots$ Cl2 <sup>iii</sup>	0.90 (2)	2.34 (2)	3.206 (3)	162 (3)
N41—H41B $\cdots$ Cl3 <sup>iv</sup>	0.89 (2)	2.49 (2)	3.326 (3)	158 (3)

Symmetry codes: (ii)  $-x+1/2, y-1/2, z$ ; (iii)  $x+1/2, y, -z+3/2$ ; (iv)  $-x+1/2, -y+1/2, -z+2$ .