

Original Paper

# Serum Cortisol-to-Cortisone Ratio and Blood Pressure in Severe Obesity before and after Weight Loss

James B. Byrd<sup>a</sup> Amy E. Rothberg<sup>b, c</sup> Robert Chomic<sup>d</sup>  
Charles F. Burant<sup>b–e</sup> Robert D. Brook<sup>a</sup> Richard J. Auchus<sup>b, d, f</sup>

Divisions of <sup>a</sup>Cardiovascular Medicine and <sup>b</sup>Metabolism, Endocrinology and Diabetes, Department of Internal Medicine, <sup>c</sup>Weight Management Program, <sup>d</sup>Metabolomics Core, and Departments of <sup>e</sup>Molecular and Integrative Physiology and <sup>f</sup>Pharmacology, University of Michigan, Ann Arbor, Mich., USA

## Key Words

Obesity · Hypertension · Cortisol · Cortisone · Adrenal steroids

## Abstract

**Background/Aims:** The pathogenesis of obesity-associated hypertension is poorly understood. Serum cortisol-to-cortisone ratio (F/E ratio) is a marker of cortisol metabolism. Our objective was to determine whether the serum F/E ratio is associated with blood pressure (BP) in patients after significant weight loss ( $\geq 15\%$  from baseline weight). **Methods:** Sera from 43 nondiabetic, severely obese males participating in a weight management program were assayed for F and E by mass spectrometry. We assessed whether changes in the F/E ratio accompanying weight loss correlate with changes in the systolic (SBP) and diastolic BP (DBP). Linear regression was used to evaluate change in the F/E ratio as a predictor of change in BP. **Results:** The body mass index decreased from  $40.8 \pm 5.6$  to  $33.7 \pm 4.8$  ( $p < 0.001$ ); also, SBP ( $133.2 \pm 13.8$  vs.  $124.1 \pm 14.3$  mm Hg;  $p < 0.001$ ) and DBP ( $69.8 \pm 8.0$  vs.  $66.6 \pm 9.4$  mm Hg;  $p = 0.026$ ) decreased during the study. The baseline F/E ratio tended to associate with baseline DBP (Spearman's  $r = -0.29$ ,  $p = 0.06$ ), and change in the serum F/E ratio correlated with change in DBP (Spearman's  $r = -0.32$ ,  $p = 0.036$ ). Change in the F/E ratio also tended to associate with change in SBP (Spearman's  $r = -0.27$ ,  $p = 0.08$ ). A multiple linear regression model adjusted for change in the F/E ratio and age explained 22% of the variance in SBP change ( $R^2 = 0.22$ ,  $p = 0.007$ ). Change in the F/E ratio independently predicted change in SBP ( $p = 0.036$ ). **Conclusion:** In our sample of nondiabetic, severely obese males, change in the serum F/E ratio was associated with change in BP after weight loss.

© 2015 S. Karger AG, Basel

James Brian Byrd, MD, MS  
Room 20-209W, North Campus Research Complex  
University of Michigan, 2800 Plymouth Road  
Ann Arbor, MI 48109-2800 (USA)  
E-Mail jbbbyrd@med.umich.edu

## Introduction

Obesity is epidemic in the United States [1] and in many other regions of the world. It is strongly associated with hypertension [2], and epidemiologic and clinical studies have demonstrated that as little as 5% weight loss lowers blood pressure (BP) [3, 4]. Obesity-associated hypertension (OAH) arises from complex mechanisms, one of which might involve dysregulation of adrenal steroid production, metabolism, or responsiveness [5]. In particular, heightened cortisol activity is a plausible candidate mechanism in the pathophysiology of OAH.

In Cushing's syndrome, excess cortisol activates the mineralocorticoid receptor (MR), increases BP, and contributes to abdominal obesity. However, even normal amounts of cortisol mediate hypertension in some contexts. 11 $\beta$ -Hydroxysteroid dehydrogenase type 2 (11 $\beta$ HSD2) is an enzyme found in mineralocorticoid target tissues, including epithelial cells of the aldosterone-sensitive distal nephron [6, 7]. 11 $\beta$ HSD2 acts principally as a dehydrogenase to convert cortisol, which binds MR with high affinity, to cortisone, which does not [8]. The importance of 11 $\beta$ HSD2 in regulating BP is demonstrated in the autosomal recessive form of hypertension called the 'apparent mineralocorticoid excess syndrome' (AME). In AME, mutations in the cognate *HSD11B2* gene severely impair 11 $\beta$ HSD2 activity, and individuals with AME have the hypertensive, hypokalemic phenotype typical of primary aldosteronism. In AME, however, aldosterone levels are low, and cortisol activates MR instead [9]. Copious ingestion of authentic licorice causes an acquired AME condition [10]. In contrast to 11 $\beta$ HSD2, 11 $\beta$ HSD1 acts predominantly in the opposite direction, reducing cortisone to cortisol in the liver, adipose, and other tissues. Previous studies have suggested altered 11 $\beta$ HSD activity in obesity [11–13]. Because of the complex interplay of these two isoenzymes in cortisol metabolism throughout the body, the contribution of 11 $\beta$ HSD isoforms to OAH remains poorly understood.

The serum cortisol-to-cortisone (F/E) ratio is a marker of net 11 $\beta$ HSD activity [14], which changes in some settings, such as inflammatory conditions [15] or after the administration of the 11 $\beta$ HSD2 inhibitor found in licorice [14]. If dysregulated cortisol metabolism contributes to OAH, then the F/E ratio might correlate with BP in obese patients, and changes in the F/E ratio might parallel BP reductions during weight loss. Previous work showed that obesity alters 11 $\beta$ HSD2 and 11 $\beta$ HSD1 expression in human adipose tissue [13], and weight loss reverses these changes [11]. It is unknown whether the serum F/E ratio correlates with a change in BP after weight loss. Therefore, we measured the serum F/E ratio using mass spectrometry in severely obese patients who enrolled in a comprehensive weight management program in which participants lost  $\geq 15\%$  of weight from baseline weight. We evaluated whether the baseline body mass index (BMI) was associated with the baseline serum F/E ratio and evaluated whether weight loss was accompanied by a change in the serum F/E ratio. Finally, we explored the association between the serum F/E ratio and BP, and changes in each after weight loss.

## Patients and Methods

### Study Population

The study was reviewed and approved by the University of Michigan Institutional Review Board. The University of Michigan's Weight Management Program (MWMP) is a 2-year intensive, multidisciplinary behavioral intervention for obesity. Program participants are given the opportunity to 'opt in' to the program's research component, which includes both baseline and interval testing (hereafter, 'phenotyping'). Procedures performed included (but were not limited to) metabolic testing (resting energy expenditure and oxidative capacity by  $\text{VO}_2\text{max}$ ); body composition by dual-energy X-ray absorptiometry; mixed-meal

tolerance testing with profiling of the metabolome and lipidome, and the measurement of steroid hormones and adipokines.

We queried the MWMP participant database for all adult male patients without diabetes mellitus. This subset of patients was selected to avoid confounding influences of female reproductive hormones and diabetes. The 22 MWMP participants who were missing a ‘phenotyping’ session were excluded. The details of the weight management protocol have been published previously [16]. Briefly, participants in the program consumed a very low-calorie diet (800 kcal/day) for the initial 12 weeks and were asked to increase their physical activity to 40 min daily at mild-to-moderate intensity. During the baseline visit, the participants’ demographics, comorbid and other health conditions, current medications, vital signs, and anthropometric measurements (hip circumference, waist circumference, height, weight, and BMI) were recorded. In addition, sera were obtained from those who consented to the research. At a follow-up visit with the MWMP approximately 4 months after the baseline, repeat measurements of the above and sera were obtained.

#### *Cortisol and Cortisone Assays*

Blood for biochemical analysis was collected in the absence of anticoagulant therapy, centrifuged immediately after clotting, and the sera were stored at  $-80^{\circ}\text{C}$  until analysis. In keeping with best practices in the measurement of adrenal steroids [17], we measured serum cortisol and cortisone using liquid chromatography-tandem mass spectrometry as described previously [18]. Serum (0.2 ml) was mixed with deuterium-labeled internal standards and deproteinated with methanol and acetonitrile. The steroids were extracted with methyl-*tert*-butylether and concentrated under nitrogen. The residue was resuspended in 50% aqueous methanol, injected into the Agilent 1290 HPLC with 6490 QQQ tandem mass spectrometer, and resolved on a Kinetex  $50 \times 2.1$  mm biphenyl column (Phenomenex, Torrance, Calif, USA) with methanol-water gradients. Cortisol and cortisone were quantitated by multiple reaction monitoring in positive ion mode using specific mass fragments derived from authentic standards and peak integrals interpolated from a standard curve with internal standard recovery correction.

#### *Statistical Methods*

Variables were evaluated for normality using the Shapiro-Wilk test. Data are presented as the mean  $\pm$  SD if normally distributed, or as the median  $\pm$  interquartile range if the Shapiro-Wilk test suggested departure from a normal distribution. We calculated the difference between the post- and preweight loss F ( $\Delta\text{F}$ ), E ( $\Delta\text{E}$ ), F/E ratio [ $\Delta(\text{F/E ratio})$ ], waist-hip ratio [ $\Delta(\text{waist-hip ratio})$ ], BMI ( $\Delta\text{BMI}$ ), systolic BP ( $\Delta\text{SBP}$ ), and diastolic BP ( $\Delta\text{DBP}$ ). The relationship between continuous variables was analyzed using the Pearson correlation coefficient, or the Spearman correlation coefficient, as appropriate. Comparison of means before and after weight loss was performed using the paired *t* test or Wilcoxon’s signed-rank test, as appropriate. The independent effects of age and  $\Delta(\text{F/E ratio})$  on  $\Delta\text{SBP}$  and  $\Delta\text{DBP}$  (model 1) were analyzed using linear regression. In an additional analysis, we modeled the independent effects of age and  $\Delta(\text{F/E ratio})$ , adjusting for  $\Delta\text{BMI}$  (model 2). A 2-sided  $p < 0.05$  was considered significant. Data were analyzed using SPSS version 22.

## Results

Of 65 screened participants, the 43 with complete data sets were included in the analysis (table 1). By design, all participants were nondiabetic males. At baseline, 67% of the participants had hypertension. Fourteen participants were not taking an antihypertensive medication at baseline, 20 participants’ antihypertensive medications were continued throughout the follow-up period, and 9 participants stopped taking one or more antihypertensive medications during follow-up. During the baseline assessment, participants’ antihypertensive medication included ACE inhibitors ( $n = 22$ ),  $\beta$ -blockers ( $n = 12$ ), diuretics ( $n = 16$ ), calcium channel blockers ( $n = 9$ ), angiotensin receptor blockers ( $n = 7$ ), or triamterene ( $n = 2$ ) in some cases. No participant was taking an MR antagonist, and no participant had been diagnosed with primary aldosteronism, Cushing’s syndrome, or adrenal insufficiency.

At baseline, F, but not E or the F/E ratio, was higher in participants with hypertension compared to those without hypertension ( $29.9 \pm 17.8$  vs.  $19.7 \pm 52.1$   $\mu\text{g/dl}$ ;  $p = 0.038$ , Mann-Whitney U test). By Spearman correlation analysis, baseline BMI did not correlate with

**Table 1.** Characteristics of the study participants (n = 43)

	Before	After	p
Age, years (median ± IQR)	53.0±9.0		N/A
Race/ethnicity, n (%)			
Non-Hispanic white	40 (93)		N/A
Other	3 (7)		N/A
Hypertension at baseline, n (%)	29 (67)		N/A
BMI (median ± IQR)	39.4±8	32.1±5.9	<0.001*
Hip circumference, cm (mean ± SD)	127.0±14.4	113.9±11.9	<0.001
Waist circumference, cm (mean ± SD)	127.5±13.6	110.5±13.3	<0.001
Waist-hip ratio (mean ± SD)	1.0±0.04	0.97±0.05	<0.001
Weight, kg (median ± IQR)	127.9±23.8	105.1±16.4	<0.001*
Systolic BP, mm Hg (mean ± SD)	133±13	124±14	<0.001
Diastolic BP, mm Hg (mean ± SD)	69.0±15.0	64.0±12.0	0.021*
Heart rate, bpm (mean ± SD)	77.5±11.8	63.1±11.7	<0.001
Serum cortisol, µg/dl (mean ± SD)	22.4±13.6	25.9±10.8	0.88*
Serum cortisone, µg/dl (mean ± SD)	4.61±2.10	4.04±2.12	0.93*
F/E ratio (mean ± SD)	5.07±2.42	5.43±2.58	0.96*

IQR = Interquartile range.

\* Wilcoxon signed-rank test.

baseline F ( $r = -0.15$ ,  $p = 0.34$ ), E ( $r = -0.14$ ,  $p = 0.39$ ), or F/E ratio ( $r = -0.13$ ,  $p = 0.039$ ). The baseline F/E ratio did not correlate with baseline SBP (Spearman's  $r = -0.21$ ,  $p = 0.18$ ), but there was a trend toward an association with baseline DBP (Spearman's  $r = -0.29$ ,  $p = 0.06$ ).

#### *Changes with Weight Loss*


After a mean follow-up time of 120 days, BMI as well as SBP and DBP decreased significantly (table 1). In univariate analysis, the mean F/E ratio did not change with weight loss. By Spearman correlation analysis,  $\Delta$ BMI did not associate with  $\Delta$ F ( $r = 0.14$ ,  $p = 0.34$ ),  $\Delta$ E ( $r = 0.07$ ,  $p = 0.65$ ), or  $\Delta$ (F/E ratio) ( $r = 0.02$ ,  $p = 0.92$ ). Nor did  $\Delta$ (waist-hip ratio) associate with  $\Delta$ F ( $r = 0.13$ ,  $p = 0.39$ ),  $\Delta$ E ( $r = 0.08$ ,  $p = 0.62$ ), or  $\Delta$ (F/E ratio) ( $r = -0.01$ ,  $p = 0.94$ ).

We evaluated whether baseline F, E, or the F/E ratio or change in these parameters after weight loss was associated with change in BP after weight loss. Baseline cortisol correlated with  $\Delta$ SBP (Spearman's  $r = 0.32$ ,  $p = 0.034$ ) and  $\Delta$ DBP (Spearman's  $r = 0.32$ ,  $p = 0.039$ ). However, baseline cortisone did not correlate with either  $\Delta$ SBP (Spearman's  $r = 0.22$ ,  $p = 0.15$ ) or  $\Delta$ DBP (Spearman's  $r = 0.23$ ,  $p = 0.13$ ). The baseline F/E ratio tended to correlate with  $\Delta$ DBP (Spearman's  $r = 0.27$ ,  $p = 0.08$ ), but not with  $\Delta$ SBP (Spearman's  $r = 0.24$ ,  $p = 0.13$ ).  $\Delta$ F tended to inversely correlate with  $\Delta$ SBP (Spearman's  $r = -0.30$ ,  $p = 0.054$ ) and showed a weak tendency to inversely correlate with  $\Delta$ DBP (Spearman's  $r = -0.26$ ,  $p = 0.087$ ).  $\Delta$ E was not associated with  $\Delta$ SBP (Spearman's  $r = -0.20$ ,  $p = 0.90$ ) or  $\Delta$ DBP (Spearman's  $r = -0.86$ ,  $p = 0.58$ ). The  $\Delta$ (F/E ratio) inversely correlated with  $\Delta$ DBP (Spearman's  $r = -0.32$ ,  $p = 0.036$ ; Pearson's  $R = -0.21$ ,  $p = 0.17$ ; fig. 1), and tended to inversely correlate with  $\Delta$ SBP (Spearman's  $r = -0.27$ ,  $p = 0.076$ ).

Using multiple linear regression, model 1 (adjusting for age; table 2) explained 22% of the variance in  $\Delta$ SBP ( $R^2 = 0.22$ ,  $p = 0.007$  for the overall model). Model 2 (adjusting for age and change in BMI) provided similar results for  $\Delta$ SBP ( $R^2 = 0.22$ ,  $p = 0.021$  for the overall model). Neither model 1 nor model 2 reached statistical significance for  $\Delta$ DBP (model 1,  $p = 0.08$ ; model 2,  $p = 0.14$ ).  $\Delta$ (F/E ratio) independently predicted  $\Delta$ SBP (model 1,  $p = 0.036$ ; model 2,  $p = 0.039$ ) but not  $\Delta$ DBP (models 1 and 2,  $p = 0.10$ ).

**Table 2.** Association between change in the serum F/E ratio and change in BP after weight loss

	Model 1 (adjusted for age)		Model 2 (adjusted for age and change in BMI)	
	$\beta$ (95% CI)	p	$\beta$ (95% CI)	p
$\Delta$ SBP	-2.204 (-4.259 to -0.150)	0.036	-2.206 (-4.28 to -0.123)	0.039
$\Delta$ DBP	-1.139 (-2.491 to 0.213)	0.096	-1.128 (-2.491 to 0.236)	0.236

**Fig. 1.** The figure shows the  $\Delta$ F/E ratio versus  $\Delta$ SBP and  $\Delta$ DBP.

## Discussion

In severely obese patients participating in a weight management program, F was higher at baseline in hypertensive compared to normotensive participants. The baseline F/E ratio tended to associate with baseline DBP. Change in the F/E ratio after weight loss correlated with change in DBP. In addition, change in the F/E ratio after weight loss predicted change in SBP after adjustment for age, or after adjustment for age and change in BMI. These findings provide additional support for a contribution of adrenal steroids to the relationship between obesity and elevated BP.

Emerging evidence suggests an important role for MR in at least one common obesity-associated cardiovascular phenotype [19]. Increased activation of MR by adrenal steroids has been hypothesized to contribute to OAH [5, 11]. Elevated aldosterone has been associated with visceral obesity in women [20] and with hypertension and the metabolic syndrome in black Americans [21]. In a smaller study of 12 patients with a lower average BMI than in the current study, the serum F/E ratio increased after significant weight loss [11]. The relationship between F/E ratio and BP was not evaluated in that study. In keeping with our findings, a recent study found no change in the serum F/E ratio after weight loss due to bariatric surgery [12]. Our findings that the  $\Delta$ (F/E ratio) correlates with  $\Delta$ SBP and independently predicts changes in BP after weight loss support the hypothesis that altered  $11\beta$ HSD activity contributes to elevated BP in obesity. The direction of the change in SBP with respect

to the change in the F/E ratio was surprising. Additional investigation will be required to further clarify the relationship between  $11\beta$ HSD enzymes and blood pressure in obesity.

This study should be interpreted in view of certain limitations. The study was designed as a pilot study with a relatively small number of participants, and our findings await confirmation in a larger population. In addition, the participants were selected from a larger population. Thus, the possibility of selection bias exists. We studied only men, so we do not know whether similar results would have been observed in women. Some of the variables were not normally distributed. Although we made an effort to minimize the impact of the non-normality (such as using nonparametric statistics for many analyses), it is possible that this limitation impacted our results. Model 2 may be prone to ‘overadjustment’ if BMI change is a mediator of F, E, or F/E change. However, we also included model 1 for comparison. In some instances, participants were treated with antihypertensive medications throughout the study, and other participants underwent changes to their antihypertensive regimen prior to the follow-up visit. Cardiomyocytes express MR but not  $11\beta$ HSD2, and glucocorticoids activate MR in the heart in an animal model [22]. Therefore, changes in the serum F/E ratio may have different implications for cardiac tissue compared to other MR-expressing tissues involved in BP regulation, such as the distal nephron. In addition, this was an observational study about hormones and BP, a dynamically changing emergent trait regulated via many factors. Therefore, conclusions about a causal relationship between changes in the F/E ratio and BP after weight loss must be drawn cautiously.

Obesity and hypertension are widespread and interrelated problems. Further insights into the adrenal contribution to OAH offer the potential to tailor therapy for this common condition. For example, if our findings are confirmed in larger studies, MR antagonists would warrant close investigation for the treatment of OAH. In addition, the role of the glucocorticoid receptor in the regulation of BP merits additional investigation.

## Acknowledgement

We thank Dr. Heidi IglayRager for her invaluable assistance with the laboratory evaluation of the samples. Funding for this study was provided by the University of Michigan McKay award (Dr. Byrd, PI), the Michigan Institute for Clinical and Health Research Translational Research Award (Dr. Byrd, PI), the University of Michigan Cardiovascular Center Inaugural Award (Dr. Byrd, PI), and the A. Alfred Taubman Research Institute (Dr. Burant). This work utilized Core Services of the Michigan Nutrition Obesity Research Center (P30 DK089503).

## Disclosure Statement

The authors have no conflicts of interest to disclose.

## References

- 1 Flegal KM, Carroll MD, Kit BK, Ogden CL: Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA* 2012;307:483–490.
- 2 Wilson PW, D’Agostino RB, Sullivan L, Parise H, Kannel WB: Overweight and obesity as determinants of cardiovascular risk: the Framingham experience. *Arch Intern Med* 2002;162:1867–1872.
- 3 Orchard TJ, Temprosa M, Barrett-Connor E, Fowler SE, Goldberg RB, Mather KJ, Marcovina SM, Montez M, Ratner RE, Saudek CD, Sherif H, Watson KE: Long-term effects of the Diabetes Prevention Program interventions on cardiovascular risk factors: a report from the DPP Outcomes Study. *Diabet Med* 2013;30:46–55.
- 4 Wing RR, Bolin P, Brancati FL, Bray GA, et al: Cardiovascular effects of intensive lifestyle intervention in type 2 diabetes. *N Engl J Med* 2013;369:145–154.



- 5 Byrd JB, Brook RD: A critical review of the evidence supporting aldosterone in the etiology and its blockade in the treatment of obesity-associated hypertension. *J Hum Hypertens* 2014;28:3–9.
- 6 Albiston AL, Obeyesekere VR, Smith RE, Krozowski ZS: Cloning and tissue distribution of the human 11beta-hydroxysteroid dehydrogenase type 2 enzyme. *Mol Cell Endocrinol* 1994;105:R11–R17.
- 7 Walker BR, Campbell JC, Williams BC, Edwards CR: Tissue-specific distribution of the NAD(+)-dependent isoform of 11beta-hydroxysteroid dehydrogenase. *Endocrinology* 1992;131:970–972.
- 8 Funder JW, Pearce PT, Smith R, Smith AI: Mineralocorticoid action: target tissue specificity is enzyme, not receptor, mediated. *Science* 1988;242:583–585.
- 9 Stewart PM, Corrie JE, Shackleton CH, Edwards CR: Syndrome of apparent mineralocorticoid excess. A defect in the cortisol-cortisone shuttle. *J Clin Invest* 1988;82:340–349.
- 10 Conn JW, Rovner DR, Cohen EL: Licorice-induced pseudoaldosteronism. Hypertension, hypokalemia, aldosteronopenia, and suppressed plasma renin activity. *JAMA* 1968;205:492–496.
- 11 Tomlinson JW, Moore JS, Clark PM, Holder G, Shakespeare L, Stewart PM: Weight loss increases 11beta-hydroxysteroid dehydrogenase type 1 expression in human adipose tissue. *J Clin Endocrinol Metab* 2004;89:2711–2716.
- 12 Woods CP, Corrigan M, Gathercole L, Taylor A, Hughes B, Gaoatswe G, Manolopoulos K, Hogan AE, O'Connell J, Stewart PM, Tomlinson JW, O'Shea D, Sherlock M: Tissue Specific Regulation of Glucocorticoids in Severe Obesity and the Response to Significant Weight Loss Following Bariatric Surgery (BARICORT). *J Clin Endocrinol Metab* 2015;100:1434–1444.
- 13 Engeli S, Bohnke J, Feldpausch M, Gorzelnik K, Heintze U, Janke J, Luft FC, Sharma AM: Regulation of 11beta-HSD genes in human adipose tissue: influence of central obesity and weight loss. *Obes Res* 2004;12:9–17.
- 14 Heilmann P, Heide J, Hundertmark S, Schoneshofer M: Administration of glycyrrhetic acid: significant correlation between serum levels and the cortisol/cortisone-ratio in serum and urine. *Exp Clin Endocrinol Diabetes* 1999;107:370–378.
- 15 Vogeser M, Zachoval R, Felbinger TW, Jacob K: Increased ratio of serum cortisol to cortisone in acute-phase response. *Horm Res* 2002;58:172–175.
- 16 Rothberg AE, McEwen LN, Fraser T, Burant CF, Herman WH: The impact of a managed care obesity intervention on clinical outcomes and costs: a prospective observational study. *Obesity (Silver Spring)* 2013;21:2157–2162.
- 17 Monaghan PJ, Keevil BG, Stewart PM, Trainer PJ: Case for the wider adoption of mass spectrometry-based adrenal steroid testing, and beyond. *J Clin Endocrinol Metab* 2014;99:4434–4437.
- 18 Turcu AF, Rege J, Chomic R, Liu J, Nishimoto HK, Else T, Moraitis AG, Palapattu GS, Rainey WE, Auchus RJ: Profiles of 21-carbon steroids in 21-hydroxylase deficiency. *J Clin Endocrinol Metab* 2015;100:2283–2290.
- 19 Bender SB, DeMarco VG, Padilla J, Jenkins NT, Habibi J, Garro M, Pulakat L, Aroor AR, Jaffe IZ, Sowers JR: Mineralocorticoid receptor antagonism treats obesity-associated cardiac diastolic dysfunction. *Hypertension* 2015;65:1082–1088.
- 20 Goodfriend TL, Kelley DE, Goodpaster BH, Winters SJ: Visceral obesity and insulin resistance are associated with plasma aldosterone levels in women. *Obes Res* 1999;7:355–362.
- 21 Kidambi S, Kotchen JM, Grim CE, Raff H, Mao J, Singh RJ, Kotchen TA: Association of adrenal steroids with hypertension and the metabolic syndrome in blacks. *Hypertension* 2007;49:704–711.
- 22 Mihailidou AS, Loan Le TY, Mardini M, Funder JW: Glucocorticoids activate cardiac mineralocorticoid receptors during experimental myocardial infarction. *Hypertension* 2009;54:1306–1312.