

SBAT: A Tool for Estimating Metal Bioaccessibility in Soils

April 2004

S. A. Heuscher, C. C. Brandt, P. M. Jardine

Environmental Sciences Division



DOCUMENT AVAILABILITY

Reports produced after January 1, 1996, are generally available free via the U.S. Department of Energy (DOE) Information Bridge.

Web site <http://www.osti.gov/bridge>

Reports produced before January 1, 1996, may be purchased by members of the public from the following source.

National Technical Information Service

5285 Port Royal Road

Springfield, VA 22161

Telephone 703-605-6000 (1-800-553-6847)

TDD 703-487-4639

Fax 703-605-6900

E-mail info@ntis.fedworld.gov

Web site <http://www.ntis.gov/support/ordernowabout.htm>

Reports are available to DOE employees, DOE contractors, Energy Technology Data Exchange (ETDE) representatives, and International Nuclear Information System (INIS) representatives from the following source.

Office of Scientific and Technical Information

P.O. Box 62

Oak Ridge, TN 37831

Telephone 865-576-8401

Fax 865-576-5728

E-mail reports@adonis.osti.gov

Web site <http://www.osti.gov/contact.html>

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Environmental Sciences Division

SBAT: A TOOL FOR ESTIMATING METAL BIOACCESSIBILITY IN SOILS

S. A. Heuscher, C. C. Brandt, P. M. Jardine
Environmental Sciences Division
Oak Ridge National Laboratory

Date Published: April 2004

Prepared for
U.S. Department of Defense
Strategic Environmental Research and Development Program
Budget Activity Number 43WQ30801

Prepared by
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831
managed by
UT-BATTELLE, LLC
for the
U.S. DEPARTMENT OF ENERGY
under contract DE-AC05-00OR22725

CONTENTS

ACKNOWLEDGEMENTS.....	v
DISCLAIMER.....	vii
1. INTRODUCTION.....	1
2. USAGE INSTRUCTIONS.....	3
2.1 USING HISTORICAL DATA	3
2.2 USING SITE-SPECIFIC DATA (OPTIONAL)	5
2.3 REVIEWING THE BIOACCESSIBILITY ESTIMATES.....	5
2.4 TEST RESULTS	7
3. PREPARATION OF THE AGGREGATED HISTORICAL DATA.....	8
4. ANALYSIS PROTOCOLS.....	10
4.1 SOIL CHARACTERIZATION	10
4.2 SAMPLE PREPARATION.....	10
4.3 DETERMINATION OF TOTAL CHROMIUM AND ARSENIC ON SOIL	10
4.4 IN VITRO BIOACCESSIBILITY	11
4. REFERENCES.....	12
APPENDIX A: SBAT QUICK GUIDE	A-1
APPENDIX B: INFLUENCE OF SOIL GEOCHEMICAL AND PHYSICAL PROPERTIES ON THE SORPTION AND BIOACCESSIBILITY OF CHROMIUM(III) (Stewart et al. 2003a)	B-1
APPENDIX C: ADSORPTION SEQUESTRATION, AND BIOACCESSIBILITY OF AS(V) IN SOILS (Yang et al. 2002)	C-1
APPENDIX D: EFFECTS OF CONTAMINANT CONCENTRATION, AGING, AND SOIL PROPERTIES ON THE BIOACCESSIBILITY OF CR(III) AND CR(IV) IN SOIL (Stewart et al. 2003b).....	D-1
APPENDIX E: FACTORS CONTROLLING THE BIOACCESSIBILITY OF ARSENIC(V) AND LEAD(II) IN SOIL (Yang et al. 2003).....	E-1

ACKNOWLEDGMENTS

This research was sponsored by the U.S. Department of Defense's Strategic Environmental Research and Development Program (Dr. Andrea Leeson, Program Manager for Cleanup) as supplemental funding to CU-1166 ("Quantifying the Bioavailability of Toxic Metals in Soils"). We would like to acknowledge Dr. Jack Parker, Oak Ridge National Laboratory, for his suggestions on improving the application design, Dr. Mark Barnett, Auburn University, for providing arsenic bioaccessibility data, and Ms. Melanie Stewart, Oak Ridge National Laboratory for providing details of analytical protocols.

DISCLAIMER

Documents available from the web server were prepared as an account of work sponsored by an agency of the U.S. Government. Neither the U.S. Government nor any agency thereof, or any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Further, Oak Ridge National Laboratory is not responsible for the contents of any off-site pages referenced.

1. INTRODUCTION

Heavy metals such as chromium and arsenic are widespread in the environment due to their usage in many industrial processes. These metals may pose significant health risks to humans, especially children, due to their mutagenic and carcinogenic properties. Typically, the health risks associated with the ingestion of soil-bound metals are estimated by assuming that the metals are completely absorbed through the human intestinal tract (100% bioavailable). This assumption potentially overestimates the risk since soils are known to strongly sequester metals thereby potentially lowering their bioavailability.

Beginning in 2000, researchers at Oak Ridge National Laboratory, with funding from the Strategic Environmental Research and Development Program (SERDP), studied the effect of soil properties on the bioaccessibility of soil-bound arsenic and chromium. Representative A and upper-B horizons from seven major U.S. soil orders were obtained from the U.S. Department of Agriculture's National Resources Conservation Service and the U.S. Department of Energy's Oak Ridge Reservation. The soils were spiked with known concentrations of arsenic (As(III) and As(V)) and chromium (Cr(III) and Cr(VI)), and the bioaccessibility was measured using a physiologically based extraction test that mimics the gastric activity of children. Linear regression models were then developed to relate the bioaccessibility measurements to the soil properties (Yang et al. 2002; Stewart et al. 2003a). Important results from these publications and other studies include:

- Cr(VI) and As(III) are more toxic and bioavailable than Cr(III) and As(V) respectively.
- Several favorable processes can occur in soils that promote the oxidation of As(III) to As(V) and the reduction of Cr(VI) to Cr(III), thereby lowering bioaccessibility. Iron and manganese oxides are capable of oxidizing As(III) to As(V), whereas organic matter and Fe(II)-bearing minerals are capable of reducing Cr(VI) to Cr(III).
- The ubiquitous metal-sequestering properties of soils significantly lower the bioaccessibility of arsenic and chromium upon ingestion relative to the currently used 100% default values.
- Key soil physical and chemical properties (particle size, pH, mineral oxide, clay, and organic matter contents) govern the extent of toxic metal bioaccessibility thus providing the necessary conceptual understanding for building accurate predictive models.
- The As(V) regression model was able to predict the *in vivo* bioavailability in ten contaminated soils within a root mean square error of <10%.
- Metal bioaccessibility is controlled by molecular-level speciation, where metal sequestration and solid phase stability are enhanced by increased soil-metal contact time.

Using the results obtained from the SERDP-funded research, we have created an Excel® 2000 application called [SBAT \(Soil BioAccessability Tool\)](#) to estimate the bioaccessibility of soil bound arsenic (As(III) and As(V)) and chromium (Cr(III) and Cr(VI)) from soil properties. The tool combines the previously developed regression models with an extensive set of summarized historical data on soil properties. When a soil series or great group name is entered, SBAT will retrieve the associated historical data and calculate an estimate of bioaccessibility. Alternatively, a user can enter site-specific soils data that will be used in calculating the estimate, or a combination of historical and site-specific data can be used. Uncertainty estimates are also calculated to provide the user with a confidence measure.

Instructions for using the tool are provided in Chapter 2. For estimation, a user can enter either site-specific soils data or use the extensive set of summarized historical data provided with

SBAT. Chapter 3 provides background information on the source and methods used to summarize the historical data. Information on sampling and analysis protocols for obtaining site-specific measurements are given in Chapter 4. An abbreviated set of instructions for using the tool is provided in Appendix A, and reprints of relevant publications are included in Appendices B through E (Stewart et al. 2003a; Yang et al. 2002; Stewart et al. 2003b; Yang et al. 2003).

SBAT is designed to help managers identify sites that pose the greatest threat to human health and are thus most deserving of remediation or additional study. Since SBAT provides estimates of bioaccessibility, it is only intended for screening work. Site-specific measurements of bioaccessibility should be made when accurate results are needed for risk assessment.

2. USAGE INSTRUCTIONS

SBAT is a standalone Microsoft Excel® 2000 application (*SBAT.xls*). Prior to use, the application should be copied to your computer. Before opening SBAT, set the macro security level of Excel® to medium. To accomplish this, open Excel®, click on the “Tools” pull down menu and then on “Options”. Once the Options box appears, click on the “Security” tab, and then click on the “Macro Security” button. Set the security level to medium and click “OK”. Next, open *SBAT.xls* and click on the “Enable Macro” button when asked by Excel® to enable or disable macros.

The SBAT interface screen is shown in Figure 1. The yellow highlighted cells are used for input. SBAT consists of the following six worksheets:

1. *Form* – Application interface. This worksheet is used to input the soil series or great group name and display the bioaccessibility estimates. The other worksheets in the workbook are linked to this worksheet.
2. *Taxonomy* – Series names and their great group assignment.
3. *Chem* – Aggregated historical soil characterization data.
4. *Metals* – Bioaccessibility regression models.
5. *Prediction Intervals* – Data for prediction interval calculation.
6. *Measurement* – Assists the user in inputting site-specific data on the *Form* worksheet.

Each worksheet is password protected to prevent users from accidentally editing the formulas or data. Initially, the worksheet tabs are not displayed when the workbook is opened. To display the tabs, click on the “Tools” pull down menu and then on “Options”. Once the Options box appears, click on the “View” tab and make sure the “Sheet Tabs” option is selected. Click on “OK” to save.

SBAT can use two different sources of data for estimating bioaccessibility. If available, a user can enter site-specific soil properties (Section 2.2) in the **Site-Specific Parameters** section of the interface screen. If site-specific data are unavailable, the user can specify the name of a soil series or great group. The tool will use soil properties data stored in the *Chem* worksheet (Section 2.1) and display the results in the **Estimated Parameters** portion of the interface screen. In either case, SBAT calculates the bioaccessibility estimate and uncertainty based on the models stored in the *Metals* worksheet (Section 2.3) and outputs these estimates to the **Bioaccessibility Estimates** section.

2.1 USING HISTORICAL DATA

At the top of the interface screen is an area in which the user can enter the name of the soil series or great group for which bioaccessibility estimates will be calculated. If the name of soil series is known (e.g., Fullerton), the user should type this name into the yellow-highlighted cell adjacent to the words *Enter Series* and press the Enter key. The series name may be upper or lower case, but it must be spelled correctly. If the user is unsure of the series spelling, she or he can click on the *Taxonomy* worksheet tab to view an alphabetic list of soil series in column B. If only the great group is known, the user should type this name into the yellow-filled cell adjacent to the words *Select Great Group* and press the Enter key. Clicking on the button on the right side of the cell will enable the user to select a great group name from a scroll list. If either the soil series or great group is unknown, leave the appropriate yellow-highlighted cell blank.

Once a soil series name is entered, SBAT retrieves the great group name from the *Taxonomy* worksheet and displays this information in cell B9. The **Estimated Parameters** section displays the aggregated historical data for this great group. The cells in the **Estimated Parameters** section will contain #N/A (not available) if this great group is not in the aggregated data (*Chem* worksheet) or if the series entered is not found in the *Taxonomy* worksheet.

	A	B	C	D	E	F	G	H
1	SBAT: Soil BioAccessibility Tool							
2	<i>Note: input fields are yellow</i>							
3								
4								
5	<i>For the soil of interest, if the soil's Great Group is known, select it from list in cell B7 or if the</i>							
6	<i>soil's Series is known enter its name in cell E7; if either is unknown leave the corresponding cell(s) blank</i>							
7	Select Great Group:		Dystrudepts	or	Enter Series:			
8								
9	Great Group: Dystrudepts							
10								
11	Estimated Parameters							
12								
13	Horizon:							
14			A			B		
15			Value	Coverage	Value	Coverage		
16	Organic carbon (TOC), wt%		3.38	64.8	0.71	66.4		
17	Carbonate content (TIC), wt%		0.572	2.0	0.986	2.3		
18	Clay content, wt%		15.0	64.5	15.6	66.1		
19	Iron content, wt%		1.751	54.3	1.847	54.0		
20	Soil pH		5.0	64.8	5.0	66.6		
21								
22	<i>Enter soil properties if site data are available: note: blank fields will be estimated from the great group properties above</i>							
23	Site-Specific Parameters							
24	Help with this Section							
25	Horizon:							
26			A			B		
27			Mean	StDev	Mean	StDev		
28	Organic carbon (TOC), wt%							
29	Carbonate content (TIC), wt%							
30	Clay content, wt%							
31	Iron content, wt%							
32	Soil pH							
33								
34	<i>note: bioaccessibility values calculated from parameters beyond the range of the model (extrapolated values) are highlighted in pink</i>							
35	Bioaccessibility Estimates							
36	P value: 0.05							
37								
38	Bioaccessibility (%) for Horizon							
39	Metal	Lower Pred. Limit	Mean	Upper Pred. Limit	Lower Pred. Limit	Mean	Upper Pred. Limit	Parameters Used
40	Cr (III) model 1	0.0	8.9	20.3	8.1	19.2	30.3	TOC Clay
41	Cr (III) model 2	6.8	17.0	27.1	3.0	13.2	23.5	TIC Clay
42	Cr (VI)	0.0	14.6	39.8	5.3	30.0	54.7	TOC pH
43	As (V)	0.0	14.4	45.2	0.0	13.5	44.3	Iron pH
44	As (III)	0.0	15.0	43.7	0.0	14.2	43.0	Iron pH
45								
46								
47								
48								
49								

4/7/2004

Fig. 1. SBAT interface.

When the great group name is retrieved, the average measurements for organic carbon, carbonate carbon, clay content, iron, and pH are displayed under the *Value* columns and the coverage of each of these soil properties is displayed under the *Coverage* columns. See Section 3 for a definition of coverage and how it was calculated.

If a measurement for a soil property is unavailable due to lack of appropriate data, the word *missing* is displayed. There are two ways in which this can result. First, there may not be any measurements for the great group and soil property in the historical data set. Possible reasons for missing measurements include: (i) the soil property was not measured in the laboratory, (ii) the horizon thickness or bulk density was missing, or (iii) quality control resulted in deletion of measurements. A second cause of missing data is that the area(s) of the aggregated series in a great group are missing from the geographic database used to identify soil series in the U.S.

2.2 USING SITE-SPECIFIC DATA (OPTIONAL)

If site-specific soils data are available, they may be entered in the **Site-Specific Parameters** section. Enter the site-specific measurements for the soil of interest in the appropriate yellow-highlighted cells. If a measurement is unavailable, leave the corresponding cell(s) blank. Soil pH should be the measured in distilled water at a 1:1 soil to solution ratio. Iron content should be measured with a dithionite citrate extraction procedure. Total organic carbon (TOC) and carbonate carbon (TIC) should be measured by combustion on a total carbon analyzer. Section 4 gives additional information about the proper analytical protocols for generating the site-specific data.

If multiple measurements are available, the user should enter the average of the measurements. For help in calculating the average, click [*Help with this section*](#). Clicking this link will take the user to the *Measurement* worksheet. Multiple measurements for soil properties can be entered here, and the average and standard deviation of these measurements will be calculated and displayed in the Results section. To copy these results to the *Form* worksheet press Ctrl and the “s” key simultaneously.

A combination of site-specific and aggregated data can be used to calculate bioaccessibility. If site-specific soils data are available but measurements are missing for some soil properties, then the aggregated data measurement will automatically be substituted for these soil properties in the bioaccessibility calculation.

2.3 REVIEWING THE BIOACCESSIBILITY ESTIMATES

Multiple regression equations were developed from data collected in previous studies (Yang et al. 2002; Stewart et al. 2003a). Reprints of these publications are provided in Appendices B and C. These equations, which form the basis of the estimates generated by SBAT, are displayed in Table 1.

Table 1. Regression models for estimating arsenic and chromium bioaccessibility (BA) in soils.

Metal	Model[†]	R²	P
As(V)	BA = -22.37 - 36.22 log Fe + 9.11 pH	0.720	<0.001
As(III)	BA = -1.41 - 30.64 log Fe + 4.76 pH	0.574	<0.001
Cr(III) model 1	BA = 15.53 - 3.78 TOC + 0.408 Clay	0.674	<0.001
Cr(III) model 2	BA = 16.02 - 9.56 TIC + 0.426 Clay	0.722	<0.001
Cr(VI)	BA = 57.34 - 22.55 log TOC - 6.15 pH	0.601	<0.001

[†]Units: BA (%), Fe (wt %), pH (standard units), Clay (wt %), TOC (wt %), TIC (wt %).

There are two models for Cr(III) for reasons described in Stewart et al. (2003a). Briefly, clay, organic carbon (TOC) and carbonate carbon (TIC) are thought to be important predictors of Cr(III) bioaccessibility. However, when clay, TIC, and TOC were used in the same model, the contribution of TOC was not significant at the 90% confidence level. This may be an artifact of the limited data from which the models were derived. A more extensive data set is necessary to test this hypothesis (Stewart et al. 2003a). For completeness, SBAT includes models with TOC and TIC.

The **Bioaccessibility Estimates** section displays the estimates and upper and lower prediction limits of Cr(III), Cr(VI), As(V), and As(III) bioaccessibility for the A and B horizons of the soil of interest. This section also displays the soil properties used to estimate bioaccessibility for the metal of interest under *Parameters Used*. Bioaccessibility values displayed in this section have been restricted between 0 and 100 (their physical limits). #VALUE! and #N/A are errors that may occasionally be displayed in this section. If #VALUE! is displayed, data were missing for an aggregated soil property necessary for modeling. Look in the **Estimated Parameters** section to see where the word *missing* is displayed. #N/A is displayed if aggregated data are not available and site-specific data for the soil properties necessary for modeling are not available.

The user may enter a probability value for the prediction interval in the upper right corner of this section adjacent to the word *P value*. For example, if a p value of 0.10 is entered, the 90% prediction interval is displayed, and the user can say that she or he is 90% confident that the bioaccessibility for the particular soil of interest lies between the upper and lower prediction limits. It is important to note that prediction intervals are displayed, not confidence intervals. Both types of intervals reflect the uncertainty of a bioaccessibility estimate. Confidence intervals quantify the uncertainty in estimating the mean bioaccessibility of all soils whose physical and chemical properties are equal to numeric values of interest. Prediction intervals are wider than confidence intervals because prediction intervals include the additional uncertainty associated with estimating the bioaccessibility of a particular soil.

Bioaccessibility estimates are highlighted in pink if they are calculated from soil measurements that are outside the range of data used to build the original bioaccessibility regression models. These estimates should be used with caution because they are extrapolations of the model. In this case, the regression model may not properly describe the relationship between bioaccessibility and soil properties.

There are several factors to consider in interpreting the output of SBAT. Foremost, the user should evaluate the data used for predicting bioaccessibility. The parameters (soil properties) necessary for modeling the bioaccessibility are listed under *Parameters Used* in the **Bioaccessibility Estimates** section. If site-specific data were not entered for these soil properties, then the corresponding aggregated soil property data in the **Estimated Parameters**

section were automatically used to predict bioaccessibility. Site specific measurements for the soil of interest are obviously better than using aggregated historical data to estimate bioaccessibility. However, if data from the **Estimated Parameters** section were used, the user should evaluate the *Coverage* values of the parameters used for modeling. If the coverage for a parameter is low, then the aggregated value is based on only a few soil series and should not be trusted to accurately predict bioaccessibility for the great group of interest. If no data were available for the aggregated soil property, then the word *missing* is displayed and the coverage value is zero.

2.4 TEST RESULTS

To test the accuracy of using aggregated soil properties, bioaccessibility estimates for nineteen soil series were calculated using SBAT. These bioaccessibility estimates were compared to laboratory measurements made on the same samples. Table 2 shows the proportion of the variance in bioaccessibility explained (R^2) by the regression equations using aggregated and measured soil properties.

Table 2. Bioaccessibility predictions using aggregated and measured soil data.

Metal	Variance Explained (R^2)	
	Aggregated Soil Data	Measured Soil Data
As(III)	0.55	0.57
As(V)	0.77	0.72
Cr(III) model 1	0.57	0.67
Cr(III) model 2	0.05	0.72
Cr(VI)	0.65	0.60

These results indicate that, with the exception of the Cr (III) model 2, soil data aggregated to the great group level provides almost as good a bioaccessibility estimate as is obtained from measured soil data. Bioaccessibility estimates used in this comparison were not restricted by coverage values. In other words, estimates based on soil properties with low coverage (<30%) were not deleted. Coverage values indicate how well the historical data represents a given great group (See Section 3). Many of the aggregated TIC values used in the Cr (III) model 2 were based on low coverage. This is a possible explanation for the low R^2 of this model. Another explanation is that there could be a large variance in TIC within great groups.

3. PREPARATION OF THE AGGREGATED HISTORICAL DATA

The aggregated historical data included with the bioaccessibility tool is derived from the Soil Survey Laboratory Soil Characterization database compiled by the U.S. Department of Agriculture's Natural Resources Conservation Service (NRCS). This database contains physical, chemical, engineering, mineralogical, and descriptive data for more than 21,000 pedons collected from all fifty states (Soil Survey Staff 1997). Analytical procedures used to measure the soil physical and chemical properties are described in the Soil Survey Laboratory Investigations Report No. 42 (Soil Survey Staff 1996).

The NRCS data were reviewed and suspicious values were removed. In order to provide more extensive spatial coverage, the NRCS data were aggregated to the great group taxonomic level. The aggregation process consists of several steps which are summarized in Figure 2. First, measurements for multiple subhorizons within a pedon were averaged to the master horizons for the pedon. The sample measurements were weighted by the subhorizon mass which was calculated by multiplying the subhorizon's thickness by its bulk density. The measurements were restricted to the top meter of soil since this represents the maximum volume of soil likely to be ingested. Next, all of the pedons belonging to a series were averaged resulting in aggregated measurements for the series. Finally, this series data was averaged to great group. This last step was weighted by series area. Series areas were obtained from the State Soil Geographic (STATSGO) Database (U.S. Department of Agriculture 1991).




Fig. 2. Data aggregation process.

The great group level aggregated data were incorporated into SBAT. However, before aggregating to great group level, the series level aggregated data were evaluated. The series aggregated data were unsuitable due to lack of sufficient data. Approximately 55% of aggregated measurements for carbonate are missing and 45% of the aggregated data are missing iron measurements. Approximately 51% of series level aggregated measurements for clay, pH, and organic carbon content were based on only one pedon.

Since approximately 60% of the NRCS database is missing bulk density measurements, only 40% of the database would be available for data aggregation. To remedy this problem, multiple regression equations were developed to estimate bulk density from common soil properties present in the database. Using these models to add predicted bulk density values to the database, only 17% of the database was missing bulk density measurements (Heuscher et al. 2004).

Since both the NRCS and STASGO databases are consistently being updated and revised, the aggregated data is essentially a snapshot in time. In other words, every existing soil series is not mapped and analytical data have not been recorded for every sample. To indicate how well the aggregated soil properties represent a given great group, percent aerial coverage was calculated as follows for each type of measurement or soil property:

$$\text{Coverage (\%)} = 100 \times \frac{\text{total area of aggregated series in the great group}}{\text{total area of all series in the great group}}$$

If the coverage value for a soil property is low, then the aggregated value is based on only a few soil series in that great group. Thus, it is probable that the aggregated soil property is not representative of the given great group.

4. ANALYSIS PROTOCOLS

Bioaccessibility is a measure of the mobilization of contaminants from soil during digestion. The bioavailability of ingested contaminated soils is defined as the contaminant fraction that reaches systemic circulation (Oomen et al. 2002). Thus, bioaccessibility is a precursor measurement to bioavailability since bioaccessibility is measured *in vitro* and bioavailability is measured *in vivo* with test animals. Bioavailability protocols are often not feasible for site-specific risk assessment studies due to this fact.

Several *in vitro* digestion models based on human physiology have been developed as cheap and reproducible tools to investigate bioaccessibility of soil contaminants (Oomen et al. 2002). A modified version of a physiologically based extraction test (PBET) developed by Ruby et al. (1996) was used to generate the data used for SBAT since it has been cross-correlated with *in vivo* bioavailability studies involving arsenic, lead, and cadmium (Ruby et al. 1996; Rodriguez et al. 1999; Schroder et al. 2003). Using this method, the reported bioaccessibility is a measure of the amount of ingested contaminant that is soluble due to simulated human gastric functions and has the potential to cross the intestinal wall. Protocols for obtaining bioaccessibility measurements using the modified PBET method are described in the attached publications (Appendices B and C) but are repeated here for convenience.

4.1 SOIL CHARACTERIZATION

Soil pH was determined using double deionized (DDI) water and 5mM CaCl₂ in a 2:1 solution to solid ratio. The pH of the clear supernatant was measured with a microprocessor ionalyzer/901 (Orion Research, Beverly, MA) using a combination glass and Calomel electrode (Beckman, Fullerton, CA). Extractable iron and manganese oxides were determined with dithionite-citrate-bicarbonate (DCB) using the methods of Mehra and Jackson (1960). Total organic carbon (TOC) and total inorganic carbon (TIC) were measured by combustion on a Perkin-Elmer 2400 Series II CHNS/O analyzer. Soil TOC was determined on pretreated samples to remove TIC which involved a near-boiling, 3 M HCl extraction method on agitated samples. Soil TIC was computed from the difference between total soil carbon (no pretreatment) and TOC.

4.2 SAMPLE PREPARATION

Soils were disaggregated with gentle grinding using a mortar and pestle and sieved to provide a soil fraction <250 μm . It is this smaller size material that is more commonly ingested by children since it adheres more readily to the hand (Sheppard et al. 1995; Rodriguez et al. 1999). Soils were spiked with various mass-loadings of Cr(III), Cr(VI), As(III), or As(V) as described in Stewart et al. (2003a) and Yang et al. (2002).

4.3 DETERMINATION OF TOTAL CHROMIUM AND ARSENIC ON SOIL

Total chromium and arsenic on the soil was determined using a modification of EPA method 3052 (<http://www.epa.gov/SW-846/pdfs/3052.pdf>). The soil was digested in a CEM microwave, model MDS-81D, with hydrofluoric and nitric acid. Boric acid was added before sample analysis in order to facilitate the removal of hydrofluoric acid from solution through the formation of fluoroboric acid. Samples were stored and analyzed for total chromium and arsenic using Inductively Coupled Plasma. Soils from the National Institute of Standards, with known concentrations of solid phase chromium or arsenic, were also analyzed with each block of analysis.

4.4 IN VITRO BIOACCESSIBILITY

Triplicate 0.39 g moist samples (0.3 g dry weight) were placed in 50 ml polyethylene tubes to which 30 ml of PBET solution was added. The PBET solution is 0.4 M glycine at pH 1.5 and 37°C. It is important to note that the quantity of soil can vary as long as the soil dry weight to PBET solution ratio of 1 to 100 is maintained. The slurries were then quickly placed in a rotating water bath of 37°C and agitated at 30±2 rpm for 1 hour. Supernatant was separated from the solid via centrifugation. The pH of the supernatant was measured to ensure that the final pH was within ±0.5 pH units of the initial pH. This scenario held for all cases. The PBET pH of 1.5 simulates the most aggressive stomach digestive scenario which is a condition indicative of human fasting.

For arsenic, the supernatant was analyzed using graphite furnace atomic adsorption spectrometry. For chromium, the supernatant was first analyzed for Cr(VI) using a modified *s*-diphenyl-carbohydride colormetric method (Bartlett and James 1979) with a Hewlett-Packard model 8453 UV-VIS spectrophotometer at wavelength 540 nm (Palo Alto, CA). Analysis of Cr(VI) was performed immediately on rapidly cooled PBET supernatant solutions to avoid possible reduction of Cr(VI) to Cr(III) by glycine (Jardine et al. 1999). Total Cr (Cr_T) in the supernatant was measured on a Perkin Elmer AAnalyst 800 atomic absorption spectrometer (Wellseley, PA). Chromium (III) was calculated as the difference between Cr_T and Cr(VI). Bioaccessibility was then calculated as:

$$\% \text{ Bioaccessibility} = \left(\frac{\text{Cr or As in PBET supernatant } (\mu\text{g/mL}) \times 30.0 \text{ mL (vol. of PBET solution)}}{\text{Cr or As on soil surface } (\mu\text{g/g}) \times 0.3 \text{ g (dry soil weight)}} \right) \times 100$$

5. REFERENCES

- Bartlett, R. and B. James. 1979. Behavior of chromium in soils: III. Oxidation. *Journal of Environmental Quality*. 8:31-35.
- Heuscher, S.A., C.C. Brandt, and P.M. Jardine. 2004. Using soil properties to estimate bulk density. *Soil Science Society of America Journal*. (submitted).
- Jardine, P.M., S.E. Fendorf, M.A. Mayes, S.C. Brooks, and W.B. Bailey. 1999. Fate and transport of hexavalent chromium in undisturbed heterogeneous soil. *Environmental Science and Technology*. 33:2939-2944.
- Mehra, O.P. and M.L. Johnson. 1960. Iron oxide removed from soils and clays by a dithionite-citrate system buffered with sodium bicarbonate. *Clays and Clay Minerals*. 7:317-327.
- Oomen, A.G., A. Hack, M. Minekus, E. Zeijdner, C. Cornelis, G. Schoeters, W. Verstraete, T. Van De Wiele, J. Wragg, C. Rompelberg, A. Sips, and J. Van Wijnen. 2002. Comparison of five in vitro digestion models to study the bioaccessibility of soil contaminants. *Environmental Science and Technology*. 36:3326-3334.
- Rodriguez, R.R., N.T. Basta, S.W. Casteel, and L.W. Pace. 1999. An in vitro gastrointestinal method to estimate bioavailable arsenic in contaminated soils and solid media. *Environmental Science and Technology*. 33:642-649.
- Ruby, M.V., A. Davis, R. Schoof, S. Eberle, and C.M. Sellstone. 1996. Estimation of lead and arsenic bioavailability using a physiologically based extraction test. *Environmental Science and Technology*. 30:422-430.
- Schroder, J.L., N.T. Basta, J. Si, S.W. Casteel, T. Evans, and M. Payton. 2003. In vitro gastrointestinal method to estimate relative bioavailable cadmium in contaminated soil. *Environmental Science and Technology*. 37:1365-1370.
- Sheppard, S.C., W.G. Evenden, and W.J. Schwartz. 1995. Ingested soil: bioavailability of sorbed lead, cadmium, cesium, iodine and mercury. *Journal of Environmental Quality*. 24:498-505.
- Soil Survey Staff. 1996. *Soil Survey Laboratory Methods Manual*. Soil Survey Laboratory Investigations Report No. 42. U.S. Department of Agriculture. 716 p.
- Soil Survey Staff. 1997. *National Soil Survey Characterization Data*. Soil Survey Laboratory, National Soil Survey Center, Natural Resources Conservation Service, U.S. Department of Agriculture. Lincoln, NE.
- Stewart, M.A., P.M. Jardine, M.O. Barnett, T.L. Mehlhorn, L.K. Hyder, and L.D. McKay. 2003a. Influence of soil geochemical and physical properties on the sorption and bioaccessibility of chromium(III). *Journal of Environmental Quality*. 32:129-137 (reprint in Appendix B).
- Stewart, M.A., P.M. Jardine, C.C. Brandt, M.O. Barnett, S.E. Fendorf, L.D. McKay, T.L. Mehlhorn, and K. Paul. 2003b. Effects of contaminant concentration, aging and soil properties on the bioaccessibility of Cr(III) and Cr(VI) in soil. *Soil and Sediment Contamination*. 12:1-21 (reprint in Appendix D).

U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, 1991. State Soil Geographic (STATSGO) Database.

Yang, J-K., M.O. Barnett, P.M. Jardine, N.T. Basta, and S.W. Casteel. 2002. Adsorption, sequestration, and bioaccessibility of As(V) in soils. *Environmental Science and Technology*. 36:4562-4569 (abstract in Appendix C).

Yang, J-K., M.O. Barnett, P.M. Jardine, and S.C. Brooks. 2003. Factors controlling the bioaccessibility of arsenic (V) and lead(II) in soils. *Soil and Sediment Contamination*. 12:165-179 (reprint in Appendix E).

APPENDIX A
SBAT QUICK GUIDE

SBAT QUICK GUIDE

1. Prior to opening SBAT, set the macro security level on Excel® to medium. To accomplish this, open Excel®, click on the “Tools” pull down menu and then on “Options”. Once the Options box appears, click on the “Security” tab, and then click on the “Macro Security” button. Set the security level to medium and click “OK”. Open the Excel® file *SBAT.xls*, and, when prompted, click the “Enable Macros” button. The worksheet tabs are not displayed when the workbook is initially opened. If the user would like to display the tabs, click Tools → Options, click the “View” tab, click the box adjacent to the words “Sheet tabs”, and click OK. Each worksheet is password protected to prevent users from accidentally editing formulas or reference data.
2. The interface worksheet (*Form*) will appear when the application is initially opened. This worksheet contains three boxed sections (top to bottom): **Estimated Parameters**, **Site-Specific Parameters**, and **Bioaccessibility Estimates**. The yellow cells are available for user input.
3. Immediately above the **Estimated Parameters** section, the user can enter the soil series or great group of interest in either of the two yellow cells. If the great group or series is not known, ignore this section and proceed to step (4). If the soil’s great group is known, click on the yellow colored cell adjacent to the words *Select Great Group*. Either type in the great group name or click on the button that appears on the right side of the cell and use the scroll bar to select the great group name from the list. If the soil great group is not known, but the soil series is known, enter the series name by clicking on the yellow colored cell adjacent to the words *Enter Series*, typing the soil series name and pressing enter. The **Estimated Parameters** section displays the great group aggregated data. Upon entering a soil series name or picking a great group from the list, the great group name is retrieved and the great group’s averages for organic carbon, inorganic carbon, clay content, iron content, and pH are displayed under the *Value* columns. If a measurement for one of these soil properties is unavailable due to lack of appropriate data, the word *missing* is displayed. If the great group of interest is not in the aggregated data or if the series entered is not listed in the most current taxonomy, then the cells in the **Estimated Parameters** section will fill with #N/A.
4. If site-specific soils data are available, they may be entered in the **Site-Specific Parameters** section. In the yellow colored cells, enter site specific measurements for the soil of interest. If a measurement is unavailable, leave the corresponding cell(s) blank.
5. If the site has multiple measurements for soil properties, then enter the average of the measurements. For help in calculating this average click [*Help with this section*](#). Clicking this link will take you to the *Measurement* worksheet. Multiple measurements for soil properties can be entered here, and the average and standard deviation of these measurements are calculated and displayed in the Results section. To copy these results to the interface worksheet press Ctrl and the “s” key simultaneously.
6. In the **Bioaccessibility Estimates** section, look under *Bioaccessibility (%) for Horizon* to see an estimate of As(V), As(III), Cr(III), and Cr(VI) bioaccessibility as well as the upper and lower prediction limits for the A and B horizons of the soil of interest. #VALUE! and #N/A are errors that may be displayed under *Bioaccessibility (%) for Horizon* in **Bioaccessibility Estimates** section. If #VALUE! is displayed, data were missing for an aggregated soil property necessary for modeling; look in the **Estimated Parameters** section to see where the

word *missing* is displayed. #N/A is displayed if aggregated data are not available and site-specific data for the soil properties necessary for modeling are not available.

7. Enter the p value of interest for the prediction interval in the upper right corner of this section adjacent to the word *P value*. For example, if a p value of 0.10 is entered, the 90% prediction interval is displayed under *Bioaccessibility (%) for Horizon*. Bioaccessibility estimates are highlighted in pink if they are calculated from soil properties that are outside the range of data used to build the bioaccessibility models. These estimates should not be trusted because they are extrapolating the model outside the range of data. Therefore, the model may not properly describe the relationship between bioaccessibility and soil properties. The **Bioaccessibility Estimates** section also displays the soil properties used to estimate bioaccessibility under *Parameters Used*.
8. If site specific data were not entered for soil properties listed under *Parameters Used*, then the corresponding aggregated soil property data in the **Estimated Parameters** section were automatically used to predict bioaccessibility. If data from the **Estimated Parameters** section were used, evaluate the coverage column(s) of the parameters used for modeling. If the coverage for a parameter is low, then the aggregated value is based on only a few soil series and should not be trusted to accurately predict bioaccessibility for the great group of interest.

APPENDIX B

**STEWART, M.A., P.M. JARDINE, M.O. BARNETT, T.L. MEHLHORN, L.K. HYDER,
AND L.D. MCKAY. 2003a. INFLUENCE OF SOIL GEOCHEMICAL AND PHYSICAL
PROPERTIES ON THE SORPTION AND BIOACCESSIBILITY OF CHROMIUM(III).
JOURNAL OF ENVIRONMENTAL QUALITY. 32:129-137.**

Influence of Soil Geochemical and Physical Properties on the Sorption and Bioaccessibility of Chromium(III)

M. A. Stewart, P. M. Jardine,* M. O. Barnett, T. L. Mehlhorn, L. K. Hyder, and L. D. McKay

ABSTRACT

There are numerous Cr(III)-contaminated sites on Department of Defense (DoD) and Department of Energy (DOE) lands that are awaiting possible clean up and closure. Ingestion of contaminated soil by children is the risk driver that generally motivates the likelihood of site remediation. The purpose of this study was to develop a simple statistical model based on common soil properties to estimate the bioaccessibility of Cr(III)-contaminated soil upon ingestion. Thirty-five uncontaminated soils from seven major soil orders, whose properties were similar to numerous U.S. DoD contaminated sites, were treated with Cr(III) and aged. Statistical analysis revealed that Cr(III) sorption (e.g., adsorption and surface precipitation) by the soils was strongly correlated with the clay content, total inorganic C, pH, and the cation exchange capacity of the soils. Soils with higher quantities of clay, inorganic C (i.e., carbonates), higher pH, and higher cation exchange capacity generally sequestered more Cr(III). The amount of Cr(III) bioaccessible from the treated soils was determined with a physiologically based extraction test (PBET) that was designed to simulate the digestive process of the stomach. The bioaccessibility of Cr(III) varied widely as a function of soil type with most soils limiting bioaccessibility to <45 and <30% after 1 and 100 d soil-Cr aging, respectively. Statistical analysis showed the bioaccessibility of Cr(III) on soil was again related to the clay and total inorganic carbon (TIC) content of the soil. Bioaccessibility decreased as the soil TIC content increased and as the clay content decreased. The model yielded an equation based on common soil properties that could be used to predict the Cr(III) bioaccessibility in soils with a reasonable level of confidence.

THE PRESENCE of chromium (Cr) in the environment is widespread due to its usage in many industrial processes. The metallurgic, tanning, and plating industries are just a few examples of very common applications, large and small, which use Cr on a daily basis (Nriagu and Nieboer, 1988). Chromium itself is thermodynamically stable in two oxidative states: cationic Cr with a valence of three, Cr(III), and anionic Cr with a valence of six, Cr(VI). Chromium(VI) is often considered to be mobile in the environment while the more environmentally stable Cr(III) is considered less mobile (Chung et al., 1994; Patterson et al., 1997). There are several factors that contribute to the decreased mobility of Cr(III) in soil: (i) strong adsorption onto the nega-

tively charged soil surfaces, (ii) the ability to form complex molecules with organics found in the soil, and (iii) the formation of oxides and hydroxides and other insoluble minerals in soil (Fendorf and Zasoski, 1992; Losi et al., 1994; Dragun, 1998).

When assessing the risks posed by Cr(VI) and Cr(III), the exposure pathway of most concern is ingestion by children (Paustenbach, 1989; Davis et al., 1990; Sheehan et al., 1991; Skowronski et al., 2001). Chromium(VI) is considered the most harmful of the oxidative states since it is both a mutagen and a carcinogen at low sub-ppm levels (Levis and Bianchi, 1982). Although Cr(III) is generally considered less harmful to human health than its oxidized counterpart, it may be of concern due to its potential to oxidize to Cr(VI) and its ability to accumulate to very high solid phase concentrations in some soils (Fendorf et al., 1992). The bioaccessibility of organic contaminants in soils has been relatively well studied (Linz and Nakles, 1997); however, the bioaccessibility of soil-bound metals such as Cr has received less attention (Ruby et al., 1996; Rodriguez et al., 1999; Skowronski et al., 2001; Stewart et al., 2003), where the bioaccessibility is defined as that amount of contaminant that is soluble due to simulated *in vitro* gastric functions and has the potential to cross the intestinal wall (Hamel et al., 1998). Typically, calculated health risks are inappropriately based on a reference dose derived from studies that use soluble aqueous metal species. The ubiquitous metal-sequestering properties of soil may significantly lower the bioaccessibility of Cr upon digestion, which, in turn may influence the decision for remediation at contaminated sites. Thus, action levels set by state regulators concerning the bioaccessibility of Cr in soil may need to consider specific soil properties instead of using generic guidelines (Proctor et al., 1997).

The intent of this paper is to show that Cr(III) can be strongly sequestered by soil, which in turn influences its bioaccessibility. We developed a simple statistical model based on measured soil properties to estimate the bioaccessibility of Cr(III)-contaminated soils upon ingestion. We show that common soil properties, which are easily obtainable from the National Resource Conservation Service (NRCS) database, can be used to assess Cr(III) bioaccessibility at contaminated sites.

M.A. Stewart, P.M. Jardine, T.L. Mehlhorn, and L.K. Hyder, Environ. Sci. Div., Oak Ridge, TN 37831-6038; M.O. Barnett, Dep. of Civil Engineering, 208 Harbert Engineering Center, Auburn Univ., AL 36849-5337; and L.D. McKay, Dep. of Geological Sciences, Univ. of Tennessee, Knoxville, TN 37996-1410. This research was sponsored by the U.S. Department of Defense (DoD) Strategic Environmental Research and Development Program. Oak Ridge National Lab is managed by the University of Tennessee– Battelle LLC, under contract DE-AC05-00OR22725 with the U.S. Department of Energy. Received 7 Nov. 2001. *Corresponding author (jardinepm@ornl.gov).

Abbreviations: DoD, Department of Defense; DOE, Department of Energy; PBET, physiologically based extraction test; NRCS, National Resource Conservation Service; CEC, cation exchange capacity; DDI, double deionized; TOC, total organic carbon; TIC, total inorganic carbon; VIF, Variance Inflation Factor; XAS, x-ray adsorption spectroscopy.

METHODS

Soil Type and Characterization

A database of metal-contaminated Department of Defense (DoD) sites was obtained from the U.S. Army Environmental Center, Aberdeen Proving Ground, Maryland. Twenty (20) DoD Army facilities throughout the USA were chosen for consideration based on the high concentration of Cr in their soils and the possible need for remediation (Table 1). Because of the difficulty in obtaining actual contaminated soils from these sites, uncontaminated soils whose properties were similar to the contaminated soils were acquired and treated with Cr(III). The soil series present at the DoD sites of interest were identified using Soil Conservation Survey documents. The USDA-NRCS database was then utilized to locate pedon numbers associated with each soil series. The NRCS was contacted and in most cases 200 g of the A-horizon and the upper B-horizon soil were obtained for each soil series (Table 1). Two additional soils were obtained from the Oak Ridge Reservation in eastern Tennessee, which also had properties similar to DoD sites in the southeast USA. Thirty-five soils were acquired and these encompassed seven major soil orders (Table 1).

Soils were disaggregated with gentle grinding using a mortar and pestle and sieved to provide a soil fraction <250 μm . It is this smaller size material that is more commonly ingested by children since it adheres more readily to the hand (Sheppard et al., 1995; Rodriguez et al., 1999). Soil properties were obtained from (i) the NRCS database and (ii) repeated or additional measurements in our laboratory. Soil properties included pH, cation exchange capacity (CEC), Fe- and Mn-oxide content, particle size distribution, and total organic and inorganic C (Table 2). Repetitive or additional measurements of soil pH, Fe- and Mn-oxide content, and total organic and inorganic C on all soils were performed to verify the quality of, and provide

missing information to the NRCS database. In general, data generated in our laboratory was in excellent agreement with the NRCS database. Soil pH was determined using double deionized (DDI) water and 5 mM CaCl_2 in a 2:1 solution/solid ratio. The pH of the clear supernatant was measured with a microprocessor ionalyzer/901 (Orion Research, Beverly, MA) using a combination glass and Calomel electrode (Beckman, Fullerton, CA). Extractable iron and manganese oxides were determined with dithionite–citrate–bicarbonate (DCB) using the methods of Mehra and Jackson (1960). Total organic carbon (TOC) and total inorganic carbon (TIC) were measured by combustion on a Perkin-Elmer 2400 Series II CHNS/O analyzer. Soil TOC was determined on pretreated samples to remove TIC, which involved a near-boiling, 3 M HCl extraction method on agitated samples. Soil TIC was computed from the difference between total soil C (no pretreatment) and TOC.

Contaminant Addition to Soil

Ten grams of soil was placed in a 200-mL glass centrifuge vessel along with 100 mL of 500 ppm Cr(III) as CrCl_3 , pH 4.0. The slurry was agitated on a reciprocal shaker for 2 d, centrifuged, and the supernatant decanted for analysis. This was repeated three more times. After the fourth addition of Cr, the soils were washed three times with DDI water and allowed to air dry. Once the soils were dry, they were gently crushed, homogenized, and then wetted with DDI water to achieve a 30% moisture content. The soils were kept in a container out of direct light and maintained at 30% water content in a moisture saturated environment. Soils were incubated in this manner for the duration of the study (i.e., at least 100 d).

Determination of Chromium on Soil

Total Cr on the soil was determined using a modification of EPA method 3052. The soil was digested in a CEM microwave, model MDS-81D, with hydrofluoric and nitric acid. Boric acid was added before sample analysis to facilitate the removal of hydrofluoric acid from solution through the formation of fluoroboric acid. Soils from the National Institute of Standards, with known concentrations of solid phase Cr, were also analyzed with each block of analysis. Samples were stored and analyzed for total Cr using inductively coupled plasma.

In Vitro Bioaccessibility

A physiologically based extraction test (PBET) was adapted from Ruby et al. (1996, 1999; Ruby, personal communication, 2000) to assess the in vitro bioaccessibility of Cr(III) from contaminated soils in humans. The method is designed to simulate the stomach digestive system in humans. The PBET method has been shown to agree with in vivo studies involving Pb-contaminated soils (Ruby et al., 1996) as well as As-contaminated soils (Rodriguez et al., 1999); however, limited data is currently available in the literature that evaluates Cr bioavailability in contaminated soils using in vivo methods (Witmer et al., 1989, 1991; Gargas et al., 1994), and this data does not appear useful for cross-correlating with the results of the current study. Nevertheless, the PBET method can serve as a useful approximation of Cr bioavailability until in vivo studies become available to validate the methods credibility with regard to Cr.

In the current study, triplicate 0.39 g moist samples (0.3 g dry wt) were placed in 50 mL polyethylene tubes to which 30 mL 0.4 M glycine at pH 1.5 and 37°C was added. The slurries were quickly placed in a rotating water bath of 37°C

Table 1. U.S. Department of Defense Army bases with their associated soil series designations.

Army bases by soil order	Facility location by state	Soil series
Ultisol		
Holston AAP	Tennessee	Allen
Fort Gillem	Georgia	Cecil
ORNL†	Tennessee	Minvale
Alfisol		
Seneca AD	New York	Angola
Indiana AAP	Indiana	Crider
Bluegrass Facility	Kentucky	Lawrence
Ft. Knox	Kentucky	Lenberg
Lexington Facility—LBAD	Kentucky	Lenberg
Inceptisol		
Letterkenny AD	Pennsylvania	Berks
ARDEC (Picatinny Arsenal)	New Jersey	Rockaway
Letterkenny	Pennsylvania	Weikert
ORNL†	Tennessee	Montevello
Spodosol		
Stratford Army Engine Plant	Connecticut	Charlton
Mollisol		
Kansas AAP	Kansas	Dennis
Lake City AAP	Missouri	Sibley
Aridisol		
Ft. Wingate	New Mexico	Doakum
Tolle Army Depot	Utah	Kzin
Desert Chem. Depot	Utah	Kzin
Dugway	Utah	Kzin
Hawthorne	Nevada	Oricto
Pueblo Chem. Depot	Colorado	Stoneham
Entisol		
Savanna Depot Activity	Illinois	Wakeland

† Department of Energy sites at the Oak Ridge National Laboratory.

Table 2. Select soil chemical and physical properties.†

	TOC	TIC	Clay	Silt	Fe	Mn	CEC	pH	pH
	%				g/kg		cmol/kg	5 mM CaCl ₂	DDI
Ultisol									
Allen A	1.55	0.56	8.7	29.5	6.95	0.31	7.7	4.59	5.05
Allen Ba	0.19	0.09	14.9	28.4	18.96	0.10	1.3	4.30	4.74
Cecil Ap	1.64	0.39	10.2	23.0	6.01	0.06	5.8	4.04	4.47
Cecil Bt1	0.29	0.21	44.8	15.5	32.56	0.11	1.6	4.44	4.48
Minvale Ap	1.89	0.99	6.1	59.0	7.71	1.51	6.0	6.01	6.61
Minvale Bt1	0.10	0.07	23.6	44.2	19.55	0.16	4.0	4.30	5.17
Alfisol									
Lawrence Apl	0.91	0.59	19.5	48.5	11.17	1.35	5.8	4.97	5.27
Lawrence Btl	0.11	0.10	25.8	38.3	17.53	0.29	3.7	4.28	4.91
Angola Ap	3.72	0.96	32.1	56.1	23.28	1.23	6.7	5.29	5.48
Crider Ap	0.55	0.39	22.5	75.8	13.34	0.72	5.6	6.57	6.84
Crider B2lt	0.21	0.13	30.9	67.2	13.38	0.30	5.4	5.27	5.63
Lenberg A	3.41	1.01	49.1	44.5	12.94	1.37	7.9	5.92	6.06
Lenberg Btl	0.36	0.25	64.7	29.5	15.69	0.12	5.5	4.35	4.77
Inceptisol									
Berks A	2.72	1.01	15.7	46.6	13.18	0.15	9.1	3.65	3.91
Rockaway A1	3.54	1.49	12.4	34.8	14.03	0.52	10.6	3.86	3.98
Rockaway B2t	0.21	0.18	12.6	32.1	17.34	0.16	3.7	4.10	4.41
Weikert Ap	3.97	2.37	24.4	56.2	21.41	6.47	13.3	4.44	4.70
Weikert Be	2.01	1.15	23.9	54.3	28.98	5.42	8.0	4.28	4.65
Montevello A	3.55	0.62	6.0	69.0	10.68	1.42	8.0	6.91	7.18
Montevello B	0.42	0.26	19.0	42.2	22.07	0.17	14.0	4.23	4.87
Spodosol									
Charlton A2	2.30	0.40	2.9	28.7	1.33	0.00	11.7	3.15	3.57
Mollisol									
Dennis Ap	1.32	0.89	15.9	66.1	15.11	0.60	8.7	5.82	6.08
Dennis Ba	0.38	0.41	29.7	57.5	24.29	0.59	4.4	4.77	5.28
Sibley A	1.06	0.49	23.5	69.7	8.23	0.67	7.1	6.36	6.66
Sibley B1	0.72	0.52	26.9	68.0	9.11	0.59	6.8	6.36	6.76
Aridisol									
Doakum Ab	0.28	0.08	10.8	24.8	4.74	0.19	6.9	6.94	7.42
Doakum Bt	0.39	0.18	29.3	15.0	6.86	0.16	7.0	6.87	7.39
Kzin A2	3.27	1.35	22.2	44.2	4.07	0.29	13.3	7.74	7.87
Kzin Bk	3.40	1.88	27.0	38.5	3.26	0.18	10.0	7.80	7.88
Oricto A2	0.09	0.94	10.2	34.7	2.92	0.34	13.7	8.72	9.60
Oricto Bt	0.16	1.10	23.2	27.5	3.16	0.29	8.6	9.01	9.60
Stoneham A	1.45	0.71	16.2	41.4	3.40	0.26	10.1	6.43	6.83
Stoneham Bt1	0.66	0.32	21.4	23.2	2.20	0.20	7.8	6.80	7.15
Entisol									
Wakeland Ap	0.92	0.00	23.8	64.7	8.82	0.71	6.1	5.86	6.09
Wakeland Cg1	0.56	0.25	21.1	66.4	9.18	0.80	5.7	5.77	6.07

† TOC, total organic carbon; TIC, total inorganic carbon; CEC, cation exchange capacity; DDI, double deionized.

and agitated at 30 ± 2 rpm for 1 h. Supernatant was separated from the solid via centrifugation. The pH of the supernatant was measured to ensure that the final pH was within ± 0.5 pH units of the initial pH. This scenario held for all cases. Thus, bioaccessibility was calculated as:

% Bioaccessibility =

$$\left(\frac{\text{Cr in PBET supernatant } (\mu\text{g/mL}) \times 30.0 \text{ mL} \div 0.3 \text{ g dry soil}}{\text{Cr on soil surface (mg/kg)}} \right) \times 100$$

The PBET pH of 1.5 simulates the most aggressive stomach digestive scenario, which is a condition indicative of human fasting. Conditions of higher pH, as a result of food intake, would most likely decrease Cr bioaccessibility even more profoundly than the results presented in the current study, thus offering a potential avenue for future research. Both Ruby et al. (1996) and Yang et al. (2002) found that soil Pb bioaccessibility was strongly pH dependent with soluble Pb decreasing profoundly over a pH range of 1.5 to 4.0.

Chromium Analysis

The PBET supernatant, soil spiking solution, and equilibrium solution were measured for Cr(VI) and total Cr (Cr_T).

Chromium(VI) was measured using a modified *s*-diphenylcarbohydrazide colorimetric method (Bartlett and James, 1979) using a UV-VIS spectrophotometer at wavelength 540 nm (HP model 8453, Palo Alto, CA). Analysis of Cr(VI) was performed immediately on rapidly cooled PBET solutions to avoid possible reduction of Cr(VI) to Cr(III) by glycine (Jardine et al., 1999). Independent studies revealed that Cr(VI) reduction by glycine at 37°C and 1 h was insignificant. Total Cr was measured on a Perkin Elmer AAnalyst 800 atomic absorption spectrophotometer (Wellseley, PA). Standards were made using an atomic absorption Cr standard (EM Industries, Hawthorne, NY). Chromium(III) was calculated as the difference between Cr_T and Cr(VI).

Modeling

A multiple regression technique in the statistical software package SigmaStat 2.0 (Jandel Scientific) was used to derive an expression that related Cr(III) sorption and bioaccessibility to common soil properties. The model was run using forward stepwise regression to determine the most salient soil properties for calculating sorption or bioaccessibility. Multiple linear regression was then employed to determine the linear equation to use when computing the Cr(III) sorption or bioaccessibility based on the important soil properties previously ascertained.

RESULTS AND DISCUSSION

Influence of Soil Properties on Chromium Sorption

Chromium sorption (i.e., adsorption and surface precipitation) by the 35 soils varied markedly with values ranging from 736 mg/kg to 17 460 mg/kg (Table 3). Sorption of Cr(III) was independent of horizon type where no distinct trend between A- and B-horizons was evident. The majority of the soils adsorbed between approximately 3000 mg/kg to approximately 6000 mg/kg with four soils as high as approximately 18 000 mg/kg. These four soils were all Aridisols and are noted for their high soil pH and for their high TIC content. Observed Cr(III) loading levels on many of these different soil types were similar to those measured on actual contaminated soils from the DoD sites. For example, actual contaminated Kzin soil (Xeric Torriorthents) from the Desert Chemical Depot contained 27 000 mg Cr/kg soil. Artificially contaminated Kzin soils in this study contained approximately 18 000 mg Cr/kg soil.

The large contrast in Cr(III) sorption by the various

soils can be explained by the differences in soil properties. Multiple linear regression showed that four soil properties were important in determining the amount of Cr adsorbed by the soils: pH, total inorganic carbon (TIC) content, clay content, and cation exchange capacity (CEC). The relationship describing Cr adsorption was:

$$\begin{aligned} \text{Cr(III) (mg/kg on soil)} = & -12\,666.3 + \\ & (113.8 \times \% \text{ clay}) + (364.6 \times \text{CEC}) + \\ & (1743.2 \times \% \text{ TIC}) + (1916.7 \times \text{soil pH}) \end{aligned}$$

Chromium(III) sorption by the soils was strongly correlated with these soil properties ($r^2 = 0.794$) suggesting that nearly 80% of the variability in Cr(III) sorption could be described by pH, TIC, clay, and CEC (Table 4). Incorporating the other measured soil properties from Table 2 (e.g., Fe-oxide content, TOC, etc.) did not improve the model fit. In fact, TIC could have been removed from the model if necessary, since the other three independent soil variables could describe approximately 77% of the variability in Cr(III) sorption. The four-parameter model above was statistically rigorous at the 95% confidence level since P values for the independent variables were all <0.05 (Table 4). Thus, it can be concluded that the independent variables, the soil properties, significantly contribute to predicting the dependent variable, Cr sorption. The Variance inflation factor (VIF) also suggested that collinearity between independent variables was not significant (Table 4). Values for VIF that are 1.0 or slightly larger suggest that the variables do not show multicollinearity and that the parameter estimates are reliable. Collinearity becomes an issue when values of VIF exceed 4.0. This model also passed the Normality Test (indicating that the data was normally distributed) and the Constant Variance Test (suggesting that the variance of the dependent variables was constant). One of the most important criteria of a successful model, however, is the true physical significance of the model parameters. Our model suggests that Cr(III) sorption is enhanced by higher soil pH, more TIC (i.e., carbonates), more clay, and higher CEC. For a sparingly soluble cation, such as Cr(III), these soil conditions should enhance sequestration as the model suggests.

The pH of the soil affects the solubility and form of Cr and therefore affects sorption. As the soil pH increases, the amount of Cr on the soil increases. At low pH, Cr(III) is adsorbed or complexed on soil negative charges; at higher soil pH values, >5.5 , Cr precipitates

Table 3. Chromium(III) solid phase concentrations on the various soils and their corresponding bioaccessibility after 1 and 100 d aging.

	C_T on soil	1 day % Cr(III) bioaccessible	100 day % Cr(III) bioaccessible
Ultisol	mg/kg		
Allen A	940.32	16.37	8.13
Allen Ba	736.15	31.11	17.98
Cecil Ap	1 342.49	18.84	9.90
Cecil Bt1	2 333.76	41.77	28.34
Minvale Ap	2 261.67	15.88	8.55
Minvale Bt1	1 294.09	54.65	35.52
Alfisol			
Lawrence Ap1	2 586.96	26.03	11.62
Lawrence Bt1	2 359.18	41.48	28.10
Angola Ap	9 408.00	32.40	16.58
Crider Ap	3 719.38	33.90	22.88
Crider B2t	4 247.30	50.30	32.35
Lenberg A	8 169.92	30.27	20.28
Lenberg Bt1	7 254.84	50.89	41.63
Inceptisol			
Berks A	2 275.20	18.77	7.67
Rockaway A1	2 482.08	11.62	6.46
Rockaway B2t	1 525.58	32.96	22.36
Weikert Ap	5 561.77	12.21	5.62
Weikert Be	3 229.97	19.73	10.35
Montevello A	5 925.66	19.03	7.03
Montevello B	2 751.57	47.71	26.23
Spodosol			
Charlton A2	1 721.95	27.65	21.26
Mollisol			
Dennis Ap	3 577.05	19.43	13.67
Dennis Ba	3 521.90	33.61	26.68
Sibley A	4 436.16	29.78	20.50
Sibley B1	4 689.17	36.36	25.37
Aridisol			
Doakum Ab	2 507.82	31.19	16.60
Doakum Bt	5 964.29	39.40	32.77
Kzin A2	16 306.33	17.22	14.00
Kzin Bk	12 452.82	24.03	19.81
Oricto A2	17 460.00	13.66	10.26
Oricto Bt	15 964.28	18.45	16.44
Stoneham A	4 377.44	29.27	18.97
Stoneham Bt1	4 599.44	33.70	24.82
Entisol			
Wakeland Ap	4 262.61	32.33	21.08
Wakeland Cg1	3 802.32	37.68	24.79

Table 4. Parameter estimates, standard errors, and statistics obtained from a multiple linear regression analysis that related soil properties to Cr(III) sorption.[†]

Parameter	Value	SE	P	VIF
Intercept	-12 666.3	1 794.5	<0.001	-
% Clay	113.8	30.4	<0.001	1.119
pH in DDI	1 916.7	250.7	<0.001	1.079
CEC, cmol/kg	364.6	155.7	0.026	1.902
% TIC	1 743.2	850.1	0.049	1.670
r^2	0.794		<0.001	

[†] DDI, double deionized; CEC, cation exchange capacity; TIC, total inorganic carbon; VIF, Variance Inflation Factor.

as hydroxides covering the surface of the soil (Bartlett and Kimble, 1976). It was presumed by Bartlett and Kimble (1976) and James and Bartlett (1983) that the Cr(III) precipitate consisted of macromolecules with Cr ions in six coordination with water and hydroxy groups. Studies by Fendorf et al. (1994) and Fendorf and Sparks (1994), using x-ray adsorption spectroscopy (XAS), showed that with a low Cr(III) surface coverage the principle mechanism was adsorption with an inner-sphere monodentate complex on the silica. With increased surface coverage (>20%), precipitation likely occurred and became the dominant sorption mechanism.

As with pH, TIC or carbonate content in soils enhanced Cr(III) sorption. The mechanism of increased sequestration is most likely a localized pH effect at the carbonate surface, which promotes the formation of $\text{Cr}(\text{OH})_3$ species. The localized pH effect is the most plausible scenario since there was no correlation between soil pH and soil TIC, thus explaining why collinearity was not a problem for these parameters when the model was fit to the Cr(III) sorption data. Several acidic Inceptisols derived from interbedded limey shales and limestone have relatively large residual carbonate contents (Table 2), due to the slow dissolution of local scale dolomite, and this may serve to enhance Cr(III) sequestration in these systems, even though the overall bulk soil pH is acidic.

The model also shows a positive correlation between the amount of Cr adsorbed and the soil clay content and CEC. This was expected since clay minerals tend to be dominated by negatively charged sites on the surface due to isomorphic substitution (Klein and Hurlbut, 1993). These negatively charged sites attract the cation Cr^{3+} and a weak, electrostatic bond is formed. The more negatively charged sites that are available (i.e., larger CEC), the greater propensity for Cr(III) sorption. Further, clay minerals typically have a large surface area that is capable of accommodating large quantities of Cr^{3+} and $\text{Cr}(\text{OH})_3$ precipitated phases. The more surface area a soil has, the more reactive sites the soil has, and consequently the more Cr that will adsorb to the soil.

Influences of Soil Properties on Chromium Bioaccessibility

The bioaccessibility of Cr(III), as measured by the PBET method, varied widely as a function of soil type with most soils limiting bioaccessibility to <45% and <30% after 1 and 100 d soil-Cr aging, respectively (Table 3, Fig. 1a–e). Bioaccessibility values were consistently higher for 1 d aging vs. 100 d aging. For all soils the percent bioaccessibility ranged from 3.0 to 54.7% at Day 1 and 1.5 to 35.5% at 100 d (Table 3, Fig. 1a–e). The aging effect is related to the enhanced stability of Cr on the soil surface with time. Structural reorientation of Cr surface bonds or slow precipitation reactions can account for the stronger sorption of Cr at longer times (Karthein et al., 1991). Previous studies by Stewart et al. (2003) have shown that aging effects are insignificant after 100 d and that the 100 d data are most relevant

to actual DoD-contaminated soils. In general, the A-horizon soils had the lowest percent bioaccessible values, even when they adsorb more Cr(III) on the soil vs. the B horizons (Table 3). Bioaccessibility did not appear to be a function of soil order, suggesting that detailed soil series data, as is used in the current study, was necessary for predictive purposes (Table 3, Fig. 1a–e). Chromium(VI) was also measured in the PBET extractant to monitor for oxidation of Cr(III) to Cr(VI). The proportion of bioaccessible Cr that was Cr(VI) was always <1%, suggesting that oxidation reactions were minimal or that any oxidation products of Cr(VI) were tightly held by the soil. These results are consistent with the data presented by Stewart et al. (2003), which showed limited bioaccessibility of Cr(VI) in several soils.

As demonstrated by Stewart et al. (2003), bioaccessibility values leveled off and reached near equilibrium after the first 50 to 100 d. Thus, the 100 d bioaccessibility data is most appropriate for use in the modeling endeavor. Stepwise multiple regression indicated two combinations of variables considered instrumental in predicting the bioaccessibility of Cr(III) in soils: (i) % clay and % TIC and (ii) % clay and % TOC. Using the independent variables from Table 2, the most significant model revealed that the bioaccessibility of Cr(III) on the soils was correlated with clay and TIC of the soil (Table 5). The relationship describing Cr(III) bioaccessibility was:

$$\begin{aligned} \% \text{ Cr(III) bioaccessible} = \\ 16.02 + (0.426 \times \% \text{ clay}) - (9.56 \times \% \text{ TIC}) \end{aligned}$$

with an r^2 value of 0.722, which indicated that as much as 72% of the variability in Cr bioaccessibility was explained by the model (Fig. 2). The model was statistically rigorous at the 99% confidence level since P values for the independent variables were well below 0.01, indicating that they all contributed to predicting the % bioaccessible Cr(III) (Table 5). Values for VIF were all nearly 1.000, indicating that there was no redundant information in the other independent variables, i.e., soil properties, and that collinearity between independent variables was not of concern. This indicated that parameter estimates in the model were reliable, which is in agreement with the low standard errors on the estimated values (Table 5). The model also passed the Normality Test and the Constant Variance Test, suggesting that the data was normally distributed around the regression line and that the variance present in the dependent variable is constant. Most important, however, is the true physical significance of the model parameters. The model suggests that Cr(III) bioaccessibility decreases as the TIC content increases and as the clay content decreases. As shown with the Cr sorption data, Cr(III) sequestration is enhanced by soils with high levels of TIC. The presence of TIC promotes the formation of solid phase Cr(III)–hydroxides that are sparingly soluble, even under acidic conditions. These hydroxides [i.e., $\text{Cr}(\text{OH})_3$] precipitate and cover the surface of the soil and are not easily bioaccessible even in the presence of the low pH in the simulated stomach fluid of the PBET. Conse-




Fig. 1. Percentage Cr(III) bioaccessibility after 1 and 100 d Cr-soil aging for (a) Ultisols; (b) Alfisols; (c) Inceptisols; (d) Spodosols, Mollisols, Entisols; and (e) Aridisols.




Fig. 1. Continued.

quently as the TIC content increases the bioaccessibility of Cr(III) in soil decreases. As shown with the Cr sorption data the clay content on the soil was also correlated with the amount of Cr sequestration and thus should be important in determining bioaccessibility. The bioaccessibility model suggested that, as the clay content of the soils increased, the percent of Cr on the soil that is bioaccessible also increased. Since the mechanism of Cr retardation on clay minerals is primarily weak electrostatic bonds, these bonds are easily broken under the conditions of the PBET, allowing Cr to desorb from the soil and be released into solution during the simulated digestion.

Stepwise multiple regression analysis also indicated that Cr(III) bioaccessibility was significantly correlated with clay and TOC content of the soil (Table 6). The relationship describing Cr(III) bioaccessibility was:

$$\% \text{ Cr(III) bioaccessible} =$$

$$15.54 + (0.408 \times \% \text{ clay}) - (3.78 \times \% \text{ TOC})$$

with an r^2 value of 0.674. This relationship was similar to the clay/TIC model where higher quantities of TIC and TOC resulted in decreased Cr(III) bioaccessibility. When clay, TIC, and TOC were used in the same model, the contribution of TOC was not significant at the 90%

Table 5. Parameter estimates, standard errors, and statistics obtained from a multiple linear regression analysis that related soil properties (clay and TIC) to percent Cr(III) bioaccessibility.†

Parameter	Value	SE	P	VIF
Intercept	16.02	1.99	<0.001	—
% Clay	0.426	0.0671	<0.001	1.002
% TIC	−9.56	1.54	<0.001	1.002
r^2	0.722		<0.001	

† TIC, total inorganic carbon; VIF, Variance Inflation Factor.




Fig. 2. The observed (data points) and model fitted (grid surface) relationship between the two most significant independent variables (% clay and TIC) and % Cr(III) bioaccessibility using the model: % Cr(III) bioaccessible = $16.02 + (0.426 \times \% \text{ clay}) - (9.56 \times \% \text{ TIC})$ and an r^2 value of 0.722.

confidence level ($P = 0.115$). This scenario may be an artifact of our limited data set, where the most appropriate model, in fact, includes both TIC and TOC along with clay content. A more extensive data set will be necessary to test this hypothesis. Nevertheless, the model using clay and TOC was statistically rigorous at the 99% confidence level since P values for the estimated parameters were <0.01 and the VIF values are approximately 1.000, indicating that the variables all contribute significantly to the equation and that no multicollinearity was present among the independent variables. This model passed the Normality Test and the Constant Variance Test. The model suggested that as the clay content decreased and the TOC content increased, the % Cr(III) bioaccessible decreased (Fig. 3). The trend regarding clay content is consistent with the previous model and the limited bioaccessibility of Cr in the presence of higher system organic C is conceptually correct. Organic matter found in soil is a major contributor to the overall negative charge in soils and thus is an important sorbent for heavy metal cations (Sparks, 1995). Organic matter has the ability to form strong bonds with the Cr(III) with the metal not readily released during the PBET process. As Cr(III) is considered a Lewis hard acid, it forms stable complexes with the carboxyl group of the organic matter (Sparks, 1995). These bonds are stable

Table 6. Parameter estimates, standard errors, and statistics obtained from a multiple linear regression analysis that related soil properties (clay and TOC) to percent Cr(III) bioaccessibility.[†]

Parameter	Value	SE	P	VIF
Intercept	15.54	2.16	<0.001	—
% Clay	0.408	0.073	<0.001	1.010
% TOC	-3.78	0.711	<0.001	1.010
r^2	0.674		<0.001	

[†] TOC, total organic carbon; VIF, Variance Inflation Factor.




Fig. 3. The observed (data points) and model fitted (grid surface) relationship between the two independent variables (% clay and TOC) and % Cr(III) bioaccessibility using the model: % Cr(III) bioaccessible = $15.54 + (0.408 \times \% \text{ clay}) - (3.78 \times \% \text{ TOC})$ with an r^2 value of 0.674.

and not easily broken. The current model again explains more than 67% of the variability in Cr(III) bioaccessibility and should be useful for soils low in carbonate (TIC).

ENVIRONMENTAL SIGNIFICANCE

This study has shown that site assessments of soil metal bioaccessibility based solely on total soil metal concentrations may not accurately reflect the risk posed by the soils. The sequestering properties of soil significantly lower the percent of Cr bioaccessible upon ingestion of the otherwise labile Cr. Chromium(III) can be immobilized as strongly bound species on clay and organic matter, and Cr-hydroxide precipitates on soil mineral surfaces. It has been shown that common soil properties are strongly correlated with Cr(III) bioaccessibility. The availability of these soil properties is commonplace (e.g., NRCS database), which allows the percent bioaccessibility of Cr(III) to be estimated for a variety of contaminated sites whose remediation is pending. The ability to rapidly assess metal bioaccessibility in soils will facilitate decision making strategies regarding the need for more detailed and expensive site-specific bioavailability (e.g., animal feeding) studies, which are designed to assess actual clean-up needs at contaminated DoD sites and other sites to a level safe for human use. Such in vivo studies are lacking with regard to Cr, but research in this area is currently underway (M.V. Ruby, personal communication, 2002).

ACKNOWLEDGMENTS

We appreciate the efforts of Ms. Cathy Vogel and Dr. Andrea Leeson, the contract officers for the U.S. DoD who supported this work and the efforts of Mr. Warren Lynn of the National Resource Conservation Service (NRCS) who provided us with most of the soils for this study.

REFERENCES

- Bartlett, R., and B. James. 1979. Behavior of chromium in soils: III. Oxidation. *J. Environ. Qual.* 8:31–35.
- Bartlett, R.J., and J.M. Kimble. 1976. Behavior of chromium in soils: I. Trivalent forms. *J. Environ. Qual.* 5:379–383.
- Chung, J., R.J. Zasoski, and S. Lim. 1994. Kinetics of chromium(III) oxidation by various manganese oxides. *Korean J. Agric. Chem. Biotechnol.* 37:414–420.
- Davis, S., P. Waller, R. Buschbom, J. Ballow, and P. White. 1990. Quantitative estimates of soil ingestion in normal children between the ages of 2 and 7 years. Population-based estimates using aluminum, silicon, and titanium as soil tracer elements. *Arch. Environ. Health* 45:112–122.
- Dragun, J. 1998. The soil chemistry of hazardous materials. Amherst Scientific Publ., Amherst, MA.
- Fendorf, S.E., M. Fendorf, D.L. Sparks, and R. Gronsky. 1992. Inhibitory mechanisms of Cr(III) oxidation by δ -MnO₂. *J. Colloid Interface Sci.* 153:37–54.
- Fendorf, S.E., G.M. Lamble, M.G. Stapleton, M.J. Kelly, and D.L. Sparks. 1994. Mechanisms of chromium (III) sorption on silica: I. Cr(III) surface structure derived by extended x-ray adsorption fine structure spectroscopy. *Environ. Sci. Technol.* 28:284–289.
- Fendorf, S.E., and D.L. Sparks. 1994. Mechanisms of chromium (III) sorption on silica: II. Effects of reaction conditions. *Environ. Sci. Technol.* 28:290–297.
- Fendorf, S.E., and R.J. Zasoski. 1992. Chromium(III) oxidation by δ -MnO₂: I. Characterization. *Environ. Sci. Technol.* 26:79–85.
- Gargas, M.L., R.L. Norton, M.A. Harris, D.J. Paustenbach, and B.L. Finley. 1994. Urinary excretion of chromium following ingestion of chromite-ore processing residues in humans: Implications for biomonitoring. *Risk Anal.* 14:1019–1024.
- Hamel, S.C., B. Buckley, and P.J. Lioy. 1998. Bioaccessibility of metals in soils for different liquid to solid ratios in synthetic gastric fluid. *Environ. Sci. Technol.* 32:358–362.
- James, B.J., and R.J. Bartlett. 1983. Behavior of chromium in soils: VII. Adsorption and reduction of hexavalent forms. *J. Environ. Qual.* 12:177–181.
- Jardine, P.M., S.E. Fendorf, M.A. Mayes, S.C. Brooks, and W.B. Bailey. 1999. Fate and transport of hexavalent chromium in undisturbed heterogeneous soil. *Environ. Sci. Technol.* 33:2939–2944.
- Karthein, R., M. Motschi, A. Schweiger, S. Ibric, B. Sulzberger, and W. Stumm. 1991. Interactions of chromium(III) complexes with hydrous delta-Al₂O₃ rearrangements in the coordination sphere studied by electron-spin-resonance and electron spin-echo spectroscopies. *Inorg. Chem.* 30:1606–1611.
- Klein, C., and C. Hurlbut, Jr. 1993. Manual of mineralogy. 21st ed. Wiley, New York.
- Levis, A.G., and V. Bianchi. 1982. Mutagenic and cytogenic effects of chromium compounds. p. 171–208. *In* S. Langjard (ed.) Biological and environmental aspects of chromium. Elsevier Biomedical Press, New York.
- Linz, D.G., and D.V. Nakles (ed.) 1997. Environmentally acceptable endpoints in soil. Am. Academy of Environ. Eng., New York.
- Losi, M.E., C. Amrhein, and W.T. Frankenberger, Jr. 1994. Bioremediation of chromate-contaminated groundwater by reduction and precipitation in surface soils. *J. Environ. Qual.* 23:1141–1150.
- Mehra, O.P., and M.L. Jackson. 1960. Iron oxide removed from soils and clays by a dithionite–citrate system buffered with sodium bicarbonate. *Clays Clay Miner.* 7:317–327.
- Nriagu, J.O., and E. Nieboer. 1988. Chromium in the natural and human environments. John Wiley & Sons, New York.
- Patterson, R.R., S. Fendorf, and M. Fendorf. 1997. Reduction of hexavalent chromium by amorphous iron sulfide. *Environ. Sci. Technol.* 31:2039–2044.
- Paustenbach, D.J. (ed.) 1989. The risk assessment of environmental and human health hazards: A textbook of case studies. John Wiley & Sons, New York.
- Proctor, D.M., E.C. Shay, and P.K. Scott. 1997. Health-based soil action levels for trivalent and hexavalent chromium: A comparison with state and federal standards. *J. Soil Contam.* 6:595–648.
- Rodriguez, R.R., N.T. Basta, S.W. Casteel, and L.W. Pace. 1999. An in vitro gastrointestinal method to estimate bioavailable arsenic in contaminated soils and solid media. *Environ. Sci. Technol.* 33:642–649.
- Ruby, M.V., A. Davis, R. Schoof, S. Eberle, and C.M. Sellstone. 1996. Estimation of lead and arsenic bioavailability using a physiologically based extraction test. *Environ. Sci. Technol.* 30:422–430.
- Ruby, M.V., R. Schoof, W. Brattin, M. Goldade, G. Post, M. Harnois, D.E. Mosby, S.W. Casteel, W. Berti, M. Carpenter, D. Edwards, D. Cragin, and W. Chappell. 1999. Advances in evaluating the oral bioavailability of inorganics in soil for use in human health risk assessment. *Environ. Sci. Technol.* 33:3697–3705.
- Sheehan, P.J., D.M. Meyer, M.M. Sauer, and D.J. Paustenbach. 1991. Assessment of the human health risks posed by exposure to chromium contaminated soils. *J. Toxicol. Environ. Health* 32:161–201.
- Sheppard, S.C., W.G. Evanden, and W.J. Achwartz. 1995. Ingested soil: Bioavailability of sorbed lead, cadmium, iodine, and mercury. *J. Environ. Qual.* 24:498–505.
- Skowronski, G.A., M. Seide, and M.S. Abdel-Rahman. 2001. Oral bioaccessibility of trivalent and hexavalent chromium in soil by simulated gastric fluid. *J. Toxicol. Environ. Health Part A* 63:351–362.
- Sparks, D.L. 1995. Environmental soil chemistry. Academic Press, New York.
- Stewart, M.A., P.M. Jardine, M.O. Barnett, L.D. McKay, T.L. Mehlhorn, S.E. Fendorf, and K. Paul. 2003. Effects of contaminant concentration, aging, and soil properties on the bioaccessibility of Cr(III) and Cr(VI) contaminated soil. *Soil Sediment Contam.* (in press).
- U.S. Environmental Protection Agency. 1996. USEPA Method 3052. Microwave assisted acid digestion of siliceous and organically based matrices. [Online.] [20 p.] <http://www.epa.gov/epaoswer/hazwaste/test/pdfs/3052.pdf>. USEPA, Washington, DC.
- Witmer, C.M., R. Harris, and S.I. Shupack. 1991. Oral bioavailability of chromium from a specific site. *Environ. Health Perspect.* 92: 105–110.
- Witmer, C.M., H.S. Park, and S.I. Shupack. 1989. Mutagenicity and disposition of chromium. *Sci. Total Environ.* 86:131–138.
- Yang, J.K., M.O. Barnett, P.M. Jardine, and S.C. Brooks. 2002. Factors controlling the bioaccessibility of arsenic(V) and lead(II) in soil. *Soil Sediment Contam.* (In press).

APPENDIX C

**YANG, J-K., M.O. BARNETT, P.M. JARDINE, N.T. BASTA, AND S.W. CASTEEL. 2002.
ADSORPTION, SEQUESTRATION, AND BIOACCESSIBILITY OF AS(V) IN SOILS.
ENVIRONMENTAL SCIENCE AND TECHNOLOGY. 36:4562-4569.**

Adsorption, Sequestration, and Bioaccessibility of As(V) in Soils

Jae-Kyu Yang,[†] Mark O. Barnett,^{*†} Philip M. Jardine,[‡] Nicholas T. Basta,[§] and Stan W. Casteel[Ⓛ]

Department of Civil Engineering, 238 Harbert Engineering Center, Auburn University, Auburn, Alabama 36849, Environmental Sciences Division, Oak Ridge National Laboratory, P.O. Box 2008, Oak Ridge, Tennessee 37831, Department of Plant and Soil Sciences, Oklahoma State University, Stillwater, Oklahoma 74078, and Veterinary Medical Diagnostic Laboratory, University of Missouri-Columbia, Columbia, Missouri 65211

Received for review December 30, 2001

Revised manuscript received August 12, 2002

Accepted August 21, 2002

Abstract:

The influence of various soil physical and chemical properties (Fe and Mn oxides, pH, cation exchange capacity, total inorganic and organic carbon, and particle size) on As(V) adsorption, sequestration, and relative bioaccessibility (as a surrogate for oral bioavailability) was investigated in a wide range of well-characterized soils over a 6-month period. Arsenic(V) bioaccessibility was measured using a streamlined version of a physiologically based extraction test (PBET), designed to replicate the solubility-limiting conditions in a child's digestive tract. The soil's dithionite-citrate-bicarbonate (DCB) extractable Fe oxide content was the most important (and only statistically significant) soil property controlling the initial degree of adsorption. Sequestration, as measured by the reduction in bioaccessibility over time, occurred to a significant extent in 17 of 36 (47.2%) soils over the first 3 months. In contrast, only 4 of 36 (11.1%) soils exhibited a significant reduction in bioaccessibility from 3 to 6 months. Soil pH was the most important (and only statistically significant) soil property affecting the decrease in bioaccessibility upon aging for 6 months. Soils with pH < 6 generally sequestered As(V) more strongly over time, whereas those with pH > 6 generally did not. The Fe oxide content and pH were the most important soil properties governing the steady-state bioaccessibility of As(V) in soil. Two multivariable linear regression models of steady-state As(V) bioaccessibility were developed using soil properties as independent variables. Generally, soils having higher Fe oxide content and lower soil pH exhibited lower bioaccessibility. These models were able to account for ~75-80% of the variability in steady-state bioaccessibility and independently predict bioaccessibility in five soils within a root-mean-square error (RMSE) of 8.2-10.9%. One of these models was also able to predict within an RMSE of 9.5% the in vivo bioavailability of As in nine contaminated soils previously used in swine dosing trials. These results indicate the

bioaccessibility, and thus, potentially the bioavailability of otherwise soluble As(V) added to soils (i.e., the worst-case bioavailability scenario) is significantly reduced in some soils over time, particularly those with lower pH and higher Fe oxide content. These results also provide a means of estimating As(V) bioaccessibility and bioavailability on the basis of soil properties.

DOI: [10.1021/es011507s](https://doi.org/10.1021/es011507s)

Reprints are available from:

http://pubs3.acs.org/acs/journals/doilookup?in_doi=10.1021/es011507s

or contact Philip Jardine at jardinepm@ornl.gov

APPENDIX D

STEWART, M.A., P.M. JARDINE, C.C. BRANDT, M.O. BARNETT, S.E. FENDORF, L.D. MCKAY, T.L. MEHLHORN AND K. PAUL. 2003b. EFFECTS OF CONTAMINANT CONCENTRATION, AGING, AND SOIL PROPERTIES ON THE BIOACCESSIBILITY OF CR(III) AND CR(IV) IN SOIL. SOIL AND SEDIMENT CONTAMINATION. 12:1-21.

Effects of Contaminant Concentration, Aging, and Soil Properties on the Bioaccessibility of Cr(III) and Cr(VI) in Soil*

M. A. Stewart,¹ P. M. Jardine,^{1*} C. Brandt,¹ M. O. Barnett,² S. E. Fendorf,³ L. D. McKay,⁴ T. L. Mehlhorn,¹ and K. Paul¹

¹Environ. Sci. Div., Oak Ridge National Lab., P.O. Box 2008, Oak Ridge, TN 37831-6038;

²Dept. of Civil Engineering, 208 Harbert Engineering Center, Auburn Univ., AL 36849-5337; ³Stanford Univ., Dept. Geol. and Environ. Sci., Stanford, CA 94305; ⁴Dept. of Geological Sciences, Univ. of Tennessee, Knoxville, TN 37996-1410

Contaminated soils at numerous U.S. Department of Defense, Department of Energy, and other industrial facilities often contain huge inventories of toxic metals such as chromium. Ingestion of soil by children is often the primary risk factor that drives the need for remediation. Site assessments are typically based solely on total soil-metal concentrations and do not consider the potential for decreased bioaccessibility due to metal sequestration by soil. The objectives of this research are to investigate the effect of soil properties on the bioaccessibility of Cr(III) and Cr(VI) as a function of contaminant concentration and aging. The A and upper B horizons of two well-characterized soils, representative of Cr-contaminated soils in the southeastern United States, were treated with varying concentration of Cr(III) and Cr(VI) and allowed to age. The bioaccessibility of the contaminated soils was measured over a 200-d time period using a physiologically based extraction test (PBET) that was de-

signed to simulate the digestive process of the stomach. The sorption of Cr(III) and Cr(VI) varied significantly as a function of soil type and horizon, and the oxidation state of the contaminant. Solid phase concentrations with Cr(III) were significantly greater than Cr(VI) for any given initial Cr concentration. This is consistent with the mechanisms of Cr(III) vs. Cr(VI) sequestration by the soils, where the formation of Cr(III)-hydroxides can result in the accumulation of large mass fractions of contaminant on mineral surfaces. Overall, Cr bioaccessibility decreased with duration of exposure for all soils and at all solid phase concentrations, with aging effects being more pronounced for Cr(III). The decrease in Cr bioaccessibility was rapid for the first 50 d and then slowed dramatically between 50 and 200 d. In general, the effects of Cr solid phase concentration on bioaccessibility was small, with Cr(III) showing the most pronounced effect; higher solid phase concentrations resulted in a decrease in bioaccessibility. Chemical extraction methods and X-ray Adsorption Spectroscopy analyses suggested that the bioaccessibility of Cr(VI) was significantly influenced by reduction processes catalyzed by soil organic carbon. Soils with sufficient organic carbon had lower Cr bioaccessibility values (~10 to 20%) due to an enhanced reduction of Cr(VI) to Cr(III). In soils where organic carbon was limited and reduction processes were minimal, the bioaccessibility of Cr(VI) dramatically increased (~60 to 70%).

* This research was sponsored by the U.S. Department of Defense (DoD) Strategic Environmental Research and Development Program. Oak Ridge National Lab is managed by the University of Tennessee – Battelle LLC, under contract DE – AC05 – 00OR22725 with the U.S. Department of Energy.

** Corresponding author (jardinepm@ornl.gov).

Key Words: metal bioavailability, metal sequestration by soil, redox transformations, X-ray absorption spectroscopy.

INTRODUCTION

Chromium is used in many industrial processes, including electroplating, leather tanning, pulp production, and wood preservation, and, consequently, can be found throughout the environment (Nriagu and Nieboer, 1988). There are two main oxidation states of chromium found in the environment, anionic Cr(VI) and cationic Cr(III). The two forms of chromium have distinct behaviors in subsurface environments. The anionic Cr(VI) is considered to be highly mobile in soils, while the Cr(III) cation is believed to be significantly less mobile (Chung *et al.*, 1994; Fendorf *et al.*, 1997; Jardine *et al.*, 1999). In regards to human health, the two forms of Cr also have major differences, with Cr(VI) considered carcinogenic and mutagenic even at low concentrations, while Cr(III) is considered potentially harmful only at high concentrations (Levis and Bianchi, 1982).

Human health is the usual risk driver that motivates the likelihood of remediation at Cr- contaminated sites. The exposure pathway of concern is usually the ingestion of contaminated soil, especially by children who traditionally have greater hand-to-mouth contact (Paustenbach, 1989; Davis *et al.*, 1990; Sheehan *et al.*, 1991; Skowronski *et al.*, 2001). U.S. EPA soil action levels for Cr(III) and Cr(VI) are 78,000 and 390 mg/kg, respectively, which are protective of soil-ingestion exposures for children in residential sites. However, certain states within the U.S. have designated action levels as low as 310 and 0.2 mg/kg for Cr(III) and Cr(VI), respectively (Proctor *et al.*, 1997). When regulators establish clean-up criteria for chromium-contaminated soils, the ubiquitous metal-sequestering properties of the soils are typically not taken into account (Proctor *et al.*, 1997). Instead, the standards are generally universal for all soils and are usually based on that of a soluble salt of the metal and the assumption that 100% of metal present will be absorbed into the body (Ruby *et al.*, 1999). In order to accurately assess the health risk posed by metal-contaminated sites, an improved understanding of the influence of soil sequestration on the bioaccessibility of Cr is needed, where bioaccessibility is defined as that amount of contaminant, which is soluble due to gastric function and has the potential to cross the intestinal wall (Hamel *et al.*, 1998).

Chromium adsorption in soil occurs under different conditions based on the oxidative state of the Cr ion. The Cr anion, Cr(VI), generally adsorbs to positively charged mineral surfaces via electrostatic attraction. Thus, conditions of decreasing pH result in enhanced adsorption of Cr(VI) (Zachara *et al.*, 1989). Surfaces with proton specific sites, particularly iron oxides, are mostly responsible for Cr(VI) adsorption (Davis and Leckie, 1980; Zachara *et al.*, 1987, 1988). Factors

interfering with Cr(VI) adsorption include the presence of SO_4^{2-} , the presence of dissolved inorganic carbon (DIC), and Al substitution for Fe in oxides. With a limited number of positive surface sites in soil, there is often competition from SO_4^{2-} and DIC for those sites (Leckie *et al.*, 1980; James and Bartlett, 1983; Zachara *et al.*, 1987, 1988, 1989). Ainsworth *et al.* (1989) concluded that Al substitution in oxides reduces the amount of chromate adsorbed due to the difference in the charge characteristics of the surface sites.

Another important mechanism of Cr(VI) sequestration by soils is the reduction of Cr(VI) to sparingly soluble Cr(III). Electron donors such as organic matter and Fe(II) are capable of reducing Cr(VI). Organic matter and surface bound organics are extremely effective at reducing Cr(VI) to Cr(III) under acidic conditions (Bartlett and Kimble, 1976b; Jardine *et al.*, 1999). Likewise, Fe(II) bearing minerals are known to rapidly reduce Cr in soils (Anderson *et al.*, 1994; Peterson *et al.*, 1997). Low soil pH facilitates the reduction reaction through the release of Fe(II) from soils (Eary and Rai, 1991). Iron sulfides also have the ability to rapidly reduce Cr(VI) to Cr(III), suggesting that complete dissolution of Fe(II) does not have to occur before the Cr can be reduced (Patterson *et al.*, 1997). These results imply that the reduction is taking place at the solid-solution interface making, FeS an effective reductant of Cr(VI).

Cationic Cr(III) also sorbs to soil through a variety of mechanisms. The pH of the soil has a strong influence on Cr(III) adsorption because changes in pH affect the variable charge on minerals and organic matter. Conditions of higher pH creates more negative surface sites on soil mineral surface and organic matter to which Cr(III) can sorb (Sparks, 1995). Further, at pH conditions above 5.5, Cr(III) rapidly precipitates from solution and forms hydroxides on the soil surface (Bartlett and Kimble, 1976a). These hydroxides have low solubility and therefore are not likely to dissolve and reenter the soil solution (Losi *et al.*, 1994).

With all the highly variable factors influencing chromium's ability to sorb to the soil surface, blanket clean-up regulations that ignore the importance of individual soil properties may not be accurate with regard to human health risk. The objective of this research was to investigate the effect of soil properties on the bioaccessibility of Cr(III) and Cr(VI) as a function of contaminant concentration and aging. We show that soils can strongly sequester both anionic and cationic forms of Cr, which, under certain circumstances, dramatically decreases toxic metal bioaccessibility.

METHODS

Soil Type and Characterization

The A and upper B horizons of two soils were obtained from the Melton Valley and Walker Branch watersheds on the Oak Ridge Reservation (ORR) in eastern Tennessee. The soils are representative of Cr-contaminated sites common to the

southeastern U.S. Selected physical and geochemical properties of these soils are listed in Table 1. The Melton Valley soil is an acidic Inceptisol derived from interbedded shales and limestone (Kooner *et al.*, 1995; Jardine *et al.*, 1999; Driese *et al.*, 2001). The soils are extensively weathered and devoid of carbonates. Illites dominate the $< 2 \mu\text{m}$ clay fraction, and the clays are heavily coated with amorphous Fe – oxides and goethite. The pH and cation exchange capacity (CEC) of these soils range from 4 to 7 and 10 to 20 $\text{cmol}_\text{c} \text{ kg}^{-1}$, respectively (Jardine *et al.*, 1989). Walker Branch soils are an acidic Ultisol that has been weathered from the Knox Group (Arnseth and Turner, 1988), a dolostone sequence with occasional interbeds of limestone and shale. The soils are also extensively weathered and devoid of carbonates. Kaolinite dominates the $< 2 \mu\text{m}$ clay fraction, and the clays are heavily coated with hematite and maghemite. The pH and cation exchange capacity (CEC) of these soils range from 4 to 6 and 4 to 6 $\text{cmol}_\text{c} \text{ kg}^{-1}$, respectively (Jardine *et al.*, 1989). All soils were dried in an oven at 40°C and gently crushed with a mortar and pestle to pass a 250- μm sieve.

Contaminant Addition to Soil

Ten grams of the soil and 100 ml of chromium solution were placed in a 200-ml glass centrifuge vessel, shaken, and allowed to equilibrate for 2 days. The spiking concentrations (dose rates) for Cr(VI), as K_2CrO_4 , were 1000, 250, and 50 ppm at a pH of 6.0 and for Cr(III), as CrCl_3 , were 500, 200, and 50 ppm at a pH of 4.0. After a 2-d equilibration period, the slurries were centrifuged and the supernatant was discarded. The soils were then rinsed with double deionized (DDI) water three times to remove chromium in the pore water and allowed to air dry. Once the soils were dry, they were gently crushed, homogenized, and then rewetted with DDI water to 30% moisture. The soils were kept in a container out of direct light and maintained at 30% water content in a moisture-saturated environment.

***In Vitro* Bioaccessibility**

A physiologically based extraction test (PBET) was adapted from Ruby *et al.* (1996, 1999; Ruby, 2000, personal communication) to assess the *in vitro* bioaccessibility of Cr(III) and Cr(VI) from contaminated soils in humans. Sampling was conducted on the treated soils that had been allowed to age in the storage container for 1, 21, 50, 100, and 200 d after the initial treatment and subsequent wetting of the treated soils. Triplicate moist samples (~0.3 g dry weight) were placed in 50-ml polyethylene tubes to which 30 ml of 0.4 M glycine at pH 1.5 and 37°C was added. The slurries were quickly placed in a rotating water bath at 37°C and agitated at 30 ± 2 rpm for 1 h. The method was designed to simulate the stomach digestive system in humans and has also been used by Skowronski *et al.*

Table 1 Select soil physical and geochemical properties

	Particle size analysis			Organic matter content (%)	pH (5mM CaCl ₂)	pH (DDI)	Fe (g/kg)	Mineralogy of <2 µm clay fraction *
	Sand (%)	Silt (%)	Clay (%)					
Melton A	56.2	30.0	13.8	3.55	6.91	7.18	10.68	
Melton B	30.8	50.4	18.8	0.42	4.23	4.87	22.07	I ₄₅ IS ₂₀ V ₁₀ K ₉ VC ₆ M ₅ Q ₃ F ₁
Walker A	34.9	58.9	6.2	1.89	6.01	6.61	7.71	
Walker B	32.2	44.2	23.6	0.10	4.30	5.17	19.55	K ₂₇ V ₂₇ VC ₁₄ Q ₁₃ I ₁₀ IS ₅ G ₃ F ₁

* K = kaolinite; V = vermiculite; VC = chloritized vermiculite; I = illite (soil mica); IS = interstratified 2:1; Q = quartz; G = gibbsite; M = montmorillonite; F = feldspar. Subscripts refer to the percent by weight of each mineral.

(2001) to assess Cr bioaccessibility in a sandy and a clayey soil. Supernatant was separated from the solid via centrifugation. The pH of the supernatant was measured to ensure that the final pH was within ± 0.5 pH units of the initial pH. This scenario held for all cases. Bioaccessibility was calculated as:

$$\% \text{ Bioaccessibility} = \left(\frac{\text{Cr in PBET supernatant } (\mu\text{g} / \text{mL}) \times 30.0 \text{ mL} \div 0.3\text{g dry soil}}{\text{Cr on soil surface } (\mu\text{g} / \text{g})} \right) \times 100$$

Standard deviations on computed %Cr(III) and Cr(VI) bioaccessibility values following triplicate PBET analyses ranged from 0.03 to 2.01 with the average standard deviation of all values being 0.52.

Chromium Analysis

The PBET supernatant was measured for Cr(VI) and Cr total (Cr_T). Cr(VI) was measured using a modified *s*-diphenylcarbohydrazide colorimetric method (Bartlett and James, 1979) with a UV-VIS spectrophotometer at a wavelength of 540 μm (HP model 8453, Palo Alto, CA). Analysis of Cr(VI) was performed immediately on rapidly cooled PBET solutions to avoid possible reduction of Cr(VI) to Cr(III) by glycine (Jardine *et al.*, 1999). Independent studies revealed that Cr(VI) reduction by glycine at 37°C and 1 h was insignificant. Total chromium was measured on a Perkin Elmer AAnalyst 800 atomic absorption spectrophotometer (Wellseley, PA). All standards used were made from an atomic absorption chromium standard (EM Industries, Hawthorne, NY). Cr(III) was calculated as the difference between Cr_T and Cr(VI).

Determination of Chromium on Soil

Total chromium on the soil was determined using a modification of EPA method 3052. The soil was digested in a CEM microwave, model MDS-81D, with hydrofluoric and nitric acid. Boric acid was added before sample analysis in order to facilitate the removal of hydrofluoric acid from solution through the formation of fluoroboric acid. Soils from the National Institute of Standards, with known concentrations of solid phase Cr, were also analyzed with each block of analyses. Samples were stored and analyzed for total chromium using Inductively Coupled Plasma.

Chromium Solid Phase Speciation

X-ray Adsorption Spectroscopy (XAS). Solid phase Cr was speciated using X-ray adsorption near-edge structure (XANES) spectroscopy, which was conducted at

the Stanford Synchrotron Radiation Laboratory (SSRL) under dedicated running conditions. Scans were recorded from -200 to 300 eV around the K-edge of chromium (5989 eV), with 0.2-eV steps across the white-line and main-edge region. Energy selection was accomplished with a Si(220) double-crystal monochromator, with a 1-mm (h) x 20-mm (w) beam. Adsorption was measured by a proportional fluorescent X-ray production using a 13-element Ge detector (Cramer *et al.*, 1988). Mass fractions of Cr(III) and Cr(VI) were determined for each soil using XANES spectroscopy by placing the soil in a 4- x 4- x 40-mm slot cut in an acrylic plate that was sealed with Kapton. The proportion of Cr(VI) relative to total chromium was then determined by the ratio of the white-line amplitude to the total atomic cross section and comparison to standard curves as described by Patterson *et al.* (1997).

Chemical Extraction. In an effort to indirectly quantify Cr(VI) reduction processes on the soils, sorption isotherms were constructed and the solid phase extracted with SO_4^{2-} . Because SO_4^{2-} competes well for Cr(VI) sorption sites, but does not compete well for Cr(III) sorption sites, an indirect measure of the reduction of Cr(VI) to Cr(III) should be possible. Approximately 1 g soil was placed in preweighed centrifuge tubes, and the soils treated with 15 ml of varying concentrations of Cr(VI) in 5 mM CaCl_2 that were adjusted to the pH of the soil. Samples were allowed to equilibrate on the shaker for 48 h. Soils were centrifuged and supernatant was saved for analysis. The Cr(VI) was extracted from soils with three sequential washings of 0.05 M Na_2SO_4 . The equilibrium solutions and extraction solutions were analyzed for both Cr(VI) and Cr(III). The chromium extract was corrected for pore water Cr of the equilibration step.

RESULTS AND DISCUSSION

Influence of Soil Properties on Cr Sorption

As expected, soils treated with solutions containing Cr(III) adsorbed 2 to 10 times more Cr than those treated with Cr(VI) (Table 2). This results from a larger cation exchange capacity vs. anion exchange capacity and the propensity for Cr(III) to precipitate on mineral surfaces at pH values above 5.5. The adsorption of both Cr species became more similar on the WB B-horizon soil because acidic conditions and abundant Fe-oxides provided positive surface charges, thereby enhancing Cr(VI) sorption. Thus, mineral phases, particularly iron oxides, with proton-specific surface sites may effectively adsorb Cr(VI) at low to medium soil pHs (Zachara *et al.*, 1987, 1988, 1989; Leckie *et al.*, 1980; Davis and Leckie, 1980; Mayer and Schick, 1981). The A-horizon soils had a higher pH and organic matter content, creating an environment that was not conducive to Cr(VI) adsorption.

Table 2 Mass loadings of Cr(III) and Cr(VI) on soil (mg/kg) for various Cr treatment concentrations (ppm)


Cr(III)	500 ppm	200 ppm	50 ppm
Melton A	4479.42	1823.26	426.42
Melton B	2002.91	1430.68	452.02
Walker A	2421.67	1779.01	451.00
Walker B	1276.05	1070.32	445.20

Cr(VI)	1000ppm	250ppm	50ppm
Melton A	386.47	199.21	91.28
Melton B	269.14	219.86	150.00
Walker A	391.83	244.24	100.22
Walker B	423.48	330.53	218.25


In the case of Cr(III) the patterns of adsorption were reversed, where the A-horizon soils typically adsorbed more Cr than the B-horizon soils. The A-horizon soils characteristically had higher pH creating an environment that favored Cr(III) adsorption. Deprotonation of oxides and organic matter occurs in soils with higher soil pH values, which results in more negatively charged sites that attract cations such as Cr(III). Also, when the soil pH is above 5.5, as with the two A-horizon soils used here, the Cr(III) most likely precipitates from solution as hydroxides creating a surface coating on a variety of soil mineral surfaces (Bartlett and Kimball, 1976a). This suggests that larger solid phase concentrations of Cr(III) can often be expected in soils with higher pH and abundant inorganic and organic carbon as shown by Stewart *et al.* (2003).

Influence of Aging on Cr Bioaccessibility

Chromium bioaccessibility, as measured by the PBET method, decreased with time for all soils tested and at all solid phase concentrations (with the exception of the 50 ppm Cr(VI) treated MV-A soil), with aging effects being most pronounced for Cr(III) (Figures 1 and 2). Standard deviations on computed % Cr bioaccessibility values were on average 0.52, which were too small to show error bars on the triplicate-measured values of Figures 1 and 2. Analysis of variance (ANOVA) t-test on day 1 vs. 200 for each of the four soils, two contaminants (Cr(III/VI)), and

**FIGURE 1**

Aging and solid phase concentration effects on the percent Cr bioaccessibility for soils treated with varying concentrations of Cr(III) (50, 200, and 500 mg/L). (a) Melton Valley A-horizon soil, (b) Melton Valley B-horizon soil, (c) Walker Branch A-horizon soil, and (d) Walker Branch B-horizon soil.

**FIGURE 2**

Aging and solid phase concentration effects on the percent Cr bioaccessibility for soils treated with varying concentrations of Cr(VI) (50, 250, and 1000 mg/L). (a) Melton Valley A-horizon soil, (b) Melton Valley B-horizon soil, (c) Walker Branch A-horizon soil, and (d) Walker Branch B-horizon soil.

three treatment concentrations (dose) confirmed that the aging effect was significant at the 95% level (i.e., most p values <0.0001) and was most pronounced on Cr(III)- treated soils (results not shown). The decrease in bioaccessibility was rapid for the first 50 d and slowed dramatically as the aging period approached 200 d. As the soils age, they most likely approach a state of equilibrium between the solution phase Cr and the surface of the soil. The aging effect is related to the enhanced stability of Cr on the soil surface with time. Structural reorientation of Cr surface bonds or, in the case of Cr(III), slow precipitation reactions can account for the stronger sorption of Cr at longer times. The greater aging effect observed for Cr(III) vs. Cr(VI) is most likely related to the time-dependent formation of solid phase $\text{Cr}(\text{OH})_3$, which is not easily dissolved under the acidic conditions of the PBET.

Influence of Solid Phase Concentration on Cr Bioaccessibility

In general, the effect of Cr solid phase concentration (dose effect) on bioaccessibility was small, with Cr(III) showing the most pronounced effect. A comprehensive ANOVA test, discussed later in the manuscript, confirmed that the dose level exerts only a minor influence on Cr bioaccessibility (see Table 6). No obvious trends were noted for Cr(VI), whose bioaccessibility remained relatively constant at different solid phase concentrations on any given soil (Figure 2). For the Cr(III) system, particularly for A-horizon soils, higher bioaccessibility was noted for soils that were treated with 50 ppm Cr(III) relative to the higher concentration treatments. This is most likely related to the fact that at low surface coverage (< 20%) adsorption is the dominant process where Cr(III) forms inner-sphere complexes with the soil, while at higher surface coverages (> 20%) surface precipitation occurs and becomes the dominant process (Fendorf *et al.*, 1994; Fendorf and Sparks, 1994). The soils that were treated with 50 ppm Cr(III) have significantly lower Cr on the soil than the other soils treated with higher concentrations. Thus, the mechanism of Cr sequestration has a higher proportion of inner-sphere bonds related to precipitated phases, which most likely causes a higher percent of Cr(III) that is bioaccessible at lower solid phase concentrations.

Influence of Soil Properties on Cr Bioaccessibility

The bioaccessibility of Cr(III) and Cr(VI) varied significantly as a function of soil type and horizon, and the oxidation state of the contaminant. Statistical analysis using the ANOVA t-test confirmed that Cr bioaccessibility was significantly influenced by these effects at the 95% level with p values typically <0.0001 (results not shown). In general, A-horizon soils exhibited less Cr bioaccessibility relative to B-horizon soils. In the Cr(III) system, the higher organic matter content and

higher pH of the A-horizon soils are probably the main factors responsible for this difference. The Walker Branch B-horizon (WB-B) soil is a good example of how soil properties effect the degree of bioaccessibility because it is the most acidic of the soils and has the lowest organic carbon content, and consequently it shows the highest percent of Cr(III) bioaccessibility (Figure 1d). Both the Melton Valley A-horizon (MV-A) and Walker Branch A-horizon (WB-A) soils have a high pH and high organic carbon content and an equally low Cr(III) bioaccessibility. These results are consistent with observations in Stewart *et al.* (2003) that showed that Cr(III) bioaccessibility was limited in systems with high levels of inorganic and organic carbon. Skowronski *et al.* (2001) also noted that Cr(III) bioaccessibility was lower on an organic-rich sandy soil vs. a clay soil that had significantly less organic carbon.

In the Cr(VI) system, the two A-horizon soils and the Melton Valley B-horizon (MV-B) showed statistically significant lower Cr bioaccessibility than WB-B for all treatment concentrations and aging times. Although the WB-B soil adsorbed the most Cr(VI), its tendency to release Cr under the acidic conditions of the PBET is due to the soil's inability to maintain the weak bond between the Cr and the surface. The Cr(VI) ion is probably electrostatically bound to mineral oxides through outer sphere complexes, which are unstable during the conditions of the PBET. This leads to the question of why is it that both A-horizon soils and even the Melton Valley Inceptisol B horizon soils (MV-B) have such low Cr(VI) bioaccessibility when soil properties are such as to discourage strong sorption?

To address the above question, both direct and indirect solid phase Cr speciation methods using X-ray Absorption Spectroscopy (XAS) and a chemical extraction technique, respectively, were employed. Analysis with XAS of the 250 and 1000 ppm Cr(VI) treated soils after 200 d aging suggested that all soils, except the WB-B soil, had Cr surface coverages that were > 95% Cr(III) (Table 3). The 250 and 1000 ppm Cr(VI) treated WB B-horizon soils contained only 30 and 53% surface bound Cr(III), respectively. Thus, the bioaccessibility of Cr(VI) was significantly influenced by the reduction of Cr(VI) to Cr(III). Skowronski *et al.* (2001) also suggested that Cr(VI) bioaccessibility in their soils was influenced by oxidation-reduction processes. In order for reduction to occur, there needs to be a source of electrons. Both organic matter and the Fe(II)-bearing minerals are able to supply electrons to catalyze the reduction of Cr(VI) to Cr(III). Because the soils used in this study were highly oxidized and most likely devoid of Fe(II)-bearing minerals, the reduction of Cr(VI) to Cr(III) was most likely catalyzed by soil organic matter or surface-bound organic carbon (Adriano, 1986; Sparks, 1995; Deng and Stone, 1996; Jardine *et al.*, 1999). Thus, extensive reduction processes for the A-horizon soils and the MV B-horizon soils are most likely related to the ample supply of organic carbon in these soils (Table 1). Even the WB B-horizon soil showed Cr(VI) reduction to Cr(III) with a solid phase carbon mass of 0.1%. Jardine *et al.* (1999) showed that in acidic soils the availability of even small amounts of surface-bound natural organic carbon (0.05% w/w on the solid) can result in significant reduction

Table 3 Percentage of soil solid phase Cr(III) and Cr(VI) quantified by X-ray Adsorption Spectroscopy (XAS) *

	Soild phase Cr(VI)	Solid phase Cr(III)
<u>250 ppm</u>	~~~~~ % ~~~~~	
MV-A	< 5	> 95
MV-B	< 5	> 95
WB-A	< 5	> 95
WB-B	70	30
<u>1000 ppm</u>		
MV-A	< 5	> 95
MV-B	< 5	> 95
WB-A	< 5	> 95
WB-B	47	53

* 200 d aged samples

of Cr(VI) to Cr(III). Therefore, Cr(VI) reduction decreases Cr bioaccessibility because the Cr(III) product is more tightly bound to the solid phase. The Cr(III) probably adsorbs to the surface through strong covalent bonds or precipitates as hydroxide complexes on mineral surfaces. Thus, the percent of Cr that is bioaccessible decreases during the PBET.

The XAS data are in agreement with aqueous Cr speciation measurements on the PBET solutions (Table 4). A significant portion of the total bioaccessible Cr was found to be Cr(III), with the WB B-horizon soil having the lowest total amount of extractable Cr(III) as indicated by the high Cr(VI) in Table 4. For all soils except WB-B, the trends in the data suggest an increasing percentage of Cr(VI) in the PBET extraction solution up to ~100 d followed by an abrupt decrease with continued Cr-soil aging to 200 d. These trends are consistent with the enhanced reduction of Cr(VI) to Cr(III) by the A-horizon soils and the MV-B soil relative to the WB-B soil. Using the 200-d aqueous speciation data coupled with the XAS solid speciation results (analyzed on 200 d aged soils), one can calculate the mass fraction of Cr(III) and Cr(VI) that are bioaccessible in each soil (Table 5). In all soils, the bioaccessibility of surface-bound Cr(VI) was significantly greater than that for Cr(III). Between 42 and 108% of the total adsorbed Cr(VI) was bioaccessible when compared with total adsorbed Cr(III), which was only 3 and 14% bioaccessible. Although Cr(III) may dominate total Cr in the PBET, surface-bound Cr(VI) is significantly more bioaccessible. Thus, the reduction of Cr(VI) to Cr(III) by soil

Table 4 Percentage of Cr(VI) in PBET extractant for soils treated with 250 and 1000 ppm Cr(VI)

	<u>1 day</u>	<u>21 day</u>	<u>50 day</u>	<u>100 day</u>	<u>200 day</u>
<u>250 ppm</u>	~~~~~ % ~~~~~				
MV-A	28.73	61.90	78.02	64.91	46.26
MV-B	21.64	59.09	71.92	67.75	32.88
WB-A	16.54	22.86	57.28	28.61	26.38
WB-B	76.92	81.16	99.10	76.07	94.27
<u>1000 ppm</u>					
MV-A	3.81	15.29	6.31	44.09	21.89
MV-B	14.59	13.26	22.09	90.46	24.07
WB-A	4.23	8.15	4.31	31.53	21.25
WB-B	60.28	84.72	63.57	73.00	78.91

organic matter significantly decreases total Cr bioaccessibility. These results are important from a human health perspective because Cr(VI) is believed to be much more toxic than Cr(III), with even sub-ppm levels considered lethal. Thus, under certain circumstances, soils that contain sufficient organic carbon or Fe(II)-bearing minerals may be capable of decreasing Cr bioaccessibility through reduction of labile Cr(VI) to the more sparingly soluble Cr(III) species.

An indirect chemical extraction method was also used to show that Cr(VI) was being reduced to Cr(III). Chromium (VI) was adsorbed onto the soils using different treatment solution concentrations, allowed to equilibrate for 2 d, and then the solid phase was treated with 0.05 M Na₂SO₄ to remove the Cr(VI) (Figure 3 a-d). The SO₄²⁻ anion should be a sufficient competitor for surface sites occupied by HCrO₄⁻ because the latter is typically sorbed to the solid phase through weak outer-sphere electrostatic bonds. Thus, if Cr reduction processes are minimal, the SO₄²⁻ should be able to recover nearly all of the initial adsorbed Cr(VI). The extractant Na₂HPO₄ was also utilized on select soils since the HPO₄²⁻ anion can aggressively compete for Cr(VI) that is bound to the soil by either inner- or outer-sphere surface complexes. The results compared favorably with the SO₄²⁻ system; however, the HPO₄²⁻ results were somewhat more erratic for reasons unknown to the authors, and thus the SO₄²⁻ system was preferred. In this study, the chemical extraction method can only be qualitatively compared with the XAS results because the latter technique was employed on 200 d aged samples, whereas the extraction method

Table 5 Percentage of total surface bound Cr(III) and Cr(VI) that was bioaccessible after 200 d aging

	<u>MV-A</u>		<u>MV-B</u>		<u>WB-A</u>		<u>WB-B</u>	
	Cr(VI) treatment conc. (ppm)							
	250	1000	250	1000	250	1000	250	1000
% Cr(III) bioaccessible	6.0	9.9	7.1	8.0	7.8	8.3	3.2	14.2
% Cr(VI) bioaccessible	101.0	51.5	65.5	48.1	52.5	42.3	71.1	107.5




FIGURE 3

Adsorbed and SO_4^{2-} extractable Cr(VI) on soil as a function of solution concentration. (a) Melton Valley A-horizon soil, (b) Melton Valley B-horizon soil, (c) Walker Branch A-horizon soil, and (d) Walker Branch B-horizon soil.

was employed on samples aged for only 2 d. Jardine *et al.* (1999) previously measured a half-life of 85 h for Cr(VI) reduction by organic carbon, so samples analyzed after 200 d of aging should have more Cr(III) product than samples analyzed after 2 d of aging. Nevertheless, the chemical extraction method agreed well with the XAS results and the quantity of organic C in the soils. With the exception of the WB B-horizon soil, the quantity of Cr(VI) extracted from the solid phase was significantly lower than the initial Cr(VI) sorbed, implying that Cr(III) is being formed and remains sorbed to the soil (Figure 3 a-d). A-horizon soils had significantly more Cr(III) production when compared with B-horizon soils, which is consistent with the larger organic carbon content of the former. The WB B-horizon, which had as little as 0.1% organic carbon, showed no Cr(VI) reduction after 2 d (Figure 3b). The low organic content of this soil does not lend itself to the rapid reduction of Cr(VI) or is the source of iron, hematite (Fe_2O_3), and maghemite ($\gamma\text{Fe}_2\text{O}_3$) conducive to Cr(VI) reduction. The presence of Fe(III) suggests that the iron is already oxidized and therefore not in the correct state to facilitate the reduction of Cr(VI). This further explains why the percent of Cr that is bioaccessible in the WB-B soil remains so high compared with the other three soils examined. These results are consistent with the XAS findings that showed Cr(VI) reduction was nearly complete on all soils after 200 d, with the exception of the WB-B-horizon soil.

Factors Influencing Cr Bioaccessibility

The entire data set was analyzed using an ANOVA model that incorporated three qualitative factors (oxidation state, soil type, and dose level) and one quantitative factor (age). The original dose amounts were converted to low (50 ppm Cr(III) and Cr(VI)), medium (200 ppm Cr(III) and 250 ppm Cr(VI)), or high (500 ppm Cr(III) and 1000 ppm Cr(VI)) categories to simplify the statistical analysis. The complete four-factor ANOVA model explained more than 95% of the variance in bioaccessibility ($r^2 = 0.952$, $F = 127.74$, $p < 0.0001$) with a summary of the ANOVA results shown in Table 6. The oxidation state, soil type, and dose main effects were all significant as were the two-way and three-way interactions among these effects. Age and its interactions with oxidation state and soil type were also significant. However, age and its interactions with dose were only marginally significant. Thus, it is thought that these marginally significant results indicate that the dose level exerts only a minor influence on the relationship between age and bioavailability. It is also important to realize that some of the significance noted in Table 6 is driven by the high analytical precision of the bioaccessibility results. Thus, in certain cases it may be difficult to tease out statistical significance from geochemical and physical significance.

Table 6. Summary of analysis of variance results showing the significance of various factors on the bioaccessibility of Cr in soils.

Factor	SS	df	MS	F	Prob.
Oxidation	1630.7	1	1630.7	73.76	<0.0001
Soil	87926.5	3	29308.8	1325.70	<0.0001
Oxidation-Soil	21324.3	3	7108.1	321.51	<0.0001
Dose	264.4	2	132.2	5.98	0.0028
Oxidation-Dose	3459.0	2	1729.5	78.23	<0.0001
Soil-Dose	1091.5	6	181.9	8.23	<0.0001
Oxidation-Soil-Dose	1317.9	6	219.6	9.94	<0.0001
Age	9378.9	1	9378.9	424.23	<0.0001
Age-Oxidation	1753.8	1	1753.8	79.33	<0.0001
Age-Soil	2956.7	3	985.6	44.58	<0.0001
Age-Oxidation-Soil	621.8	3	207.3	9.38	<0.0001
Age-Dose	126.4	2	63.2	2.86	0.0589
Age-Oxidation-Dose	224.1	2	112.0	5.07	0.0068
Age-Soil-Dose	355.5	6	59.2	2.68	0.0151
Age-Oxidation-Soil-Dose	297.7	6	49.6	2.24	0.0391
Error	6632.4	300	22.1		

SS = sum of squares, df = degrees of freedom, MS = mean squares, F = F-value statistic, Prob. = probability that one obtains the F-value other than by chance.

ENVIRONMENTAL SIGNIFICANCE

This study has shown that the metal-sequestering properties of soil significantly lower the percent of Cr(III) and Cr(VI) bioaccessible after ingestion. The percent of bioaccessible Cr is largely independent of the initial solid phase concentration of Cr prior to the PBET simulated digestion. Sorption and bioaccessibility of Cr(III) and Cr(VI) vary significantly as a function of soil type and horizon, and the oxidation state of the contaminant. Soils with higher pH and abundant inorganic and organic carbon can often be expected to have higher solid phase concentrations of Cr(III), while for Cr(VI) the patterns are reversed, with Cr(VI) adsorption favored by lower soil pH and soil minerals with amphoteric charge. Aging effects show Cr bioaccessibility decreases after the first 50 d, and this is related to the enhanced stability of Cr on the soil surface followed by stable bioaccessibility to 200 d. Bioaccessibility of Cr(III) can be significantly reduced by its ability to bind strongly to organic matter and also to Cr – hydroxide precipitates on the soil surface, even under the conditions present in the PBET. Soil sequestration of Cr(VI) significantly lowers its bioaccessibility. Organic-rich soils and/or soils with Fe(II)-bearing minerals present enhance Cr(VI) reduction to Cr(III), with the latter being strongly adsorbed and less bioaccessible. This is important from a human health perspective because Cr(VI) is believed to be much more toxic than Cr(III).

ACKNOWLEDGMENTS

We appreciate the efforts of Ms. Cathy Vogel and Dr. Andrea Leeson, the contract officers for the U.S. DoD SERDP who supported this work.

REFERENCES

- Adriano, D.C. 1986. *Trace Elements in the Terrestrial Environment*. New York, Springer-Verlag.
- Ainsworth, C.C., Girvin, D.C., Zachara, J.M., and Smith, S.C. 1989. Chromate adsorption on goethite: effects of aluminum substitution. *Soil Sci. Soc. Am. J.* **53**, 411–418.
- Anderson, L.D., Kent, D.B., and Davis, J.A. 1994. Batch experiments characterizing the reduction of Cr(VI) using suboxic material from a mildly reducing sand and gravel aquifer. *Environ. Sci. Technol.* **28**, 178–185.
- Arnseth, R.W. and Turner, R.S. 1988. Sequential extraction of iron, manganese, aluminum, and silicon in soils from two contrasting watersheds. *Soil Sci. Soc. Am. J.* **52**, 1801–1807.
- Bartlett, R.J. and Kimble, J.M. 1976a. Behavior of chromium in soils. I. Trivalent forms. *J. Environ. Qual.* **5**, 379–383.
- Bartlett, R.J. and Kimble, J.M. 1976b. Behavior of chromium in soils. II. Hexavalent forms. *J. Environ. Qual.* **5**, 383–386.
- Bartlett, R.J. and Kimble, J.M. 1979. Behavior of chromium in soils. III. Oxidation. *J. Environ. Qual.* **8**, 31–35.
- Chung J., Zasoski, R.J., and Lim, S. 1994. Kinetics of chromium(III) oxidation by various manganese oxides. *Korean J Agric Chem Biotechnol.* **37**, 414–420.
- Cramer, S.P., Tench, O., Yocum, M., and George, G.N. 1988. A 13-element GE detector for fluorescence EXAFS. *Nucl. Instrum. Meth.* **A266**, 586–591.
- Davis, J.A. and Leckie, J.O. 1980. Surface ionization and complexation at the oxide/water interface 3. Adsorption on anions. *J. Colloid Interface Sci.* **74**, 32–43.
- Davis, S., Waller, P., Buschbom, R., Ballow, J., and White, P. 1990. Quantitative estimates of soil ingestion in normal children between the ages of 2 and 7 years. Population-based estimates using aluminum, silicon, and titanium as soil tracer elements. *Arch. Environ. Health.* **45**, 112–122.
- Deng, B. and Stone, A.T. 1996. Surface catalyzed chromium(VI) reduction: reactivity comparisons of different organic reductants and different oxide surfaces. *Environ. Sci. Technol.* **30**, 2484–2494.
- Driese, S.G., McKay, L.D., and Penfield, C.P. 2001. Lithologic and pedogenic influences on porosity distribution and groundwater flow fractured sedimentary saprolite: a new application of environmental sedimentology. *J. Sedimentary Res.* **71**, 843–857.
- Eary, L.E. and Rai, D. 1991. Chromate reduction by subsurface soils under acidic conditions. *Soil Sci. Soc. Am. J.* **55**, 676–683.
- EPA Method 3052. <http://www.epa.gov/epaoswer/hazwaste/test/pdfs/3052.pdf>.
- Fendorf, S.E., Lamble, G.M., Stapleton, M.G., Kelly, M.J., and Sparks, D.L. 1994. Mechanisms of chromium (III) sorption on silica. 1. Cr(III) surface structure derived by extended x-ray adsorption fine structure spectroscopy. *Environ. Sci. Technol.* **28**, 284–289.
- Fendorf, S.E. and Sparks, D.L. 1994. Mechanisms of chromium (III) sorption on silica: 2. Effects of reaction conditions. *Environ. Sci. Technol.* **28**, 290–297.
- Fendorf, S., Eick, M.J., Grossl, P., and Sparks, D.L. 1997. Arsenate and chromate retention mechanisms on goethite. 1. Surface structure. *Environ. Sci. Technol.* **31**, 315–320.
- Hamel, S.C., Buckley, B., and Liroy, P.J. 1998. Bioaccessibility of metals in soils for different liquid to solid ratios in synthetic gastric fluid. *Environ. Sci. Technol.* **32**, 358–362.

- James, B.J. and Bartlett, R.J. 1983. Behavior of chromium in soils. VII. Adsorption and reduction of hexavalent forms. *J. Environ. Qual.* **12**, 177–181.
- Jardine, P.M., Fendorf, S.E., Mayes, M.A., Brooks, S.C., and Bailey, W.B. 1999. Fate and transport of hexavalent chromium in undisturbed heterogeneous soil. *Environ. Sci. Technol.* **33**, 2939–2944.
- Jardine, P.M., Weber, N.L., and McCarthy, J.F. 1989. Mechanisms of dissolved organic carbon adsorption on soil. *Soil. Sci. Soc. Am. J.* **53**, 1378–1385.
- Kooner, Z.S., Jardine, P.M., and Feldman, S. 1995. Competitive surface complexation reactions of sulfate and natural organic carbon on soil. *J. Environ. Qual.* **24**, 656–662.
- Leckie, J.O., Benjamin, M.M., Hayes, K., Kaufman, G., and Altman, S. 1980. Adsorption/coprecipitation of trace elements from water with iron oxyhydroxide. *Electric Power Res. Inst. Rept.* Palo Alto, Ca., EPRI-RP-910.
- Levis, A.G. and V. Bianchi. 1982. Mutagenic and cytogenic effects of chromium compounds. In: *Biological and Environmental Aspects of Chromium*. pp. 171–208 (Sverre Langjard, Ed.) New York, Elsevier Biomedical Press.
- Losi, M.E., Amrhein, C., and Frankenberger, W.T. Jr. 1994. Bioremediation of chromate-contaminated groundwater by reduction and precipitation in surface soils. *J. Environ. Qual.* **23**, 1141–1150.
- Mayer, L.M., and Schick, L.L. 1981. Removal of hexavalent chromium from estuarine waters by model substrates and natural sediments. *Environ. Sci. Technol.* **15**, 1482–1484.
- Nriagu, J. O. and Nieboer, E. 1988. *Chromium in the Natural and Human Environments*. New York, John Wiley & Sons.
- Patterson, R.R., Fendorf, S., and Fendorf, M. 1997. Reduction of hexavalent chromium by amorphous iron sulfide. *Environ. Sci. Technol.* **31**, 2039–2044.
- Paustenbach, D. J. 1989. *The Risk Assessment of Environmental and Human Health Hazards: A Textbook of Case Studies*. New York, John Wiley & Sons.
- Peterson, M.L., Brown, G.E. Jr., Parks, G.A., and Stein, C.L. 1997. Differential redox and sorption of Cr(III/VI) on natural silicate and oxide minerals: EXAFS and XANES results. *Geochim. Cosmochim. Acta.* **61**, 3399–3412.
- Proctor, D.M., Shay, E.C., and Scott, P.K. 1997. Health-based soil action levels for trivalent and hexavalent chromium: a comparison with state and federal standards. *J. Soil Contamin.* **6**, 595–648.
- Ruby, M.V., Davis, A., Schoof, R., Eberle, S., Sellstone, S.M. 1996. Estimation of lead and arsenic bioavailability using a physiologically based extraction test. *Environ. Sci. Technol.* **30**, 422–430.
- Ruby, M.V., Schoof, R., Brattin, W., Goldade, M., Post, G., Harnois, M., Mosby, D.E., Casteel, S.W., Berti, W., Carpenter, M., Edwards, D., Cragin, D., and Chappell, W. 1999. Advances in evaluating the oral bioavailability of inorganics in soil for use in human health risk assessment. *Environ. Sci. Technol.* **33**, 3697–3705.
- Sheehan, P.J., Meyer, D.M., Sauer, M.M., and Paustenbach, D.J. 1991. Assessment of the human health risks posed by exposure to chromium contaminated soils. *J. Toxicol. Environ. Health* **32**, 161–201.
- Skowronski, G.A., Seide, M., and Abdel-Rahman, M.S. 2001. Oral bioaccessibility of trivalent and hexavalent chromium in soil by simulated gastric fluid. *J. Toxicol. Environ. Health, Part A.* **63**, 351–362.
- Sparks, D.L. 1995. *Environmental Soil Chemistry*. New York, Academic Press.
- Stewart, M.A., Jardine, P.M., Barnett, M.O., Mehlhorn, T.L., Hyder, K., and McKay, L.D. 2003. Influence of soil geochemical and physical properties on the sorption and bioaccessibility of Cr(III). *J. Environ. Qual.* **32**, 129–137.

- Zachara, J.M., Girvin, D.C., Schmidt, R.L., and Resch, C.T. 1987. Chromate adsorption on amorphous iron oxyhydroxide in the presence of major groundwater ions. *Environ. Sci. Technol.* **21**, 589-594.
- Zachara, J.M., Cowan, C.E., Schmidt, R.L., and Ainsworth, C.C. 1988. Chromate adsorption by kaolinite. *Clays Clay Miner.* **36**, 317-326.
- Zachara, J.M., Ainsworth, C.C., Cowan, C.E., Resch, C.T. 1989. Adsorption of chromate by subsurface soil horizons. *Soil Sci. Soc. Am. J.* **53**, 418-428.
-

APPENDIX E

YANG, J-K., M.O. BARNETT, P.M. JARDINE, AND S.C. BROOKS. 2003. FACTORS CONTROLLING THE BIOACCESSIBILITY OF ARSENIC(V) AND LEAD(II) IN SOIL. SOIL AND SEDIMENT CONTAMINATION. 12:165-179.

Factors Controlling the Bioaccessibility of Arsenic(V) and Lead(II) in Soil

Jae-Kyu Yang,¹ Mark O. Barnett,^{1*}
Philip M. Jardine,² and Scott C.
Brooks²

¹Department of Civil Engineering, 238 Harbert Engineering Center, Auburn University, Auburn, AL 36849; ²Environmental Sciences Division, Oak Ridge National Laboratory, P. O. Box 2008, Oak Ridge, TN 37831

The relative oral bioaccessibility of labile Pb(II) and As(V) added to soils was investigated in a well-characterized soil using a physiologically based extraction test (PBET) to simulate metal solubility in a child's digestive sys-

tem. The effect of soil and PBET (i.e., simulated stomach and small intestine) pH, soil metal concentration, soil to solution ratio, and soil-metal aging time were investigated. Arsenic bioaccessibility was relatively unaffected by a variation in simulated stomach and small intestine pH over the range 2 to 7 and soil pH over the range 4.5 to 9.4. In contrast, Pb(II) bioaccessibility was strongly dependent on both the simulated stomach, small intestine, and soil pH, showing enhanced sequestration and decreased bioaccessibility at higher pH values in all cases. Although the bioaccessibility of Pb(II) was constant over the concentration range of approximately 10 to 10,000 mg/kg, the As(V) bioaccessibility significantly increased over this concentration range. The bioaccessibility of both arsenic and lead increased as the soil-to-solution ratio decreased from 1:40 to 1:100. Additional lead sequestration was not observed during 6 months of soil aging, but As(V) bioaccessibility decreased significantly during this period.

* Corresponding author phone: (334) 844-6291; fax: (334)844-6290; email: barnettm@eng.auburn.edu

Key Words: bioaccessibility, bioavailability, extraction, arsenic, lead, soil.

INTRODUCTION

Soil ingestion is typically the primary human health exposure pathway at metal-contaminated sites. For residential or recreational land use scenarios, for example, the ingestion of soil by children is almost always the critical exposure pathway. The calculated health risk due to the incidental ingestion of a metal-contaminated soil is a function of several variables: the soil-metal concentration, soil ingestion rate, body weight, exposure frequency and duration, and the oral toxicity (cancer slope factor for carcinogens or the reference dose for non-carcinogens). However, the oral toxicity of metals is often based on toxicological studies where the metal is potentially more bioavailable than metals in soils (e.g., from animal feeding studies with soluble metal salts). Thus, with the exception of Pb, risk assessments implicitly assume a default value of 100% relative bioavailability. In other words, the bioavailability of the metal in the soil is implicitly assumed to be the same as in the dosing medium (e.g., water or food) in the critical toxicity study. The risk assessment methodology for Pb in soils is unique; Pb is the only metal that has an explicit soil bioavailability adjustment.

Metals in soils, however, are often relatively insoluble, requiring aggressive digestion procedures for complete analytical metal recovery. As a result, an oral toxicity value developed from studies using soluble metal species may overstate the risk posed by less-soluble metals in soils. The generally low bioavailability of Pb and As in soils in mining areas has been well documented, and risk assessments based on data from studies using soluble metal salts overestimate the risk posed by these soils (Davis *et al.*, 1992). Numerous studies, for example, have shown that Pb in soil (Freeman *et al.*, 1994; Casteel *et al.*, 1997), mining waste (Dieter *et al.*, 1993; Polak *et al.*, 1996) and aggregate (Cheng *et al.*, 1991; Preslan *et al.*, 1996), is much less bioavailable than more soluble Pb species, such as Pb oxide, nitrate, or acetate used in toxicological studies. Relatively low Pb bioavailability is a consequence of Pb speciation and the corresponding solubility constraints (Davis *et al.*, 1993) and kinetic limitations to dissolution in the limited residence time of the GI tract (Ruby *et al.*, 1992). Similarly, the oral toxicity for As is based on a human epidemiological study of As in drinking water. However, soluble As in drinking water is much more bioavailable than insoluble As in soils, the latter of which is primarily excreted through the feces without being absorbed through the GI tract (Freeman *et al.*, 1995). Estimates of risk due to As ingestion in soils in mining areas would overstate the risk unless the lower bioavailability of As in these soils is considered (Davis *et al.*, 1996; Davis *et al.*, 2001).

In mining-impacted areas, low soil-metal bioavailability might be due to the presence of residual low solubility metal sulfides from the ore body. However, even in non-mining areas, soil metal bioavailability may be lower than for soluble metal species because soils typically bind metals due to sorption to the solid phase and the formation of other secondary solid phases with lower solubility, including authigenic metal sulfides (Barnett *et al.*, 1997). For example, the presence of the

soil matrix significantly reduced the absorption of soluble CdCl_2 from the GI tract in rat studies (Schilderman *et al.*, 1997). In fact, animals are believed to instinctively consume soils when exposed to contaminants in their diets as a way of decreasing the bioavailability and the effect of these contaminants (Sheppard *et al.*, 1995).

The purpose of this article is to describe the results of an investigation into the bioaccessibility of Pb(II) and As(V) added to soils using a physiologically based extraction test (PBET) to simulate soil ingestion. As(V) and Pb(II)-spiked soils were used because (1) the initial metal concentration and speciation could be controlled, (2) changes in bioaccessibility from the initial labile metal could be followed with time, and (3) beginning with labile metals provided insight into the ability of soils themselves to limit metal bioaccessibility, without regard to any unique site-specific speciation. The effects of soil and PBET pH, soil-to-solution ratio, soil-metal concentration, and soil-metal aging time were investigated.

II. EXPERIMENTAL METHODS

A. Materials

All chemicals employed in this research were analytical grade or above, and solutions were prepared with deionized water ($18 \text{ M}\Omega\text{-cm}$) from a reverse osmosis/ion exchange apparatus (Milli-Q™ Water System). Soil samples were collected from the B- and C-horizon of a weakly developed Inceptisol on the Department of Energy Oak Ridge (Tenn.) Reservation. The soils were air dried and passed through a $250\text{-}\mu\text{m}$ (B-horizon) or 2-mm (C-horizon) sieve. The $<250\text{-}\mu\text{m}$ fraction represents the soil fraction most likely ingested as a result of children's hand-to-mouth activities and was adopted after the initial experiments with the C-horizon material were begun. These soils are acidic (pH ~ 4.2 in a $1:2 \text{ g/mL}$ suspension) and heavily coated with Fe-oxides. Some physical and chemical properties of the two soil samples are shown in Table 1.

B. Soil Spiking

Arsenic(V) and Pb(II) were added to the soil from a small volume of concentrated metal stock solution to a $1:10 \text{ g/mL}$ suspension in 10^{-3} M CaCl_2 solution. In most experiments, the soil slurry was maintained at the natural soil pH (~ 4.5 in a $1:10 \text{ g/mL}$ suspension) by immediately neutralizing the acidity from the metal stock solution with dilute NaOH. The pH of some slurries was changed by adding additional dilute NaOH to study the effect of soil pH on the bioaccessibility of As(V) and Pb(II). After mixing for 48 h, the soil suspension was centrifuged and the supernatant was decanted. The remaining soil was washed twice with distilled

TABLE 1
Some Physical and Chemical Properties of Inceptisol Soils Used in Study

Property	B-horizon	C-horizon
Sand (%)	31	31
Silt (%)	50	50
Clay (%)	19	19
pH*	4.2	4.1
Mn (g/kg)**	0.17	0.36
Fe (g/kg)**	22.1	25.8
Organic Matter (%)	0.42	0.55
Inorganic Carbon (%)	0.26	not measured
CEC (cmol/kg)	14	not measured
Mineralogy (%)***	I(45)IS(20)V(10)K(9) VC(6)M(5)Q(3)F(1)	I(30)IS(25)K(20)S(10) V(10)Q(5)

* The pH of the soil solution was measured in 5 mM CaCl₂ in a 1:2 g/mL suspension.

** Dithionite/citrate/bicarbonate extractable.

*** K=kaolinite, V=vermiculite, VC=chloritized vermiculite, I=illite, IS=interstratified 2:1, Q=quartz, G=gibbsite, M=montmorillonite, F=feldspar, S = smectite.

water to remove any traces of the original soluble As(V) or Pb(II) spike. The decanted supernatant and rinse water were filtered through 0.45- μ m membrane filter, and the concentration of As(V) and Pb(II) in the filtrate was analyzed using an atomic absorption spectrophotometer equipped with an electrodeless discharge lamp (EDL) for As and a hollow cathode lamp for Pb. The difference between the amount of As(V) or Pb(II) added and that remaining in the supernatant was used to calculate the initial soil concentration. The soil residues from the PBET extraction (below) were also analyzed for Pb and As using EPA Method 3050B to verify a mass balance of $\pm 10\%$.

The soils were then air-dried and homogenized by mixing. Initial subsamples were taken representing the conditions at the beginning of the aging experiment (i.e., $t = 0$). The remaining soil was placed in a weighing dish, and deionized water was added to bring the soil to field capacity (30% moisture). The open containers were then aged in a larger container through which a steady flow of 100% relative humidity air was passed. The moisture content of the soils was monitored periodically by weight, with deionized water added as necessary to maintain a constant moisture content of 30%. Periodically, subsamples were removed and analyzed as described below.

C. Adsorption

The degree of adsorption As(V) and Pb(II) to the soil was measured by adding 5 g/L B-horizon soil and 1 mg/L Pb(II) or As(V) concentrations in 10^{-2} M NaNO₃

solution. After adjusting the pH of the initial suspension to between 2 and 12 using dilute HNO_3 or NaOH solutions, the samples were shaken for 48 h at normal room temperature (22 to 25°C). After 48 h, the suspension pH of each sample was measured, and the suspensions were filtered using 0.45- μm filters (Gelman). The concentration of As(V) or Pb(II) in the filtrate was measured using an atomic absorption spectrophotometer as described above.

D. Extractions

The physiologically based extraction test (PBET) used here was adopted from a modification to the original PBET described by Ruby *et al.* (1996). This extraction test has been shown to be predictive of Pb bioavailability in two animal models and is currently being validated for As (Ruby *et al.*, 1999). The extraction device consisted of a sample holder that held 16 wide-mouth, high-density polyethylene bottles (125 mL) and a motor that rotated the sample holder at variable speed. The sample holder was located in a temperature-controlled water bath. During the extraction, the water temperature in the bath was maintained at body temperature ($37 \pm 2^\circ\text{C}$). The extraction solution consisted of 30 g/L glycine (0.4 M) with the pH adjusted to 1.5, 2, 3, or 4 with HCl. These conditions simulated the stomach, because recent research has suggested that Pb and As dissolution in the simulated stomach environment is predictive of Pb and As bioavailability in animals (Ruby *et al.*, 1999).

One gram of each air-dried soil was placed in a 125-mL HDPE bottle. Then 40 or 100 mL of 37°C simulated gastric solution was poured in each bottle. After capping, each bottle was placed in the sample holder and rotated end over end at 30 ± 2 rpm for 1 h. After 1 h, the bottles were immediately removed and stood up right for approximately 5 min before taking a portion of the supernatant, which was then filtered with 0.45- μm filter. For all experiments, duplicate or triplicate samples were run and the results were reported as \pm one standard deviation unless otherwise noted. The dissolved metal concentration in the filtrate was measured with an atomic absorption spectrophotometer, with the fraction of metal dissolved representing the bioaccessibility (see below). Although the stomach may be important in solubilizing soil-bound metals, systemic absorption occurs in the small intestine, where chemical conditions (especially pH) are significantly different. To examine these effects, the pH of the remaining PBET solution was adjusted to 7 by adding 4 mL of 0.5 M NaHCO_3 , maintaining a constant soil-solution ratio. The bottles were returned to the extractor and rotated end over end at 30 ± 2 rpm for 3 h, when they were sampled and analyzed as described previously. The remaining soil sample was analyzed for As or Pb using acid digestion (see below) to verify mass balance within $\pm 10\%$.

To measure the pH and the readily soluble and exchangeable concentrations of As and Pb, 1 g of each soil was mixed with 2 mL of 5×10^{-3} M CaCl_2 solution for

2 h. After centrifugation (10 min at 2000 rpm), the supernatant was filtered with 0.45- μ m filter. Then the pH and metal content of the supernatant were measured. For all soils, blanks (no metal added) were used to correct all data obtained from CaCl_2 and PBET extractions.

The absolute oral bioavailability is the fraction of an administered metal dose that reaches systemic circulation from the gastrointestinal tract (Ruby *et al.*, 1999). The relative bioavailability is the bioavailability of a metal in one form or media compared with another (e.g., the bioavailability of a metal in soil relative to the bioavailability of the metal in water). In *in vitro* extraction tests, the fraction of metal solubilized and available for absorption is termed the bioaccessibility and is an indicator of the bioavailability of soil-bound metals relative to the soluble metal species on which the oral toxicity is based (Ruby *et al.*, 1996). The bioaccessibility of soluble As_2O_5 and $\text{Pb}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ at the same concentration as in the soils was $96.1 \pm 0.1\%$ and $99.8 \pm 1.1\%$, respectively. *In vitro* extraction procedures are a more useful tool than expensive and time-consuming animal feeding studies for investigating the effect a number of variables on bioaccessibility/bioavailability.

E. Soil Analysis

In order to verify the mass balance, the residual soil Pb or As was determined using a strong acid extraction method (EPA 3050B; 10 mL of 50% HNO_3 , 5 mL of concentrated HNO_3 , 2 mL of water + 3 mL of 30% H_2O_2 at $95 \pm 5^\circ\text{C}$) after each PBET extraction. After digestion, the samples were filtered using a Whatman filter paper, and the filtrate was measured with AAS to obtain the total metal amounts remaining on the soil. An analysis of the soil residues from the procedure yielded a mass recovery of $100 \pm 10\%$.

III. RESULTS AND DISCUSSION

A. Effect of Aging Time

Figure 1 shows the water-soluble/exchangeable and bioaccessible concentrations of Pb(II) and As(V) in contact with soils from the C-horizon as a function of aging time. The soil rapidly and strongly sequestered both Pb(II) and As(V). The CaCl_2 -extractable Pb(II) and As(V) was less than 3% of the total soil concentration over all time periods. The bioaccessibility of arsenic was rapidly and dramatically reduced, decreasing from $11.3 \pm 0.7\%$ initially to $5.8 \pm 0.2\%$ after 6 months, a significant decrease ($p < 0.001$). The Pb(II) bioaccessibility was greater than that of arsenic, $62.6 \pm 3.2\%$ initially with no further significant sequestration over 6 months. The reductions in bioaccessibility are due to metal-soil interactions rather than preexisting solid phase speciation, as soluble metals were added to the soil




FIGURE 1

Simulated stomach bioaccessibility of As(V) and Pb(II) with aging times (C-horizon soil, pH 2, 1:40 soil/solution ratio). Error bars are \pm one standard deviation ($n = 2-4$).

initially. This is important because it implies a long-term reduction in bioaccessibility as long as the soil properties governing metal sequestration do not change. This is in contrast to long-term changes in metal speciation (e.g., metal sulfide oxidation) that may be a concern in situations where metal speciation, as opposed to soil-metal interactions, is controlling bioaccessibility.

B. Effect of Simulated Gastrointestinal pH

The pH of the stomach is variable, ranging from approximately 2 (fasting) to 4 to 5 after eating (Ruby *et al.* 1996). To examine the potential effects of different pH conditions in a simulated stomach, the bioaccessibility of As(V) and Pb(II) were measured at four different pH values. Figure 2a shows a comparison of the bioaccessibility of As(V) and Pb(II) in freshly spiked B-horizon soil at four different simulated stomach pH values. The bioaccessibility of arsenic was constant at $25.9 \pm 6.8\%$ with pH over the range 1.5 to 4. The differences between the bioaccessibility in the B-horizon (Figure 2a) and C-horizon (Figure 1) may be due




FIGURE 2A

Simulated stomach bioaccessibility of As(V) and Pb(II) from fresh soil with variation of pH of PBET solution (B-horizon soil, 1:40 soil/solution ratio). Error bars are \pm one standard deviation ($n = 2-3$).

to differences in particle size used ($<250 \mu\text{m}$ vs. $<2000 \mu\text{m}$) or small changes in the amount and reactivity of Fe or other metal-sequestering solid phases. The Pb(II) bioaccessibility, in contrast to As(V), exhibited a greater pH dependence. At pH 1.5, $81.1 \pm 1.3\%$ of lead was bioaccessible, while only $11.1 \pm 0.7\%$ was bioaccessible at pH 4. This result suggests that the bioaccessibility of lead is strongly affected by the stomach pH. Thus, an eightfold variation in bioaccessibility is possible due to a daily variation in stomach pH. This phenomenon illustrates another source of uncertainty that must be considered in conducting a risk assessment. From studies of the bioavailability of soil-borne lead in adults, Maddaloni *et al.* (1998) reported a great difference in lead absorption between fasting (26.1%) and after eating (2.5%).

After approximately 2 h of residence time in the stomach, food enters the small intestine where the pH increases to approximately 7 (Ruby *et al.*, 1996). Although the overall dissolution of Pb(II) and As(V) may be controlled by the stomach, it is not clear that all the metals dissolved in the stomach may be absorbed, because the proximal area of the small intestine is known as the primary region of heavy metal




FIGURE 2B

Simulated stomach (pH 2) and small intestine (pH 7) bioaccessibility of As(V) and Pb(II) from fresh B-horizon soil (1:40 soil/solution ratio). Error bars are \pm one standard deviation ($n = 2-3$).

absorption (Ashmead *et al.*, 1985). To simulate the small intestine, the pH of the extraction solution was increased to 7 by the addition of NaHCO_3 . Metal bioaccessibility in the simulated small intestine following digestion is shown in Figure 2b. The bioaccessible As was not significantly affected as the pH of the extraction solution was changed from 2 to 7, suggesting that the pH is not a major controlling parameter for the dissolution of As(V) from this soil at this pH range. However, lead bioaccessibility decreased significantly ($p < 0.01$) from $76.7 \pm 3.1\%$ to $37.4 \pm 2.3\%$. As the small intestine is the major region of heavy metal absorption, the bioaccessibility of Pb in the stomach may be greater than the actual bioavailability.

The pH-dependent bioaccessibility of As(V) and Pb(II) can be understood in terms of standard geochemical phenomena. For example, cationic metals (e.g., Pb(II)) typically partition to solids to a greater degree at higher pH, while anionic metals (e.g., As(V)) exhibit the opposite behavior, as shown in Figure 2c. Therefore, the lower bioaccessibility of Pb(II) at higher pH may be due to the same factors (e.g., pH-dependent sorption) that favor Pb(II) adsorption at higher pH. In contrast, As(V) bioaccessibility was relatively independent of simulated stomach




FIGURE 2C

Pb(II) and As(V) adsorption onto B-horizon soil as a function of pH (5 g/L soil; 1 mg/L Pb(II) and As(V); I = 0.01 M NaNO₃).

and small intestine pH, which is consistent with a relatively little variation in adsorption from pH 2 to 7 (Figure 2c).

C. Effect of Soil pH

B-horizon soil was used to study initial soil pH effects on metal bioaccessibility. As shown in Figure 3, the effect of soil pH on As(V) and **Pb(II) bioaccessibility was different**. Although As(V) sorption increased sharply from pH 7 to 9 (Figure 2c), the variation of As(V) bioaccessibility was relatively small over the pH range 4.5 to 9. These results indicate that the As(V) bioaccessibility in this soil is controlled by the simulated stomach and small intestine pH rather than the initial soil pH, possibly reflecting relatively rapid pH-dependent partitioning in the solution phase (i.e., As(V) partitioning responds relatively rapidly to solution pH independent of initial soil pH). In contrast, Pb(II) bioaccessibility significantly ($p < 0.02$) decreased from $76.7 \pm 3.1\%$ to $47.2 \pm 3.2\%$ at higher soil pH, reflecting




FIGURE 3

Simulated stomach bioaccessibility of As(V) and Pb(II) with variation of soil pH (fresh B-horizon soil, pH 2, 1:40 soil/solution ratio). Error bars are \pm one standard deviation ($n = 2-3$).

the same pattern as typical cationic-type adsorption (Figure 2c). These results indicate that the binding of Pb(II) in the soil is influenced by the initial soil pH, and that the Pb(II) bioaccessibility depended on both the simulated stomach (Figure 2a) and soil (Figure 3) pH. In contrast, the As(V) bioaccessibility was relatively independent of both the simulated stomach (Figure 2a) and soil (Figure 3) pH.

D. Effect of Concentration

Figure 4 shows the bioaccessibility of Pb(II) and As(V) as a function of soil metal concentration. The current risk assessment methodology implicitly assumes that the bioavailability is independent of the concentration by using a constant relative bioavailability adjustment factor. However, metals often partition to the solid phase in a nonlinear manner (i.e., the fraction of metal sorbed decreases with increasing concentration). In order to investigate the accuracy of using a constant bioaccessibility, Pb(II) and As(V) bioaccessibility were measured over almost three orders of magnitude of concentration (10 to 10,000 mg/kg). Pb(II)




FIGURE 4

Simulated stomach bioaccessibility of As(V) and Pb(II) with variation of initial concentration (fresh B-horizon soil, pH 2, 1:40 soil/solution ratio). Error bars are \pm one standard deviation ($n = 2-3$).

bioaccessibility was relatively insensitive to the concentration of lead, yielding $78.7 \pm 6.8\%$ bioaccessibility over all concentration ranges. However, As(V) bioaccessibility significantly ($p < 0.05$) increased from $23.0 \pm 4.2\%$ to approximately $42.8 \pm 2.7\%$ as the concentration of As(V) increased from approximately 10 to 1000 mg/kg, illustrating another potential source of uncertainty introduced in risk assessments by using a constant bioavailability adjustment.

E. Effect of Soil to Solution Ratio

The soil to solution ratio in the stomach will not be constant over time due to the ingestion of different amounts of soil on varying occasions and because the volume of fluid in the stomach depends on the fasting condition of the child. As a result, the bioaccessibility of As(V) and Pb(II) was investigated as a function of soil to solution ratio (Figure 5). The bioaccessibility increased from 30.8 ± 13.8 to 53.2 ± 0.9 for As(V) and from 81.8 ± 1.8 to 91.0 ± 2.1 for Pb(II) with decreasing soil




FIGURE 5

Simulated stomach bioaccessibility of As(V) and Pb(II) from fresh B-horizon soil (pH 1.5, 1:40 and 1:100 soil/solution ratio). Error bars are \pm one standard deviation ($n = 2$).

to solution ratio. The same trend was observed in soluble and exchangeable Pb(II) in the CaCl_2 solution, especially for aged soil samples. Hamel *et al.* (1998) reported that the effect of the soil to solution ratio on lead and arsenic bioaccessibility depended on the soil sample. Although a higher bioaccessibility of Pb and As from Jersey City soil was observed as the soil to solution ratio decreased from 1:100 to 1:5000, a relatively constant bioaccessibility of Pb(II) and As(V) was observed with Montana soils over all soil to solution ratios tested.

IV. SUMMARY AND CONCLUSIONS

These results have illustrated several salient aspects of Pb(II) and As(V) bioaccessibility. First, the soils decreased both Pb(II) and especially As(V) bioaccessibility solely as a result of soil-metal interactions and not as a result of any specific preexisting metal speciation. Reduced Pb(II) and As(V) bioaccessibility then can be a result of the fundamental nature of soil-metal interactions rather than site-specific speciation (e.g., metal sulfides from ore bodies). These results also promote greater confidence in the long-term ability of soil to lower Pb(II) and

As(V) bioaccessibility as long as the soil properties governing metal sequestration remain constant. This might not be the case if the reduced Pb(II) or As(V) bioaccessibility was due to unique metal speciation that was subject to change over time (e.g., oxidation of metal sulfides in surface soils). In fact, the As(V) bioaccessibility significantly decreased over a 6-month aging period. Second, Pb(II) bioaccessibility significantly depended on the pH of both the simulated GI fluid and the soil, showing enhanced sequestration and reduced bioaccessibility at higher pH values. In contrast, neither soil nor GI pH significantly affected As(V) bioaccessibility over the range of GI pH from 2 to 7 and soil pH from 4.5 to 9.4. Third, although Pb(II) bioaccessibility was not significantly influenced by soil-metal concentration over the range 10 to 10,000 mg/kg, the As(V) bioaccessibility significantly increased over this same concentration range. Thus, the use of a concentration-independent bioaccessibility/bioavailability factor in a risk assessment for As(V) may not be warranted. Finally, both Pb(II) and As(V) bioaccessibility increased with decreasing soil to solution ratio, illustrating another degree of uncertainty in estimating the risk of soil ingestion at metal-contaminated sites.

ACKNOWLEDGMENTS

The authors acknowledge the comments of two anonymous reviewers that greatly improved the paper. This research was sponsored by the Strategic Environmental Research and Development Program (SERDP) under the direction of Ms. Cathy Vogel and Dr. Andrea Leeson. We appreciate the assistance of Beth Derrick-Williams in measuring soil-metal partitioning.

REFERENCES

- Ashmead, H. D., Graff, D. J., and Ashmead, H. H. 1985. *Intestinal Absorption of Metal Ions and Chelates*, pp. 76–77. (Ed.) Charles C Thomas, Springfield, IL.
- Barnett, M. O., Harris, L. A., Turner, R. R., Stevenson R. J., Henson, T. J., Melton, R. C., and Hoffman, D. P. 1997. Formation of mercuric sulfide in soil. *Environ. Sci. Technol.* **31**, 3037–3043.
- Casteel, S. W., Cowart, R. P., Weis, C. P., Henningsen, G. M., Hoffman, E., Brattin, W. J., Guzman, R. E., Starost, M. F., Payne, J. T., Stockham, S. L., Becker, S. V., Drexler, J. W., and Turk, J. R. 1997. Bioavailability of lead to juvenile swine dosed with soil from the smuggler mountain NPL site of Aspen, Colorado. *Fundam. Appl. Toxicol.* **36**, 177–187.
- Cheng, Y. L., Preslan, J. E., Anderson, M. B., and George, W. J. 1991. Solubility and bioavailability of lead following oral ingestion of vitrified slagged aggregate. *J. Hazard. Mater.* **27**, 137–147.
- Davis, A., Drexler, J. W., Ruby, M. V., and Nicholson, A. 1993. Micromineralogy of mine wastes in relation to lead bioavailability, Butte, Montana. *Environ. Sci. Technol.* **27**, 1415–1425.
- Davis, A., Ruby, M. V., and Bergstrom, P. D. 1992. Bioavailability of arsenic and lead in soils from the Butte, Montana, mining district. *Environ. Sci. Technol.* **26**, 461–468.

- Davis, A., Ruby, M. V., Bloom, M., Schoof, R., Freeman, G., and Bergstrom, P. D. 1996. Mineralogic constraints on the bioavailability of arsenic in smelter-impacted soils. *Environ. Sci. Technol.* **30**, 392–399.
- Davis, A., Sherwin, D., Ditmars, R., and Hoenke, K. A. 2001. An analysis of soil arsenic records of decision. *Environ. Sci. Technol.* **35**, 2401–2406.
- Dieter, M. P., Matthews, H. B., Jeffcont, R. A., and Mosemers, R. F. J. 1993. Comparison of lead bioavailability in F344 rats fed lead acetate, lead-oxide, lead sulfide, or lead ore concentrate from Skagway, Alaska. *J. Toxicol. Environ. Health* **39**, 79–93.
- Freeman, G. B., Johnson, J. D., Liao, S. C., Feder, P. I., Davis, A. O., Ruby, M. V., Schoof, R. A., Chaney, R. L., and Bergstrom, P. D. 1994. Absolute bioavailability of lead acetate and mining waste lead in rats. *Toxicology* **91**, 151–163.
- Freeman, G. B., Schoof, R. A., Ruby, M. V., Davis, A. O., Dill, J. A., Liao, S. C., Lapin, C. A., and Bergstrom, P. D. 1995. Bioavailability of arsenic in soil and house dust impacted by smelter activities following oral administration in cynomolgus monkeys. *Fundam. Appl. Toxicol.* **28**, 215–222.
- Hamel, S. C., Buckley, B., and Lioy, P. J. 1998. Bioaccessibility of metals in soils for different liquid to solid ratios in synthetic gastric fluid. *Environ. Sci. Technol.* **32**, 358–362.
- Maddaloni, M., Lolocono, N., Manton, W., Blum, C., Drexler, J., and Graziano, J. 1998. Bioavailability of soilborne lead in adults, by stable isotope dilution. *Environ. Health Perspect.* **106**, 1589–1594.
- Polak, J. E., Oflaherty, E. J., Freeman, G. B., Johnson, J. D., Liao, S. C., and Bergstrom, P. D. 1996. Evaluating lead bioavailability data by means of a physiologically based lead kinetic model. *Fundam. Appl. Toxicol.* **29**, 63–70.
- Preslan, J. E., Chang, C. Y., Schiller, N. K., and George, W. J. 1996. Bioavailability of lead from vitrified slagged aggregate. *J. Hazard. Mater.* **48**, 207–218.
- Ruby, M. V., Davis, A., Kempton, J. H., Drexler, J. W., and Bergstrom, P. D. 1992. Lead bioavailability-dissolution kinetics under simulated gastric conditions. *Environ. Sci. Technol.* **26**, 1242–1248.
- Ruby, M. V., Davis, A., Schoof, R., Eberle, S., and Sellstone, C. M. 1996. Estimation of lead and arsenic bioavailability using a physiologically based extraction test. *Environ. Sci. Technol.* **30**, 422–430.
- Ruby, M. V., Schoof, R., Brattin, W., Goldade, M., Post, G., Harnois, M., Mosby, D. E., Casteel, S. W., Berti, W., Carpenter, M., Edwards, D., Cragin, D., and Chappell, W. 1999. Advances in evaluating the oral bioavailability of inorganics in soil for use in human health risk assessment. *Environ. Sci. Technol.* **33**, 3697–3705.
- Schilderman, P. E., Moonen, J. C., Kempkers, P., and Kleinjans, J. C. S. 1997. Bioavailability of soil-adsorbed cadmium in orally exposed male rats. *Environ. Health Perspect.* **105**, 234–238.
- Sheppard, S. C., Evenden, W. G., and Schwatz, W. J. 1995. Ingested soil bioavailability of sorbed lead, cadmium, cesium, iodine, and mercury. *J. Environ. Qual.* **24**, 498–505.

INTERNAL DISTRIBUTION

- | | |
|--------------------------------|----------------------------------|
| 1. G. K. Jacobs, 1505, MS-6037 | 3. – 4. ESD Library |
| 2. D. E. Fowler, 1505, MS-6037 | 5. ORNL Central Research Library |
| | 6. ORNL Laboratory Records-RC |
| | 7. P. M. Jardine, 1505, MS-6038 |
| | 8. C. C. Brandt, 1505, MS-6038 |
| | 9. S. A. Heuscher, 1505, MS-6036 |

ELECTRONIC NOTIFICATION

10. Scott Dockum (sdockum@hgl.com)
11. Andrea Leeson (andrea.leeson@osd.mil)
12. Richard Mach (MachRG@navfac.navy.mil)
13. Alicia Shepard (aanderson@hgl.com)
14. Annette Gatchett (gatchett.annette@epa.gov)
15. Beth Moore (beth.moore@em.doe.gov)
16. Chuck Coyle (charles.g.coyle@nwd02.usace.army.mil)
17. Don Ficklen (Holmes.Ficklen@brooks.af.mil)
18. Erica Becvar (Erica.Becvar@brooks.af.mil)
19. Erik Hangeland (erik.hangeland@aec.apgea.army.mil)
20. Grover Chamberlain (grover.chamberlain@em.doe.gov)
21. Hans Stroo (hstroo@retec.com)
22. Ivette O'Brien (ivette.obrien@brooks.af.mil)
23. Jeff Breckenridge (jeff.l.breckenridge@usace.army.mil)
24. Jeff Cornell (Jeffrey.Cornell@pentagon.af.mil)
25. John Cullinane (cullinm@wes.army.mil)
26. Linda Chrissey (chrisel@onr.navy.mil)
27. Mark Hampton (mark.hampton@aec.apgea.army.mil)
28. Martin Nguyen (mnguyen@comdt.uscg.mil)
29. Marvin Unger (MUnger@retec.com)
30. Michelle Simon (Simon.Michelle@epamail.epa.gov)
31. Rajat Ghosh (rghosh@retec.com)
32. Rebecca Biggers (rbigger@nfesc.navy.mil)
33. Katie Houff (khouff@hgl.com)
34. Katherine Perdue (KPerdue@hgl.com)
35. Mark Barnett (barnettm@eng.auburn.edu)
36. Rebecca Bigger (biggersr@nfesc.navy.mil)
37. Nick Basta (basta.4@osu.edu)
38. Amy Hawkins (amy.hawkins@navy.mil)
39. Roman Lanno (lanno.1@osu.edu)
40. Stan Casteel (CasteelS@missouri.edu)
41. Jeff Marqusee (jeffery.marqusee@osd.mil)
42. Valerie Eisenstein (vke@hgl.com)