

**Determinations of the Matrix Element V(Ub) From Inclusive Semileptonic B
Decays With Reduced Model Dependency
for the Enriched Xenon Observatory**

Edward J. Hill

Stanford Linear Accelerator Center
Stanford University
Stanford, CA 94309

SLAC-Report-814

Prepared for the Department of Energy
under contract number DE-AC02-76SF00515

Printed in the United States of America. Available from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161.

This document, and the material and data contained therein, was developed under sponsorship of the United States Government. Neither the United States nor the Department of Energy, nor the Leland Stanford Junior University, nor their employees, nor their respective contractors, subcontractors, or their employees, makes an warranty, express or implied, or assumes any liability of responsibility for accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use will not infringe privately owned rights. Mention of any product, its manufacturer, or suppliers shall not, nor is it intended to, imply approval, disapproval, or fitness of any particular use. A royalty-free, nonexclusive right to use and disseminate same of whatsoever, is expressly reserved to the United States and the University.

UNIVERSITY OF CALIFORNIA, SAN DIEGO

Determinations of $|V_{ub}|$ from Inclusive Semileptonic B Decays
with Reduced Model Dependency

A dissertation submitted in partial satisfaction of the
requirements for the degree Doctor of Philosophy

in

Physics

by

Edward J. Hill

Committee in charge:

Professor David MacFarlane, Chair
Professor James Branson
Professor Bruce Driver
Professor Douglas Magde
Professor Aneesh Manohar

2005

TABLE OF CONTENTS

Signature Page	iii
Dedication	iv
Table of Contents	v
List of Figures	viii
List of Tables	xix
Acknowledgments	xxii
Vita	xxiii
Abstract	xxiv
I Introduction	1
1.1 Outline of Dissertation	2
2 The Theory of Weak Interactions and the Extraction of V_{ub}	5
2.1 The Standard Model	5
2.2 The CKM Picture of Weak Interactions	8
2.3 Semileptonic Decays	13
2.4 Inclusive Decays of the B Meson	16
2.5 Extracting $ V_{ub} $ from Charmless Inclusive Semileptonic Decays	17
2.6 Extracting $ V_{ub} $ in the Real World	18
2.7 Extracting $ V_{ub} $ with Reduced Model Dependency	22
3 The BABAR Experiment	28
3.1 PEP-II B Factory	28
3.2 BABAR Overview	32
3.3 Charged Particle Tracking System	35
3.3.1 Silicon Vertex Tracker	35
3.3.2 Drift Chamber	39
3.3.3 Tracking Summary	41
3.4 Detector of Internally Reflected Cherenkov Light	43
3.5 Electromagnetic Calorimeter	46
3.6 Instrumented Flux Return	49
3.7 Trigger System	52
3.7.1 The Level 1 Trigger System	52
3.7.2 The Level 3 Trigger System	53
3.8 Data Acquisition System	55

4	Particle Reconstruction	57
4.1	Charged Particle Reconstruction	57
4.2	Charged Particle Identification	59
4.2.1	Electron Identification	60
4.2.2	Muon Identification	63
4.2.3	Kaon Identification	64
4.3	Neutral Particle Reconstruction	65
4.4	π^0 and K^0_S Reconstruction	67
4.4.1	π^0 Reconstruction	67
4.4.2	K^0_S Reconstruction	68
4.5	D Meson Reconstruction	69
4.5.1	D^+ Meson Selection	70
4.5.2	D^* Meson Selection	71
4.5.3	D^{**} Meson Selection	73
4.5.4	D^{*0} Meson Selection	73
4.6	B Meson Reconstruction	74
4.6.1	Energy conservation and ΔE variable	75
4.6.2	B mass reconstruction and m_{tag} variable	76
5	Semi-Exclusive Reconstruction	80
5.1	Introduction	80
5.2	Reconstruction Method	81
5.2.1	Selection of B candidates	82
5.3	Categorization and Summary of B Modes	83
5.4	Signal Event Selection	84
5.5	Selection of the Best B	85
6	Analysis on the Recoil	89
6.1	Introduction	89
6.2	Data and Monte Carlo Samples	91
6.2.1	Data Sample	91
6.2.2	Monte Carlo Samples	91
6.2.3	Summary of Samples	97
6.3	Recoil Reconstruction	99
6.3.1	Recovery of Bremsstrahlung Photons	99
6.3.2	Reconstruction of the Hadronic System	100
6.3.3	Event Based Selection	101
6.3.4	Comparison of Data and Monte Carlo	112
7	Signal Extraction and Results	120
7.1	Extraction of $\delta\Gamma(c)$	120
7.1.1	Extraction of N_c, the Number of $b - u\bar{d}\nu$ Signal Events	121
7.1.2	Extraction of f	123

7.1.3	Extraction of N_{SL} , the Number of Semileptonic Events	125
7.1.4	Efficiency Corrections	126
7.1.5	Fit Validation	126
7.1.6	Summary of $\delta\Gamma(c)$ Results	127
7.2	Extraction of I_0 and I_s from the differential $b \rightarrow s\gamma$ Photon Spectrum	128
7.3	Determination of $ V_{ub} $ and $\mathcal{B}(\bar{B} \rightarrow X_u \ell \bar{\nu})$	129
8	Systematic Uncertainties	138
8.1	$\delta\Gamma(c)$ Systematics	138
8.1.1	E_{reco} Reconstruction	138
8.1.2	Fit to the m_{CS} Distributions	139
8.1.3	Floating the Other Component	140
8.1.4	B_{track} Tagging Efficiency ($\epsilon_t^d/\epsilon_t^u$)	140
8.1.5	Binning of the m_X Distribution	140
8.1.6	Tracking	141
8.1.7	Lepton Identification	142
8.1.8	Charged Kaon Identification	142
8.1.9	Neutral Reconstruction	142
8.1.10	K_L^0 Reconstruction	143
8.1.11	Semileptonic B Branching Fractions	143
8.1.12	$B \rightarrow D^* l \bar{\nu}$ Form Factors	144
8.1.13	Charm Decay Branching Fractions	145
8.1.14	Signal Model and Parameterization	145
8.1.15	Fermi Motion	146
8.1.16	Modeling of $\bar{B} \rightarrow X_u \ell \bar{\nu}$ Decays	147
8.1.17	Hadronization Uncertainties	148
8.1.18	$s\bar{s}$ Popping	149
8.1.19	Monte Carlo Statistics	149
8.1.20	Stability Checks	149
8.2	Statistical and Systematic Uncertainties from the $b \rightarrow s\gamma$ Photon Energy Spectrum	155
8.3	Theoretical Uncertainties on the Extraction of $ V_{ub} $	155
8.3.1	Scale Uncertainty of α	156
8.3.2	Perturbative Error	157
8.3.3	Non-perturbative Error	157
9	Conclusions	158
A	Fit to the m_{CS} Distribution	162
B	Fit Results on Various Subsamples	175
	References	179

LIST OF FIGURES

2.1	The three unitary triangles detailed in Eq. 2.13 (a), 2.14 (b), and 2.15 (c)	11
2.2	The unitary triangle.	12
2.3	The rescaled unitary triangle, drawn in the (ρ, η) plane.	13
2.4	The experimental constraints for the sides and angles of the unitary triangle in the (β, η) plane.	14
2.5	The semileptonic decay of a B meson. The complicated QCD effects carried by the gluons (depicted by the curly lines) are located entirely within the hadronic current.	15
2.6	The contributing terms to the transition operator \mathbf{T} (left), and the corresponding operators in the OPE (right). The open squares represent the four-fermion interaction in the weak Lagrangian \mathcal{L}_{eff} and the black dots represent local operators of the $1/m_b$ expansion	16
2.7	The lepton energy spectra of semileptonic decays. The charmless $b \rightarrow u\bar{\nu}$ decays are shown in blue while the $b \rightarrow c\bar{l}\bar{\nu}$ decays are shown in red. The high momentum region above 2.3 GeV/c is dominated by $b \rightarrow u\bar{\nu}$ decays.	19
2.8	The hadronic mass spectra from semileptonic decays before the effects of detector resolution. The charmless $b \rightarrow u\bar{\nu}$ decays are shown in blue while the $b \rightarrow c\bar{l}\bar{\nu}$ decays are shown in red. Note the logarithmic scale for the vertical axis. The low mass region is dominated by $b \rightarrow u\bar{\nu}$ decays and the region above m_D is dominated by $b \rightarrow u\bar{\nu}$ decays.	20
3.1	The Stanford Linear Accelerator Center.	28
3.2	The polar angle relationship between the CM frame (expressed as the inscribed $\cos \theta_{\text{CM}}$ lines) and the laboratory frame (outer markings on the protractor). The detector active region lies between 350 mrad in the forward direction and 400 mrad in the backward direction.	29
3.3	The total integrated luminosity delivered by PEP-II (blue) and recorded by BABAR (red). The delivered off-peak data is shown in green.	31

3.4	The daily recorded luminosity delivered by PEP-II (blue) and recorded by <i>BABAR</i> (red).	32
3.5	A longitudinal cutaway (top) and a horizontal cutaway (bottom) view of the <i>BABAR</i> detector.	34
3.6	The Silicon Vertex Tracker (SVT).	37
3.7	A longitudinal view of the SVT. The roman numerals indicate the six different types of sensor modules.	37
3.8	A cross-sectional view of the SVT. The orientation of the five layers and the individual strip sensors is shown.	38
3.9	A longitudinal cutaway view of the DCH. The dimensions shown are in units of mm. Note that the interaction point, labeled IP, is offset 370 nm from the center of the DCH.	40
3.10	The ten DCH superlayers (left) and the layout of the drift cells for the first four superlayers (right). The numbers on the right side of the right plot give the stereo angle of the layer in units of mrad and the lines drawn between field wires illustrate the boundaries of the drift cells.	41
3.11	Drift cell 100 ns isochrones in the third and fourth layer of an axial superlayer. The isochrones are quite circular near the sense wires but become non-circular near the edges of a cell.	42
3.12	The track reconstruction efficiency in the DCH at operating voltages of 1900 V and 1960 V, as a function of transverse momentum (top), and polar angle (bottom). The efficiency is measured using multi-hadron events.	43
3.13	Track parameter differences between upper and lower tracks for tracks with $p_T > 3 \text{ GeV}/c$ as determined from cosmic ray muons.	44
3.14	The p_T resolution as a function of p_T as determined from cosmic ray muons. The resolution is well described by a linear function.	44
3.15	The DCH dE/dx measurement vs. the track momentum. The lines display the Bethe-Bloch predictions for the dE/dx measurement.	45
3.16	A longitudinal cutaway view of the DRC.	46
3.17	DRC schematic. The radiator bar reflects Cherenkov photons and the PMTs provide the imaging to determine the Cherenkov angle θ_C	47

3.18	The fitted Cherenkov angle versus track momentum from a sample of multi-hadron events. The lines drawn indicate the predicted values for the various particles.	48
3.19	The expected π - K separation in $B^0 \rightarrow \pi^+\pi^-$ events versus the track momentum.	49
3.20	A longitudinal cutaway view of the top half of the EMC. Cross-sections of the 56 crystal rings are visible. Dimensions are listed in mm.	50
3.21	A schematic diagram of a single CsI(Tl) crystal. Please note that this figure is not to scale.	51
3.22	The energy resolution (left) and angular resolution (right) of the EMC. The solid lines are fits described in Eqs. 3.4 and 3.5.	52
3.23	The electron identification efficiency and the pion misidentification probability as a function of the momentum (a) and the polar angle (b). The efficiency is indicated by the axis labels on the left and the misidentification probability is indicated by the axis labels on the right.	53
3.24	An overview of the IFR. The Barrel is pictured on the left and the end doors are pictured on the right. The dimensions listed are in mm.	54
3.25	A cross sectional view of a planar RPC.	54
3.26	The muon identification efficiency and pion misidentification probability as a function of the momentum (a) and the polar angle (b). The efficiency is indicated by the axis labels on the left and the misidentification probability is indicated by the axis labels on the right.	55
3.27	A schematic of the BABAR DAQ system.	56
4.1	Event with looping tracks and ghost tracks. Note that this is not a common event: Loopers affect a relatively small number of events and to have the coincidence with ghost tracks (seen in the lower right quadrant) is very rare.	59
4.2	Distribution for E/p (top left) using a control sample of electrons and E/p versus momentum (top right), polar angle (bottom left) and azimuthal angle (bottom right).	61

4.3	Electron identification and hadron misidentification probability for the likelihood-based electron selector as a function of momentum (left) and polar angle (right). Note the different scales for identification and misidentification on the left and right ordinates, respectively. The measurements are for luminosity-averaged rates for Run-1 and Run-2.	63
4.4	Muon identification and hadron misidentification probability for the tight muon selector as a function of momentum (left) and polar angle (right). The solid markers indicate the efficiency in 2000, the empty markers the efficiency in 2001. Note the different scales for identification and misidentification on the left and right ordinates, respectively.	65
4.5	Charged kaon identification and pion misidentification probability for the tight kaon micro selector as a function of momentum (left) and polar angle (right). The solid markers indicate the efficiency for positive particles, the empty markers the efficiency for negative particles. Note the different scales for identification and misidentification on the left and right ordinates.	66
4.6	Definition of the variables r_0 , φ_0 and R_0	67
4.7	π^0 peaks for simulated events and for data.	68
4.8	Mass distributions for $K_S^0 \rightarrow \pi^+ \pi^-$. The distribution is fitted with a sum of a double Gaussian and a first-order polynomial function.	69
4.9	K_S momentum (left) and polar angle (right) distributions in data (solid markers) and Monte Carlo simulation (hatched histogram).	69
4.10	D^0 candidates selected for $D^0 \rightarrow K\pi$, $D^0 \rightarrow K\pi\pi^0$, $D^0 \rightarrow K_S^0\pi\pi$, and $D^0 \rightarrow K3\pi$ modes.	71
4.11	Distribution of soft pion momentum in the $T(4S)$ frame (left) and $m(D^0\pi^+) - m(D^0)$ mass distribution for D^{*+} candidates in the $B \rightarrow D^{*+}\pi^+$, $D^0 \rightarrow K\pi$ mode. Vertical lines indicate the signal windows used in the selection.	73
4.12	Δm distribution for $D^{*0} \rightarrow D^0\pi^0$ decays, where $p^*(D^0) < 2.5$ GeV/c	75
4.13	An example of ΔE distribution for $B \rightarrow D^{*+}\pi^-$ with $D^0 \rightarrow K\pi$.	76
4.14	ΔE versus m_{ES} for the decay, $B \rightarrow D^{*+}\pi^-$ with $D^0 \rightarrow K\pi$.	77

4.15	Left: m_{ES} distribution for candidates in the off-resonance data (40 MeV below the $\Upsilon(4S)$ mass). Right: m_{ES} distribution for $b\bar{b}$ background (B^0 reconstructed as B^+). ARGUS shape fit is superimposed in both cases.	78
4.16	MC m_{ES} distributions for reconstructed B modes with (left) no π^0 in the final state, (middle) 1 π^0 in the final state and (right) 2 π^0 's in the final state. The fit function is a sum of crystal ball and ARGUS function.	79
5.1	Definition of the $m_{\text{ES}}, \Delta E$ regions. In each iteration of the semi-exclusive reconstruction a combination is used (A) only as candidate, (B) both as candidate and as seed, (C) used as seed but not as candidate and (D) discarded.	83
5.2	ΔE distributions for the four D^0 decay modes in $B \rightarrow D^* \pi \pi^0$, $D^* \rightarrow D^0 \pi$: a) $D^0 \rightarrow K\pi$, b) $D^0 \rightarrow K\pi\pi^0$, c) $D^0 \rightarrow K3\pi$, and d) $D^0 \rightarrow K_s^0 \pi\pi$	87
5.3	Dependence of the quality factor $S/\sqrt{S+B}$ as a function of the yield when adding modes for the $B^0 \rightarrow D^{**} X$ case. Statistics corresponds to 80 fb^{-1}	88
6.1	Semileptonic $B_{\text{ recoil}}$ decay recoiling from a fully reconstructed B meson.	90
6.2	m_X distributions at generator level for pure resonant (left) and pure non-resonant $b \rightarrow u\ell\bar{\nu}$ MC simulation (right).	93
6.3	Parton-level distributions for the generator without Fermi motion. The upper left plot shows the scaled lepton energy $x = 2E_\ell/m_b$, the upper right plot the scaled hadron energy $z = 2(E_h - \vec{p})/m_b$, the lower left plot shows the scaled hadron mass squared $s_h = m_h^2/m_b^2$ and the lower right plot shows the scaled virtual W mass squared $Q^2 = q^2/m_b^2$. The red lines denote analytical single differential functions with the same input quantities and the green lines show the tree results without α_s corrections.	94
6.4	m_X distribution at the generator-level for the hybrid signal Monte Carlo for $b \rightarrow u\ell\bar{\nu}$ events. The resonant and non-resonant contributions to the hybrid are shown in magenta and red respectively along with the purely non-resonant distribution in blue.	96

6.5	Event yields for all seeds combined without additional requirement on the recoil system (left) and after requiring a lepton with $p_{\text{ons}} > 1 \text{ GeV}/c$ (right). The numbers printed on each plot indicate signal yield (S), background yield (B), and purity (P) all in the signal region defined by $m_{\text{gg}} > 5.27 \text{ GeV}/c^2$. Only the Argus background is fitted in these plots with the signal yield taken as the difference between the histogram and the fitted Argus background. These plots serve as an illustration of the event yields and are not used in the analysis.	98
6.6	Signal MC: a) p^* resolution for all electrons, b) p^* resolution for electrons that are known through generator truth-matching to contain at least one Bremsstrahlung photon, c) m_X distribution. The blue line corresponds to no recovery and the red points correspond to recovery of photons within $\alpha_{\text{brems}} < 0.08$	100
6.7	Left: Missing-mass squared of the event before (upper plot) and after (lower plot) the kinematic fit. Due to the zero-mass hypothesis for the neutrino, the missing-mass of the event after the fit is within the precision compatible with zero. Right: Mass resolution of the B_{reco} (left side) before (upper plots) and after (lower plots) the kinematic fit. Due to the imposed equal-mass constraint, the masses of the two B mesons are, within precision, identical after the fit.	102
6.8	Cocktail MC: The resolution on the kinematically fitted hadronic mass with all analysis cuts applied for $b \rightarrow u\ell\nu$ events (left) and $b \rightarrow c\ell\nu$ events (right).	103
6.9	Cocktail MC: The mean (left) and RMS (right) of the resolution on the kinematically fitted hadronic mass with all analysis cuts applied in bins of true m_X (upper) and m_{miss}^2 (lower). The blue points are $b \rightarrow u\ell\nu$ events and the red points are $b \rightarrow c\ell\nu$ events.	104
6.10	Generic MC: Statistical significance ($S/\sqrt{S+B}$) as a function of the purity of the mode of the reconstructed B sample for the four charm seeds after all cuts: clockwise (starting from the upper left plot) the seeds are $B^0 \rightarrow D^{**}X$, $B^0 \rightarrow D^+X$, $B^+ \rightarrow D^0X$, $B^+ \rightarrow D^{*0}X$	106
6.11	Cocktail MC: The lepton momentum spectrum with all analysis cuts applied	107
6.12	Cocktail MC: The number of leptons observed with all analysis cuts applied	107
6.13	Cocktail MC: The total event charge with all analysis cuts applied.	108
6.14	Cocktail MC: The missing-mass squared with all analysis cuts applied . . .	108

6.15	Cocktail MC: Left: the number of charged kaons with all analysis cuts applied Right: the number of K_S with all analysis cuts applied	109
6.16	Cocktail MC: The missing momentum with all analysis cuts applied	109
6.17	Cocktail MC: The cosine of the angle of the missing system with all analysis cuts applied	110
6.18	Cocktail MC: The momentum of the slowest track with all analysis cuts applied	110
6.19	Cocktail MC: Missing-mass distribution m_{miss, p_R}^2 for a) $D^* \rightarrow D^0\pi$, b) $D^* \rightarrow D\pi^0$, c) $D^{*0} \rightarrow D^0\pi^0$	111
6.20	B^0 (left) and B^+ (right) lepton spectra (side-band subtracted) in generic/cocktail MC and data for $b \rightarrow u\ell\nu$ enhanced (top row) and depleted (bottom row) event samples	113
6.21	Electron (left) and muon (right) spectra (side-band subtracted) in generic/cocktail MC and data for $b \rightarrow u\ell\nu$ enhanced (top row) and depleted (bottom row) event samples	114
6.22	Reconstructed (left) and kinematically fitted (right) hadronic recoil invariant mass spectra (side-band subtracted) in generic/cocktail MC and data for $b \rightarrow u\ell\nu$ enhanced (top row) and depleted (bottom row) event samples	115
6.23	Missing-mass squared (left) and Q^2 (right) distributions (side-band subtracted) in generic/cocktail MC and data for $b \rightarrow u\ell\nu$ enhanced (top row) and depleted (bottom row) event samples	116
6.24	Missing momentum (left) and Θ_{miss} (right) distributions (side-band subtracted) in generic/cocktail MC and data for $b \rightarrow u\ell\nu$ enhanced (top row) and depleted (bottom row) event samples	117
6.25	Charged (left) and neutral (right) multiplicity distributions (side-band subtracted) in generic/cocktail MC and data for $b \rightarrow u\ell\nu$ enhanced (top row) and depleted (bottom row) event samples	118
6.26	Total charge (left) distributions (side-band subtracted) in generic/cocktail MC and data for $b \rightarrow u\ell\nu$ enhanced (top row) and depleted (bottom row) event samples	119
7.1	The fit results for $m_X < 1.55 \text{ GeV}/c^2$	124

7.2	The fit results for $m_X < 1.59 \text{ GeV}/c^2$	124
7.3	The fit results for $m_X < 1.67 \text{ GeV}/c^2$	124
7.4	The fit results for $m_X < 1.75 \text{ GeV}/c^2$	125
7.5	The fit results for $m_X < 1.83 \text{ GeV}/c^2$	125
7.6	The fit results for $m_X < 2.50 \text{ GeV}/c^2$	125
7.7	The fit results for the depleted sample.	127
7.8	Photon energy spectrum for the decay $b \rightarrow s\gamma$ measured by BABAR [69]. This measurement is based on the sum of exclusive modes and results in an excellent photon energy resolution.	129
7.9	$b \rightarrow s\gamma$ photon energy spectrum as a function of $u = 2E_\gamma/m_B$, weighted with the weight functions $i_0(u)$ and $i_+(u)$, calculated for a cut on the hadronic mass $m_X < 1.67 \text{ GeV}/c^2$. The integral of these functions corre- spond to I_0 and I_+ , respectively.	130
7.10	$ V_{ub} $ vs. the upper cut on m_X . The error bars illustrate (from innermost to outermost) the experimental (statistical plus detector systematics), back- ground and signal modeling, $b \rightarrow s\gamma$ errors. Points without the inner error bars have only the statistical error displayed. The shaded error band illustrate the perturbative error (innermost yellow band) and the total the- oretical error (outermost bluish band) where the QED scale uncertainty, the perturbative and non-perturbative error have been added in quadrature.	131
7.11	Enlarged view of $ V_{ub} $ vs. the upper cut on m_X with special emphasis on the statistical error from the $b \rightarrow s\gamma$ photon energy spectrum. The error bars illustrate (from innermost to outermost) the statistical error from the $b \rightarrow s\gamma$ photon energy spectrum and the total error (statistical, experimen- tal systematic, signal and background modeling). The shaded error band illustrate the perturbative error (innermost yellow band) and the total the- oretical error (outermost bluish band) where the QED scale uncertainty, the perturbative and non-perturbative error have been added in quadrature.	132
8.1	Integrated purity for cocktail MC (left), generic MC (middle) and data (right), as an indicator of the sample composition.	139
8.2	The $b \rightarrow c\ell\nu m_X$ distribution (all cuts applied) for cross-feed events com- pared with the total sample. Left plot corresponds to B^0 s, right to $B^+ s$	140

8.3	Relative efficiency for Monte Carlo (histogram) and for data (dots) for the soft pions from $D^{**} \rightarrow D^0\pi^+$, $D^0 \rightarrow K^-\pi^+$ decays, as a function of the momentum.	141
8.4	Measurement of $R_{u/d}$ as a function of the m_X cut applied. The left column displays the results with correlated errors. The right column shows the difference to the default analysis working point with uncorrelated errors shown. The errors are statistical only.	151
8.5	Measurement of $R_{u/d}$ as a function of the $m_{\mu\mu\gamma}^2$ cut applied. The left column displays the results with correlated errors. The right column shows the difference to the default analysis working point with uncorrelated errors shown. The errors are statistical only.	152
8.6	Measurement of $R_{u/d}$ as a function the cut on the lepton momentum applied. The left column displays the results with correlated errors. The right column shows the difference to the default analysis working point with uncorrelated errors shown. The errors are statistical only.	153
8.7	Relative error estimation for $ V_{ub} $ due to the combined statistical and systematic error in the $b \rightarrow s\gamma$ photon energy spectrum.	156
9.1	Results for $ V_{ub} $ extractions from other experiments. The world average is calculated by HFAG [83]. The inclusive results are obtained with shape function parameters obtained from fits to the Belle $b \rightarrow s\gamma$ photon spectrum [82].	161
A.1	DATA: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for B^\pm s.	163
A.2	DATA: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for right sign B^0 s.	164
A.3	DATA: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for wrong sign B^0 s.	165
A.4	GENERIC MC, $b \rightarrow c\ell\bar{\nu}$: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for B^\pm s.	166
A.5	GENERIC MC, $b \rightarrow c\ell\bar{\nu}$: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for right sign B^0 s.	167

A.6	GENERIC MC, $b \rightarrow c\bar{\nu}$: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for wrong sign B^0 s.	168
A.7	GENERIC MC, other: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for B^\pm s.	169
A.8	GENERIC MC, other: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for right sign B^0 s.	170
A.9	GENERIC MC, other: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for wrong sign B^0 s.	171
A.10	SIGNAL MC: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for B^\pm s.	172
A.11	SIGNAL MC: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for right sign B^0 s.	173
A.12	SIGNAL MC: The total m_{ES} distribution (top left) and the individual m_{ES} distributions in each bin of m_X for wrong sign B^0 s.	174
B.1	Fit results for $m_X < 1.55 \text{ GeV}/c^2$: The left plot is neutral B and the right plot is charged B .	175
B.2	Fit results for $m_X < 1.59 \text{ GeV}/c^2$: The left plot is neutral B and the right plot is charged B .	175
B.3	Fit results for $m_X < 1.67 \text{ GeV}/c^2$: The left plot is neutral B and the right plot is charged B .	176
B.4	Fit results for $m_X < 1.75 \text{ GeV}/c^2$: The left plot is neutral B and the right plot is charged B .	176
B.5	Fit results for $m_X < 1.83 \text{ GeV}/c^2$: The left plot is neutral B and the right plot is charged B .	176
B.6	Fit results for $m_X < 2.50 \text{ GeV}/c^2$: The left plot is neutral B and the right plot is charged B .	176
B.7	Fit results for $m_X < 1.55 \text{ GeV}/c^2$: The left plot is electrons and the right plot is muons.	177
B.8	Fit results for $m_X < 1.59 \text{ GeV}/c^2$: The left plot is electrons and the right plot is muons.	177

B.9	Fit results for $m_X < 1.67 \text{ GeV}/c^2$: The left plot is electrons and the right plot is muons.	177
B.10	Fit results for $m_X < 1.75 \text{ GeV}/c^2$: The left plot is electrons and the right plot is muons.	177
B.11	Fit results for $m_X < 1.83 \text{ GeV}/c^2$: The left plot is electrons and the right plot is muons.	177
B.12	Fit results for $m_X < 2.50 \text{ GeV}/c^2$: The left plot is electrons and the right plot is muons.	178

LIST OF TABLES

2.1	The six quarks and six leptons and their properties are presented in this table. They are split into three generations. u is the “up” quark, d is the “down” quark, c is the “charm” quark, s is the “strange” quark, t is the “top” quark, b is the “bottom” quark, e is the electron, μ is the muon, τ is the tau lepton, and ν_X are the corresponding neutrinos.	6
2.2	The fundamental forces of nature and their associated force-carrying gauge bosons, spins, masses, effective range, sources and couplings.	7
3.1	The PEP-II beam design parameters and the corresponding achieved values as of July 2004. σ_{Lx} , σ_{Ly} , σ_{Lz} refer to the size of the interaction region of the collision.	30
3.2	The cross-sections for the production of various states at the $T(4S)$ mass $\sqrt{s} = M(T(4S))$	30
3.3	Overview of the coverage, segmentation, and performance of the BABAR detector systems. The notation (C), (F), and (B) refers to the central barrel, forward and backward components of the system, respectively. The detector coverage in the laboratory frame is specified in terms of the polar angles θ_1 (forward) and θ_2 (backward). The number of readout channels is listed. Performance numbers are quoted for 1 GeV/c particles, except where noted. The performances for the SVT and DCH are quoted for a combined Kalman fit (for the definition of the track parameters, see section 3.3)	36
4.1	Summary of track selection criteria.	60
4.2	Summary of selection criteria for the D^0 selection	70
4.3	Summary of selection criteria for the D^+ selection	72
4.4	Summary of the selection criteria for the D^{*+} selection	74
4.5	Summary of selection criteria for the D^{*0} selection	74

5.1	Some of the inclusive and exclusive branching fractions relevant to the semi-exclusive reconstruction [6]. The reason that the branching fractions sum to a value greater than unity is that for inclusive $B \rightarrow D^{(*)}Y$ decays, there can be a significant amount of overlap as the Y can include $D^{(*)}$ mesons.	81
5.2	Summary of the number of semi-exclusive modes.	84
6.1	Monte Carlo event samples used in this analysis. Equivalent statistics in $ V_{ub} $ MC assumes $\mathcal{B}(b \rightarrow u\ell\nu) = 1.7 \times 10^{-3}$	91
6.2	Branching ratios used in the resonant and non-resonant $\bar{B} \rightarrow X_s \ell\bar{\nu}$ signal MC (before the hybrid reweighting). The hadron masses m_X are those used in the generator.	92
6.3	Branching fractions used in the $\bar{B} \rightarrow X_s/\ell$ hybrid signal MC after the reweighting.	95
6.4	Signal yield, S, and background, B, per charm seed mode of the B_{reco} candidate. The numbers are obtained from fits to the m_{gg} distributions.	105
6.5	Selection criteria for $b \rightarrow q\ell\nu$ and $b \rightarrow u\ell\nu$ events.	112
7.1	The χ^2/DOF results from the fits to the hadronic mass spectrum	123
7.2	Analysis validation and fit results for high-statistics cocktail MC. The generated value corresponds to $R_u \equiv \mathcal{B}(b \rightarrow u\ell\nu)/\mathcal{B}(b \rightarrow c\ell\nu) = 0.0116$.	127
7.3	The fraction of signal events below a cut-off in the hadronic mass spectrum. The error shown is statistical. This number is only used in the extraction of the total charmless branching fraction (Eq. 7.2) for the full rate measurement.	128
7.4	Summary of the fit parameters for data, on the full (top) and depleted (bottom) samples.	133
7.5	Summary of the fit parameters for data, on the neutral B (top) and charged B (bottom) samples.	134
7.6	Summary of the fit parameters for data, on the electron (top) and muon (bottom) samples.	135

7.7	Differential $b \rightarrow s\gamma$ branching fraction in bins of photon energy E_γ . Note that these numbers are normalized to the bin width and not to 100 MeV as in Fig. 7.8. The systematic error has been symmetrized.	136
7.8	The results for the determination of $ V_{cb} $ and summary of the relative uncertainties. The first part of the table shows the results obtained in the framework of Leibovich, Low, and Rothstein and the right-most column provides the results based on the full rate.	137
8.1	Smearing factor in different neutral energy bins.	143
8.2	The current best measurements for the branching fractions for $B \rightarrow X_c \ell \nu$ decays and values used in MC simulation. The non resonant $B \rightarrow D \ell \nu X$ is obtained by difference of the inclusive rate and the other 4 components.	144
8.3	D^+ branching fractions, current best measurements and values used in the MC.	146
8.4	D^0 branching fractions, current best measurements and values used in the MC.	147
8.5	$\delta\Gamma(c) \times 10^3$ results and errors for various m_X cuts for the signal modeling studies in sections 8.1.14–8.1.17. The error is determined from the largest deviations to the nominal value.	148
8.6	Summary of errors on the partial $b \rightarrow u\ell\bar{\nu}$ rate $\delta\Gamma(c)$. The errors are listed in units of % of $\delta\Gamma(c)$.	154

Acknowledgments

The work in this thesis would not be possible if not for the instruction and support I have received from my colleagues along the way. There are many people to thank, and I will attempt to list them below. I apologize to anyone who I may have forgotten.

Rolf-Dieter Heinz and Urs Langenegger. Thanks for working with me on this analysis. It was a pleasure working with both of you.

Adam Leibovich, Ian Low, and Ira Rothstein. Thanks for consistently and patiently answering my questions.

The members of the Semileptonic AWG, in particular Daniele Del Re, Riccardo Faccini, Virginia Azzolini, Concezio Bozzi, Kerstin Tackmann, and Dominique Fortin. The work in this thesis has benefited greatly from your work.

The members of *BABAR* who devoted time to reviewing this analysis and helping prepare it for publication.

The UC San Diego *BABAR* group of past and present. In particular, I would like to thank my advisor, David MacFarlane, for his support.

Lastly, I would like to offer a special thank you to Urs Langenegger. This analysis would not have been possible without you. Thanks for being limitlessly patient and always finding time to speak with me from the other side of the Atlantic, regardless of the time. You've been a true mentor and friend.