



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

UCRL-TR-217981

# Large-Scale First-Principles Molecular Dynamics Simulations on the BlueGene/L Platform using the Qbox Code

F. Gygi, E. Draeger, B. de Supinski, R. K. Yates, F. Franchetti, S. Kral, J. Lorenz, C. Ueberhueber, J. A. Gunnels, J. Sexton

January 5, 2006

## Disclaimer

---

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

# Large-Scale First-Principles Molecular Dynamics Simulations on the BlueGene/L Platform using the Qbox Code

## 2005 Gordon Bell Award - Finalist

*François Gygi, Erik W. Draeger, Bronis de Supinski, and Robert K. Yates (CASC), Franz Franchetti (Carnegie Mellon University), Stefan Kral, Jeorgen Lorenz and Christoph Ueberhueber (Vienna University of Technology), John A. Gunnels and James C. Sexton (IBM)*

First-Principles Molecular Dynamics (FPMD) is an accurate, atomistic simulation approach that is routinely applied to a variety of areas including solid-state physics, chemistry, biochemistry and nanotechnology. FPMD enables one to perform predictive materials simulations, as no empirical or adjustable parameters are used to describe a given system. Instead, a quantum mechanical description of electrons is obtained by solving the Kohn–Sham equations within a pseudopotential plane-wave formalism. This rigorous first-principles treatment of electronic structure is computationally expensive and limits the size of tractable systems to a few hundred atoms on most currently available parallel computers. Developed specifically for large parallel systems at LLNL's Center for Applied Scientific Computing, the Qbox implementation of the FPMD method shows unprecedented performance and scaling on BlueGene/L.



*Team members (L-R) Kim Yates (CASC), Christoph Ueberhueber (Vienna University of Technology), Franz Franchetti (Carnegie Mellon University), Erik Draeger, Francois Gygi, and Bronis de Supinski (CASC), as well as Bill Gropp (ANL).*

Predicting the properties of metals under extreme conditions of temperature and pressure has been a longstanding goal of materials science and high-energy density physics. Under extreme conditions, some important properties of metals require simulations with a significantly larger number of atoms than have previously been feasible, for example the calculation of melting temperatures using the two-phase simulation technique and the calculation of defect energies and defect migration processes. Qbox simulations of 1000 molybdenum atoms (12,000 electrons) showed that a strong scaling parallel efficiency of 86% can be obtained between 1024 and 32,768 processors. Using 65,536 processors, we observed a sustained performance of up to 64 TFlops. This system is considerably larger than any previously feasible FPMD simulation and represents the next generation of predictive materials simulations.