

Counting trees in graphs

Dhruv Mubayi*

Jacques Verstraëte†

Submitted: Apr 19, 2016; Accepted: Aug 11, 2016; Published: Aug 19, 2016

Mathematics Subject Classifications: 05C

Abstract

Erdős and Simonovits proved that the number of paths of length t in an n -vertex graph of average degree d is at least $(1 - \delta)nd(d - 1) \cdots (d - t + 1)$, where $\delta = (\log d)^{-1/2+o(1)}$ as $d \rightarrow \infty$. In this paper, we strengthen and generalize this result as follows. Let T be a tree with t edges. We prove that for any n -vertex graph G of average degree $d \geq t$, the number of labelled copies of T in G is at least

$$(1 - \varepsilon)nd(d - 1) \cdots (d - t + 1)$$

where $\varepsilon = (4t)^5/d^2$. This bound is tight except for the term $1 - \varepsilon$, as shown by a disjoint union of cliques. Our proof is obtained by first showing a lower bound that is a convex function of the degree sequence of G , and this answers a question of Dellamonica et. al.

1 Introduction

If H and G are graphs, let $\mathbf{N}_H(G)$ denote the number of labelled copies of H in G . More precisely, $\mathbf{N}_H(G)$ is the number of injections $\phi : V(H) \rightarrow V(G)$ such that $\phi(u)\phi(v) \in E(G)$ for every edge uv of H . Let $N_G(v)$ denote the neighborhood of vertex v in a graph G and let $d_G(v) = |N_G(v)|$. Let $(a)_b = a(a - 1)(a - 2) \cdots (a - b + 1)$ when $a \geq b \geq 1$ and a is real and b is an integer, and let $(a)_0 = 1$. Let T be a t -edge tree and $G = (V, E)$ be a graph with minimum degree at least t . In this paper, we obtain lower bounds on $\mathbf{N}_T(G)$. This is a basic question in combinatorics, for example, the simple lower bound $\sum_{v \in V} (d(v))_t$ in the case when T is a star is the main inequality needed for a variety of fundamental problems in extremal graph theory. Counting paths and walks in graphs has numerous applications, such as finding bounds on the spectral radius of the graph and the energy of a graph (see Täubig and Weihmann [6] and the references therein), as well as

*Department of Mathematics, Statistics, and Computer Science, University of Illinois, Chicago, IL, 60607 USA. Research partially supported by NSF grant DMS-1300138. Email: mubayi@uic.edu

†Department of Mathematics, UCSD. E-mail: jacques@ucsd.edu

in standard graph theoretic reductions of combinatorial problems such as counting words over alphabets with forbidden patterns.

Blakley and Roy [2] used elementary linear algebraic methods to show that if A is any symmetric pointwise non-negative matrix and v is any pointwise non-negative unit vector, then for any $t \in \mathbb{N}$, $\langle A^t v, v \rangle \geq \langle Av, v \rangle^t$. If t is even, this is a simple consequence of the fact that A is diagonalizable and Jensen's Inequality applied to a convex combination of the t -th powers of the eigenvalues of A . When A is the adjacency matrix of an n -vertex graph of average degree d , this shows that the number of walks of length t is at least nd^t . There are now a number of proofs of this result, for instance another approach to counting walks was used by Sidorenko, using an analytic method and the tensor power trick [5], as well as generalizations to various other inequalities involving walks (see [6] and the references therein).

Alon, Hoory and Linial [1] showed that the number of non-backtracking walks in an n -vertex graph G of average degree d and minimum degree at least two is at least $nd(d-1)^{t-1}$, which is again tight for d -regular graphs. A key fact in their proof is that if one considers a random walk of length t on the graph, starting with a uniformly randomly chosen edge, then the distribution of every edge of the walk is also uniform. The method of Alon, Hoory and Linial [1] was generalized in Dellamonica et. al. [3] to show that the number of homomorphisms of a t -edge tree T in G is at least $nd \prod_{v \in V} d(v)^{(t-1)d(v)/nd}$. By Jensen's Inequality, this is at least nd^t when G has no isolated vertices.

In Problem 4 of [3], the question of finding similar lower bounds for $\mathbf{N}_T(G)$ when T is a tree is raised. In the case T is a path with three edges, it was shown in [3] that $\mathbf{N}_T(G) \geq nd \prod_{v \in V} (d-2)^{2d(v)/dn}$ when G has minimum degree at least three. If P is a t -edge path, and G is a d -regular graph, then clearly $\mathbf{N}_P(G) \geq n(d)_t$ with equality for $d \geq t$ if and only if every component of G is a clique. Erdős and Simonovits [4] showed that if G is any graph of average degree d , then $\mathbf{N}_P(G) \geq n(d - o(d))_t$ as $d \rightarrow \infty$. In this paper, we answer Problem 4 of [3], extend the result in [4] to arbitrary trees and strengthen their lower bound to $n(d - O(1/d))_t$ as $d \rightarrow \infty$:

Theorem 1. *If T is any t -edge tree and G is an n -vertex graph of average degree $d \geq t$, and $\varepsilon_d = (4t)^5/d^2$, then*

$$\mathbf{N}_T(G) \geq (1 - \varepsilon_d)n(d)_t.$$

This result is best possible up to the factor $1 - \varepsilon_d$ in view of the graph comprising $n/(d+1)$ cliques of order $d+1 \geq t+1$ when $d+1$ is a positive integer dividing n , since these graphs have exactly $n(d)_t$ copies of T . Note that since $\varepsilon = O(1/d^2)$, Theorem 1 implies $\mathbf{N}_T(G) \geq n(d - O(1/d))_t$ as $d \rightarrow \infty$.

Since we do not have an example of a graph G with n vertices and average degree d for which $\mathbf{N}_T(G) < n(d)_t$, we propose the following conjecture:

Conjecture 2. For every $t > 1$, there exists $d_0 = d_0(t)$ such that for any tree T with t edges and any n -vertex graph G with average degree $d \geq d_0$, $\mathbf{N}_T(G) \geq n(d)_t$.

Once more the example of disjoint complete graphs of order $d+1$ shows that this conjecture if true is best possible. Perhaps it is even true that if d is an integer, then this is the only example where equality holds.

This paper is organized as follows. In the next section, we determine a lower bound on $N_T(G)$ for a t -edge tree T , as a function of the degree sequence of G . This answers Problem 4 of [3]. The proof of this result extends the method of [1] by carefully controlling the distribution of the edges in a random embedding of a tree in a graph of average degree d . The function of degrees is “almost” convex, and in Section 3 we give a suitable approximate form of Jensen’s Inequality, which is used in Section 4 to derive Theorem 1.

2 Degree sequences and counting trees

As a stepping stone to Theorem 1, we are going to prove the following theorem, which gives a lower bound for $N_T(G)$ in terms of the degree sequence of G , and addresses Problem 4 in [3]:

Theorem 3. *Let T be a t -edge tree, and let $G = (V, E)$ be an n -vertex graph with average degree d and minimum degree $k > t > 1$. Then*

$$N_T(G) \geq nd \prod_{v \in V} \prod_{i=3}^{t+1} (d(v) - i + 2)^{\frac{d(v)-i+2}{(d-i+2)n} \theta_i(v)} \quad (1)$$

where for $3 \leq i \leq t+1$,

$$\frac{1}{n} \sum_{v \in V} (d(v) - i + 2) \theta_i(v) = d - i + 2 \quad (2)$$

and for $3 \leq i \leq t+1$,

$$\left(\frac{k-t}{k}\right)^t \leq \theta_i(v) \leq \left(\frac{k}{k-t}\right)^t \quad (3)$$

Proof. Let Ω be the set of all labelled copies of T in the graph G . Fix a breadth-first search ordering $\vec{x} = (x_1, x_2, \dots, x_{t+1})$ of the vertices of T , starting with a leaf x_1 of T . For $2 \leq i \leq t+1$, let $a(i)$ be the unique $h < i$ such that x_h is the parent of x_i in the ordering \vec{x} . Note that $a(2) = 1$ and $a(3) = 2$. Let $v_1 v_2$ be a randomly and uniformly chosen (oriented) edge of G and define $\phi(x_1) = v_1$ and $\phi(x_2) = v_2$. Having defined $\phi(x_j) = v_j$ for all $j < i$, let x_i be mapped to a uniformly chosen vertex of

$$N_+(v_{a(i)}) := N(v_{a(i)}) \setminus \{v_1, v_2, \dots, v_{i-1}\}.$$

One can always define ϕ in this manner since $d(v_{a(i)}) \geq k \geq t+1 \geq i$ and therefore

$$\begin{aligned} |N_+(v_{a(i)})| &\geq |N(v_{a(i)})| - |\{v_1, v_2, \dots, v_{i-1}\} \cap N(v_{a(i)})| \\ &\geq d(v_{a(i)}) - i + 2 > 0. \end{aligned}$$

Here we noted that $v_{a(i)} \in \{v_1, v_2, \dots, v_{i-1}\}$. Then ϕ gives a probability measure P on the sample space Ω . Let \vec{v} denote a vector $(v_1, v_2, \dots, v_{t+1})$ of vertices of G . We write $\phi(\vec{x}) = \vec{v}$ to denote the event that $\phi(x_i) = v_i$ for all $i \in [t+1]$. Then

$$P(\phi(\vec{x}) = \vec{v}) = \frac{1}{nd} \prod_{i=3}^{t+1} \frac{1}{|N_+(v_{a(i)})|}. \quad (4)$$

By the inequality of arithmetic and geometric means:

$$\begin{aligned} N_T(G) &= \sum_{\vec{v} \in \Omega} P(\phi(\vec{x}) = \vec{v}) \cdot \frac{1}{P(\phi(\vec{x}) = \vec{v})} \\ &\geq \prod_{\vec{v} \in \Omega} P(\phi(\vec{x}) = \vec{v})^{-P(\phi(\vec{x}) = \vec{v})} \\ &= \prod_{\vec{v} \in \Omega} (nd)^{P(\phi(\vec{x}) = \vec{v})} \cdot \prod_{\vec{v} \in \Omega} \prod_{i=3}^{t+1} |N_+(v_{a(i)})|^{P(\phi(\vec{x}) = \vec{v})} \\ &= nd \prod_{\vec{v} \in \Omega} \prod_{i=3}^{t+1} |N_+(v_{a(i)})|^{P(\phi(\vec{x}) = \vec{v})}. \end{aligned}$$

Let $\Omega_{iv} = \{\vec{v} \in \Omega : v_{a(i)} = v\}$. Since $|N_+(v_{a(i)})| \geq d(v_{a(i)}) - i + 2$ for $3 \leq i \leq t+1$,

$$\begin{aligned} N_T(G) &\geq nd \cdot \prod_{\vec{v} \in \Omega} \prod_{i=3}^{t+1} |N_+(v_{a(i)})|^{P(\phi(\vec{x}) = \vec{v})} \\ &= nd \cdot \prod_{i=3}^{t+1} \prod_{v \in V} \prod_{\vec{v} \in \Omega_{iv}} |N_+(v_{a(i)})|^{P(\phi(\vec{x}) = \vec{v})} \\ &\geq nd \cdot \prod_{i=3}^{t+1} \prod_{v \in V} \prod_{\vec{v} \in \Omega_{iv}} (d_G(v) - i + 2)^{P(\phi(\vec{x}) = \vec{v})} \\ &= nd \cdot \prod_{i=3}^{t+1} \prod_{v \in V} (d_G(v) - i + 2)^{\sum_{\vec{v} \in \Omega_{iv}} P(\phi(\vec{x}) = \vec{v})} \\ &= nd \cdot \prod_{i=3}^{t+1} \prod_{v \in V} (d_G(v) - i + 2)^{P(\phi(x_{a(i)}) = v)}. \end{aligned} \quad (5)$$

In the next two claims, we fix $v \in V$. For $i \in [t+1]$, define

$$g_i(v) = P(\phi(x_i) = v).$$

We observe that $g_1(v) = g_2(v) = d_G(v)/dn$. For $3 \leq i \leq t+1$, we determine upper and lower bounds on $g_i(v)$ relative to $d_G(v)/dn$. We write $d(v)$ instead of $d_G(v)$ in what follows.

To bound the quantities $g_i(v)$, let T_i be the subtree of T consisting of the vertices x_1, x_2, \dots, x_i in order. Now consider a breadth-first search ordering y_1, y_2, \dots, y_i of T_i such that $y_1 = x_i$ and $y_i = x_1$. For $2 \leq j \leq i$, let $b(j)$ be the unique $h < j$ such that y_h is the parent of y_j in the ordering y_1, y_2, \dots, y_i and let

$$N_+(u_{b(j)}) = N_G(u_{b(j)}) \setminus \{u_1, u_2, \dots, u_{j-1}\}.$$

As an example, if T_i is a path, then $b(j) = j - 1$ and $N_+(u_j) = N_G(u_j) \setminus \{u_1, u_2, \dots, u_{j-1}\}$.

Claim 1. Let $3 \leq i \leq t + 1$ and let $v = u_{b(2)} = u_1$. Then

$$\frac{1}{nd} \sum_{u_2 \in N_+(u_{b(2)})} \cdots \sum_{u_i \in N_+(u_{b(i)})} \prod_{j=3}^i \frac{1}{d(u_{b(j)})} \leq g_i(v) \leq \frac{1}{nd} \sum_{u_2 \in N_+(u_{b(2)})} \cdots \sum_{u_i \in N_+(u_{b(i)})} \prod_{j=3}^i \frac{1}{d(u_{b(j)}) - t} \quad (6)$$

Proof of Claim 1. The number of choices of an embedding of T_i with x_i mapping to v is

$$\sum_{u_2 \in N_+(u_{b(2)})} \sum_{u_3 \in N_+(u_{b(3)})} \cdots \sum_{u_i \in N_+(u_{b(i)})} 1. \quad (7)$$

Here we are counting the images of T_i according to the breadth-first search ordering y_1, y_2, \dots, y_i of T_i with $y_1 = x_i$ and $y_i = x_1$, so that the image of y_j is u_j for $j \in [i]$. At the stage where y_j is to be embedded, we have to select a vertex $u_j \in N_G(u_{b(j)})$ such that u_j is not equal to any of $v = u_1, u_2, \dots, u_{j-1}$, which have already been embedded. By definition of $N_+(u_{b(j)})$, this is equivalent to $u_j \in N_+(u_{b(j)})$.

Suppose we fix such an embedding with vertices u_1, u_2, \dots, u_i . Let v_1, v_2, \dots, v_i be the re-ordering of u_1, u_2, \dots, u_i such that x_j is mapped to v_j for $j \in [i]$. This fixed embedding has probability

$$P(\phi(\vec{x}) = \vec{v}) = \frac{1}{nd} \prod_{j=3}^i \frac{1}{|N_+(v_{a(j)})|}. \quad (8)$$

Since $d(v_{a(j)}) - t \leq |N_+(v_{a(j)})| \leq d(v_{a(j)})$,

$$\frac{1}{nd} \prod_{j=3}^i \frac{1}{d(v_{a(j)})} \leq P(\phi(\vec{x}) = \vec{v}) \leq \frac{1}{nd} \prod_{j=3}^i \frac{1}{d(v_{a(j)}) - t}. \quad (9)$$

We now prove $\{v_{a(j)} : 3 \leq j \leq i\}$ and $\{u_{b(j)} : 3 \leq j \leq i\}$ are equal as multisets. For example, if T_i is a path, then these multisets are sets, and $\{v_{a(j)} : 3 \leq j \leq i\} = \{v_2, v_3, \dots, v_{i-1}\} = \{u_2, u_3, \dots, u_{i-1}\}$ since $u_j = v_{i-j+1}$ for $j \in [i]$. For $2 \leq \ell \leq i - 1$, the number of times u_ℓ appears in the multiset $\{u_{b(j)} : 3 \leq j \leq i\}$ is the number of j with $3 \leq j \leq i$ such that $b(j) = \ell$. In other words, it is the number of times u_ℓ is a parent of some vertex u_j in the ordering u_1, \dots, u_i . This is precisely $d_T(y_\ell) - 1$. Similarly, the number of times $u_\ell \in \{v_2, v_3, \dots, v_{i-1}\}$ appears in the multiset $\{v_{a(j)} : 3 \leq j \leq i\}$ is the

number of times u_ℓ is a parent of some vertex v_j in the ordering v_1, \dots, v_i , and this is again $d_T(y_\ell) - 1$. We conclude $\{v_{a(j)} : 3 \leq j \leq i\}$ and $\{u_{b(j)} : 3 \leq j \leq i\}$ are equal as multisets. Therefore by (9),

$$\frac{1}{nd} \prod_{j=3}^i \frac{1}{d(u_{b(j)})} \leq \mathbf{P}(\phi(\vec{x}) = \vec{v}) \leq \frac{1}{nd} \prod_{j=3}^i \frac{1}{d(u_{b(j)}) - t}.$$

Replacing the summand 1 in (7) with $\mathbf{P}(\phi(\vec{x}) = \vec{v})$, we obtain Claim 1. \square

For each $v \in V$ and $3 \leq i \leq t+1$, define $\theta_i(v)$ by

$$g_{a(i)}(v) = \frac{d(v) - i + 2}{(d - i + 2)n} \theta_i(v). \quad (10)$$

Note that since $\sum_{v \in V} g_\ell(v) = \sum_{v \in V} \mathbf{P}(\phi(x_\ell) = v) = 1$ for all $\ell \leq t+1$, we have

$$\frac{1}{n} \sum_{v \in V} (d(v) - i + 2) \theta_i(v) = d - i + 2$$

and this verifies (2). The goal of the next claim is to verify (3).

Claim 2. For $3 \leq i \leq t+1$,

$$\left(\frac{k-t}{k}\right)^t \leq \theta_i(v) \leq \left(\frac{k}{k-t}\right)^t.$$

Proof of Claim 2. We only prove the upper bound, since the lower bound is similar. For convenience, for $3 \leq \ell \leq i$, and $v = u_1 = u_{b(2)}$, define

$$f(\ell) = \frac{1}{nd} \sum_{u_2 \in N_+(u_{b(2)})} \sum_{u_3 \in N_+(u_{b(3)})} \cdots \sum_{u_\ell \in N_+(u_{b(\ell)})} \prod_{j=3}^{\ell} \frac{1}{d(u_{b(j)}) - t}.$$

Then $g_i(v) \leq f(i)$ by Claim 1. Since $|N_+(u_{b(i)})| \leq d(u_{b(i)})$ and $d(u_{b(i)}) \geq k$,

$$\sum_{u_i \in N_+(u_{b(i)})} \frac{1}{d(u_{b(i)}) - t} = \frac{|N_+(u_{b(i)})|}{d(u_{b(i)}) - t} \leq \frac{d(u_{b(i)})}{d(u_{b(i)}) - t} \leq \frac{k}{k-t}.$$

This shows $f(i) \leq \frac{k}{k-t} f(i-1)$. Continuing in this way we obtain

$$\begin{aligned} g_i(v) \leq f(i) &\leq \left(\frac{k}{k-t}\right)^{i-2} f(2) \\ &= \frac{1}{nd} \left(\frac{k}{k-t}\right)^{i-2} \sum_{u_2 \in N(v)} 1 \\ &= \frac{d(v)}{nd} \left(\frac{k}{k-t}\right)^{i-2} \\ &\leq \frac{d(v)}{nd} \left(\frac{k}{k-t}\right)^{t-2}. \end{aligned}$$

For any j such that $3 \leq j \leq i \leq t+1$, $d(v)/(d(v)-j+2) \leq k/(k-j+2) \leq k/(k-t)$ since $d(v) \geq k \geq t+1 \geq j$. Therefore

$$\frac{d(v)}{nd} \leq \frac{d(v)-j+2}{(d-j+2)n} \cdot \frac{d(v)}{d(v)-j+2} \leq \frac{d(v)-j+2}{(d-j+2)n} \cdot \frac{k}{k-t}.$$

Inserting this in the preceding upper bound for $g_i(v)$, we find for any j with $3 \leq j \leq i \leq t+1$,

$$g_i(v) \leq \frac{d(v)-j+2}{(d-j+2)n} \cdot \frac{k}{k-t} \cdot \left(\frac{k}{k-t}\right)^{t-2} \leq \frac{d(v)-j+2}{(d-j+2)n} \cdot \left(\frac{k}{k-t}\right)^t.$$

This also holds for $i=2$ as

$$g_2(v) = \frac{d(v)}{dn} \leq \frac{d(v)-j+2}{(d-j+2)n} \cdot \frac{d(v)}{d(v)-j+2} \leq \frac{d(v)-j+2}{(d-j+2)n} \left(\frac{k}{k-t}\right).$$

Replacing i with $a(i) \geq 2$ and selecting $j=i$,

$$g_{a(i)}(v) \leq \frac{d(v)-i+2}{(d-i+2)n} \left(\frac{k}{k-t}\right)^t.$$

By the definition (10), this gives $\theta_i(v) \leq (k/(k-t))^t$. The lower bound is similar: we use the lower bound on $g_i(v)$ from Claim 1, together with the observation $|N_+(u_{b(i)})| \geq d(u_{b(i)})-t$ and $d(u_{b(i)}) \geq k$ to obtain

$$g_i(v) \geq \frac{d(v)}{nd} \left(\frac{k-t}{k}\right)^{t-2}.$$

Then we use

$$\frac{d(v)}{dn} \geq \frac{d(v)-j+2}{(d-j+2)n} \cdot \frac{d-j+2}{d} \geq \frac{d(v)-j+2}{(d-j+2)n} \cdot \frac{k-t}{k}$$

for $3 \leq j \leq t+1$, and as above this gives the required lower bound on $\theta_i(v)$. \square

Now we finish the proof of Theorem 3: (5) becomes

$$N_T(G) \geq nd \prod_{v \in V} \prod_{i=3}^{t+1} (d(v)-i+2)^{g_{a(i)}(v)} = nd \prod_{v \in V} \prod_{i=3}^{t+1} (d(v)-i+2)^{\frac{d(v)-i+2}{(d-i+2)n} \theta_i(v)}$$

where $\theta_i(v)$ satisfies (2) and (3), by Claims 1 and 2. This proves (1). \square

3 An Approximate Jensen's Inequality

An approximate form of Jensen's Inequality is used to prove Theorem 1 from Theorem 3:

Lemma 4. Let $\delta, x_i, y_i \in [0, \infty)$ for all $i \in [n]$, and suppose $\frac{1}{n} \sum_{i=1}^n x_i = \frac{1}{n} \sum_{i=1}^n y_i = \gamma$ and $|x_i/y_i - 1| \leq \delta$ for all $i \in [n]$. Then

$$\prod_{i=1}^n x_i^{\frac{y_i}{\gamma^n}} \geq (1 - \delta)e^\delta \gamma.$$

Proof. The lemma is clear if $\delta \geq 1$ so we assume $\delta \in [0, 1)$. This implies $x_i, y_i \in (0, \infty)$. Let $f(y) = y \log y$ and $\frac{x_i}{y_i} = c_i$. Since f is convex on $(0, \infty)$, Jensen's Inequality gives:

$$\frac{1}{n} \sum_{i=1}^n f(y_i) \geq f\left(\frac{1}{n} \sum_{i=1}^n y_i\right) = \gamma \log \gamma.$$

Furthermore, $y_i \log x_i = f(y_i) + y_i \log c_i$. Therefore

$$\begin{aligned} \log \prod_{i=1}^n x_i^{\frac{y_i}{\gamma^n}} &= \frac{1}{\gamma^n} \sum_{i=1}^n y_i \log x_i \\ &= \frac{1}{\gamma} \cdot \frac{1}{n} \sum_{i=1}^n (f(y_i) + y_i \log c_i) \\ &= \frac{1}{\gamma} \left(\frac{1}{n} \sum_{i=1}^n f(y_i) \right) + \frac{1}{\gamma n} \sum_{i=1}^n y_i \log c_i \\ &\geq \log \gamma + \frac{1}{\gamma n} \sum_{i=1}^n y_i \log c_i \\ &= \log \gamma + \frac{1}{\gamma n} \sum_{i=1}^n y_i (c_i - 1) + \frac{1}{\gamma n} \sum_{i=1}^n y_i (\log c_i - c_i + 1). \end{aligned}$$

Since $\sum c_i y_i = \sum x_i = \gamma n$, $\sum y_i (c_i - 1) = 0$. The function $h(x) = \log x - x$ on the interval $[1 - \delta, 1 + \delta]$ has one critical point, a maximum at $x = 1$, and therefore its minimum in this interval occurs at one of its endpoints. We claim that $h(1 - \delta) < h(1 + \delta)$. Indeed, this is equivalent to $e^{2\delta} < (1 + \delta)/(1 - \delta) = 1 + \sum_{j=1}^{\infty} 2\delta^j$. Since $e^{2\delta} = 1 + \sum_{j=1}^{\infty} (2\delta)^j/j!$ and $(2\delta)^j/j! \leq 2\delta^j$ for all $j \geq 1$ with strict inequality for $j > 2$, this proves that $h(1 - \delta) < h(1 + \delta)$ for $\delta \in [0, 1)$. Therefore $\log c_i - c_i + 1 \geq \log(1 - \delta) + \delta$ for all $i \in [n]$, which gives

$$\log \prod_{i=1}^n x_i^{\frac{y_i}{\gamma^n}} \geq \log \gamma + \log(1 - \delta) + \delta.$$

Exponentiating gives the lemma. □

Remarks. As $\delta \rightarrow 0$, $(1 - \delta)e^\delta = 1 - \frac{1}{2}\delta^2 + O(\delta^3)$. If n is even, $y_i = \gamma$ for all $i \leq n$ and $x_i = (1 + \delta)\gamma$ for $i \leq \frac{1}{2}n$ and $x_i = (1 - \delta)\gamma$ for $i > \frac{1}{2}n$, then as $\delta \rightarrow 0$,

$$\prod_{i=1}^n x_i^{\frac{y_i}{\gamma^n}} = (1 - \delta^2)^{\frac{1}{2}} \gamma = 1 - \frac{1}{2}\delta^2 + O(\delta^3).$$

This shows the bound in Lemma 4 is asymptotically tight as $\delta \rightarrow 0$. Note that if $\delta = 0$, then $\prod_{i=1}^n x_i^{y_i/\gamma^n} \geq \gamma$ directly from Jensen's Inequality.

4 Proof of Theorem 1

We prove Theorem 1 by induction on n . Let G be an n -vertex graph with average degree $d \geq t$. Theorem 1 states $\mathbf{N}_T(G) \geq (1 - \varepsilon_d)n(d)_t$. If $t \leq d < 16t^2$, then $1 - \varepsilon_d \leq 0$ and so Theorem 1 is true in this case. This also proves the theorem when $n < 16t^2$. Suppose $d \geq 16t^2$. If k is the minimum degree of G , we consider the two cases $k < d/8$ and $k \geq d/8$ separately.

Case 1. $k < d/8$. In this case, remove from G a vertex of degree k to get a graph H with $n - 1$ vertices and average degree at least

$$\frac{2}{n-1} \left(\frac{dn}{2} - \frac{d}{8} \right) > d + \frac{d}{n-1} - \frac{d}{4(n-1)} > d + \frac{3d}{4n} := d'.$$

By definition of d' ,

$$\begin{aligned} (n-1)d'(d'-1) &= (n-1)\left(d + \frac{3d}{4n}\right)\left(d - 1 + \frac{3d}{4n}\right) \\ &= nd(d-1) + \frac{1}{4}d + \frac{1}{2}d^2 - \frac{9d^2}{16n^2} + \frac{3d}{4n} - \frac{15d^2}{16n} \\ &> nd(d-1) + \frac{1}{2}d^2 - \frac{1}{n}d^2 \\ &> nd(d-1). \end{aligned}$$

From the second to the third line, we used $3d/4n > 9d^2/16n^2$ since $d < n$, and $15d^2/16n < d^2/n$. Recall $\varepsilon_d = (4t)^5/d^2$. Note also $d' > d \geq t$, so by induction, and since $1 - \varepsilon_{d'} \geq 1 - \varepsilon_d$ for $d' \geq d$,

$$\begin{aligned} \mathbf{N}_T(H) &\geq (1 - \varepsilon_{d'}) \cdot (n-1)(d')_t \\ &\geq (1 - \varepsilon_d) \cdot (n-1)d'(d'-1) \cdot (d-2)_{t-2} \\ &> (1 - \varepsilon_d) \cdot nd(d-1) \cdot (d-2)_{t-2} \\ &= (1 - \varepsilon_d) \cdot n(d)_t. \end{aligned}$$

The proof is complete in this case.

Case 2. $k \geq d/8$. We apply Lemma 4 and Theorem 3. By Theorem 3, shifting the range of summation from $3 \leq i \leq t+1$ to $j \in [t-1]$,

$$\mathbf{N}_T(G) \geq nd \prod_{v \in V} \prod_{j=1}^{t-1} (d(v) - j)^{\frac{d(v)-j}{(d-j)n} \theta_{j+2}(v)}. \quad (11)$$

By (3), for all $j \in [t-1]$,

$$\left(\frac{k-t}{k} \right)^t \leq \frac{1}{\theta_{j+2}(v)} \leq \left(\frac{k}{k-t} \right)^t.$$

Since $(1-x)^t \geq 1-xt$ for $0 \leq x \leq 1$ and $(1+x)^t \leq 1+2xt$ for $0 \leq x \leq 1/t$,

$$1 - \frac{t^2}{k} \leq \frac{1}{\theta_{j+2}(v)} \leq 1 + \frac{2t^2}{k-t}.$$

Since $2t^2/(k-t) \leq 4t^2/k$ is easily true for $k \geq d/8 \geq 2t^2$, letting $\delta = 4t^2/k$ we obtain

$$\left| \frac{1}{\theta_{j+2}(v)} - 1 \right| \leq \delta$$

for all $j \in [t-1]$. Now we use Lemma 4. Fixing $j \in [t-1]$, let $x_{v,j} = d(v) - j$ and let $y_{v,j} = (d(v) - j)\theta_{j+2}(v)$ for $v \in V$. Then

$$\left| \frac{x_{v,j}}{y_{v,j}} - 1 \right| = \left| \frac{1}{\theta_{j+2}(v)} - 1 \right| \leq \delta.$$

Furthermore, from (2) in Theorem 3,

$$\frac{1}{n} \sum_{v \in V} y_{v,j} = \frac{1}{n} \sum_{v \in V} (d(v) - j)\theta_{j+2}(v) = d - j = \frac{1}{n} \sum_{v \in V} (d(v) - j) = \frac{1}{n} \sum_{v \in V} x_{v,j}.$$

Also $x_{v,j} \geq d(v) - j \geq k - t \geq 0$ and $y_{v,j} = \theta_{j+2}(v)x_{v,j} \geq 0$. Therefore all the conditions of Lemma 4 are satisfied, and so for each $j \in [t-1]$, Lemma 4 with $\gamma = d - j$ gives

$$\prod_{v \in V} x_{v,j}^{\frac{y_{v,j}}{(d-j)n}} \geq (1 - \delta)e^\delta(d - j).$$

Applying this for each $j \in [t-1]$ to (11) gives

$$\mathbf{N}_T(G) \geq nd \prod_{j=1}^{t-1} (1 - \delta)e^\delta(d - j) = n(d)_t \cdot (1 - \delta)^{t-1}e^{(t-1)\delta}.$$

Finally, using $(1 - \delta)e^\delta \geq 1 - \delta^2$ and $(1 - \delta^2)^{t-1} \geq 1 - t\delta^2$,

$$\begin{aligned} (1 - \delta)^{t-1}e^{(t-1)\delta} &\geq (1 - \delta^2)^{(t-1)} \\ &\geq 1 - t\delta^2 \\ &= 1 - \frac{16t^5}{k^2} \\ &\geq 1 - \varepsilon_d. \end{aligned}$$

From the third line to the fourth, we used $k \geq d/8$. This completes the proof. \square

5 Concluding remarks

• Theorem 1 can be improved if structural information on G is given. For instance, if G is an n -vertex triangle-free graph of average degree d , and T is the t -edge path, then one can adapt the proof of Theorem 1 to obtain

$$N_T(G) \geq (1 - O(\frac{t^5}{d^2}))nd \prod_{i=1}^{t-1} (d - \lceil \frac{i}{2} \rceil).$$

A disjoint union of complete bipartite graphs $K_{d,d}$ shows this is tight up to the factor $1 - O(\frac{t^5}{d^2})$. It is plausible as in Conjecture 1 that this construction has the fewest copies of T amongst all triangle-free graphs of average degree d , when d is large enough.

• Fix a tree T with t edges and degrees $1, d_1, d_2, \dots, d_t$ in a breadth-first labelling starting with a leaf. A *local isomorphism* of T in a graph G is a neighbourhood preserving embedding of T in G i.e. a map $\phi : V(T) \rightarrow V(G)$ such that if $\phi(x) = v$ for some $x \in V(T)$, then $\phi(y) \neq \phi(z)$ whenever $y, z \in N_T(x)$ are distinct. Using the method of Theorem 3, one can show that the number of local isomorphisms of T in G is at least

$$nd \prod_{v \in V} \prod_{i=1}^{t-1} (d(v) - 1)_{d_i-1}^{\frac{d(v)}{d_i}}.$$

If G has minimum degree say at least $2t$, and average degree d , it is possible to show that the above expression is at least

$$nd \prod_{i=1}^{t-1} (d - 1)_{d_i-1}.$$

If T is a path, then we recover the main result of [1] stating that the number of non-backtracking walks of t edges in G is at least $nd(d - 1)^{t-1}$. Furthermore, the bound is tight since equality is achieved for any d -regular n -vertex graph. Finally, if G has girth at least $\text{diam}(T)$, then every local isomorphism of T in G is an isomorphism, so we obtain the lower bound

$$N_T(G) \geq nd \prod_{i=1}^{t-1} (d - 1)_{d_i-1}.$$

Again, this is tight for any d -regular graph of sufficiently large girth.

Acknowledgments

We thank Eoin Long for pointing out an error in an earlier version of this paper, and the anonymous referees for helpful comments.

References

- [1] N. Alon, S. Hoory, and N. Linial, The Moore Bound for Irregular Graphs, *Graphs Combin.* 18 (2002), no. 1, 53–57.
- [2] G. Blakley and P. Roy, Hölder type inequality for symmetrical matrices with non-negative entries, *Proc. Amer. Math. Soc.* (1965) 16, 1244–1245.
- [3] D. Dellamonica Jr., P. Haxell, T. Łuczak, D. Mubayi, B. Nagle, Y. Person, V. Rödl, M. Schacht, Tree-minimal graphs are almost regular, *J. Comb.* 3 (2012), no. 1, 49–62.
- [4] P. Erdős and M. Simonovits, Compactness results in extremal graph theory, *Combinatorica* 2 (1982) no. 3, 275–288.
- [5] A. F. Sidorenko, Inequalities for functionals generated by bipartite graphs, *Diskretnaya Matematika* 3 (1991), 50–65 (in Russian), *Discrete Math. Appl.* 2 (1992), 489–504 (in English).
- [6] H. Täubig. J. Weihmann, Matrix power inequalities and the number of walks in graphs, *Discrete Appl. Math.* 176, 122–129.