

A better lower bound on the average degree of online k -list-critical graphs

Landon Rabern
IQVIA, U.S.A.

Submitted: Aug 24, 2016; Accepted: Feb 18, 2018; Published: Mar 2, 2018
Mathematics Subject Classifications: 05C15

Abstract

We improve the best known bounds on the average degree of online k -list-critical graphs for $k \geq 6$. Specifically, for $k \geq 7$ we show that every non-complete online k -list-critical graph has average degree at least $k - 1 + \frac{(k-3)^2(2k-3)}{k^4-2k^3-11k^2+28k-14}$ and every non-complete online 6-list-critical graph has average degree at least $5 + \frac{93}{766}$. The same bounds hold for offline k -list-critical graphs.

1 Introduction

Imagine a classroom with n children and one teacher. Each child wishes to go to lunch with a group of her friends, but the teacher is dreadfully strict and will only allow certain groups of students to leave for lunch at a time. The teacher starts by giving each child k lunch tokens. Then she selects a group of students S and the children must decide on a group of friends F from S to go to lunch. The group of friends F leave for lunch and the remaining students in S each give the teacher one of their lunch tokens. The teacher keeps picking groups of students and allowing the children to decide on a subgroup of friends to go to lunch like this until either all of the students have gone to lunch with a lunch token in hand; or some student has no lunch tokens left. The teacher wants to be strict, but fair, so she desires to find the smallest number k of tokens to give each student so that there is always a way for the children to play that ends up with all the children going to lunch with a token and a group of friends. To her chagrin, she cannot find an easy way to determine the smallest k for a given classroom of children; it seems she has constructed a nearly impossible problem for herself in her perverse authoritarian zeal. We do not really want to help her solve her problem so she can inflict this game upon the children, but it is an interesting problem nonetheless. So, let us think about the problem and just not tell her of any progress we make. First off, we can abstract things a little and consider a graph G with vertex set consisting of all the children such that vertices x and y (representing children) are adjacent just in case x and y are not friends. Then a group of friends is just an independent set in G . We can think of giving a number of

lunch tokens to each student as a function from $V(G)$ to \mathbb{N} where $f(v)$ is the number of tokens child v gets. For such a function f , say that G is *online f -list-colorable* just in case $V(G)$ is empty, or for every $S \subseteq V(G)$ there is an independent set $F \subseteq S$ such that $G - F$ is online f' -list-colorable where f' is the function from $V(G) \setminus F$ to \mathbb{N} given by $f'(v) = f(v) - 1$ for $v \in S \setminus F$ and $f'(v) = f(v)$ for $v \in V(G) \setminus S$. For $k \in \mathbb{N}$, we say that G is *online k -list-colorable* just in case G is online f -list-colorable where $f(v) = k$ for all $v \in V(G)$. The smallest k for which G is online k -list-colorable is the online list-chromatic number of G . The teacher's problem then becomes the following.

Problem. Given a graph G that represents a classroom of children and their friend relationships, how can we best find the online list-chromatic number of G ?

This problem is hard (supposing other problems we think are hard are in fact hard [6]), we will not solve it here. Surely, more friendships amongst the children should make it easier for the teacher to use fewer tokens per child. This intuition is most succinctly captured by the *average degree* $d(G)$ of the vertices in the non-friend graph G defined above. Specifically, a low average degree should make it easier for the teacher to use fewer tokens per child. We improve the best known quantitative bounds for this problem. A graph G is *online k -list-critical* if G is not online $(k - 1)$ -list-colorable, but every proper subgraph of G is. For further graph theory definitions and notation, we refer the reader to Diestel's textbook [3].

Main Theorem. *For $k \geq 7$, every non-complete online k -list-critical graph has average degree at least*

$$k - 1 + \frac{(k - 3)^2(2k - 3)}{k^4 - 2k^3 - 11k^2 + 28k - 14}.$$

Every non-complete online 6-list-critical graph has average degree at least $5 + \frac{93}{766}$.

The theorem also holds, with a nearly identical proof, for offline list coloring. The proof is similar to the 4-list-critical case in [12], but now we incorporate reducibility lemmas from Kierstead and Rabern [7]. Basically, we show that the average degree of the subgraph induced on vertices of degree $k - 1$ is small, which implies that the number of edges incident to the vertices of degree at least k must be large, and hence the number of vertices of degree at least k must be large; that is, the graph must have high average degree. A tight bound on the average degree of the subgraph induced on vertices of degree $k - 1$ in an online k -list-critical graph was proved by Gallai [5]. The connected graphs in which each block is a complete graph or an odd cycle are called *Gallai trees*. Gallai [5] proved that in a k -critical graph, the vertices of degree $k - 1$ induce a disjoint union of Gallai trees. The same is true for online k -list-critical graphs [1, 4, 13]. Since Gallai's bound is tight, it may appear that there is no hope of improvement using the above method. While it is true that the upper bound on average degree of Gallai trees cannot be improved in general, it can be improved in the absence of certain *bad* properties. Let G be an online k -list-critical graph and let \mathcal{L} be the subgraph of G induced on vertices of degree $k - 1$. If the presence of bad properties in \mathcal{L} could be shown to lead to reducible configurations in G , we would have a pathway to improvement. Kostochka and Stiebitz

[9] made the first progress along these lines. Further improvements in [7], [2] and [12] follow the same general outline. As in [2] and [12], it is convenient to have a measure of how bad \mathcal{L} is. So, if b is a function measuring badness, this could be realized as an upper bound of the form:

$$2 \|\mathcal{L}\| \leq s(k) |\mathcal{L}| + t(k)b(\mathcal{L}).$$

Of course, we can measure badness along multiple axes. In our proof we use two badness measures $\beta(\mathcal{L})$ and $q(\mathcal{L})$, so the upper bound looks like:

$$2 \|\mathcal{L}\| \leq s(k) |\mathcal{L}| + h(k)\beta(\mathcal{L}) + z(k)q(\mathcal{L}).$$

High $\beta(\mathcal{L})$ badness leads to reducible configurations by kernel-perfect orientations and high $q(\mathcal{L})$ badness leads to reducible configurations by Alon-Tarsi orientations. That means the same proof shows that Main Theorem holds for offline k -list-critical graphs as well (in fact, for the larger class of OC -irreducible graphs with $\delta(G) = k - 1$ defined in section 5).

Let $c_k^*(\mathcal{L})$ be the number of components of \mathcal{L} containing a copy of K_{k-1} . Let $q_k(\mathcal{L})$ be the number of non-cut vertices in \mathcal{L} that appear in copies of K_{k-1} . Let $\beta_k(\mathcal{L})$ be the independence number of the subgraph of \mathcal{L} induced on the vertices of degree $k - 1$. When k is defined in context, we just write $c^*(\mathcal{L})$, $q(\mathcal{L})$ and $\beta(\mathcal{L})$. We need upper bounds on our badness parameters $q(\mathcal{L})$ and $\beta(\mathcal{L})$. More general versions of the following two lemmas are proved in Section 2 and Section 3, respectively. The first lemma allows us a measure of control over K_{k-1} blocks in \mathcal{L} .

Lemma 1.1. *Let G be a non-complete online k -list-critical graph where $k \geq 5$. Let \mathcal{L} be the subgraph of G induced on $(k - 1)$ -vertices, \mathcal{H}^- the subgraph of G induced on k -vertices and \mathcal{H}^+ the subgraph of G induced on $(k + 1)^+$ -vertices. Then*

$$q(\mathcal{L}) \leq c^*(\mathcal{L}) + 4 |\mathcal{H}^-| + \|\mathcal{H}^+, \mathcal{L}\|,$$

and if $k \geq 7$, then

$$q(\mathcal{L}) \leq 2c^*(\mathcal{L}) + 3 |\mathcal{H}^-| + \|\mathcal{H}^+, \mathcal{L}\|.$$

The second lemma allows us some control over the number of vertices in \mathcal{L} with no neighbors in \mathcal{H} .

Lemma 1.2. *Let G be an online k -list-critical graph. Let \mathcal{L} be the subgraph of G induced on $(k - 1)$ -vertices and \mathcal{H} the subgraph of G induced on k^+ -vertices. If $2 \leq \lambda \leq \frac{6(k-1)}{k}$, then*

$$\beta(\mathcal{L}) \leq \frac{2}{\lambda} \|\mathcal{H}\| + \frac{2 \|G\| - (k - 2) |G| - \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |\mathcal{H}| - 1}{k - 1}.$$

In Section 4, we use these two lemmas and a little careful counting to prove general lower bounds on the average degree of online k -list-critical graphs in terms of parameters p, h, z, f that are functions of k and satisfy bounds of the form

$$2 \|T\| \leq (k - 3 + p(k)) |T| + h(k)q(T) + z(k)\beta(T) + f(k),$$

k	Gallai [5] $d(G) \geq$	KS [9] $d(G) \geq$	KR [7] $d(G) \geq$	CR [2] $d(G) \geq$	R [12] $d(G) \geq$	Here $d(G) \geq$
4	3.0769	—	—	—	3.1000	3.1000
5	4.0909	—	4.0984	4.1000	4.1176	4.1176
6	5.0909	—	5.1053	5.1076	5.1153	5.1214
7	6.0870	—	6.1149	6.1192	6.1081	6.1296
8	7.0820	—	7.1128	7.1167	7.1000	7.1260
9	8.0769	8.0838	8.1094	8.1130	8.0923	8.1213
10	9.0722	9.0793	9.1055	9.1088	9.0853	9.1162
15	14.0541	14.0610	14.0864	14.0884	14.0609	14.0930
20	19.0428	19.0490	19.0719	19.0733	19.0469	19.0762

Table 1: Lower bounds on average degree $d(G)$ of a k -list-critical graph G .

for every Gallai tree T with $\Delta(T) \leq k - 1$ (and a few further caveats). So, basically Section 4 is about turning upper bounds on the average degree of Gallai trees into lower bounds on the average degree of online k -list-critical graphs.

Finally, Section 5 proves upper bounds on the average degree of Gallai trees in terms of the parameters p, h, z, f .

2 Bounding $q(\mathcal{L})$

This section is devoted to extracting the reusable Lemma 2.1 from the proof of Kierstead and Rabern [7]. All of the hard work was already done in [7].

Definition 1. A graph G is *AT-reducible* to H if H is a nonempty induced subgraph of G which is f_H -AT where $f_H(v) := \delta(G) + d_H(v) - d_G(v)$ for all $v \in V(H)$. If G is not AT-reducible to any nonempty induced subgraph, then it is *AT-irreducible*.

Lemma 2.1. *Let G be a non-complete AT-irreducible graph with $\delta(G) = k - 1$ where $k \geq 5$. Let \mathcal{L} be the subgraph of G induced on $(k - 1)$ -vertices, \mathcal{H}^- the subgraph of G induced on k -vertices and \mathcal{H}^+ the subgraph of G induced on $(k + 1)^+$ -vertices. Then*

$$q(\mathcal{L}) \leq c^*(\mathcal{L}) + 4|\mathcal{H}^-| + \|\mathcal{H}^+, \mathcal{L}\|,$$

and if $k \geq 7$, then

$$q(\mathcal{L}) \leq 2c^*(\mathcal{L}) + 3|\mathcal{H}^-| + \|\mathcal{H}^+, \mathcal{L}\|.$$

Observation. The hypotheses of Lemma 2.1 are satisfied by non-complete k -critical, k -list-critical, online k -list-critical and k -AT-critical graphs.

The proof of Lemma 2.1 requires the following four lemmas from [7].

Lemma 2.2. *Let G be a graph and $f: V(G) \rightarrow \mathbb{N}$. If $\|G\| > \sum_{v \in V(G)} f(v)$, then G has an induced subgraph H such that $d_H(v) > f(v)$ for each $v \in V(H)$.*

Proof. Suppose not and choose a counterexample G minimizing $|G|$. Then $|G| \geq 3$ and we have $x \in V(G)$ with $d_G(x) \leq f(x)$. But now $\|G - x\| > \sum_{v \in V(G-x)} f(v)$, contradicting minimality of $|G|$. \square

Let \mathcal{T}_k be the Gallai trees with maximum degree at most $k - 1$, excepting K_k . For a graph G , let $W^k(G)$ be the set of vertices of G that are contained in some K_{k-1} in G .

Lemma 2.3. *Let $k \geq 5$ and let G be a graph with $x \in V(G)$ such that:*

1. $K_k \not\subseteq G$; and
2. $G - x$ has t components H_1, H_2, \dots, H_t , and all are in \mathcal{T}_k ; and
3. $d_G(v) \leq k - 1$ for all $v \in V(G - x)$; and
4. $|N(x) \cap W^k(H_i)| \geq 1$ for $i \in [t]$; and
5. $d_G(x) \geq t + 2$.

Then G is f -AT where $f(x) = d_G(x) - 1$ and $f(v) = d_G(v)$ for all $v \in V(G - x)$.

For a graph G , $\{X, Y\}$ a partition of $V(G)$ and $k \geq 4$, let $\mathcal{B}_k(X, Y)$ be the bipartite graph with one part Y and the other part the components of $G[X]$. Put an edge between $y \in Y$ and a component T of $G[X]$ iff $N(y) \cap W^k(T) \neq \emptyset$. The next lemma tells us that we have a reducible configuration if this bipartite graph has minimum degree at least three.

Lemma 2.4. *Let $k \geq 7$ and let G be a graph with $Y \subseteq V(G)$ such that:*

1. $K_k \not\subseteq G$; and
2. the components of $G - Y$ are in \mathcal{T}_k ; and
3. $d_G(v) \leq k - 1$ for all $v \in V(G - Y)$; and
4. with $\mathcal{B} := \mathcal{B}_k(V(G - Y), Y)$ we have $\delta(\mathcal{B}) \geq 3$.

Then G has an induced subgraph G' that is f -AT where $f(y) = d_{G'}(y) - 1$ for $y \in Y$ and $f(v) = d_{G'}(v)$ for all $v \in V(G' - Y)$.

We also have the following version with asymmetric degree condition on \mathcal{B} . The point here is that this works for $k \geq 5$. The consequence is that we trade a bit in our bound for the proof to go through with $k \in \{5, 6\}$.

Lemma 2.5. *Let $k \geq 5$ and let G be a graph with $Y \subseteq V(G)$ such that:*

1. $K_k \not\subseteq G$; and
2. the components of $G - Y$ are in \mathcal{T}_k ; and

3. $d_G(v) \leq k - 1$ for all $v \in V(G - Y)$; and
4. with $\mathcal{B} := \mathcal{B}_k(V(G - Y), Y)$ we have $d_{\mathcal{B}}(y) \geq 4$ for all $y \in Y$ and $d_{\mathcal{B}}(T) \geq 2$ for all components T of $G - Y$.

Then G has an induced subgraph G' that is f -AT where $f(y) = d_{G'}(y) - 1$ for $y \in Y$ and $f(v) = d_{G'}(v)$ for all $v \in V(G' - Y)$.

Proof of Lemma 2.1. Let \mathcal{H} be the subgraph of G induced on k^+ -vertices and let \mathcal{D} be the components of \mathcal{L} containing a copy of K_{k-1} . Put $W := W^k(\mathcal{L})$ and $L' := V(\mathcal{L}) \setminus W$. Define an auxiliary bipartite graph F with parts A and B where:

1. $B = V(\mathcal{H}^-)$ and A is the disjoint union of the following sets A_1, A_2 and A_3 ,
2. $A_1 = \mathcal{D}$ and each $T \in \mathcal{D}$ is adjacent to all $y \in B$ where $N(y) \cap W^k(T) \neq \emptyset$,
3. For each $v \in L'$, let $A_2(v)$ be a set of $|N(v) \cap B|$ vertices connected to $N(v) \cap B$ by a matching in F . Let A_2 be the disjoint union of the $A_2(v)$ for $v \in L'$,
4. For each $y \in B$, let $A_3(y)$ be a set of $d_{\mathcal{H}}(y)$ vertices which are all joined to y in F . Let A_3 be the disjoint union of the $A_3(y)$ for $y \in B$.

Define $f: V(F) \rightarrow \mathbb{N}$ by $f(v) = 1$ for all $v \in A_1 \cup A_2 \cup A_3$ and $f(v) = 3$ for all $v \in B$. First, suppose $\|F\| > \sum_{v \in V(F)} f(v)$. Then by Lemma 2.2, F has an induced subgraph Q such that $d_Q(v) > f(v)$ for each $v \in V(Q)$. In particular, $V(Q) \subseteq B \cup A_1$ and $d_Q(v) \geq 4$ for $v \in B \cap V(Q)$ and $d_Q(v) \geq 2$ for $v \in A_1 \cap V(Q)$. Put $Y := B \cap V(Q)$ and let X be $\bigcup_{T \in V(Q) \cap A_1} V(T)$. Now $Z := G[X \cup Y]$ satisfies the hypotheses of Lemma 2.5, so Z has an induced subgraph G' that is f -AT where $f(y) = d_{G'}(y) - 1$ for $y \in Y$ and $f(v) = d_{G'}(v)$ for $v \in X$. Since $Y \subseteq B$ and $X \subseteq V(\mathcal{L})$, we have $f(v) = k - 1 + d_{G'}(v) - d_G(v)$ for all $v \in V(G')$. Hence, G is AT-reducible to G' , a contradiction. Therefore $\|F\| \leq \sum_{v \in V(F)} f(v) = 3|B| + |\mathcal{D}| + |A_2| + |A_3|$. By Lemma 2.3, for each $y \in B$ we have $d_F(y) \geq k - 1$. Hence $\|F\| \geq (k - 1)|B|$. This gives $(k - 4)|B| \leq |\mathcal{D}| + |A_2| + |A_3|$. Now the first inequality in the lemma follows since $B = V(\mathcal{H}^-)$, $|A_3| = \sum_{v \in V(\mathcal{H}^-)} d_{\mathcal{H}}(v)$ and

$$\begin{aligned} |A_2| &= -q(\mathcal{L}) + \|\mathcal{H}, \mathcal{L}\| \\ &= -q(\mathcal{L}) + k|\mathcal{H}^-| + \|\mathcal{H}^+, \mathcal{L}\| - \sum_{v \in V(\mathcal{H}^-)} d_{\mathcal{H}}(v). \end{aligned}$$

Suppose $k \geq 7$. Define $f: V(F) \rightarrow \mathbb{N}$ by $f(v) = 1$ for all $v \in A_2 \cup A_3$ and $f(v) = 2$ for all $v \in B \cup A_1$. First, suppose $\|F\| > \sum_{v \in V(F)} f(v)$. Then by Lemma 2.2, F has an induced subgraph Q such that $d_Q(v) > f(v)$ for each $v \in V(Q)$. In particular, $V(Q) \subseteq B \cup A_1$ and $\delta(Q) \geq 3$. Put $Y := B \cap V(Q)$ and let X be $\bigcup_{T \in V(Q) \cap A_1} V(T)$. Now $Z := G[X \cup Y]$ satisfies the hypotheses of Lemma 2.4, so Z has an induced subgraph G' that is f -AT where $f(y) = d_{G'}(y) - 1$ for $y \in Y$ and $f(v) = d_{G'}(v)$ for $v \in X$. Since $Y \subseteq B$ and $X \subseteq V(\mathcal{L})$, we have $f(v) = k - 1 + d_{G'}(v) - d_G(v)$ for all $v \in V(G')$. Hence, G is AT-reducible to G' , a contradiction.

Therefore $\|F\| \leq \sum_{v \in V(F)} f(v) = 2(|B| + |\mathcal{D}|) + |A_2| + |A_3|$. By Lemma 2.3, for each $y \in B$ we have $d_F(y) \geq k - 1$. Hence $\|F\| \geq (k - 1)|B|$. This gives $(k - 3)|B| \leq 2|\mathcal{D}| + |A_2| + |A_3|$. Now the second inequality in the lemma follows as before. \square

3 Bounding $\beta(\mathcal{L})$

This section is devoted to extracting the reusable Lemma 3.1 from the proof of Rabern [12].

Definition 2. A graph G is *OC-reducible* to H if H is a nonempty induced subgraph of G which is online f_H -choosable where $f_H(v) := \delta(G) + d_H(v) - d_G(v)$ for all $v \in V(H)$. If G is not OC-reducible to any nonempty induced subgraph, then it is *OC-irreducible*.

Lemma 3.1. *Let G be an OC-irreducible graph with $\delta(G) = k - 1$. Let \mathcal{L} be the subgraph of G induced on $(k - 1)$ -vertices and \mathcal{H} the subgraph of G induced on k^+ -vertices. If $2 \leq \lambda \leq \frac{6(k-1)}{k}$, then*

$$\beta(\mathcal{L}) \leq \frac{2}{\lambda} \|\mathcal{H}\| + \frac{2\|G\| - (k - 2)|G| - \left(\frac{k}{2} + \frac{k-1}{\lambda}\right)|\mathcal{H}| - 1}{k - 1}.$$

Observation. The hypotheses of Lemma 3.1 are satisfied by k -critical, k -list-critical and online k -list-critical graphs.

The proof of Lemma 3.1 requires the following lemma from Kierstead and Rabern [8] that generalizes a kernel technique of Kostochka and Yancey [10].

Definition. The *maximum independent cover number* of a graph G is the maximum $\text{mic}(G)$ of $\|I, V(G) \setminus I\|$ over all independent sets I of G .

Lemma 3.2 (Kernel Magic). *Every OC-irreducible graph G with $\delta(G) = k - 1$ satisfies*

$$2\|G\| \geq (k - 2)|G| + \text{mic}(G) + 1.$$

Theorem 3.3 (Löwenstein, et al. [11]). *If G is a connected graph, then*

$$\alpha(G) \geq \frac{2}{3}|G| - \frac{1}{4}\|G\| - \frac{1}{3}.$$

Corollary 3.4. *If G is a connected graph, then*

$$\alpha(G) \geq \frac{2}{3}|G| - \frac{1}{3}\|G\|.$$

Proof. By Theorem 3.3,

$$\alpha(G) \geq \frac{2}{3}|G| - \frac{1}{3}\|G\| + \frac{1}{12}\|G\| - \frac{1}{3},$$

so, the corollary holds if $\frac{1}{12}\|G\| \geq \frac{1}{3}$. If not, then $\|G\| < 4$, so G is K_1 , K_2 , P_3 or K_3 which all satisfy the desired bound. \square

Proof of Lemma 3.1. Fix λ with $2 \leq \lambda \leq \frac{6(k-1)}{k}$. Let M be the maximum of $\|I, V(G) \setminus I\|$ over all independent sets I of G with $I \subseteq \mathcal{H}$. Since the vertices in \mathcal{L} with $k-1$ neighbors in \mathcal{L} have no neighbors in \mathcal{H} ,

$$\text{mic}(G) \geq M + (k-1)\beta(\mathcal{L}). \quad (1)$$

Claim 1. *If C is a component of \mathcal{H} , then*

$$k\alpha(C) \geq \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |C| - \left(\frac{2(k-1)}{\lambda}\right) \|C\|.$$

First, suppose $\|C\| < |C|$. Then $\|C\| = |C| - 1$ and C is a tree. If $|C| \geq 2$, then

$$\begin{aligned} k\alpha(C) &\geq k \frac{|C|}{2} \\ &\geq \left(\frac{k}{2} - \frac{k-1}{\lambda}\right) |C| + \frac{2(k-1)}{\lambda} \\ &= \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |C| - \left(\frac{2(k-1)}{\lambda}\right) (|C| - 1) \\ &= \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |C| - \left(\frac{2(k-1)}{\lambda}\right) \|C\|. \end{aligned}$$

If instead, $|C| = 1$, then $k\alpha(C) = k \geq \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |C| - \left(\frac{2(k-1)}{\lambda}\right) \|C\|$ since $\lambda \geq 2$.

So, we may assume $\|C\| \geq |C|$. Applying Corollary 3.4, we conclude

$$\begin{aligned} k\alpha(C) &\geq \frac{2k}{3} |C| - \frac{k}{3} \|C\| \\ &= \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |C| - \left(\frac{2(k-1)}{\lambda}\right) \|C\| + \left(\frac{k}{6} - \frac{k-1}{\lambda}\right) |C| - \left(\frac{k}{3} - \frac{2(k-1)}{\lambda}\right) \|C\| \\ &= \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |C| - \left(\frac{2(k-1)}{\lambda}\right) \|C\| + \left(\frac{k-1}{\lambda} - \frac{k}{6}\right) |C| \\ &\geq \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |C| - \left(\frac{2(k-1)}{\lambda}\right) \|C\|, \end{aligned}$$

where in the final inequality we used $\lambda \leq \frac{6(k-1)}{k}$.

Claim 2. *Lemma 3.1 is true.*

Summing the bound in Claim 1 over all components of \mathcal{H} and plugging into (1) gives

$$\text{mic}(G) \geq \left(\frac{k}{2} + \frac{k-1}{\lambda}\right) |\mathcal{H}| - \left(\frac{2(k-1)}{\lambda}\right) \|\mathcal{H}\| + (k-1)\beta(\mathcal{L}). \quad (2)$$

Applying Lemma 3.2 using (2) and solving for $\beta(\mathcal{L})$ proves the claim. \square

4 General lower bounds on average degree

This is the counting portion of the proof, which is simpler and more general than the counting in [7] and [2].

Definition 3. A quadruple (p, h, z, f) of functions from \mathbb{N} to \mathbb{R} is r -Gallai if for every $k \geq r$ and Gallai tree $T \neq K_k$ with $\Delta(T) \leq k - 1$, the following hold:

- if $K_{k-1} \subseteq T$, then $2 \|T\| \leq (k - 3 + p(k)) |T| + h(k)q(T) + z(k)\beta(T) + f(k)$; and
- if $K_{k-1} \not\subseteq T$, then $2 \|T\| \leq (k - 3 + p(k)) |T| + z(k)\beta(T)$.

Theorem 4.1. Let (p, h, z, f) be 7-Gallai. If $k \geq 7$ and $2 \leq z(k) \leq \frac{6(k-1)}{k}$, then for any non-complete online k -list-critical graph G ,

$$d(G) \geq k - 1 + \frac{2 - p(k) - \frac{z(k)}{k-1} + \frac{\frac{z(k)}{k-1} - (2h(k) + f(k))c^*(\mathcal{L})}{|G|}}{k + 1 + 3h(k) - p(k) - \frac{(k-2)z(k)}{2(k-1)}},$$

where \mathcal{L} is the subgraph of G induced on $(k - 1)$ -vertices.

Proof. Let \mathcal{H}^- the subgraph of G induced on k -vertices, \mathcal{H} the subgraph of G induced on k^+ -vertices, \mathcal{H}^+ the subgraph of G induced on $(k + 1)^+$ -vertices and \mathcal{D} the components of \mathcal{L} containing K_{k-1} . Plainly, the following bounds hold.

$$2 \|G\| \geq k |G| - |\mathcal{L}| \tag{3}$$

$$2 \|G\| \geq (k + 1) |G| - |\mathcal{H}^-| - 2 |\mathcal{L}| \tag{4}$$

$$2 \|G\| \geq k |\mathcal{H}^-| + (k - 1) |\mathcal{L}| + \|\mathcal{H}^+, \mathcal{L}\| \tag{5}$$

$$\|\mathcal{H}, \mathcal{L}\| = (k - 1) |\mathcal{L}| - 2 \|\mathcal{L}\| \tag{6}$$

Since (p, h, z, f) is 7-Gallai,

$$2 \|\mathcal{L}\| \leq (k - 3 + p(k)) |\mathcal{L}| + f(k) |\mathcal{D}| + h(k)q(\mathcal{L}) + z(k)\beta(\mathcal{L}) \tag{7}$$

By Lemma 1.1,

$$q(\mathcal{L}) \leq 2 |\mathcal{D}| + 3 |\mathcal{H}^-| + \|\mathcal{H}^+, \mathcal{L}\|,$$

plugging this into (7) gives

$$2 \|\mathcal{L}\| \leq (k - 3 + p(k)) |\mathcal{L}| + 3h(k) |\mathcal{H}^-| + h(k) \|\mathcal{H}^+, \mathcal{L}\| + z(k)\beta(\mathcal{L}) + S_1, \tag{8}$$

where

$$S_1 := (2h(k) + f(k)) |\mathcal{D}|.$$

Now using (3) and (8),

$$\begin{aligned}
2 \|G\| &= 2 \|\mathcal{H}\| + 2 \|\mathcal{H}, \mathcal{L}\| + 2 \|\mathcal{L}\| \\
&= 2 \|\mathcal{H}\| + 2((k-1) |\mathcal{L}| - 2 \|\mathcal{L}\|) + 2 \|\mathcal{L}\| \\
&= 2 \|\mathcal{H}\| + 2(k-1) |\mathcal{L}| - 2 \|\mathcal{L}\| \\
&\geq 2 \|\mathcal{H}\| + (k+1-p(k)) |\mathcal{L}| - 3h(k) |\mathcal{H}^-| - h(k) \|\mathcal{H}^+, \mathcal{L}\| - z(k)\beta(\mathcal{L}) - S_1 \quad (9)
\end{aligned}$$

Adding $h(k)$ times (5) to (9) gives

$$2 \|G\| \geq \frac{2 \|\mathcal{H}\| + (k+1+(k-1)h(k)-p(k)) |\mathcal{L}| + (k-3)h(k) |\mathcal{H}^-| - z(k)\beta(\mathcal{L}) - S_1}{1+h(k)} \quad (10)$$

Lemma 1.2 gives

$$\beta(\mathcal{L}) \leq \frac{2}{z(k)} \|\mathcal{H}\| + \frac{2 \|G\| - (k-2) |G| - \left(\frac{k}{2} + \frac{k-1}{z(k)}\right) |\mathcal{H}| - 1}{k-1}.$$

Plugging this into (10) yields

$$2 \|G\| \geq \frac{(k+1+(k-1)h(k)-p(k)) |\mathcal{L}| + (k-3)h(k) |\mathcal{H}^-| + \frac{(k-2)z(k)}{k-1} |G| + \left(\frac{kz(k)}{2(k-1)} + 1\right) |\mathcal{H}| + S_2}{1+h(k) + \frac{z(k)}{k-1}}, \quad (11)$$

where

$$S_2 := \frac{z(k)}{k-1} - S_1.$$

Now using $|\mathcal{H}| = |G| - |\mathcal{L}|$ gives

$$2 \|G\| \geq \frac{\left(k + (k-1)h(k) - p(k) - \frac{kz(k)}{2(k-1)}\right) |\mathcal{L}| + (k-3)h(k) |\mathcal{H}^-| + \left(\frac{(3k-4)z(k)}{2(k-1)} + 1\right) |G| + S_2}{1+h(k) + \frac{z(k)}{k-1}}. \quad (12)$$

Now using (4) to get a lower bound on $|\mathcal{H}^-|$ gives

$$2 \|G\| \geq \frac{\left(k - (k-5)h(k) - p(k) - \frac{kz(k)}{2(k-1)}\right) |\mathcal{L}| + \left((k+1)(k-3)h(k) + \frac{(3k-4)z(k)}{2(k-1)} + 1\right) |G| + S_2}{1+(k-2)h(k) + \frac{z(k)}{k-1}}. \quad (13)$$

Using (3) to get a lower bound on $|\mathcal{L}|$ and simplifying gives

$$\frac{2 \|G\|}{|G|} \geq \frac{k^2 + 3(k-1)h(k) - kp(k) + 1 - \frac{k^2-3k+4}{2(k-1)} z(k) + \frac{S_2}{|G|}}{k+1+3h(k)-p(k)-\frac{(k-2)z(k)}{2(k-1)}}. \quad (14)$$

Now factoring out $k-1$ gives the desired bound. \square

A nearly identical argument, using the other inequality in Lemma 1.1, proves a bound that holds for $k \geq 5$.

Theorem 4.2. *Let (p, h, z, f) be 5-Gallai. If $k \geq 5$ and $2 \leq z(k) \leq \frac{6(k-1)}{k}$, then for any non-complete online k -list-critical graph G ,*

$$d(G) \geq k - 1 + \frac{2 - p(k) - \frac{z(k)}{k-1} + \frac{\frac{z(k)}{k-1} - (h(k) + f(k))c^*(\mathcal{L})}{|G|}}{k + 1 + 4h(k) - p(k) - \frac{(k-2)z(k)}{2(k-1)}},$$

where \mathcal{L} is the subgraph of G induced on $(k-1)$ -vertices.

When $k = 4$, we cannot apply Lemma 1.1, but using $h(k) = 0$ and running through the same argument proves the following bound for $k \geq 4$.

Theorem 4.3. *Let $(p, 0, z, f)$ be 4-Gallai. If $k \geq 4$ and $2 \leq z(k) \leq \frac{6(k-1)}{k}$, then for any non-complete online k -list-critical graph G ,*

$$d(G) \geq k - 1 + \frac{2 - p(k) - \frac{z(k)}{k-1} + \frac{\frac{z(k)}{k-1} - f(k)c^*(\mathcal{L})}{|G|}}{k + 1 - p(k) - \frac{(k-2)z(k)}{2(k-1)}},$$

where \mathcal{L} is the subgraph of G induced on $(k-1)$ -vertices.

When $z(k) < 2$, using Lemma 1.2 worsens the lower bound, so we may as well use $z(k) = 0$; that is, drop the $\beta(\mathcal{L})$ term entirely. Doing so in the above argument shows that Theorems 4.1, 4.2, 4.3 hold for $z(k) = 0$ if we replace $k + 1$ in the denominator with $k + 2$. This gives the bounds proved by discharging in Cranston and Rabern [2].

5 Gallai quadruples

All known proofs of lower bounds for average degree of list-critical graphs are essentially a counting argument combined with the fact that some quadruple is Gallai.

Lemma 5.1 (Gallai [5]). *The tuple $(\frac{k+1}{k-1}, 0, 0, -2)$ is 4-Gallai.*

Lemma 5.2 (Kostochka-Stiebitz [9]). *The tuple $(\frac{4(k-1)}{k^2-3k+4}, \frac{k^2-3k}{k^2-3k+4}, 0, \frac{-4(k^2-3k+2)}{k^2-3k+4})$ is 7-Gallai.*

Lemma 5.3 (Cranston-Rabern [2]). *The tuple $(\frac{3k-5}{k^2-4k+5}, \frac{k(k-3)}{k^2-4k+5}, 0, \frac{-2(k-1)(2k-5)}{k^2-4k+5})$ is 5-Gallai.*

Lemma 5.4 (Rabern [12]). *The tuple $(1, 0, 2, 0)$ is 4-Gallai.*

We give a list of inequalities that provide a sufficient condition for (p, h, z, f) to be 5-Gallai. These inequalities take a form quite similar to the inequalities in Cranston and Rabern [2], but now they involve $z(k)$ as well. The sufficiency proof is a small modification of the proof in [2]. To use a Gallai quadruple in Lemma 4.1, we want $2h(k) + f(k) \leq 0$ to get rid of the term involving $c^*(\mathcal{L})$. Similarly, for Lemma 4.2, we want $h(k) + f(k) \leq 0$. Finding the p, h, z, f that give the largest average degree subject to these constraints is a fractional linear program that can be converted to a linear program and solved for each k . This is useful for verification of bounds, but we want a formula in terms of k . For $k \geq 7$, we use the following quadruple.

Lemma 5.5. *The tuple $\left(\frac{3k-7}{k^2-4k+5}, \frac{(k-1)(k-4)}{k^2-4k+5}, 2, \frac{-2(k-1)(k-4)}{k^2-4k+5}\right)$ is 5-Gallai.*

For $k = 6$, we use the following quadruple. For $k = 5$, the quadruple in Lemma 5.4 is the optimal choice of p, h, z, f .

Lemma 5.6. *The tuple $\left(\frac{3k-5}{k^2-3k+3}, \frac{(k-1)(k-4)}{k^2-3k+3}, \frac{(3k-5)(k-2)}{k^2-3k+3}, \frac{-(k-1)(k-4)}{k^2-3k+3}\right)$ is 5-Gallai.*

Now on to the sufficiency proof. We prove two technical lemmas, the proof of Lemma 5.5 and Lemma 5.6 are straightforward computations from these. For an endblock B of a Gallai tree T , let x_B be the cutvertex contained in B .

Lemma 5.7. *Let $z: \mathbb{N} \rightarrow \mathbb{R}$ such that $z(k) = 0$ or $z(k) \geq 2$ for all $k \in \mathbb{N}$. For all $k \geq 5$ and Gallai trees T with $\Delta(T) \leq k-1$ and $K_{k-1} \not\subseteq T$, we have*

$$2 \|T\| \leq \left(k - 3 + \frac{\max\{2, 3 - z(k)\}}{k - 2}\right) |T| + z(k)\beta(T).$$

Proof. Suppose the lemma is false and choose a counterexample T minimizing $|T|$.

Claim 1. *T has at least two blocks.*

If T has only one block, then $2 \|T\| \leq (k-3) |T|$.

Claim 2. *Each endblock of T is K_{k-2} .*

Suppose T has an endblock B that is not K_{k-2} . Then removing $V(B) \setminus \{x_B\}$ from T to get T' and applying minimality of $|T|$ gives

$$2 \|B\| > \left(k - 3 + \frac{\max\{2, 3 - z(k)\}}{k - 2}\right) (|B| - 1).$$

This is a contradiction unless $k = 5$ and $B = K_3$, but then $B = K_{k-2}$, a contradiction.

Claim 3. *If B is an endblock of T , then $d_T(x_B) = k-1$.*

Suppose B is an endblock of T with $d_T(x_B) < k-1$. Then $B = K_{k-2}$ by Claim 2 and hence $d_T(x_B) = k-2$. Removing $V(B)$ from T to get T^* and applying minimality of $|T|$ gives the contradiction

$$(k-2)(k-3) + 6 > \left(k - 3 + \frac{\max\{2, 3 - z(k)\}}{k - 2}\right) (k-1).$$

Claim 4. *T does not exist.*

By the previous claims, we know that every endblock T is a K_{k-2} that shares a vertex with an odd cycle. Pick an endblock B that is the end of a longest path in the block-tree of T . Let C be the odd cycle sharing x_B with B . Since B is the end of a longest path in the block-tree, there is a neighbor y of x_B on C such that $d_T(y) = 2$ or y is contained in another endblock A (which must be a K_{k-2}). First, suppose $d_T(y) = 2$. Removing $V(B) \cup \{y\}$ from T to get T' and applying minimality of $|T|$ gives the contradiction (since $\beta(T') < \beta(T)$)

$$(k-2)(k-3) + 6 > \left(k - 3 + \frac{\max\{2, 3 - z(k)\}}{k - 2}\right) (k-1) + z(k)(\beta(T) - \beta(T')).$$

Hence y is contained in another K_{k-2} endblock A . Removing $V(B) \cup V(A)$ from T to get T^* and applying minimality of $|T|$ gives the contradiction (since $\beta(T^*) < \beta(T)$)

$$2(k-2)(k-3) + 6 > \left(k-3 + \frac{\max\{2, 3-z(k)\}}{k-2}\right) (2(k-2)) + z(k)(\beta(T) - \beta(T^*)). \quad \square$$

Lemma 5.8. *Let $p: \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$, $f: \mathbb{N} \rightarrow \mathbb{R}$, $h: \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$, $z: \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ such that $z(k) = 0$ or $z(k) \geq 2$. For all $k \geq 5$ and Gallai trees $T \neq K_k$ with $\Delta(T) \leq k-1$ and $K_{k-1} \subseteq T$, we have*

$$2\|T\| \leq (k-3+p(k))|T| + f(k) + h(k)q(T) + z(k)\beta(T)$$

whenever p , f , h and z satisfy all of the following conditions:

- (1) $f(k) \geq (k-1)(1-p(k)-h(k))$; and
- (2) $p(k) \geq \frac{3-\frac{z(k)}{2}}{k-2}$; and
- (3) $p(k) \geq h(k) + 5 - k$; and
- (4) $p(k) \geq \frac{2+h(k)}{k-2}$; and
- (5) $(k-1)p(k) + (k-3)h(k) + z(k) \geq k+1$.

Proof. Suppose the lemma is false and choose a counterexample T minimizing $|T|$.

Claim 1. T has at least two blocks.

Otherwise, $T = K_{k-1}$ and (1) gives a contradiction.

Claim 2. Each endblock of T is K_{k-2} or K_{k-1} .

Suppose T has an endblock B that is not K_{k-2} or K_{k-1} . Then removing $V(B) \setminus \{x_B\}$ from T to get T' and applying minimality of $|T|$ gives

$$2\|B\| > (k-3+p(k))(|B|-1) + h(k)(q(T) - q(T')) + z(k)(\beta(T) - \beta(T')).$$

If $B = K_2$, then $q(T') \leq q(T) + 1$, otherwise $q(T') = q(T)$. For $B = K_2$, we have to contradiction (to (3))

$$2 > (k-3+p(k)) - h(k).$$

Suppose $B = K_t$ for $4 \leq t \leq k-3$. Then we have the contradiction

$$t(t-1) > (k-3+p(k))(t-1).$$

Finally, suppose B is an odd cycle of length ℓ . Then, we have

$$2\ell > (k-3+p(k))(\ell-1).$$

This simplifies to

$$\ell < 1 + \frac{2}{k-5+p(k)}.$$

Since $k - 5 + p(k) \geq 1$ when $k \geq 6$, this implies that $k = 5$. Using (4), we conclude $\ell = 3$, but then $B = K_{k-2}$, a contradiction.

Claim 3. *T has at most one K_{k-1} endblock.*

Suppose T has at least two K_{k-1} endblocks. Let B be one of them. Then removing $V(B)$ from T leaves a graph T' with $K_{k-1} \subseteq T'$. So, we may apply minimality of $|T|$ to get

$$(k-1)(k-2) + 2 > (k-3+p(k))(k-1) + h(k)(q(T) - q(T')) + z(k)(\beta(T) - \beta(T')).$$

Now $\beta(T') < \beta(T)$ and $q(T') \leq q(T) - (k-2) + 1$, so we have the contradiction (to (5))

$$k+1 > (k-1)p(k) + (k-3)h(k) + z(k).$$

Claim 4. *If B is an endblock of T , then $d_T(x_B) = k-1$.*

Suppose B is an endblock of T with $d_T(x_B) < k-1$. Then $B = K_{k-2}$ by Claim 2. Removing $V(B)$ from T leaves a graph T' with $K_{k-1} \subseteq T'$. So, we may apply minimality of $|T|$ to get

$$(k-2)(k-3) + 2 > (k-3+p(k))(k-2) + h(k)(q(T) - q(T')) + z(k)(\beta(T) - \beta(T')).$$

We have $q(T') \leq q(T) + 1$, so this gives the contradiction (to (4))

$$2 > (k-1)p(k) - h(k).$$

Claim 5. *T does not exist.*

By Claims 2 and 3, all but at most one endblock of T is K_{k-2} with a cutvertex that is also in an odd cycle. Pick an endblock B that is the end of a longest path in the block-tree of T . Let C be the odd cycle sharing x_B with B . Since B is the end of a longest path in the block-tree, there is a neighbor y of x_B on C such that $d_T(y) = 2$ or y is contained in another endblock A (which must be a K_{k-2}). First, suppose $d_T(y) = 2$. Removing $V(B) \cup \{y\}$ from T to get T' and applying minimality of $|T|$ gives (since $q(T') = q(T)$ and $\beta(T') < \beta(T)$)

$$(k-2)(k-3) + 6 > (k-3+p(k))(k-1) + z(k),$$

so

$$p(k) < \frac{9-k-z(k)}{k-1},$$

contradicting (2). Hence y is contained in another K_{k-2} endblock A . Removing $V(B) \cup V(A)$ from T to get T^* and applying minimality of $|T|$ gives (since $q(T^*) = q(T)$ and $\beta(T^*) < \beta(T)$)

$$2(k-2)(k-3) + 6 > (k-3+p(k))(2(k-2)) + z(k),$$

so

$$6 > 2(k-2)p(k) + z(k),$$

contradicting (2). □

That is all we need to prove our lower bounds on average degree. If a good upper bound on $c^*(\mathcal{L})$ is known, it may be better to allow $2h(k) + f(k) > 0$. In that case, one could use the following.

Lemma 5.9. *If $z: \mathbb{N} \rightarrow \mathbb{R}$ is such that $z(k) = 0$ or $2 \leq z(k) \leq \frac{k(k-3)}{k-2}$ for all $k \in \mathbb{N}$, then (p, h, z, f) is 5-Gallai, where*

$$\begin{aligned} h(k) &:= \frac{k(k-3) - (k-2)z(k)}{k^2 - 4k + 5}, \\ p(k) &:= \frac{2 + h(k)}{k-2}, \\ f(k) &:= (k-1)(1 - h(k) - p(k)). \end{aligned}$$

References

- [1] O.V. Borodin, *Criterion of chromaticity of a degree prescription*, Abstracts of IV All-Union Conf. on Th. Cybernetics, 1977, pp. 127–128.
- [2] D. Cranston and L. Rabern, *Edge lower bounds for list critical graphs, via discharging*, *Combinatorica*, to appear [doi:10.1007/s00493-016-3584-6](https://doi.org/10.1007/s00493-016-3584-6).
- [3] R. Diestel, *Graph Theory*, 4 ed., Springer-Verlag, Heidelberg, 2010.
- [4] P. Erdős, A.L. Rubin, and H. Taylor, *Choosability in graphs*, Proc. West Coast Conf. on Combinatorics, Graph Theory and Computing, Congressus Numerantium, vol. 26, 1979, pp. 125–157.
- [5] T. Gallai, *Kritische Graphen I.*, Publ. Math. Inst. Hungar. Acad. Sci **8** (1963), 165–192 (in German).
- [6] G. Gutowski, *On-line choosability*, Ph.D. thesis, Jagiellonian University, Kraków, 2011.
- [7] H.A. Kierstead and L. Rabern, *Improved lower bounds on the number of edges in list critical and online list critical graphs*, [arXiv:1406.7355](https://arxiv.org/abs/1406.7355) (2014).
- [8] ———, *Extracting list colorings from large independent sets*, *Journal of Graph Theory*, to appear [doi:10.1002/jgt.22128](https://doi.org/10.1002/jgt.22128).
- [9] A.V. Kostochka and M. Stiebitz, *A new lower bound on the number of edges in colour-critical graphs and hypergraphs*, *Journal of Combinatorial Theory, Series B* **87** (2003), no. 2, 374–402.
- [10] A.V. Kostochka and M. Yancey, *Ore’s conjecture on color-critical graphs is almost true*, *J. Combin. Theory Ser. B* **109** (2014), 73–101.
- [11] C. Löwenstein, A. Pedersen, D. Rautenbach, and F. Regen, *Independence, odd girth, and average degree*, *Journal of Graph Theory* **67** (2011), no. 2, 96–111.
- [12] L. Rabern, *A better lower bound on average degree of 4-list-critical graphs*, *Electronic Journal of Combinatorics* **23(3)** (2016), #P3.37.
- [13] U. Schauz, *Mr. Paint and Mrs. Correct*, *Electronic Journal of Combinatorics* **16(1)** (2009), #R77.