

# Migratory birds in southern Brazil are a source of multiple avian influenza virus subtypes

Jansen Araujo<sup>1</sup>  | Maria Virgínia Petry<sup>2</sup> | Thomas Fabrizio<sup>3</sup> | David Walker<sup>3</sup> | Tatiana Ometto<sup>1</sup> | Luciano M. Thomazelli<sup>1</sup> | Angelo L. Scherer<sup>2</sup> | Patricia P. Serafini<sup>4</sup> | Isaac S. Neto<sup>4</sup> | Scott Krauss<sup>3</sup> | Robert G. Webster<sup>3</sup> | Richard J. Webby<sup>3</sup> | Edison L. Durigon<sup>1</sup>

<sup>1</sup>Laboratório de Virologia Clínica e Molecular do Instituto de Ciências Biomédicas (ICB-II), Universidade de São Paulo, São Paulo, SP, Brazil

<sup>2</sup>Laboratório de Ornitologia e Animais Marinhos (LOAM), Universidade do Vale do Rio dos Sinos, UNISINOS, São Leopoldo, RS, Brazil

<sup>3</sup>Department of Infectious Diseases, St. Jude Children's Research Hospital, Memphis, TN, USA

<sup>4</sup>Centro Nacional de Pesquisa e Conservação das Aves Silvestres (CEMAVE/ICMBio/MMA), Brazil, Florianópolis, Brazil

## Correspondence

Jansen Araujo, Institute of Biomedical Science, University of São Paulo, São Paulo, SP, Brazil.  
Email: jansentequila@usp.br

## Funding information

Fundação de Amparo à Pesquisa do Estado de São Paulo, Grant/Award Number: 13821-7; US National Institute of Allergy and Infectious Disease Centers of Excellence for Influenza Research and Surveillance (CEIRS) and by the American Lebanese Syrian Associated Charities (ALSAC) program, Grant/Award Number: (HHSN266200700005C); Wildlife Conservation Society (WCS) Project; Grant/Award Number: 2008005\_WCS\_OWOH and 2009005\_WCS\_OWOH; Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul, Grant/Award Number: 09/0574-7

**Background:** There is insufficient knowledge about the relation of avian influenza virus (AIV) to migratory birds in South America. Accordingly, we studied samples obtained over a 4-year period (2009-2012) from wild birds at a major wintering site in southern Brazil.

**Methods:** We obtained 1212 oropharyngeal/cloacal samples from wild birds at Lagoa do Peixe National Park and screened them for influenza A virus by RT-PCR amplification of the matrix gene. Virus isolates were subjected to genomic sequencing and antigenic characterization.

**Results:** Forty-eight samples of 1212 (3.96%) contained detectable influenza virus RNA. Partial viral sequences were obtained from 12 of these samples, showing the presence of H2N2 (1), H6Nx (1), H6N1 (8), H9N2 (1), and H12N5 (1) viruses. As H6 viruses predominated, we generated complete genomes from all 9 H6 viruses. Phylogenetic analyses showed that they were most similar to viruses of South American lineage. The H6N1 viruses caused no disease signs in infected ferrets and, despite genetic differences, were antigenically similar to North American isolates.

**Conclusions:** Lagoa do Peixe National Park is a source of multiple AIV subtypes, with the levels of influenza virus in birds being highest at the end of their wintering period in this region. H6N1 viruses were the predominant subtype identified. These viruses were more similar to viruses of South American lineage than to those of North American lineage.

## KEYWORDS

avian influenza virus, Brazil, clade, H6N1, migratory birds, North America, reassortment, South America, waterfowl

## 1 | INTRODUCTION

Aquatic birds, particularly the Anseriformes (waterfowl) and Charadriiformes (gulls and shorebirds), are the natural reservoirs

of avian influenza virus (AIV) and serve to perpetuate most known influenza A subtypes.<sup>1-3</sup> The major migratory bird stopover sites may create areas for enhanced interspecies contact, AIV transmission, and AIV genetic reassortment in different bird species from

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.


© 2017 The Authors. *Influenza and Other Respiratory Viruses* Published by John Wiley & Sons Ltd.

different flyways.<sup>4</sup> Every year, millions of wild birds from different North American flyways converge at wintering sites in Brazil. The Lagoa do Peixe National Park is recognized as the most important of these stopovers and acts as a wintering site for North American shorebirds.<sup>5-7</sup> The birds usually remain in Brazil from September to May, and they depend on this habitat to replenish the energy expended in their winter migration.<sup>8</sup> Lagoa do Peixe National Park is also important to resident birds and to South American migrant birds,<sup>9</sup> suggesting that there is great potential there for the spread and genetic reassortment of influenza A viruses. Despite the commingling of birds from major North and South American flyways, there is a marked phylogenetic separation of some influenza viruses residing within them. Based on genetic sequence, there are 2 major groups of influenza viruses in the Americas, one made up of viruses from North and South America, and the second only with viruses from South America.<sup>10</sup>

Influenza A viruses are composed of 8 RNA segments and are sub-typed according to their hemagglutinin (HA) and neuraminidase (NA) antigenic glycoproteins. Evidence suggests that viruses representing the 16 HA and 9 NA subtypes exist in harmony with their natural reservoirs, causing little or no disease.<sup>11</sup> AIVs in Eurasia and Australia appear to have diverged from those in North America, presumably as a

result of different flyways being established and a consequent lack of contact between birds and therefore viruses, from the different continents.<sup>12</sup> Similarly, the HA and NA genes of AIVs in Argentina differ from those of viruses circulating in North America and Eurasia.<sup>13</sup> The main subtypes of AIV consistently reported to circulate in other countries<sup>14-19</sup> have not been reported in Brazil. However, an H11N9 influenza virus was recently identified for the first time in South American migratory birds in the Amazon region, and this virus showed great similarity to viruses of North American, rather than South American, lineages.<sup>20</sup>

Compared to surveillance activities in other regions of the world, influenza virus surveillance in South American wildlife has been limited. It remains unclear whether the North American and South American AIV lineages converge and whether migrating wild birds transport AIVs of South American lineage to the United States.<sup>21</sup> Much remains unknown about AIV ecology in Brazil, including the identity of the main virus carrier species and the extent to which wild birds support virus spread between North and South America. Here, we studied samples obtained from wild birds in southern Brazil over a 4-year period to determine the extent of AIV circulation, the major host species of AIV, and the nature of any viruses present.



**FIGURE 1** Sampling site in Lagoa do Peixe National Park. The inset indicates the location of Rio Grande do Sul State in Brazil

## 2 | METHODS

### 2.1 | Ethics statement

The study was conducted in strict accordance with the guidelines of the Ethical Principles of Animal Experimentation adopted by the Brazilian Society of Laboratory Animal Science (SBCAL), and the animal protocol was approved by the Ethics Committee on Animal Experiments (EAEC) (Permit No. 105). All procedures involving wild birds were approved by the Ministério do Meio Ambiente (MMA) and the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio/SISBIO) (SISBIO numbers. 17565-1, 22976-8, 23159-3, 24381-3, and 14966-1). The ferret studies were approved by the Institutional Animal Care and Use Committee of St. Jude Children's Research Hospital.

### 2.2 | Study site and sample collection

During the years 2009 through 2012, AIV surveillance was conducted in Lagoa do Peixe National Park in Rio Grande do Sul State, Brazil (31°82'17.94"S, 51°80'52.37"W), which is a natural gathering site for migratory birds (Figure 1). Oropharyngeal and cloacal swabs were obtained from 1212 wild birds. The bird families captured in mist nets included Ardeidae, Charadriidae, Haematopodidae, Recurvirostridae, Scolopacidae, Rostratulidae, Tyrannidae, Furnariidae, and Laridae, except Kelp Gull, and were then released after sampling. The other bird families were sampled along a 120-km stretch of beach north of the Lagoa do Peixe National Park, between Balneário Pinhal (lat 30°14'55"S, long 50°13'47"W) and Mostardas (lat 31°10'52"S, long 50°50'03"W). Oropharyngeal and cloacal samples were also collected from recently stranded dead birds on the beach in monthly trips from November 2009 to June 2012. The swabs were placed in vials containing VTM transport medium composed of salt solution (PBS) supplemented with antimicrobial agents (200 U/mL penicillin G, 200 U/mL streptomycin, 25 µg/mL fungizone, and 6 µg/mL gentamycin) and 10% glycerol. All samples were immediately placed in liquid nitrogen following collection in duplicate. The duplicate sample was shipped to St. Jude Children's Research Hospital via the vapor phase of liquid nitrogen to prevent any freeze/thaw cycles.

### 2.3 | RNA extraction and AIV screening

RNA was extracted from the VTM medium using an RNA extraction kit (MagMax TM-96 RNA Isolation Kit; Ambion, Austin, TX, USA) in accordance with the manufacturer's instructions. The RNA was eluted into 80 µL of RNase-free water. All samples were screened for the presence of the AIV matrix gene by one-step real-time reverse transcriptase (RT)-PCR, using an AIV-M TaqMan Reagents Kit (Applied Biosystems, Foster City, CA, USA).<sup>20,22</sup>

### 2.4 | Virus isolation and subtyping

Virus was isolated by standard methods in 10-day-old embryonated chicken eggs, and viral RNA was amplified from samples positive

for AIV by real-time RT-PCR.<sup>23,24</sup> The Flu DETECT Avian Influenza Virus Type A Test Kit (Synbiotics, San Diego, CA, USA) was used to quickly confirm the presence of AIV protein within each sample.<sup>25</sup> Hemagglutination and neuraminidase inhibition tests were performed with a panel of polyclonal reference sera to identify the influenza subtypes as recorded in Table 1.<sup>26-28</sup>

### 2.5 | Virus sequencing

Viral RNA was extracted from 200 µL of allantoic fluid using the RNeasy Mini Kit (Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions. RNA was eluted into 80 µL of RNase-free water. The cDNA was transcribed using a High-capacity cDNA Archive Kit (Applied Biosystems) in accordance with the manufacturer's directions. The full genomes of all H6-positive samples were sequenced using a high-throughput next-generation sequencing pipeline on an Ion Torrent Personal Genome Machine (PGM) sequencer (Life Technologies Inc., New York, NY, USA)<sup>29-31</sup> in a BSL3+ laboratory at the University of São Paulo and on Roche 454 GS FLX (Roche Diagnostics Corp., Branford, CT, USA) and Illumina MiSeq platforms (Illumina, San Diego, CA, USA), as described previously,<sup>32-34</sup> at St. Jude Children's Research Hospital. Samples for MiSeq sequencing were prepared using the Illumina Nextera XT DNA Library Preparation Kit in accordance with the manufacturer's instructions. Briefly, whole influenza genomes were amplified using universal primers (sequences available upon request) and tagged into fragments of approximately 300 bp each. The fragments were enriched and amplified before sequencing on the MiSeq instrument using the 300-cycle MiSeq Reagent v2 Kit (Illumina). Sequences were deposited in the GenBank database under accession numbers KX620053-KX620135, KX670559-KX670562, KX814376-KX814382, and KX827310.

### 2.6 | Phylogenetic analysis

The complete sequences of all segments of each isolate were compared using the Basic Local Alignment Search Tool (BLAST) to identify the most closely related sequences available in public databases.<sup>35</sup> The nucleotide sequences of each gene segment were aligned using CLC GENOMICS WORKBENCH version 7.6 (Qiagen, Aarhus, Denmark), and the percentage nucleotide similarity was calculated using the MegAlign program of the DNASTAR package (DNASTAR, Inc., Madison, WI, USA). The available full-genome reference sequences of AIVs isolated from birds were used in the construction of the phylogenetic trees. The HA and NA genes were analyzed using open reading frames (ORFs); to identify the most closely related sequences, maximum likelihood trees were inferred on the basis of nucleotide alignment, using the program Bayesian Markov Chain Monte Carlo (MCMC) in v1.8.0,<sup>36</sup> with the generalized time-reversible (GTR) model and gamma-distributed rates among sites.

### 2.7 | Ferret studies

To prepare post-infection ferret antiserum, ferrets aged approximately 8 months (Triple F Farms, Sayre, PA, USA), previously

**TABLE 1** Forty-eight samples testing positive for influenza A virus. The 17 samples in which the presence of a specific subtype was confirmed are in bold. Nine H6N1-subtype viruses were confirmed, along with one virus of each of the H6Nx, H2N2, H12N5, and H9N2 subtypes

| Tubes | Sample ID       | Ct        | Scientific name                | Date              | Location              | HA | Flu detect | HI      | Subtype      |
|-------|-----------------|-----------|--------------------------------|-------------------|-----------------------|----|------------|---------|--------------|
| 1     | PNLP 233        | 38        | <i>Calidris fuscicollis</i>    | 25/03/2010        | Lagoa do Peixe        | –  |            |         | –            |
| 2     | <b>PNLP 304</b> | <b>37</b> | <b><i>Sterna hirundo</i></b>   | <b>27/03/2010</b> | <b>Lagoa do Peixe</b> | –  |            |         | <b>H6N</b>   |
| 3     | <b>PNLP 315</b> | <b>35</b> | <b><i>Calidris canutus</i></b> | <b>27/03/2010</b> | <b>Lagoa do Peixe</b> | +  | +          | 640     | <b>H12N5</b> |
| 4     | PNLP 319        | 35        | <i>C. canutus</i>              | 27/03/2010        | Lagoa do Peixe        | –  |            |         | –            |
| 5     | PNLP-320        | 37        | <i>C. canutus</i>              | 27/03/2010        | Lagoa do Peixe        | –  |            |         | –            |
| 6     | PNLP-325        | 36        | <i>C. fuscicollis</i>          | 27/03/2010        | Lagoa do Peixe        | –  |            |         | –            |
| 7     | PNLP-346        | 37        | <i>C. fuscicollis</i>          | 27/03/2010        | Lagoa do Peixe        | –  |            |         | –            |
| 8     | PNLP-395        | 39        | <i>C. fuscicollis</i>          | 29/03/2010        | Lagoa do Peixe        | –  |            |         | –            |
| 9     | RS 738          | 36        | <i>Larus dominicanus</i>       | 20/12/2010        | Lagoa do Peixe        | –  |            |         | –            |
| 10    | <b>RS 787</b>   | <b>38</b> | <b><i>C. canutus</i></b>       | <b>04/04/2011</b> | <b>Lagoa do Peixe</b> | –  |            |         | <b>H9N2</b>  |
| 11    | <b>RS 1147</b>  | <b>28</b> | <b><i>C. fuscicollis</i></b>   | <b>17/04/2012</b> | <b>Lagoa do Peixe</b> | –  |            |         | <b>H6N1</b>  |
| 12    | RS 1148         | 36        | <i>Charadrius semipalmatus</i> | 17/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 13    | <b>RS 1149</b>  | <b>25</b> | <b><i>C. fuscicollis</i></b>   | <b>17/04/2012</b> | <b>Lagoa do Peixe</b> | +  | +          | 160     | <b>H6N1</b>  |
| 14    | RS 1150         | 34        | <i>C. fuscicollis</i>          | 17/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 15    | <b>RS 1151</b>  | <b>30</b> | <b><i>C. fuscicollis</i></b>   | <b>18/04/2012</b> | <b>Lagoa do Peixe</b> | +  | +          | 640     | <b>H6N1</b>  |
| 16    | RS 1152         | 39        | <i>C. semipalmatus</i>         | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 17    | RS 1153         | 38        | <i>C. semipalmatus</i>         | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 18    | <b>RS 1154</b>  | <b>29</b> | <b><i>C. fuscicollis</i></b>   | <b>18/04/2012</b> | <b>Lagoa do Peixe</b> | –  |            |         | <b>H6N1</b>  |
| 19    | RS 1155         | 32        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 20    | RS 1156         | 34        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 21    | RS 1158         | 31        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 22    | <b>RS 1167</b>  | <b>32</b> | <b><i>C. fuscicollis</i></b>   | <b>18/04/2012</b> | <b>Lagoa do Peixe</b> | +  | –          | –       | <b>H6N1</b>  |
| 23    | <b>RS 1169</b>  | <b>33</b> | <b><i>C. fuscicollis</i></b>   | <b>18/04/2012</b> | <b>Lagoa do Peixe</b> | –  |            |         | <b>H6N1</b>  |
| 24    | RS 1170         | 38        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 25    | <b>RS 1177</b>  | <b>23</b> | <b><i>C. fuscicollis</i></b>   | <b>18/04/2012</b> | <b>Lagoa do Peixe</b> | +  | –          | –       | <b>H6N1</b>  |
| 26    | RS 1179         | 39        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 27    | RS 1183         | 38        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 28    | RS 1188         | 32        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 29    | RS 1189         | 38        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 30    | RS 1190         | 39        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 31    | <b>RS 1193</b>  | <b>30</b> | <b><i>C. fuscicollis</i></b>   | <b>18/04/2012</b> | <b>Lagoa do Peixe</b> | –  |            |         | <b>H2N2</b>  |
| 32    | <b>RS 1196</b>  | <b>30</b> | <b><i>C. fuscicollis</i></b>   | <b>18/04/2012</b> | <b>Lagoa do Peixe</b> | –  |            |         | <b>H6N1</b>  |
| 33    | RS 1197         | 39        | <i>C. semipalmatus</i>         | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 34    | RS 1198         | 39        | <i>Rynchops niger</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 35    | RS 1199         | 40        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 36    | RS 1200         | 38        | <i>C. fuscicollis</i>          | 18/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 37    | RS 1203         | 38        | <i>C. fuscicollis</i>          | 19/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 38    | RS 1204         | 35        | <i>C. fuscicollis</i>          | 19/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 39    | RS 1205         | 36        | <i>C. fuscicollis</i>          | 19/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 40    | RS 1207         | 39        | <i>C. fuscicollis</i>          | 19/04/2012        | Lagoa do Peixe        | –  |            |         | –            |
| 41    | RS 1208         | 35        | <i>C. fuscicollis</i>          | 00/01/1900        | Lagoa do Peixe        | +  | –          | NDV:160 | –            |
| 42    | RS 1210         | 38        | <i>C. fuscicollis</i>          | 19/04/2012        | Lagoa do Peixe        | –  |            |         | –            |

(Continues)

**TABLE 1** (Continued)

| Tubes | Sample ID | Ct | Scientific name             | Date       | Location       | HA | Flu detect | HI  | Subtype |
|-------|-----------|----|-----------------------------|------------|----------------|----|------------|-----|---------|
| 43    | RS 1212   | 39 | <i>Haematopus palliatus</i> | 19/04/2012 | Lagoa do Peixe |    |            |     | –       |
| 44    | RS 1216   | 39 | <i>C. fuscicollis</i>       | 19/04/2012 | Lagoa do Peixe | –  |            |     | –       |
| 45    | RS 1217   | 33 | <i>C. fuscicollis</i>       | 19/04/2012 | Lagoa do Peixe | –  |            |     | –       |
| 46    | RS 1218   | 39 | <i>Calidris alba</i>        | 19/04/2012 | Lagoa do Peixe | –  |            |     | –       |
| 47    | RS 1222   | 33 | <i>R. niger</i>             | 19/04/2012 | Lagoa do Peixe | –  |            |     | –       |
| 48    | RS 1239   | 37 | <i>C. fuscicollis</i>       | 22/04/2012 | Lagoa do Peixe | –  |            |     | –       |
| 49    | i536 DE   | –  | Shorebirds                  | 2012       | Delaware Bay   | +  | +          | 320 | H6N1    |
| 50    | i548 DE   | –  | Shorebirds                  | 2012       | Delaware Bay   | +  | +          | 640 | H6N1    |
| 51    | i556 DE   | –  | Shorebirds                  | 2012       | Delaware Bay   | +  | +          | 512 | H6N1    |
| 52    | i558 DE   | –  | Shorebirds                  | 2012       | Delaware Bay   | +  | +          | 512 | H6N1    |
| 53    | i595 DE   | –  | Shorebirds                  | 2012       | Delaware Bay   | +  | +          | 512 | H6N1    |

Ct, real-time RT-PCR cycle threshold; HA, hemagglutination assay; HI, hemagglutination inhibition assay.

verified as seronegative for influenza A (H1 and H3 subtypes) and influenza B by hemagglutinin inhibition (HI) tests, were inoculated with 1:10 dilutions of each viral stock in a 0.5-mL volume under light isoflurane anesthesia (actual doses ranged from  $10^{4.9}$  to  $10^{6.4}$ ). After 21 days, blood was collected to test for seroconversion by HI assay with turkey red blood cells.<sup>37,38</sup> The HI assay was conducted in triplicate, and the titers are reported as the geometric means. An HI titer of 40 or higher is indicative of seroconversion. All ferret studies were conducted in an ABSL-2 laboratory space in accordance with the St. Jude Children's Research Hospital Animal Care and Use Committee guidelines.

### 3 | RESULTS

#### 3.1 | Virus detection

We examined 1212 samples from wild birds. Cloacal and oropharyngeal swabs from each bird were collected and combined to facilitate the analysis (Table 2). Each year of the study featured 2 distinct events: the arrival of birds from the Northern Hemisphere in November/December (at the end of their migration south) and their departure in March/April (for their migration north). Most of the samples were collected from migratory bird species. The AIV M gene was detected in 48 of 1212 samples (3.96%), most of which came from White-rumped sandpipers (*Calidris fuscicollis*). With the exception of one sample collected in December 2010, all of the positive samples were collected during March or April, suggesting that the AIV prevalence builds up during the wintering period and that birds migrating northward carry a burden of virus with them. We detected the M gene at a higher rate in 2012 (9/163 birds, 5.5%) than in 2010 (2/371 birds, 0.53%) or 2011 (1/466 birds, 0.21%); no M gene was detected in 2009 (0/212 birds). The lack of positive samples collected during the austral summer also suggests that AIV is not maintained at high levels in resident bird populations.

#### 3.2 | Virus identification

Of the 48 samples positive for the M gene, 12 yielded enough sequence data for subtyping the virus. The subtypes identified were H2N2 (1), H6Nx (1), H6N1 (8), H9N2 (1), and H12N5 (1). The samples containing H6NX virus from the common tern (*Sterna hirundo*) and H12N5 virus from the red knot (*Calidris canutus*) were collected in the 2010 austral autumn; the sample containing H9N2 (*C. canutus*) was collected in the 2011 austral autumn, and the samples containing H6N1 and H2N2 viruses were collected in the 2012 austral autumn. All 8 of the H6N1 viruses from 2012 were isolated from samples collected from White-rumped sandpipers. We attempted to isolate virus from all M gene-positive samples but recovered only 2 of the H6N1 viruses collected in 2012 and only one H12N5 virus collected in 2010.

#### 3.3 | Genetic and antigenic analyses of isolated H6N1 viruses

In our genetic analysis, we were able to generate complete genomes from 9 samples. Eight of these genomes were obtained from swab material, and only one was obtained from an egg isolate. All 9 viruses had >99% amino acid homology with each other. A comparison of the virus surface glycoprotein gene sequences by BLAST analyses showed the HA glycoproteins of all the viruses to be similar to those of the H6 subtype (with 97% to 99% similarity), and their NA glycoproteins were similar to those of the N1 subtype (with 96% to 99% similarity). Both the HA and NA genes showed a strong relation to South American H6N1 lineages, as did their other genes (Figures 2–4).

The phylogenetic analysis used 1013 sequences available in the GenBank database, together with 5 new sequences of H6 viruses isolated the same year in Delaware Bay (North America) for comparison. The data showed that the South American isolates formed a unique clade; however, it was still similar to the North American viruses


**TABLE 2** Frequency of AIV RNA detection by bird family and by species sampled at Lagoa do Peixe National Park, southern Brazil. Status: SA, South American migrant; NA, North American migrant; R, resident

| Family            | Scientific name                    | Common name                     | Status | Sampling | Positives | AIV frequency (%) | AIV subtype         |
|-------------------|------------------------------------|---------------------------------|--------|----------|-----------|-------------------|---------------------|
| Spheniscidae      | <i>Spheniscus magellanicus</i>     | Magellanic penguin              | SA     | 151      | 0         | –                 | –                   |
| Procellariidae    | <i>Thalassarche chlororhynchos</i> | Atlantic yellow-nosed albatross | SA     | 15       | 0         | –                 | –                   |
|                   | <i>Thalassarche melanophrys</i>    | Black-browed albatross          | SA     | 21       | 0         | –                 | –                   |
|                   | <i>Macronectes giganteus</i>       | Southern giant petrel           | SA     | 6        | 0         | –                 | –                   |
|                   | <i>Pterodroma mollis</i>           | Soft-plumaged petrel            | SA     | 1        | 0         | –                 | –                   |
|                   | <i>Pterodroma incerta</i>          | Atlantic petrel                 | SA     | 2        | 0         | –                 | –                   |
|                   | <i>Pachyptila desolata</i>         | Antarctic prion                 | SA     | 1        | 0         | –                 | –                   |
|                   | <i>Pachyptila belcheri</i>         | Slender-billed prion            | SA     | 1        | 0         | –                 | –                   |
|                   | <i>Procellaria aequinoctialis</i>  | White-chinned petrel            | SA     | 18       | 0         | –                 | –                   |
|                   | <i>Calonectris borealis</i>        | Cory's shearwater               | NA     | 21       | 0         | –                 | –                   |
|                   | <i>Ardena gravis</i>               | Great shearwater                | SA     | 17       | 0         | –                 | –                   |
|                   | <i>Puffinus puffinus</i>           | Manx shearwater                 | NA     | 39       | 0         | –                 | –                   |
| Oceanitidae       | <i>Oceanites oceanicus</i>         | Wilson's storm-petrel           | SA     | 1        | 0         | –                 | –                   |
| Phalacrocoracidae | <i>Phalacrocorax brasilianus</i>   | Neotropic cormorant             | R      | 4        | 0         | –                 | –                   |
| Threskiornithidae | <i>Plegadis chihi</i>              | White-faced ibis                | R      | 1        | 0         | –                 | –                   |
| Ardeidae          | <i>Egretta thula</i>               | Snowy egret                     | R      | 2        | 0         | –                 | –                   |
| Accipitridae      | <i>Buteo</i> sp.                   | Hawks                           | –      | 1        | 0         | –                 | –                   |
|                   | <i>Buteogallus meridionalis</i>    | Savanna hawk                    | R      | 1        | 0         | –                 | –                   |
| Rallidae          | <i>Fulica leucoptera</i>           | White-winged coot               | R      | 1        | 0         | –                 | –                   |
|                   | <i>Gallinula galeata</i>           | Common gallinule                | R      | 1        | 0         | –                 | –                   |
| Charadriidae      |                                    |                                 |        |          |           | 5.26              |                     |
|                   | <i>Vanellus chilensis</i>          | Southern lapwing                | R      | 6        | 0         | –                 | –                   |
|                   | <i>Pluvialis dominica</i>          | American golden plover          | NA     | 31       | 0         | –                 | –                   |
|                   | <i>Pluvialis squatarola</i>        | Black-bellied plover            | NA     | 6        | 0         | –                 | –                   |
|                   | <i>Charadrius semipalmatus</i>     | Semipalmated plover             | NA     | 28       | 4         | 14.29             |                     |
|                   | <i>Charadrius collaris</i>         | Collared plover                 | R      | 3        | 0         | –                 | –                   |
|                   | <i>Charadrius falklandicus</i>     | Two-banded plover               | SA     | 1        | 0         | –                 | –                   |
|                   | <i>Charadrius modestus</i>         | Rufous-chested plover           | SA     | 1        | 0         | –                 | –                   |
| Haematopodidae    |                                    |                                 |        |          |           | 3.13              |                     |
|                   | <i>Haematopus palliatus</i>        | American oystercatcher          | R      | 32       | 1         | 3.13              |                     |
| Recurvitostridae  | <i>Himantopus melanurus</i>        | White-backed stilt              | R      | 35       | 0         | –                 | –                   |
| Scolopacidae      |                                    |                                 |        |          |           | 7.41              |                     |
|                   | <i>Tringa melanoleuca</i>          | Greater yellowlegs              | NA     | 5        | 0         | –                 | –                   |
|                   | <i>Tringa flavipes</i>             | Lesser yellowlegs               | NA     | 10       | 0         | –                 | –                   |
|                   | <i>Arenaria interpres</i>          | Ruddy turnstone                 | NA     | 10       | 0         | –                 | –                   |
|                   | <i>Calidris canutus</i>            | Red knot                        | NA     | 44       | 4         | 9.09              | H12N5 (1); H9N2 (1) |
|                   | <i>Calidris alba</i>               | Sanderling                      | NA     | 25       | 1         | 4.00              |                     |
|                   | <i>Calidris pusilla</i>            | Semipalmated sandpiper          | NA     | 14       | 0         | –                 | –                   |

(Continues)

TABLE 2 (Continued)


| Family        | Scientific name                  | Common name                  | Status | Sampling | Positives | AIV frequency (%) | AIV subtype        |
|---------------|----------------------------------|------------------------------|--------|----------|-----------|-------------------|--------------------|
|               | <i>Calidris fuscicollis</i>      | White-rumped sandpiper       | NA     | 370      | 34        | 9.19              | H6N1 (8); H2N2 (1) |
|               | <i>Calidris melanotos</i>        | Pectoral sandpiper           | NA     | 1        | 0         | –                 | –                  |
|               | <i>Calidris himantopus</i>       | Stilt sandpiper              | NA     | 2        | 0         | –                 | –                  |
|               | <i>Calidris subruficollis</i>    | Buff-breasted sandpiper      | NA     | 45       | 0         | –                 | –                  |
| Rostratulidae | <i>Nycticryphes semicollaris</i> | South American painted snipe | R      | 2        | 0         | –                 | –                  |
| Laridae       |                                  |                              |        |          |           | 1.71              |                    |
|               | <i>Larus maculipennis</i>        | Brown-hooded gull            | R      | 4        | 0         | –                 | –                  |
|               | <i>Larus dominicanus</i>         | Kelp gull                    | R      | 4        | 1         | 25.00             |                    |
|               | <i>Sternula superciliaris</i>    | Yellow-billed tern           | R      | 7        | 0         | –                 | –                  |
|               | <i>Sterna hirundo</i>            | Common tern                  | NA     | 83       | 1         | 1.20              | H6Nx (1)           |
|               | <i>Sterna hirundinacea</i>       | South American tern          | R      | 9        | 0         | –                 | –                  |
|               | <i>Sterna vittata</i>            | Antarctic tern               | SA     | 1        | 0         | –                 | –                  |
|               | <i>Sterna trudeaui</i>           | Snowy-crowned tern           | R      | 13       | 0         | –                 | –                  |
|               | <i>Thalasseus acutiflavus</i>    | Cabot's tern                 | R      | 3        | 0         | –                 | –                  |
|               | <i>Rynchops niger</i>            | Black skimmer                | R      | 110      | 2         | 1.82              |                    |
| Tyrannidae    | <i>Elaenia mesoleuca</i>         | Olivaceous elaenia           | R      | 1        | 0         | –                 | –                  |
| Furnariidae   | <i>Anumbius annumbi</i>          | Firewood-gatherer            | R      | 1        | 0         | –                 | –                  |
| Total         |                                  |                              |        | 1212     | 48        | 3.96              |                    |



**FIGURE 2** Phylogenetic analysis of influenza A virus based on the hemagglutinin gene. The trees are drawn to similar scales with branch lengths proportional to the evolutionary distance. The clusters of sequences from the Brazilian samples are indicated by branches colored red. Bootstraps values >50% were obtained in the analysis of 1000 replicates and are presented at the branching points

(Figures 2 and 3). Although the viruses from Brazilian samples were closely related to North American sequences, they formed a divergent clade consisting of only South American H6N1 sequences.

To assess the antigenic relatedness of our isolated H6N1 viruses to North American H6N1 viruses, we infected ferrets with each virus at various doses. Although we did not power the ferret study to assess



**FIGURE 3** Phylogenetic analysis of influenza A virus based on the neuraminidase gene. The trees are drawn to similar scales with branch lengths proportional to the evolutionary distance. Bootstraps values >50% were obtained in the analysis of 1000 replicates and are presented at the branching points. The clusters of sequences from the Brazilian samples are indicated by branches colored red. The clusters of sequences from Delaware Bay, isolated contemporaneously in the United States, are indicated by branches colored blue


pathogenicity or transmission, none of the infected ferrets showed any overt clinical signs, and all of them remained healthy and active throughout the study. HI analysis showed that the South and North American H6N1 viruses were antigenically indistinguishable (Table 3).

## 4 | DISCUSSION

In our 4 years of sampling at this important migratory bird wintering site, we found evidence of AIV circulation in consecutive years, with H6N1 viruses being predominant (mostly as a result of increased detection in the austral autumn of 2012). Rimondi et al<sup>13</sup> reported the existence of a South American H6 influenza virus lineage in Argentinian aquatic birds in 2008; most of the Argentinian virus genes were phylogenetically distinct from those of the North American lineages and had probably evolved independently and in isolation. However, one of the sequenced NA genes appeared to be derived from a virus of North American lineage, indicating that viruses of the 2 American genetic lineages do have opportunities to interact.<sup>13</sup>

The H6 subtype isolated in our study showed high sequence similarity to the Argentinian H6 virus isolated from a Rosy-Billed Pochard (*Netta peposaca*). This result is difficult to explain. The birds we sampled are known to migrate to North America, and we identified the highest prevalence of virus just before this northward migration, yet the viruses we isolated were clearly distinct from those detected in North America. These data suggest that the viruses isolated in the

austral autumn do not reach North America but are cleared from this population of birds en route. Our data also support a hypothesis that the source of AIV infection at Lagoa do Peixe National Park is local resident birds (or birds migrating to this area from other regions of South America), rather than birds migrating to the area from North America. We should note that 2012 saw the highest recorded incidence of virus detection in wild birds at Lagoa do Peixe, perhaps reflecting the great number of White-rumped sandpipers (*C. fuscicollis*) that congregated there that year. This species is among the most abundant of the shorebird species that migrate to South America. White-rumped sandpipers migrate from the Arctic to the Nova Scotia coast and then follow the Atlantic route to Patagonia, stopping over in Colombia, Venezuela, Suriname, and South Brazil. During the austral summer, flocks of up to 7000 White-rumped sandpipers use the Lagoa do Peixe National Park as a feeding, molting, and resting area. The flocks are greatest there during April/May, when individuals wintering in Patagonia fly north to join those that winter in Lagoa do Peixe National Park for feeding and remain there until migrating back to North America. In wintering and stopover areas in South America, the species may be found along with other migratory shorebirds from North America [eg, the red knot (*C. canutus*), sanderling (*Calidris alba*), and semipalmated plover (*Charadrius semipalmatus*)], as well as South American migrants [eg, the black skimmer (*Rynchops niger*) and Chilean flamingo (*Phoenicopterus chilensis*)] and resident species [eg, the American oystercatcher (*Haematopus palliatus*), southern lapwing (*Vanellus chilensis*), and collared plover (*Charadrius collaris*)].



**FIGURE 4** Phylogenetic analysis of influenza A virus based on internal gene sequences. The trees are drawn to similar scales with branch lengths proportional to evolutionary distance. Clades are broadly classified as Eurasian (green), North American (pink), and South American (red). Bootstraps values >50% were obtained in the analysis of 1000 replicates and are presented at the branching points. The phylogenetic trees were constructed using maximum likelihood methodology. Detailed phylogenetic trees with virus designations can be found in Figure S1 in the Supporting Information

**TABLE 3** HI analysis of North American and South American H6N1 viruses. The HI assays were conducted in triplicate, and the titers are reported as the geometric mean. An HI titer of 40 or higher is indicative of seroconversion

|                                                 | Sandpiper/Brazil/RS1149 |       | Sandpiper/Brazil/RS1151 |       | Shorebird/DE/548 |       | Shorebird/DE/595 |       |
|-------------------------------------------------|-------------------------|-------|-------------------------|-------|------------------|-------|------------------|-------|
|                                                 | F-930                   | F-928 | F-925                   | F-926 | F-921            | F-922 | F-919            | F-920 |
| A/White-rumped sandpiper/<br>Brazil/RS1149/2012 | 80                      | 80    | 40                      | 31.75 | 40               | 80    | 127              | 63.5  |
| A/Shorebird/DE/548/2012                         | 80                      | 80    | 80                      | 40    | 80               | 80    | 254              | 80    |
| A/Shorebird/DE/595/2012                         | 80                      | 80    | 40                      | 31.75 | 40               | 80    | 160              | 80    |

We have obtained samples from several other migratory shorebird species; however, among these others, none of the species that are most abundant in wet open-field habitats around the Lagoa do Peixe National Park [eg, the buff-breasted sandpiper (*Calidris subruficollis*), lesser yellowlegs (*Tringa flavipes*), and American golden plover (*Pluvialis dominica*)] have yielded detectable virus. Because these species prefer more open fields, they do not interact much with the other aquatic shorebird species, such as *C. canutus* and *C. fuscicollis*, in which AIVs have been detected. In the case of *S. hirundo*, we found AIV in one of the 2 individuals sampled, because this species is associated with the bodies of water where other aquatic species are found. The fact that all of our positive samples were collected in the austral autumn suggests that the infection builds up during the wintering period, leading to a local outbreak phenomenon, as has been suggested to occur at Delaware Bay.<sup>15</sup>

In 2014, researchers reported the result of a study of the blue-winged teal and AIV infections in North and South America. They found no evidence of South American lineage genes in samples from these birds, although this teal species is widely distributed in South America, and they suggested that viral gene flow may be limited in some species as a result of geographic barriers.<sup>21</sup> The major caveat to this interpretation is that we have enough genetic information for comparison purposes from only the H6N1 viruses, and it is possible that both the North and South American lineages circulate at Lagoa do Peixe National Park. Consequently, we are attempting to gather more genetic information from other samples through enrichment procedures. The H11N9 virus is of particular interest, as this subtype has been found in ruddy turnstones (*Arenaria interpres*) in the Amazon region of Brazil, and it has a strong relation to North American AIV lineages, clearly showing that virus exchange between the 2 continents does occur.<sup>20</sup>

In South America, AIV has been confirmed in wild birds in multiple years in Argentina (H13N9, H1N1, H6N2, and H9N2),<sup>13,39,40</sup> Bolivia (H7N3),<sup>41</sup> (H7N3),<sup>42</sup> Colombia (H5N2),<sup>19</sup> Peru (H3N8, H4N5, H10N9, and H13N2),<sup>42,43</sup> and Brazil (H11N9).<sup>20</sup> However, the H6N1 subtype has not previously been identified in shorebirds. Despite the predominance of H6N1 virus, other important subtypes, such as H2N2, H9N2, and H12N5, were identified in birds in the same region during the same collection period. This was the first time that the H2N2 and H12N5 subtypes were reported in South America.

Most recent outbreaks of HPAI in North American poultry clearly resulted from the introduction of low-pathogenic precursors into poultry populations and their subsequent evolution to a highly

pathogenic form in large poultry farms.<sup>44</sup> More recently, however, a HPAI virus spread from Asia, most likely through wild bird movement, into poultry in North America<sup>45</sup>. Despite Brazil being considered free of HPAI, surveillance of AIV in migratory birds remains incomplete in that country, and AIV data remain scarce. Although H6N1 viruses are considered low-pathogenic, an isolate of this subtype, albeit genetically distinct from our isolates, has been isolated from a human infection in Taiwan.<sup>46</sup> Regardless of the zoonotic potential of the viruses found in Lagoa do Peixe National Park, increased surveillance in all animal species in the country is prudent.

The aim of the present work was to assess the importance of Lagoa do Peixe National Park as a source of AIV and to determine the nature of any viruses identified. We were able to detect AIV in the avian community over multiple years, with multiple subtypes being identified. Our phylogenetic analysis demonstrated the high genomewide similarity of our H6N1 isolates and their close relation to South American H6 viruses. Further studies are needed to explain the data showing the presence of viruses of South American lineage in birds that were beginning to migrate north. The avian surveillance program in Brazil is not continuous, but it should be. Our findings are potentially important for surveillance of AIV in Brazil, and they represent a valuable contribution to the study of AIV viruses in South America.

## ACKNOWLEDGEMENTS

We thank the National Center of Research and Conservation of Wild Birds (CEMAVE-ICMBio/MMA), the Universidade Vale do Rio do Sinos (UNISINOS), the field team of the Department of Microbiology of the University of São Paulo, the Secretaria de Estado da Agricultura, Pecuária e Abastecimento (SEAPA) RS, and the Ministério da Agricultura (MAPA) for their support of the field work and logistics. We would also like to extend our appreciation to Keith Laylock for revisions and suggestions to the manuscript. We would like to thank Dr. Sook-San Wong for conducted statistical analyses an earlier version of this manuscript.

## FIELD TEAM MEMBERS

We thank the Field team: Ana L. Bezerra, Aparecida B. Basler, César R. Santos, Elisa S. Petersen, Emily T. Moura, Fernanda L. Valls, Gabriel V. Vier, Guilherme P. Cauduro, Hélio A. A. Fracasso, Janete F. M. Scherer, Jaqueline Brummelhaus, Jéssica Sartor, Julia V. G. Finger,

Laura A. Lindenmeyer de Sousa, Liana C. Rossi, Lucas Kruger, Marina M. M. Seixas, Carla Meneguín, Alexandre Farias de Lemos, Priscilla S. Kiscorski, Roberta C. Piuco, and Suzana Seibert.

## CONFLICT OF INTEREST

None of the authors have a conflict of interest.

## BIOGRAPHICAL SKETCH

Dr. Jansen Araujo has a Ph.D. in Microbiology and is the coordinator of the field team in the Virology Laboratory at the Biomedical Science Institute at the University of São Paulo. His research interests include eco-epidemiological studies of wildlife and emerging infectious diseases with zoonotic potential in Brazil.

## STATEMENT

Wild birds are the natural reservoir for avian influenza viruses (AIVs). Brazil provides stopover and wintering sites for migratory birds. We wished to investigate the potential for influenza A viruses to be transported by birds that migrate annually along with resident species and to determine the subtypes involved and their potential for reassortment. In Brazil, data about AIV subtypes are scarce, and few groups have worked with AIV in wild birds. At present, only AIV H11N9 has been isolated from the Amazon region (in 2014). This study was conducted to elucidate which AIV subtypes were present at an important stopover and wintering site of wild birds in Brazil and the genetic relatedness between viruses in the Northern and Southern Hemispheres. In addition to performing isolation and next-generation sequencing (NGS), we used a ferret model to generate influenza A antiserum to compare antigenic differences in influenza viruses from the Northern and Southern Hemispheres.

## ORCID

Jansen Araujo  <http://orcid.org/0000-0002-7250-3024>

## REFERENCES

- Webster RG, Bean WJ, Gorman OT, Chambers TM, Kawaoka Y. Evolution and ecology of influenza A viruses. *Microbiol Rev.* 1992;56:152-179.
- Stallknecht DE. Impediments to wildlife disease surveillance, research, and diagnostics. *Curr Top Microbiol Immunol.* 2007;315:445-461.
- Gaidet N, Caron A, Cappelle J, et al. Understanding the ecological drivers of avian influenza virus infection in wildfowl: a continental-scale study across Africa. *Proc Biol Sci.* 2012;279:1131-1141.
- Olsen B, Munster VJ, Wallensten A, Waldenström J, Osterhaus ADME, Fouchier RA. Global patterns of influenza A virus in wild birds. *Science.* 2006;312:384-388.
- Colwell MA. *Shorebird Ecology, Conservation, and Management.* Oakland, CA: University of California Press; 2010.
- Petry MV, Petersen EDS. Riqueza e abundância sazonal de Laridae e Sternidae no litoral médio do Rio Grande do Sul, Brasil. *Rev Bras Ornitol.* 2011;19:493-497.
- Mader A, Petry MV, Efe MA. Litoral médio do Rio Grande do Sul. In: *Conservação de Aves Migratórias Neárticas no Brasil.* Belém, Brazil: Conservação Internacional; 2011:325-330.
- CEMAVE-ICMBio. *Relatório Anual de Rotas e Áreas de Concentração de Aves Migratórias no Brasil.* Brasília: Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio); 2014.
- Harrison NM, Whitehouse MJ, Madureira LASP. List observations of the under-described avifauna of the Mostardas Peninsula, Rio Grande do Sul, Brazil. *Check List.* 2013;9:391-399.
- Xu K, Ferreri L, Rimondi A, et al. Isolation and characterization of an H9N2 influenza virus isolated in Argentina. *Virus Res.* 2012;168:41-47.
- Krauss S, Obert CA, Franks J, et al. Influenza in migratory birds and evidence of limited intercontinental virus exchange. *PLoS Pathog.* 2007;3:e167.
- Munster VJ, Fouchier RAM. Avian influenza virus: of virus and bird ecology. *Vaccine.* 2009;27:6340-6344.
- Rimondi A, Xu K, Craig MI, et al. Phylogenetic analysis of H6 influenza viruses isolated from rose-billed pochards (*Netta peposaca*) in Argentina reveals the presence of different HA gene clusters. *J Virol.* 2011;85:13354-13362.
- Pearce JM, Ramey AM, Ip HS, Gill RE. Limited evidence of trans-hemispheric movement of avian influenza viruses among contemporary North American shorebird isolates. *Virus Res.* 2010;148:44-50.
- Bahl J, Krauss S, Kühnert D, et al. Influenza A virus migration and persistence in North American wild birds. *PLoS Pathog.* 2013;9:e1003570.
- D'Amico VL, Bertellotti M, Baker AJ, Diaz La. Exposure of red knots (*Calidris canutus rufa*) to select avian pathogens; Patagonia, Argentina. *J Wildl Dis.* 2007;43:794-797.
- Ghera BM, Sovero MM, Icochea E, et al. Isolation of low-pathogenic H7N3 avian influenza from wild birds in Peru. *J Wildl Dis.* 2011;47:792-795.
- González-Reiche AS, Morales-Betoulle ME, Alvarez D, et al. Influenza A viruses from wild birds in Guatemala belong to the North American lineage. *PLoS ONE.* 2012;7:e32873.
- Karlsson EA, Ciuderas K, Freiden PJ, et al. Prevalence and characterization of influenza viruses in diverse species in Los Llanos, Colombia. *Emerg Microb Infect.* 2013;2:e20.
- De Araujo J, de Azevedo SM, Gaidet N, et al. Avian influenza virus (H11N9) in migratory shorebirds wintering in the Amazon Region, Brazil. *PLoS ONE.* 2014;9:e110141.
- Ramey aM, Walther P, Link P, et al. Optimizing surveillance for South American origin influenza A viruses along the United States Gulf Coast through genomic characterization of isolates from blue-winged teal (*Anas discors*). *Transbound Emerg Dis.* 2016;63:194-202.
- Curd E, Pollinger J, Toffelmier E, Smith T. Rapid influenza A detection and quantitation in birds using a one-step real-time reverse transcriptase PCR and High Resolution Melting. *J Virol Methods.* 2011;176:125-130.
- Shortridge KF, Butterfield WK, Webster RG, Campbell CH. Isolation and characterization of influenza A viruses from avian species in Hong Kong. *Bull World Health Organ.* 1977;55:15-20.
- Kawaoka YN. *Influenza Virus Methods and Protocols.* New York, NY: Springer; 2012.
- Das A, Spackman E, Thomas C, Swayne DE, Suarez DL. Detection of H5N1 high-pathogenicity avian influenza virus in meat and tracheal samples from experimentally infected chickens. *Avian Dis.* 2008;52:40-48.
- Aymard-Henry M, Coleman MT, Dowdle WR, Laver WG, Schild GC, Webster RG. Influenzavirus neuraminidase and neuraminidase-inhibition test procedures. *Bull World Health Organ.* 1973;48:199-202.
- Palmer DFDW, Coleman MT, Schild GC. *Advanced Laboratory Techniques for Influenza Diagnosis.* Immunology series no. 6. Atlanta: Centers for Disease Control, US Department of Health, Education and Welfare; 1975.

28. Webster RG, Isachenko VA, Carter M. A new avian influenza virus from feral birds in the USSR: recombination in nature? *Bull World Health Organ.* 1974;51:325-332.
29. Bowman AS, Sreevatsan S, Killian ML, et al. Molecular evidence for interspecies transmission of H3N2pM/H3N2v influenza A viruses at an Ohio agricultural fair, July 2012. *Emerg Microb Infect.* 2012;1:e33.
30. Delangue J, Roca Sanchez Y, Piorkowski G, et al. Viral aetiology influenza like illnesses in Santa Cruz, Bolivia (2010-2012). *Virology.* 2014;11:35.
31. Life Technologies. ION Torrent. 2014. <http://tools.lifetechnologies>.
32. Zhou B, Donnelly ME, Scholes DT, et al. Single-reaction genomic amplification accelerates sequencing and vaccine production for classical and swine origin human influenza A viruses. *J Virol.* 2009;83:10309-10313.
33. Depew J, Zhou B, McCarrison JM, et al. Sequencing viral genomes from a single isolated plaque. *Virology.* 2013;10:181.
34. Djikeng A, Halpin R, Kuzmickas R, et al. Viral genome sequencing by random priming methods. *BMC Genom.* 2008;9:5.
35. Zhang Z, Schwartz S, Wagner L, Miller W. A greedy algorithm for aligning DNA sequences. *J Comput Biol.* 2000;7:203-214.
36. Drummond AJ, Suchard MA, Xie D, Rambaut A, Bayesian Phylogenetics with BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* 2012;29:1969-1973.
37. Hurt AC, Vijaykrishna D, Butler J, et al. Detection of evolutionarily distinct avian influenza A viruses in Antarctica. *MBio.* 2014;5:e01098-14.
38. Jones JC, Baranovich T, Marathe BM, et al. Risk assessment of H2N2 influenza viruses from the avian reservoir. *J Virol.* 2014;88:1175-1188.
39. Alvarez P, Mattiello R, Rivallier P, et al. First isolation of an H1N1 avian influenza virus from wild terrestrial non-migratory birds in Argentina. *Virology.* 2010;396:76-84.
40. Pereda AJ, Uhart M, Perez AA, et al. Avian influenza virus isolated in wild waterfowl in Argentina: evidence of a potentially unique phylogenetic lineage in South America. *Virology.* 2008;378:363-370.
41. Williams RAJ, Segovia-Hinostroza K, Ghersi BM, Gonzaga V, Peterson AT, Montgomery JM. Avian Influenza infections in nonmigrant land birds in Andean Peru. *J Wildl Dis.* 2012;48:910-917.
42. Ghersi BM, Blazes DL, Icochea E, et al. Avian influenza in wild birds, central coast of Peru. *Emerg Infect Dis.* 2009;15:935-938.
43. Spackman E, McCracken KG, Winker K, Swayne DE. An avian influenza virus from waterfowl in South America contains genes from North American avian and equine lineages. *Avian Dis.* 2007;51:273-274.
44. World Organisation for Animal Health (OIE). *Highly Pathogenic Avian Influenza, United States of America.* <http://www.oie.int/animal-health-in-the-world/update-on-avian-influenza/2015/>. Accessed January 19, 2015.
45. Lee D-H, Torchetti MK, Winker K, Ip HS, Song C-S, Swayne DE. Intercontinental spread of Asian-Origin H5N8 to North America through Beringia by Migratory Birds. *J Virol.* 2015;89:6521-6524.
46. Wei S-H, Yang J-R, Wu H-S, et al. Human infection with avian influenza A H6N1 virus: an epidemiological analysis. *Lancet Respir Med.* 2013;1:771-778.

## SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

**How to cite this article:** Araujo J, Petry MV, Fabrizio T, et al. Migratory birds in southern Brazil are a source of multiple avian influenza virus subtypes. *Influenza Other Respi Viruses.* 2018;12:220-231. <https://doi.org/10.1111/irv.12519>