

Original Article

Long-term affected flat oyster (*Ostrea edulis*) haemocytes show differential gene expression profiles from naïve oysters in response to *Bonamia ostreae*

P. Ronza^a, A. Cao^b, D. Robledo^c, A. Gómez-Tato^d, J.A. Álvarez-Dios^e, A.F.M. Hasanuzzaman^f, M.I. Quiroga^a, A. Villalba^{b,g}, B.G. Pardo^{h,*}, P. Martínez^{h,*}

^a Departamento de Anatomía, Producción Animal e Ciencias Clínicas Veterinarias, Facultad de Veterinaria, Universidade de Santiago de Compostela, 27002 Lugo, Spain

^b Centro de Investigacións Mariñas (CIMA), Consellería do Mar, Xunta de Galicia, 36620 Vilanova de Arousa, Spain

^c The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Midlothian EH25 9RG, UK

^d Departamento de Matemáticas, Facultad de Matemáticas, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain

^e Departamento de Matemática Aplicada, Facultad de Matemáticas, Universidade de Santiago de Compostela, 15781 Santiago de Compostela, Spain

^f Fisheries and Marine Resource Technology Discipline, Khulna University, Khulna 9208, Bangladesh

^g Departamento de Ciencias de la Vida, Universidad de Alcalá, 28871 Alcalá de Henares, Spain

^h Departamento de Zoología, Genética y Antropología Física, Facultad de Veterinaria, Universidade de Santiago de Compostela, 27002 Lugo, Spain

ARTICLE INFO

Keywords:

Bonamiosis
Parasite
Oligo-microarray
Immune response
Mollusc
Bivalve

ABSTRACT

European flat oyster (*Ostrea edulis*) production has suffered a severe decline due to bonamiosis. The responsible parasite enters in oyster haemocytes, causing an acute inflammatory response frequently leading to death. We used an immune-enriched oligo-microarray to understand the haemocyte response to *Bonamia ostreae* by comparing expression profiles between naïve (NS) and long-term affected (AS) populations along a time series (1 d, 30 d, 90 d). AS showed a much higher response just after challenge, which might be indicative of selection for resistance. No regulated genes were detected at 30 d in both populations while a notable reactivation was observed at 90 d, suggesting parasite latency during infection. Genes related to extracellular matrix and protease inhibitors, up-regulated in AS, and those related to histones, down-regulated in NS, might play an important role along the infection. Twenty-four candidate genes related to resistance should be further validated for selection programs aimed to control bonamiosis.

1. Introduction

The European flat oyster *Ostrea edulis* has traditionally been a highly appreciated marine resource for human nutrition. Overexploitation has led to the exhaustion of most flat oyster beds through Europe in the last century, and its production has been mostly sustained through aquaculture [42]. However, flat oyster farming has been affected at specific areas by disease outbreaks, such as the “shell disease” caused by the fungus *Ostracoblabe implexa* [1], or at larger areas involving different countries, such as infections with the protist *Marteilia refringens* [7,32] and *Bonamia ostreae* [72]. The latter is a parasite widely distributed throughout the European coast, constraining European flat oyster production [11,12,20,21,43]. Thus, controlling bonamiosis is required for a successful recovery of the European flat oyster production, but its eradication from affected areas seems not to be a simple task [33,45,87], and the use of resistant oyster stocks appears to be the most promising strategy [12,19,46,63]. Successful breeding programs to produce disease resistant oysters, such as those against summer

mortality or infections with *Haplosporidium nelsoni*, *Perkinsus olseni*, *Roseobacter crassostreae*, *Marteilia sydneyi* or *B. ostreae*, have been based on selecting broodstock among survivors after long exposure in heavily affected areas [6,13,18,26,27,46,76]. This procedure relies on the cumulative benefit of successive generations of selection to achieve profitable increased resistance levels. Considering that the maximal oyster mortality due to *B. ostreae* occurs when the oysters have almost reached market-size [4,10,12], the evaluation of each generation in the field to select breeders and obtaining the next generation should take at least 4–5 years. Such procedure would need a long-term program to reach acceptable oyster survival in affected areas [46]. Availability of molecular markers associated with resistance would avoid long field evaluation of breeders, thus shortening the process and making it more efficient. Such resistance markers have been successfully found for some diseases in mollusks [64,66,68,75,86], and even a specific mutation conferring resistance to perkinsosis has been identified in the eastern oyster [37].

Mollusc immunity relies on innate cellular and humoral

* Corresponding authors.

E-mail address: paulino.martinez@usc.es (P. Martínez).

mechanisms, where haemocytes, the circulating cells present in haemolymph and infiltrating through other tissues, play a key role as responsible cells for phagocytosis of foreign particles, among other functions. *B. ostreae* is an intracellular parasite able to proliferate and survive inside the oyster haemocytes, inducing a strong inflammatory response and disturbing the immune capacity of the host [8]. Oyster response involves cytotoxic mechanisms as well as non-oxidative and oxidative pathways, including lysosomal enzymes and reactive oxygen/nitrogen intermediates. The haemolymph serum contains humoral defense factors, such as soluble lectins, hydrolytic enzymes and antimicrobial peptides [2,3]. The evaluation of immune-gene expression profiles of oyster and their interactions along the process of infection is crucial to understand the response to bonamiosis and requires the development of large-scale genomic resources and appropriate tools to identify genes and signaling networks related to disease resistance. Recently, a comprehensive transcriptomic haemocyte database has been used to design and validate an oligo-microarray to assess gene expression profiles of flat oyster in response to *B. ostreae* [70].

In this study, we used this oligo-microarray to identify genes and pathways differentially expressed along a time series after challenge with *B. ostreae* in two flat oyster stocks with differential susceptibility to bonamiosis: i) a naïve stock from a free-bonamiosis area; and ii) a stock from a long-term affected area, which had shown some resistance to *B. ostreae*. The comparison of the pattern of response along the infection process and the differentially expressed genes and pathways between both stocks provided candidate genes and pathways to be validated for their application on breeding programmes for a more efficient selection.

2. Materials and methods

2.1. Biological material and challenge

Flat oysters were collected from Limfjord (Denmark) and Ortigueira (NW Spain) in 2010. Surveillance since 1996 has shown Limfjord to be a *Bonamia*-free area until 2014, when *B. ostreae* was detected for the first time in November [48]. Thus the Limfjord oyster population was still naïve regarding bonamiosis in 2010 (NS: naïve stock onwards). Conversely, bonamiosis affects oysters in Ortigueira since early 1980's (AS: long-term affected stock onwards) and oysters from this bed are more resistant to *B. ostreae* than oysters from other geographic origins, as shown in an experiment where families from Ortigueira showed significantly less prevalence and lower infection intensity [12].

Oysters of both origins were experimentally challenged with *B. ostreae* and compared with their corresponding non-challenged controls. The challenge was performed by individually immersing oysters in beakers containing an aerated suspension of 300,000 *B. ostreae* cells in filtered seawater for 24 h at the Centro de Investigaciones Mariñas. The *B. ostreae* cells for the challenge were isolated from heavily infected oysters collected in Lough Foyle (Ireland), following the procedure described by Mialhe et al. [53]. After challenge, oysters were kept in tanks with running, filtered (1 µm) seawater plus continuous (pumped) supply of mixed cultured algae until the end of the experiment. Haemolymph samples were taken at three different times after the challenge (1, 30 and 90 days post-challenge, dpc) considering previous information on the progression of the infection [11]; as much haemolymph as possible (from 0.2 ml to 1.5 ml) was withdrawn from the adductor muscle of each oyster with a 21 gauge needle screwed to a 2 ml cold syringe. The haemolymph was immediately poured into a cold vial and kept in crushed ice to avoid haemocyte clumping. Haemolymph samples were centrifuged (800 g, 10 min, 4 °C) and the cell fractions (pellets) resuspended in RNA later; then suspensions, after one night at 4 °C, were stored at –20 °C until further processing. Five biological replicates for each condition (control and challenged), stock (NS and AS) and sample times (1, 30 and 90 dpc) were used.

2.2. RNA isolation, cDNA synthesis and library construction

RNA was individually extracted using RNeasy Mini Kit (Qiagen, Germany) following manufacturer's instructions with slight modifications, including two additional RPE washes and DNase digestion. Quantity was determined using a Nanodrop spectrophotometer (Nanodrop Technologies, USA) and quality (RNA integrity number, RIN) using a Bioanalyzer (Agilent Technologies, USA). All samples showed high RNA quality (RIN > 8.0) and consequently, they were further processed for microarray analysis. For each experimental condition, the control group was pooled by combining equimolar RNA quantities of the five biological replicates, while the RNA of each challenged oyster was processed individually (five replicates). Next, 50 ng of total RNA were labeled using the Low Input Quick Amp Labeling Kit, One-Color (Cy3; Agilent Technologies).

2.3. Microarray hybridisation and scanning

All samples (control pools and challenged oysters) were hybridized individually in a custom 8 × 15 K Agilent oligo-microarray at the Universidade de Santiago de Compostela (USC) Genomics Platform following the protocols for Agilent one-color Gene Expression Analysis. This custom microarray was mainly based on OedulisDB, an *Ostrea edulis* database obtained from oyster haemocytes challenged with *B. ostreae* [70]. A total of 36 microarrays were used for hybridizations of 6 control (RNA pools from 2 stocks × 3 sample times) and 30 challenged oysters (5 biological replicates × 2 stocks × 3 sample times). Copy RNA (cRNA) was hybridized overnight during 17 h at 65 °C and washed with the corresponding buffers on the following day. Hybridized slides were scanned using an Agilent G2565B microarray scanner (Agilent Technologies). The scanner images were segmented with the protocol GE1-v5.95 from the Agilent Feature Extraction Software (v9.5) and to avoid saturation in the highest intensity range extended dynamic range was implemented. Agilent Feature Extraction produced the raw data for further preprocessing. The processed signal (gProcessed-Signal) value was the chosen parameter for statistical analysis as recommended [55]. Feature quality filtering was performed following Millán et al. [54].

2.4. qPCR validation

Quantitative real-time PCR (qPCR) was performed to validate microarray data. A set of 16 genes was selected following the random stratified procedure proposed by Miron et al. [56] with some modifications as described by Millán et al. [55]. A total of 39 cases were selected covering the fold change (FC) range of the experiment and including at least one gene per microarray and two cases per gene. The glyceraldehyde-3-phosphate dehydrogenase (*gapdh*) was chosen as reference gene following literature recommendations [59].

Primers for qPCR were designed using *Primer Express* Software v2.0 (Applied Biosystems) with default settings (Table S1). The same RNA samples used for microarray hybridizations were used for qPCR. The concordance between microarray and qPCR data was assessed using the Pearson correlation and the null hypothesis of no differences between both methods tested by *t*-test ($P \leq 0.05$).

2.5. Data analysis

Normalization using all microarray data was done by the Quantile method implemented in the Limma R package [78]. All further statistical analyses were performed on the software Multiple Experiment Viewer (MeV) version 4.9.0, after a log₂ transformation of the fluorescence intensity values. Fold change (FC) values were obtained as the log₂ (treatment/control) after averaging technical replicates within each microarray. Principal component analysis (PCA) was computed with default parameters. Hierarchical Clustering (HCL) trees were constructed using Pearson correlation and default parameters (100

iterations). The SOTA program (Self Organizing Tree Algorithm; [17,91]) was applied to identify groups of genes with similar expression patterns using Pearson correlation and internal cell diversity at $P < 0.05$. Regulated genes (RGs) at each experimental condition were detected by Significance Analysis of Microarrays test (SAM; [85]) with a significance threshold of 5% False Discovery Rate (FDR). A two-way ANOVA considering stock and time as factors was performed to detect differentially expressed genes (DEGs) between stocks and across time ($P < 0.001$).

3. Results & discussion

3.1. Experimental challenge and microarray evaluation

The gene expression profiles of *O. edulis* haemocytes challenged with *B. ostreae* were evaluated using an *in vivo* experimental infection, by immersing the oyster in beakers containing a suspension of parasitic cells. These conditions mimic the host-parasite interactions occurring in a natural scenario, so constituting an approach aimed to a more comprehensive evaluation of haemocytes response than other published approaches consisting of injecting the parasite into the oyster adductor muscle [49], or challenging haemocytes *in vitro* [30,60–62]. This approach enabled to identify a set of 715 regulated genes (RG) in infected oysters regarding controls (FC \neq 0) along the time series evaluated (Table S2). Moreover, the availability of AS and NS stocks allowed to compare expression profiles of oysters showing some degree of resistance to the parasite with those that never had contact with it, and thus, identify 837 differentially expressed genes (DEG) between stocks (Table S3).

Gene profiles were analyzed using an Agilent custom 8×15 K oligo-microarray previously validated by Pardo et al. [70]. This microarray contains 6882 different oligos and two replicates per oligo, which enabled controlling technical noise. The microarray platform Agilent-038418 and the data presented in this publication have been deposited in the NCBI's Gene Expression Omnibus (GEO, <http://www.ncbi.nlm.nih.gov/geo/>) and are available under accession number XXX. Comparison of expression data between microarray and qPCR for selected genes was highly consistent except for two values among the 39 comparisons performed, corresponding to the genes predicted HD phosphohydrolase family protein (ID: 4160e) and tripartite motif-containing protein 2 (4677e). The Pearson correlation coefficient increased from 0.769 to 0.929 after excluding these two cases ($P < 0.001$ for both correlations), and no significant differences were detected between both data sets including all cases (Wilcoxon test $P = 0.137$). Similar results have been reported with custom oligo-microarrays in other aquaculture species [25,54,77].

3.2. Gene expression profiles

Hierarchical clustering (HCL) was applied to check the grouping of biological replicates using RGs and considering the two main experimental factors: time after challenging (T1, T2 and T3) and stock evaluated (NS and AS). The codes NT1, NT2, NT3, AT1, AT2 and AT3 combining time and stock will be used onwards to refer to each experimental condition. Most biological replicates clustered in the expected group according to the experimental condition (Fig. 1), especially for T1 and T3 in both stocks, which supports the fact that experimental factors (time and stock) override biological variation and that technical noise was kept under control in our hybridization protocol. T2 showed a slightly more erratic pattern because, as shown below, no response to the parasite was detected at this time.

A certain proportion (6.6%) of RGs or DEGs showed the same annotation (Table S4), reflecting the redundancy of the oyster database [70]. Probes with the same annotation usually showed correlated profiles along the experimental conditions tested (88.2%; Table S4). For example, the three genes annotated as peptidylglycine alpha-

hydroxylating monooxygenase ($r > 0.990$) or the two annotated as collectin-12 ($r = 0.999$) showed highly significant correlation between them, suggesting that they are part of the same gene resulting from limitations of the transcriptome assembly process [70] or they represent true duplicated genes with the same functional role. In the remaining 11.8% cases, no significant correlation or even a significant negative correlation was observed for the same annotation, suggesting that they are duplicated genes belonging to different paralogs with refined functions (neofunctionalization) or they represent alternative splice variants of the same gene. The most remarkable cases were those related to calcium and integrin-binding protein ($r = -0.647$) and complement C1q-like protein ($r = -0.493$), which showed a significant negative correlation value ($P < 0.001$). It has been suggested that stress- and immune-related genes suffered a notable genomic expansion in molluscs due to their feeding by filtering in an enriched microbial environment, as reported for *Crassostrea gigas* [93]. This might explain our observations, especially considering that these genes were modulated after a parasite challenge.

A total of 715 RGs were detected after *Bonamia* challenge in at least one of the 6 experimental conditions (2 stocks \times 3 times) tested (Table S2). While an important amount of RGs were detected at 1 dpc and 90 dpc, none was observed at 30 dpc either in NS or in AS stocks, which suggests some kind of latency during the infection progress. It is well known that the infection by *B. ostreae*, an intracellular parasite, presents a long pre-patent period, from one to several months, which is a major issue still unclear regarding the course of this parasitosis [11,19,57,84]. In general, the AS stock showed a more intense response than the NS (536 vs 206 RGs, respectively) and a sharply different temporal pattern was observed between both stocks (Table 1). The response was much more intense at 1 dpc in AS (AT1: 507 RGs vs AT3: 31 RGs), while in NS more RGs were detected at 90 dpc (NT1: 86 vs NT3: 127). The huge difference observed at 1 dpc (six times more RGs in AS than NS) suggests that natural selection in the bonamiosis-endemic area might have favored a more intense and quicker response to the parasite. Interestingly, very few RGs were shared between times (1 dpc and 90 dpc) or stocks (NS vs AS), the highest common set being detected between NT1 and AT1 (12 RGs), which suggests a specific response to the challenge at each time and stock (Fig. 2).

In general, significant FC values were moderate: mean $FC > 0 = 1.204$; mean $FC < 0 = -1.722$; 15.8% RGs showing $2 < FC < -2$ (Table S5). FC ranged between 3.943 and -7.486 for the von Willebrand factor type A domain containing protein (NT3) and leucine-rich repeat-containing G-protein coupled receptor 6 (NT1), respectively; down-regulated genes were more frequent than up-regulated among those showing the more extreme FCs (41 vs 18); and the AS stock showed a higher proportion of genes in this list than NS, especially among the up-regulated ones (14 out of 18) (Table S5).

The two-way ANOVA identified a total of 837 significant DEGs between stocks, 250 DEGs across time and 637 DEGs due to interaction of stock \times time ($P < 0.001$) (Table S3). The large number of genes regarding interaction between both factors (stock \times time) agrees with the particular behavior of each stock across time highlighted in the analysis of RGs. Accordingly, no significantly enriched pathways were detected in the 250 DEGs across time, and thus we focused our attention on the 837 DEGs between stocks as the most interesting to extract relevant information for future selection programs. Among them, 196 were in common with the RGs list (Table S2). The clustering of expression profiles of the 43 annotated genes showing the highest differences between both stocks ($SD > 0.8$) is represented in Fig. 3. Among them, several genes related to histones and extracellular matrix (ECM) proteins, up-regulated in the AS stock and down-regulated in NS, were identified.

The application of SOTA analysis to identify clusters of genes showing correlated profiles rendered four RG groups in each of the two stocks (Fig. S1, Table S6) and five DEG clusters (Fig. S2, Table S7).




Fig. 3. Hierarchical clustering of the differentially expressed genes (ANOVA, $P < 0.001$) showing the highest difference (standard deviation > 0.8) between the naïve (N) and long-term affected (A) flat oyster (*O. edulis*) stocks after challenge with *B. ostreae* along a time series (T1: 1 day post-challenge (dpc), T2: 30 dpc, T3: 90 dpc).

stimulus, reflecting the expression of several G-protein coupled receptors (GPCRs) and toll-like receptor (TLR) 1 in response to the parasite. GPCRs are essential in several signaling pathways regulating a wide range of cell responses, including the development of the immune response through activation, migration and adhesion of the cells of the immune system [36,82]. On the other hand, TLRs are well known as key components of the innate immune system that recognizes pathogen-associated molecular patterns (PAMPs), and a prominent role of TLR signaling was demonstrated in infection-induced haemocyte activation in *C. gigas* [94]. We found several GPCRs up-regulated in AT1 (Table S2), mostly classified as rhodopsin-like receptors, including orexin receptor type 2, cholecystokinin A receptor, cephalotocin receptor 1, neuropeptide FF receptor 2, growth hormone secretagogue receptor type 1 and prostaglandin E2 receptor 4. The involvement of G-protein signaling genes in the pathogenesis of bonamiosis in flat oyster was previously hypothesized [49,50]. Furthermore, in other invertebrates like the mosquito *Anopheles gambiae*, rhodopsin-like GPCRs have demonstrated an important role in response to the infection with the African malaria parasite *Plasmodium falciparum* [52], and specifically, prostaglandin E2 receptor 4 has been involved in the innate immune response of *Crassostrea hongkongensis* to different pathogens [74]. Two genes encoding alpha-adrenergic receptors were, by contrast, down-regulated in AT1 (Table S2). Interestingly, those GPCRs have been associated with immunomodulatory functions in the scallop *Chlamys farreri*, and their activation was shown to repress haemocyte phagocytic and antibacterial capacities [95]. In two recent transcriptomic studies on mollusks, GPCRs and TLRs have been associated with defense response of *C. gigas* to the ostreid herpes virus 1 microvariant and with that of *Mercenaria mercenaria* against Quahog Parasite Unknown (QPX) [38,89]. In our study, NS oysters barely presented some RGs related to

GPCRs, and one was leucine-rich repeat-containing G-protein coupled receptor 6, which was the most down-regulated gene detected in any condition ($FC = -7486$ at T1; Table S5).

We found other up-regulated genes related to cell membrane in AS oysters (cluster AS.RG2; Fig. S1, Table S6), such as integrin alpha-PS3, calcium- and integrin-binding protein (*CIB1*) and innexin, which might be involved in the early immune response. Integrins are transmembrane proteins that show a wide functional range, including spreading, adhesion and migration, and, particularly, integrin- α PS3 was reported to be activated in haemocytes of Manila clam (*Ruditapes philippinarum*) exposed to zoospores of *Perkinsus olseni* [24], and involved in phagocytosis of apoptotic cells and bacteria in *Drosophila* haemocytes [67]. Moreover, up-regulation of *CIB1*, encoding for an integrin binding partner that regulates integrin function [28], was observed in *C. gigas* oysters infected by ostreid herpes virus and related to the mechanisms of virus entry [41]. Innexins are transmembrane proteins, better characterized in insects than in bivalves, which are known to be structural components of the gap junctions in invertebrates. In arthropods, the formation of gap junction between haemocytes has been described during defense response, particularly for pathogen encapsulation, which is also a main defense mechanism of bivalves, for example against the previously cited parasites *Perkinsus* spp. and QPX [2,81].

All in all, results suggest that up-regulation of cell-surface molecules and pathogens recognition receptors occur in AS haemocytes early after challenge, possibly making them “battle ready” to efficiently fight against the antigenic stimuli.

We also found significantly enriched GO terms associated to up-regulated genes encoding protease inhibitors (cluster AS.RG2; Fig. 4 and Fig. S1, Table S8), which were especially present in AT1 (inter-alpha-trypsin inhibitor heavy chain H3, inter-alpha-trypsin inhibitor





Fig. 4. Relevant enriched GO terms of regulated genes groups showing correlated expression profiles (SOTA groups) along the time course in the long-term affected (AS) flat oyster (*O. edulis*) stock after challenge with *B. ostreae* along a time series (T1: 1 day post-challenge (dpc), T2: 30 dpc, T3: 90 dpc); A: AS.G1; B: AS.G2. In the upper right the profile pattern of the group including T1, T2 and T3 and the five replicates per time (see Fig. S1).

heavy chain H4, serine protease inhibitor dipetalogastin, pregnancy zone protein, protein AMBP, murinoglobulin-2). These molecules, often acting as acute phase proteins, represent an important defense mechanism against infectious diseases in bivalves [2,34,90], and a polymorphism in a serine protease inhibitor gene has been associated to disease resistance in *C. virginica* [92].

Genes related to histones are known to participate in mollusc immune response being up-regulated in different oyster species in response to *Perkinsus* spp. [65,73]. These genes were broadly down-regulated in NS as compared to AS oysters (Tables S2 and S3). Histones H2A, H2B, H3 and H4 were among those showing the highest differences between stocks (Fig. 3) and were clustered in the SOTA group GS2 (Table S7), where significantly enriched GO terms related to these nuclear proteins were found (Table S9, Fig. 5).

Conversely, cluster GS4 contained genes mostly up-regulated in the NS stock at 1 dpc and 90 dpc, but down-regulated in the AS stock (Fig. S2, Table S7), and included six ubiquitin- and ten proteasome-annotated genes. Accordingly, several enriched related GO terms were related to the ubiquitin-proteasome pathway (Fig. 5, Table S9). This pathway is a critical regulator of innate immunity; nonetheless, intracellular pathogens are capable of manipulating the ubiquitin-proteasome system of the host to their benefit for growth, replication or immune evasion [9,44,69]. Some indications of proteasome activation in wild flat oysters infected by *B. ostreae* were previously reported [49], and, interestingly, transcripts related to this pathway were found up-regulated in Atlantic salmon susceptible to infectious pancreatic necrosis virus (IPNV) in a study comparing susceptible and resistant specimens [79].

B. ostreae was shown to interfere with essential processes in the defense response of *O. edulis* haemocytes, such as phagocytosis and lysosomal destruction, and reactive oxygen species (ROS) production

Fig. 5. Relevant enriched GO terms of differentially expressed genes between naïve (NS) and long-term affected (AS) flat oyster (*O. edulis*) stocks showing correlated expression profiles (SOTA groups) after challenge with *B. ostreae* along a time series (T1: 1 day post-challenge (dpc), T2: 30 dpc, T3: 90 dpc). A: GS2; B: GS3; C: GS4.

[58,62]. Nevertheless, NADPH oxidase (NOX5) and cytochrome *b*-245 light chain, both involved in ROS production [5,80], were found among RGs in AT1 (Table S2). Furthermore, two genes related to antioxidant defense by avoiding the collateral effects of ROS, such as catalase and glutathione S-transferase [39], showed a significant activation in the AS stock as compared to NS (cluster GS3, Table S7), which suggests that resistant oysters might be able to circumvent parasite interference showing more active ROS production. In the same way, several cytoskeleton-related RGs were found in AS and among DEGs, mostly up-regulated in AT1. Cytoskeleton plays a key role in cell proliferation, migration and phagocytosis, but also during cell death by apoptosis, and the interaction with host cytoskeleton is essential for a successful invasion by intracellular pathogens [14,22,29]. Among those genes, we found filamin, previously detected in oyster haemocytes infected with *B. ostreae*, and hypothesized to be involved in parasite internalization [60]. Also, several genes related to the actin cytoskeleton structure,

Table 2

List of relevant immune-genes regulated (RG) at 1 day post challenge (dpc) (SAM test, FDR < 0.05) and differentially expressed (DEG) between long-term affected (AS) and naïve (NS) flat oyster (*O. edulis*) (ANOVA; P < 0.001).

Gene ^a	Regulation at 1 dpc ^b	Expression in AS vs NS ^c	Associated function/GO terms
Orexin receptor type 2	UP (AS)	UP	F: G-protein coupled receptor activity
Orexin receptor type 1	DOWN (AS)	DOWN	F: G-protein coupled receptor activity
Cephalotocin receptor 1 (4)	UP (AS)	UP	F: G-protein coupled receptor activity
Integrin alpha-PS3	UP (AS)	UP	F: receptor activity; P: integrin-mediated signaling pathway
Calcium and integrin-binding protein (2)	UP (AS)	UP*	P: positive regulation of cell adhesion mediated by integrin
Fermitin family-like protein 2	UP (AS)	UP	P: cell-matrix adhesion P: integrin-mediated signaling pathway
Vinculin	UP (AS)	UP	F: actin binding; P: cell-matrix adhesion
Catenin beta (2)	UP (AS)	UP	P: cell-matrix adhesion; P: Wnt signaling pathway
Kinesin-like protein (2)	UP (AS)	UP	F: microtubule motor activity; P: microtubule-based movement
Rho GTPase-activating protein17	UP (AS)	UP	P: regulation of small GTPase mediated signal transduction
Complement component 1 Q subcomponent-binding protein (2)	DOWN (AS)	DOWN	F: complement component C1q binding; P: Innate immune response
Complement C1q-like protein 4 (2)	UP (AS)	UP*	C: Collagen trimer
Collagen alpha-1(IV) chain (4)	UP (AS)	UP	C: collagen trimer; F:extracellular matrix structural constituent; P: extracellular matrix organization
Hemicentin 1 (2)	UP (AS)	UP	C: extracellular matrix
Tenascin-R	UP (AS)	UP	C: extracellular matrix; P: extracellular matrix organization
Inter-alpha-trypsin inhibitor heavy chain H4	UP (AS)	UP	F: endopeptidase inhibitor activity; P: negative regulation of endopeptidase activity
Pregnancy zone protein	UP (AS)	UP	F: endopeptidase inhibitor activity; P: negative regulation of endopeptidase activity
Endonuclease G	UP (AS)	UP	P: positive regulation of apoptotic process
Programmed cell death protein 10	UP (NS)	DOWN	P: negative regulation of apoptotic process
Salvador-like protein 1	UP (AS)	UP	P: apoptotic process
Histone H2A (3)	DOWN (NS)	UP*	F: protein binding
Histone H2B (2)	DOWN (NS)	UP	F: protein binding
Histone H4	DOWN (NS)	UP	F: protein binding
Ubiquitin-like modifier-activating enzyme 1	UP (NS)	DOWN	P: protein ubiquitination

^a In parentheses the number of different genes with the same annotation detected either among RG or DEG.

^b In parentheses the stock where the gene was differentially expressed.

^c Asterisks indicate genes with the same annotation but showing an opposite expression pattern.

such as alpha-actinin and vinculin, and cytoskeleton dynamics, including three members of the Rho family of GTPase, known to be involved in phagocytosis, cell spreading and motility [29], were identified. Conversely, genes encoding tubulin alpha and beta, the major proteins of microtubules, and kinesin, a motor protein participating in microtubules dynamics, were down-regulated in AT1 (Table S2). The dynamic rearrangement of cytoskeletal components is key for apoptosis, and microtubules appear to have a main role in Fas-mediated apoptosis [15]. This mechanism was previously hypothesized as important for defense in *Bonamia*-resistant oysters, where a Fas-ligand was up-regulated in haemocytes [62].

As highlighted in a recent work [30], the activation of the apoptotic machinery appears as a main defense response of *O. edulis* against *B. ostreae* infection; the controlled death of infected haemocytes would avoid the intracellular development and multiplication of the parasite decreasing the number of parasitic cells. Several genes involved in apoptosis were detected among RGs in AS oysters, such as the pro-apoptotic genes tumor protein 53-induced nuclear protein 1, caspase 7, caspase 10, growth arrest and DNA-damage-inducible protein alpha, endonuclease G and salvador-like protein 1, as well as three genes related to the Inhibitor of Apoptosis (IAP) family (Table S2).

A refined interpretation of the processes underlying the modulation of the cytoskeleton- and apoptosis-related genes is beyond the scope of this study, where the expression profiles evaluated might include heterogeneous populations of haemocytes, including different cell types. However, our results show consistently that the haemocytes of AS oysters present a wide modulation of these sets of genes at 1 dpc, reflecting cytoskeletal dynamics and remodeling, as well as activation of the apoptosis machinery in response to the parasite challenge, which were unnoticed in NS.

In general, the expression profile of the NS haemocytes at 1 dpc reflected a limited defense response, but an increased activation, reflected in the number of RGs, was found at 90 dpc. Among these RGs,

we detected several immune-related genes, such as universal stress protein (USP), macrophage migration inhibitory factor and interferon-induced protein 44, or galectin 1, shown to be involved in *P. marinus* recognition in *Crassostrea virginica* [23]. Also some genes activated in AT1, such as beta catenin and kinesin, involved in cytoskeletal organizations, and a caspase family member (caspase 2) acting in the execution-phase of apoptosis (Table S2), were detected in NT3. This might be indicative of a delayed, and probably less effective, response by NS oysters, given the lack of previous contact with the parasite.

Despite mechanisms such as apoptosis of infected haemocytes appears to be a strategy to avoid the spreading of the infection, consistent with the lower infection intensity showed by AS oysters [12], the results of our work suggest that the resistance of AS has a multifactorial nature, depending also on a more efficient defense response by preventing the haemocytes invasion by *B. ostreae*.

A list of 24 candidate genes for resistance/susceptibility of *O. edulis* to *B. ostreae* is presented in Table 2. These genes were chosen among those in common between the lists of DEGs and RGs at the most significant time point for AS response (1 dpc), but also considering their involvement in the functions here hypothesized as relevant after data analysis and interpretation. G-protein, integrin-mediated and ubiquitin signaling pathways are represented, as well as cell-matrix interactions, apoptosis and immune-related genes, such as histones, protease inhibitors and complement. This set of genes represents one of the main outcomes of this work and the basis for further investigation and validation of potential markers for resistance to bonamiosis. This could be tackled either by testing SNP associated allelic variants in resistant/susceptible individuals or by testing the use of target genes (mRNA, proteins) on hemolymph samples from infected oysters as intermediate phenotypes.

4. Conclusions

The application of a validated haemocyte flat oyster oligo-microarray has contributed to a better understanding of the response against the threatening parasite *B. ostreae* and to the identification of candidate genes related to resistance/susceptibility. The expression profiles observed in the present study revealed that the haemocytes of AS presents a more intense and quicker response as compared to the NS, with activation of several signal transduction pathways, early after challenge with *B. ostreae*. This aspect might represent a key factor for the resistance characteristics observed for this stock. Also, a main role was found for cell-ECM interactions and proteases inhibitors in the defense response of AS, while the broad down-regulation of histones-related genes in challenged haemocytes from NS suggests that these proteins might have an important part during the infection process. Further work is needed to deepen into these aspects and to investigate the validity and potential of the candidate genes for genetic selection.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ygeno.2018.04.002>.

Acknowledgements

David Iglesias, Niall O'Boyle, Grace Flannery, Sarah Culloty and Carsten Fomsgaard helped to get oysters from Ria of Ortigueira, Lough Foyle and Limfjord. Lucía Insua, Sonia Gómez, Victoria Gregorio, Juan C. Pazos, Rocío Renda, Maribel Meléndez and Elena Penas provided technical support. This study has been supported by the OYSTERECOVER project (FP7-SM3-2008-2-243583) from the European Community's Seventh Framework Programme and Xunta de Galicia Local Government (GRC2014/010, R2014/046).

References

- [1] D.J. Alderman, E.B.G. Jones, Shell disease of oysters, *Fish. Investig. Lond. (Ser. 2)* 26 (8) (1971) 1–19.
- [2] B. Allam, D. Raftos, Immune responses to infectious diseases in bivalves, *J. Invertebr. Pathol.* 131 (2015) 121–136.
- [3] E. Bachère, R.D. Rosa, P. Schmitt, A.C. Poirier, N. Merou, G.M. Charrière, D. Destoumieux-Garzón, The new insights into the oyster antimicrobial defense: cellular, molecular and genetic view, *Fish Shellfish Immunol.* 46 (2015) 50–64.
- [4] J. Baud, A. Gerard, Y. Naciri-Graven, Comparative growth and mortality of *Bonamia ostreae*-resistant and wild flat oysters, *Ostrea edulis*, in an intensive system. I. First year of experiment, *Mar. Biol.* 130 (1997) 71–79.
- [5] K. Bedard, K.-H. Krause, The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology, *Physiol. Rev.* 87 (2007) 245.
- [6] J.H. Beattie, J.P. Davis, S.L. Downing, K.K. Chew, Summer mortality of Pacific oysters, *Am. Fish. Soc. Spec. Publ.* 18 (1988) 265–268.
- [7] F.C.J. Berthe, F. Le Roux, R.D. Adlard, A. Figueras, Marteiliopsis in molluscs: a review, *Aquat. Living Resour.* 17 (2004) 433–448.
- [8] D. Bucke, Pathology of bonamiasis, *Parasitol. Today* 4 (1988) 174–176.
- [9] C.A. Collins, E.J. Brown, Cytosol as battleground: ubiquitin as a weapon for both host and pathogen, *Trends Cell Biol.* 20 (2010) 205–213.
- [10] S.C. Culloty, M.F. Mulcahy, Season-, age-, and sex-related variation in the prevalence of bonamiasis in flat oysters (*Ostrea edulis* L) on the south coast of Ireland, *Aquaculture* 144 (1996) 53–63.
- [11] S.C. Culloty, M.F. Mulcahy, *Bonamia ostreae* in the Native Oyster *Ostrea edulis*: A Review vol Marine Environment & Health Series, (2007).
- [12] P.M. da Silva, J. Fuentes, A. Villalba, Growth, mortality and disease susceptibility of oyster *Ostrea edulis* families obtained from brood stocks of different geographical origins, through on-growing in the Ria de Arousa (Galicia, NW Spain), *Mar. Biol.* 147 (2005) 965–977.
- [13] L. Dégremont, M. Nourry, E. Maurouard, Mass selection for survival and resistance to OsHV-1 infection in *Crassostrea gigas* sat in field conditions: response to selection after four generations, *Aquaculture* 446 (2015) 111–121.
- [14] M. Desouza, P.W. Gunning, J.R. Stehn, The actin cytoskeleton as a sensor and mediator of apoptosis, *BioArchitecture* 2 (2012) 75–87.
- [15] E. Doma, K. Chakrabandhu, A.O. Hueber, A novel role of microtubular cytoskeleton in the dynamics of caspase-dependent Fas/CD95 death receptor complexes during apoptosis, *FEBS Lett.* 584 (2010) 1033–1040.
- [16] Y. Dong, G. Dimopoulos, Anopheles fibrinogen-related proteins provide expanded pattern recognition capacity against bacteria and malaria parasites, *J. Biol. Chem.* 284 (2009) 9835–9844.
- [17] J. Dopazo, J.M. Carazo, Phylogenetic reconstruction using an unsupervised growing neural network that adopts the topology of a phylogenetic tree, *J. Mol. Evol.* 44 (1997) 226–233.
- [18] M.C. Dove, J.A. Nell, W.A. O'Connor, Evaluation of the progenie of the fourth-generation Sydney rock oyster *Saccostrea glomerata* (Gold, 1850) breeding lines for resistance to QX (*Marteilia sydneyi*) and winter mortality (*Bonamia roughleyi*), *Aquat. Res.* 44 (2013) 1791–1800.
- [19] R.A. Elston, M.L. Kent, M.T. Wilkinson, Resistance of *Ostrea edulis* to *Bonamia ostreae* infection, *Aquaculture* 64 (1987) 237–242.
- [20] M.Y. Engelsma, S. Kerhoff, I. Roozenburg, O.L.M. Haenen, A. van Gool, W. Sijstermans, S. Wijnhoven, H. Hummel, Epidemiology of *Bonamia ostreae* infecting European flat oysters *Ostrea edulis* from Lake Grevelinjen, the Netherlands, *Mar. Ecol. Prog. Ser.* 409 (2010) 131–142.
- [21] M.Y. Engelsma, S.C. Culloty, S.A. Lynch, I. Arzul, R.B. Carnegie, *Bonamia* parasites: a rapidly changing perspective on a genus of important mollusc pathogens, *Dis. Aquat. Org.* 110 (2014) 5–23.
- [22] C.L. Epting, B.M. Coates, D.M. Engman, Molecular mechanisms of host cell invasion by *Trypanosoma cruzi*, *Exp. Parasitol.* 126 (2010) 283–291.
- [23] C. Feng, A. Ghosh, M.N. Amin, B. Giomarelli, S. Shridhar, A. Banerjee, J.A. Fernández-Robledo, M.A. Bianchet, L.-X. Wang, I.B.H. Wilson, G.R. Vasta, The galectin CvGal1 from the eastern oyster (*Crassostrea virginica*) binds to blood group a oligosaccharides on the hemocyte surface, *J. Biol. Chem.* 288 (2013) 24394–24409.
- [24] S. Fernández-Boo, A. Villalba, A. Cao, Protein expression profiling in haemocytes and plasma of the Manila clam *Ruditapes philippinarum* in response to infection with *Perkinsus olseni*, *J. Fish Dis.* 39 (2016) 1369–1385.
- [25] S. Ferrareso, N. Vitulo, A.N. Mininni, C. Romualdi, B. Cardazzo, E. Negrisolo, R. Reinhardt, A.V. Canario, T. Patarnello, L. Bargelloni, Development and validation of a gene expression oligo microarray for the gilthead sea bream (*Sparus aurata*), *BMC Genomics* 9 (2008) 580.
- [26] S.E. Ford, H.H. Haskin, Infection and mortality patterns in strains of oysters *Crassostrea virginica* selected for resistance to the parasite *Haplosporidium nelsoni* MSX, *J. Parasitol.* 73 (1987) 386–396.
- [27] A. Frank-Lawale, S.K. Allen, L. Dégremont, Breeding and domestication of eastern oyster (*Crassostrea virginica*) lines for culture in the mid-Atlantic, USA: line development and mass selection for disease resistance, *J. Shellfish Res.* 33 (2014) 153–165.
- [28] T.C. Freeman, J.L. Black, H.G. Bray, O. Dagliyan, Y.I. Wu, A. Tripathy, N.V. Dokholyan, T.M. Leisner, L.V. Parise, Identification of novel integrin binding partners for CIB1: structural and thermodynamic basis of CIB1 promiscuity, *Biochemistry* 52 (2013) 7082–7090.
- [29] S.A. Freeman, S. Grinstein, Phagocytosis: receptors, signal integration, and the cytoskeleton, *Immunol. Rev.* 262 (2014) 193–215.
- [30] O. Gervais, B. Chollet, T. Renault, I. Arzul, Flat oyster follows the apoptosis pathway to defend against the protozoan parasite *Bonamia ostreae*, *Fish Shellfish Immunol.* 56 (2016) 322–329.
- [31] M.A. Gordy, E.A. Pila, P.C. Hanington, The role of fibrinogen-related proteins in the gastropod immune response, *Fish Shellfish Immunol.* 46 (2015) 39–49.
- [32] H. Grizel, M. Comps, J.R. Bonami, F. Cousserans, J.L. Duthoit, M.A. Le Penne, Recherche sur l'agent de la maladie de la glande digestive de *Ostrea edulis* Linne, *Sci. Pêche. Bull. Inst. Pêches. Marit.* 240 (1974) 7–29.
- [33] H. Grizel, E. Bachère, E. Mialhe, G. Tigé, Solving Parasite-Related Problems in Cultured Molluscs, in: M.J. Howell (Ed.), *Parasitology, Quo vadit?* Proc 6th Int Congr Parasitol, Australian Academy of Science, Canberra, 1986, pp. 301–308.
- [34] A.F.M. Hasanuzzaman, D. Robledo, A. Gómez-Tato, J.A. Álvarez Dios, P.W. Harrison, A. Cao, S. Fernández-Boo, A. Villalba, B.G. Pardo, P. Martínez, Transcriptomic profile of Manila clam (*Ruditapes philippinarum*) haemocytes in response to *Perkinsus olseni* infection, *Aquaculture* 467 (2017) 170–181.
- [35] P.C. Hanington, S.-M. Zhang, The primary role of fibrinogen-related proteins in invertebrates is defense, not coagulation, *J. Innate Immun.* 3 (2010) 17–27.
- [36] C.D. Hanlon, D.J. Andrew, Outside-in signaling – a brief review of GPCR signaling with a focus on the *Drosophila* GPCR family, *J. Cell Sci.* 128 (2015) 3533.
- [37] Y. He, H. Yu, Z. Bao, Q. Zhang, X. Guo, Mutation in promoter region of a serine protease inhibitor confers *Perkinsus marinus* resistance in the eastern oyster (*Crassostrea virginica*), *Fish Shellfish Immunol.* 33 (2012) 411–417.
- [38] Y. He, A. Jouaux, S.E. Ford, C. Lelong, P. Sourdain, M. Mathieu, X. Guo, Transcriptome analysis reveals strong and complex antiviral response in a mollusc, *Fish Shellfish Immunol.* 46 (2015) 131–144.
- [39] M. Hermes-Lima, Oxygen in biology and biochemistry: role of free radicals, in: Kenneth B. Storey (Ed.), *Functional Metabolism*, John Wiley & Sons, Inc, 2004, p. 319e368.
- [40] B. Huang, L. Zhang, L. Li, X. Tang, G. Zhang, Highly diverse fibrinogen-related proteins in the Pacific oyster *Crassostrea gigas*, *Fish Shellfish Immunol.* 43 (2015) 485–490.
- [41] A. Jouaux, M. Lafont, J.L. Blin, M. Houssin, M. Mathieu, C. Lelong, Physiological change under OsHV-1 contamination in Pacific oyster *Crassostrea gigas* through massive mortality events on fields, *BMC Genomics* 14 (2013) 590.
- [42] P. Korringa, Farming the Flat Oysters of the Genus *Ostrea*: A Multidisciplinary Treatise, Elsevier Publishing company, Amsterdam, 1976.
- [43] I. Laing, P. Dunn, E.J. Peeler, S.W. Feist, M. Longshaw, Epidemiology of *Bonamia* in the UK, 1982–2012, *Dis. Aquat. Org.* 110 (2014) 101–111.
- [44] J. Laliberte, V.B. Carruthers, Host cell manipulation by the human pathogen *Toxoplasma gondii*, *Cell. Mol. Life Sci.* 65 (2008) 1900–1915.
- [45] S.A. Lynch, D.V. Armitage, J. Coughlan, M.F. Mulcahy, S.C. Culloty, Investigating the possible role of benthic macroinvertebrates and zooplankton in the life cycle of the haplosporidian *Bonamia ostreae*, *Exp. Parasitol.* 115 (2007) 359–368.
- [46] S.A. Lynch, G. Flannery, T. Hugh-Jones, D. Hugh-Jones, S.C. Culloty, Thirty-year history of Irish (Rossmore) *Ostrea edulis* selectively bred for disease resistance to *Bonamia ostreae*, *Dis. Aquat. Org.* 110 (2014) 113–121.

- [47] Y.J. Ma, E. Hein, L. Munthe-Fog, M.O. Skjoed, R. Bayarri-Olmos, L. Romani, P. Garred, Soluble Collectin-12 (CL-12) is a pattern recognition molecule initiating complement activation via the alternative pathway, *J. Immunol.* 195 (2015) 3365–3373.
- [48] L. Madsen, Thomassen H (2015) First detection of *Bonamia ostreae* in native flat oysters from the Limfjord in Denmark, 17th International Conference on Diseases of Fish and Shellfish: Abstract Book, European Association of Fish Pathologists, Las Palmas de Gran Canaria, Spain, 2015, p. 92.
- [49] L. Martín-Gómez, A. Villalba, E. Abollo, Identification and expression of immune genes in the flat oyster *Ostrea edulis* in response to bonamiosis, *Gene* 492 (2012) 81–93.
- [50] L. Martín-Gómez, A. Villalba, R.H. Kerkhoven, E. Abollo, Role of microRNAs in the immunity process of the flat oyster *Ostrea edulis* against bonamiosis, *Infect. Genet. Evol.* 27 (2014) 40–50.
- [51] I.C. McDowell, C. Nikapitiya, D. Aguiar, C.E. Lane, S. Istrail, M. Gomez-Chiarri, Transcriptome of American oysters, *Crassostrea virginica*, in response to bacterial challenge: insights into potential mechanisms of disease resistance, *PLoS One* 9 (2014) e105097.
- [52] A.M. Mendes, P.H. Awono-Ambene, S.E. Nsango, A. Cohuet, D. Fontenille, F.C. Kafatos, G.K. Christophides, I. Morlais, D. Vlachou, Infection intensity-dependent responses of *Anopheles gambiae* to the African malaria parasite *Plasmodium falciparum*, *Infect. Immun.* 79 (2011) 4708–4715.
- [53] E. Mialhe, E. Bachere, D. Chagot, H. Grizel, Isolation and purification of the protozoan *Bonamia ostreae* (Pichot et al. 1980), a parasite affecting the flat oyster *Ostrea edulis* L. *Aquaculture* 71 (1988) 293–299.
- [54] A. Millán, A. Gómez-Tato, C. Fernández, B.G. Pardo, J.A. Alvarez-Dios, M. Calaza, C. Bouza, M. Vázquez, S. Cabaleiro, P. Martínez, Design and performance of a turbot (*Scophthalmus maximus*) oligo-microarray based on ESTs from immune tissues, *Mar. Biotechnol.* 12 (2010) 452–465.
- [55] A. Millán, A. Gómez-Tato, B.G. Pardo, C. Fernández, C. Bouza, M. Vera, J.A. Alvarez-Dios, S. Cabaleiro, J. Lamas, M.L. Lemos, P. Martínez, Gene expression profiles of the spleen, liver, and head kidney in turbot (*Scophthalmus maximus*) along the infection process with *Aeromonas salmonicida* using an immune-enriched oligo-microarray, *Mar. Biotechnol.* 13 (2011) 1099–1114.
- [56] M. Miron, O.Z. Woody, A. Marcil, C. Murie, R. Sladek, R. Nadon, A methodology for global validation of microarray experiments, *BMC Bioinforma.* 7 (2006) 333.
- [57] J. Montes, A. Villalba, M.C. López, M.J. Carballed, S.G. Mourelle, Bonamiosis in native flat oysters (*Ostrea edulis* L.) from two intertidal beds of the Ortigueira estuary (Galicia, N.W. Spain) with different histories of oyster culture, *Aquaculture* 93 (1991) 213–224.
- [58] B. Morga, I. Arzul, B. Chollet, T. Renault, Infection with the protozoan parasite *Bonamia ostreae* modifies in vitro haemocyte activities of flat oyster *Ostrea edulis*, *Fish Shellfish Immunol.* 26 (2009) 836–842.
- [59] B. Morga, I. Arzul, N. Faury, T. Renault, Identification of genes from flat oyster *Ostrea edulis* as suitable housekeeping genes for quantitative real time PCR, *Fish Shellfish Immunol.* 29 (2010) 937–945.
- [60] B. Morga, I. Arzul, N. Faury, A. Segarra, B. Chollet, T. Renault, Molecular responses of *Ostrea edulis* haemocytes to an in vitro infection with *Bonamia ostreae*, *Dev. Comp. Immunol.* 35 (2011) 323–333.
- [61] B. Morga, T. Renault, N. Faury, B. Chollet, I. Arzul, Cellular and molecular responses of haemocytes from *Ostrea edulis* during in vitro infection by the parasite *Bonamia ostreae*, *Int. J. Parasitol.* 41 (2011) 755–764.
- [62] B. Morga, T. Renault, N. Faury, I. Arzul, New insights in flat oyster *Ostrea edulis* resistance against the parasite *Bonamia ostreae*, *Fish Shellfish Immunol.* 32 (2012) 958–968.
- [63] Y. Naciri-Graven, J. Haure, A. Gérard, J.P. Baud, Comparative growth and mortality of *Bonamia ostreae*-resistant and wild flat oysters, *Ostrea edulis*, in an intensive system. II. Second year of the experiment, *Aquaculture* 171 (1999) 195–208.
- [64] Q. Nie, X. Yue, B. Liu, Development of *Vibrio* spp. infection resistance related SNP markers using multiplex SNaPshot genotyping method in the clam *Meretrix meretrix*, *Fish Shellfish Immunol.* 43 (2015) 469–471.
- [65] C. Nikapitiya, T. Dorrington, M. Gómez-Chiarri, The role of histones in the immune responses of aquatic invertebrates, *Invertebr. Surviv. J.* 10 (2013) 94–101.
- [66] C. Nikapitiya, I.C. McDowell, L. Villamil, P. Muñoz, S.B. Sohn, M. Gomez-Chiarri, Identification of potential general markers of disease resistance in American oysters, *Crassostrea virginica* through gene expression studies, *Fish Shellfish Immunol.* 41 (2014) 27–36.
- [67] S. Nonaka, K. Nagaosa, T. Mori, A. Shiratsuchi, Y. Nakanishi, Integrin α PS3/ β -mediated phagocytosis of apoptotic cells and bacteria in *Drosophila*, *J. Biol. Chem.* 288 (2013) 10374–10380.
- [68] J. Normand, R. Li, V. Quillien, J.L. Nicolas, P. Boudry, F. Pernet, A. Huvet, Contrasted survival under field or controlled conditions displays associations between mRNA levels of candidate genes and response to OsHV-1 infection in the Pacific oyster *Crassostrea gigas*, *Mar. Genomics* 15 (2014) 95–102.
- [69] M. Olivier, D.J. Gregory, G. Forget, Subversion mechanisms by which leishmania parasites can escape the host immune response: a signaling point of view, *Clin. Microbiol. Rev.* 18 (2005) 293–305.
- [70] B.G. Pardo, J.A. Alvarez-Dios, A. Cao, A. Ramilo, A. Gomez-Tato, J.V. Planas, A. Villalba, P. Martinez, Construction of an *Ostrea edulis* database from genomic and expressed sequence tags (ESTs) obtained from *Bonamia ostreae* infected haemocytes: development of an immune-enriched oligo-microarray, *Fish Shellfish Immunol.* 59 (2016) 331–344.
- [71] A.M. Piccinini, K.S. Midwood, Endogenous control of immunity against infection: tenascin-C regulates TLR4-mediated inflammation via microRNA-155, *Cell Rep.* 2 (2012) 914–926.
- [72] Y. Pichot, M. Comps, G. Tigé, H. Grizel, M.A. Rabouin, Recherches sur *Bonamia ostreae* gen. n., sp. n., parasite nouveau de l'huître plate *Ostrea edulis* L., *Rev. Trav. Inst. Pêches. Marit.* 43 (1980) 131–140.
- [73] A.C. Poirier, P. Schmitt, R.D. Rosa, A.S. Vanhove, S. Kieffer-Jaquinod, T.P. Rubio, G.M. Charriere, D. Destoumieux-Garzon, Antimicrobial histones and DNA traps in invertebrate immunity: evidences in *Crassostrea gigas*, *J. Biol. Chem.* 289 (2014) 24821–24831.
- [74] F. Qu, Z. Xiang, F. Wang, L. Qi, F. Xu, S. Xiao, Z. Yu, Prostaglandin E receptor 4 (PTGER4) involved in host protection against immune challenge in oyster, *Crassostrea hongkongensis*, *Fish Shellfish Immunol.* 42 (2015) 316–324.
- [75] D.A. Raftos, R. Kuchel, S. Aladaile, D. Butt, Infectious microbial diseases and host defense responses in Sydney rock oysters, *Front. Microbiol.* 5 (2014) 135.
- [76] L.M. Ragone Calvo, G.W. Calvo, E.M. Burreson, Dual disease resistance in a selectively bred eastern oyster, *Crassostrea virginica*, strain tested in Chesapeake Bay, *Aquaculture* 220 (2003) 69–87.
- [77] L. Ribas, D. Robledo, A. Gómez-Tato, A. Viñas, P. Martínez, F. Piferrer, Comprehensive transcriptomic analysis of the process of gonadal sex differentiation in the turbot (*Scophthalmus maximus*), *Mol. Cell. Endocrinol.* 422 (2016) 132–149.
- [78] M.E. Ritchie, B. Phipson, D. Wu, Y. Hu, C.W. Law, W. Shi, G.K. Smyth, Limma powers differential expression analyses for RNA-sequencing and microarray studies, *Nucleic Acids Res.* 43 (2015) e47.
- [79] D. Robledo, J.B. Taggart, J.H. Ireland, B.J. McAndrew, W.G. Starkey, C.S. Haley, A. Hamilton, D.R. Guy, J.C. Mota-Velasco, A.A. Gheyas, A.E. Tinch, D.W. Verner-Jeffrey, R.K. Paley, G.S. Rimmer, L.J. Tew, S.C. Bishop, J.E. Bron, R.D. Houston, Gene expression comparison of resistant and susceptible Atlantic salmon fry challenged with Infectious Pancreatic Necrosis virus reveals a marked contrast in immune response, *BMC Genomics* 17 (2016) 279.
- [80] A.W. Segal, et al., Cytochrome b245 is a flavocytochrome containing FAD and the NADPH-binding site of the microbicidal oxidase of phagocytes, *Biochem. J.* 284 (1992) 781–788.
- [81] P. Soudant, R. Leite, F. Chu, A. Villalba, L. Cancela, Bivalves-*Perkinsus* spp. interactions, Workshop for the analysis of the impact of Perkinsosis to the European Shellfish Industry, Vilanova de Arousa, Spain, Citeaser, 2008, pp. 78–109.
- [82] L. Sun, R.D. Ye, Role of G protein-coupled receptors in inflammation, *Acta Pharmacol. Sin.* 33 (2012) 342–350.
- [83] T. Thomsen, A. Schlosser, U. Holmskov, G.L. Sorensen, Ficolins and FIBCD1: soluble and membrane bound pattern recognition molecules with acetyl group selectivity, *Mol. Immunol.* 48 (2011) 369–381.
- [84] G. Tigé, H. Grizel, Essai de contamination d'*Ostrea edulis* Linne par *Bonamia ostreae* (Pichot et al., 1979) en riviere de Crach (Morbinhan), *Rev. Trav. Inst. Pêch. Marit.* 46 (1982) 307–314.
- [85] V.G. Tusher, R. Tibshirani, G. Chu, Significance analysis of microarrays applied to the ionizing radiation response, *Proc. Natl. Acad. Sci. U. S. A.* 98 (2001) 5116–5121.
- [86] V. Vaibhav, M. Lepretre, E. Thompson, D. Raftos, P.A. Haynes, Biomarkers of Winter Mortality resistance in selectively bred Sydney rock oysters (*Saccostrea glomerata*), *Aquaculture* 465 (2016) 323–329.
- [87] P. van Banning, Observations on bonamiosis in the stock of the European flat oyster, *Ostrea edulis*, in the Netherlands, with special reference to the recent developments in Lake Grevelingen, *Aquaculture* 93 (1991) 205–211.
- [88] K. Wang, C. del Castillo, E. Corre, E. Pales Espinosa, B. Allam, Clam focal and systemic immune responses to QPX infection revealed by RNA-seq technology, *BMC Genomics* 17 (2016) 146.
- [89] K. Wang, E. Pales Espinosa, A. Tanguy, B. Allam, Alterations of the immune transcriptome in resistant and susceptible hard clams (*Mercentaria mercenaria*) in response to Quahog Parasite Unknown (QPX) and temperature, *Fish Shellfish Immunol.* 49 (2016) 163–176.
- [90] Q. Xue, J.P. Beguel, J. Gauthier, J. La Peyre, Identification of cvSI-3 and evidence for the wide distribution and active evolution of the 184 family of protease inhibitors in mollusks, *Fish Shellfish Immunol.* 62 (2017) 332–340.
- [91] L. Yin, C.H. Huang, J. Ni, Clustering of gene expression data: performance and similarity analysis, *BMC Bioinforma.* 7 (Suppl. 4) (2006) S19.
- [92] H. Yu, Y. He, X. Wang, Q. Zhang, Z. Bao, X. Guo, Polymorphism in a serine protease inhibitor gene and its association with disease resistance in the eastern oyster (*Crassostrea virginica* Gmelin), *Fish Shellfish Immunol.* 30 (2011) 757–762.
- [93] G. Zhang, X. Fang, X. Guo, L. Li, R. Luo, et al., The oyster genome reveals stress adaptation and complexity of shell formation, *Nature* 490 (2012) 49–54.
- [94] Y. Zhang, X. He, F. Yu, Z. Xiang, J. Li, K.L. Thorpe, Z. Yu, Characteristic and functional analysis of Toll-like receptors (TLRs) in the lophotrocozoan, *Crassostrea gigas*, reveals ancient origin of TLR-mediated innate immunity, *PLoS One* 8 (2013) e76464.
- [95] Z. Zhou, Q. Jiang, M. Wang, F. Yue, L. Wang, L. Wang, F. Li, R. Liu, L. Song, Modulation of haemocyte phagocytic and antibacterial activity by alpha-adrenergic receptor in scallop *Chlamys farreri*, *Fish Shellfish Immunol.* 35 (2013) 825–832.