

Nocturnal enuresis in children is associated with differences in autonomic control

Alexandra Bascom, Mary Ann McMaster, R Todd Alexander, Joanna E MacLean

Sleep, Volume 42, Issue 3, March 2019, zsy239, <https://doi.org/10.1093/sleep/zsy239>

Published: 26 November 2018 **Article history** ▼

 PDF  Split View  Cite  Permissions  Share ▼

Abstract

Study Objectives

To assess the relationship between urine osmolality, cardiovascular parameters, and nocturnal enuresis in a population of children undergoing polysomnographic assessment.

Methods

This prospective observational study included consecutive children aged 5–17 years presenting for overnight polysomnography. Children were evaluated using continuous ambulatory blood pressure monitoring to assess heart rate and blood pressure. Urine samples were collected throughout the night to determine urine sodium excretion and osmolality. Comparisons of results were made between children with and without a history of nocturnal enuresis.

Results

A total of 61 children were included for analysis; 13 had a history of nocturnal enuresis. Children with nocturnal enuresis had greater disruption in respiratory parameters including higher apnea–hypopnea index (mean difference 12.2 ± 8.8 events/h, $p < 0.05$), attributable to more central respiratory events (mean difference 5.4 ± 4.9 , $p < 0.05$), and higher variability in both oxygen and carbon dioxide parameters compared to those without nocturnal enuresis. Sleep parameters, urine osmolality, and blood pressure did not differ between groups. Children with nocturnal enuresis showed an increase, rather than a decrease, in heart rate across the night ($+5.4 \pm 19.1$ vs. -6.0 ± 14.8 beats/min, $p < 0.05$).

Conclusions

Children with a history of nocturnal enuresis have greater respiratory abnormalities, no differences in urine osmolality or blood pressure, and loss of normal heart rate decrease across the night. This pattern suggests that autonomic control, rather than renal or hemodynamic abnormalities, may contribute to the pathophysiology of nocturnal enuresis.

[polysomnography](#), [blood pressure](#), [pediatrics](#), [urine analysis](#)

Statement of Significance

Nocturnal enuresis and obstructive sleep apnea commonly co-occur in children; understanding the link between these conditions will shed light on their shared pathophysiology and potential for novel treatment strategies. Our results highlight autonomic disruption as a potential contributor

to nocturnal enuresis and a plausible link between nocturnal enuresis and obstructive sleep apnea in children. Exploring pathways linked to autonomic function may lead to novel treatments for both nocturnal enuresis and obstructive sleep apnea.

Introduction

Nocturnal enuresis (NE), commonly referred to as bedwetting, is defined as inappropriate urine outflow during sleep and affects approximately 15% of 5-year-old and 5% of 10-year-old children [1]. Enuresis can have dramatic psychological and emotional impact on young children and severely affect their quality of life [2]. The most dominant theories with respect to proposed mechanisms for NE include maturational delay in bladder response, increased threshold for arousal from sleep, abnormal bladder dynamics, and impaired release of renin–angiotensin–aldosterone system hormones, atrial natriuretic peptide, or antidiuretic hormone (ADH) [3, 4]. With a strong relationship between obstructive sleep apnea (OSA) and NE in young children, understanding the mechanisms associated with NE in the context of OSA may shed new light on the mechanisms of NE overall.

OSA is defined by repetitive complete or partial airway closure that may disrupt breathing during sleep, sleep patterns, or both. Several retrospective studies have demonstrated a positive correlation between NE and OSA in children including resolution of NE after surgical management of OSA by adenotonsillectomy. A systematic review which included data from 3550 children reported a prevalence of 33% for NE in children with reported sleep disordered breathing (SDB), a broader term that includes OSA; the prevalence of NE dropped to 16% after adenotonsillectomy [5]. NE occurs more commonly in children with higher apnea–hypopnea index (AHI) [5, 6] and children with NE have a high rate of snoring and symptoms of SDB [7, 8], suggesting a common pathway contributing to both NE and SDB. Treatment for OSA with adenotonsillectomy leads to resolution or improvements of NE in 33%–74% and 23%–31% of children, respectively [1, 9–11], showing that OSA alone cannot explain NE in children with OSA. Pathways postulated to be involved in the association between NE and OSA include heightened sympathetic output leading to increased blood pressure and secondary natriuresis, heightened arousal threshold leading to a failure of arousal from sleep, and swings in intrathoracic pressure resulting in the release of cardiac hormones including brain-derived neurotrophic factor leading to increased urine production [12, 13]. There is, however, a paucity of physiological data available to support these hypothesized mechanisms.

The objective of this study was to determine the association between polysomnography results, physiologic variables, including hemodynamics and urine osmolality, and NE in a population of children undergoing polysomnography. We hypothesize that children with NE will have greater respiratory disturbance resulting in increases in heart rate and blood pressure as well as higher urine volume throughout the night when compared to children without NE.

Methods

The study design was a prospective observational cohort study. Consecutive children aged 5–17 years undergoing overnight diagnostic polysomnography for the investigation of sleep-related symptoms were recruited through the pediatric sleep laboratory at a tertiary care children's hospital. The lower limit of 5 years of age for inclusion is consistent with international consensus guidelines on NE [14]. Exclusion criteria included children with known neurological disorders, significant congenital bladder anomalies, or conditions associated with alterations in cardiac or renal function. Written informed consent from parents/guardians and assent from children was obtained prior to participation. The study protocol was approved by the institutional health research ethics board (Pro00034514).

Patient demographics, including age, height, weight, body mass index, past medical history, reason for the polysomnography, and medications were collected from sleep laboratory records. Information about daily sodium and fluid intake was collected using a dietary recall sheet completed by the parents/guardians and analyzed using Food Processor Nutrition software (www.ehsa.com). Children and parents were asked about previous history of NE, defined for the purpose of this study as greater than three wet nights per month for the past 6 months.

Polysomnography was conducted in accordance with standard laboratory protocols and scored according to the criteria of the American Academy of Sleep Medicine [15] by an experienced polysomnography technologist. The AHI includes the total number of apneas and hypopneas per hour of total sleep time. The obstructive-mixed AHI and central AHI include obstructive/mixed and central apnea and hypopneas per hour of total sleep time, respectively. Oxygen desaturation index (ODI) was defined by the number of oxygen desaturation events $\geq 3\%$ per hour of total sleep time.

During the polysomnography, children were monitored with an ambulatory blood pressure monitor (Spacelabs Medical Ambulatory Blood Pressure Cuff 90217-1Q). Heart rate and blood pressure were checked in the supine position when the child first went to bed. Additional readings were taken automatically throughout the night on an hourly basis and at the time of waking (between 0600 and 0700) while the child was in a supine position.

Blood pressure and heart rate data were extracted from the analysis software (Spacelabs software On Track). Measurements were converted to z-scores using established reference equations and thresholds [16, 17].

Children were asked to provide urine samples overnight, collected using a toilet hat. Each sample was measured for volume and then divided into two separate containers: the first for urine osmolality, and the second for urine sodium. The first urine collection occurred just prior to going to bed and represented the pre-sleep urine collection. Further urine specimens from spontaneous voids were collected throughout the night. Children were then asked to fully empty their bladder in the morning after awakening. Enuresis overnight was documented by the parent or sleep technician. Urine samples were then deposited at the laboratory for analysis. Samples were kept refrigerated according to laboratory protocols. Urine sodium was assessed by the University of Alberta Hospital Laboratory, while urine osmolality was measured with an Advanced Osmometer (model 3D3, Advanced Instruments, Norwood, MA).

Children were grouped into those with and without a history of NE for the purpose of analysis. Non-parametric data were transformed, where possible, prior to analysis. Student *t*-tests and Mann–Whitney *U* were used to compare group results for continuous variables as appropriate. Chi-square was used to compare distribution of categorical variables. Spearman correlation was used to assess bivariate non-parametric relationships. Linear regression was used for multi-variable analysis of parametric data. A *p*-value of <0.05 indicated statistically significant group differences.

Results

Sixty-eight children agreed to participate in the study, of which 61 children completed the protocol and were included for analysis. Mean age for this sample of children was 9.9 ± 3.8 years (range 5–17 years) and 13 (21%) patients had a history of NE. Age was the only descriptive factor that differed between those with and without a history of NE where children with enuresis were younger (Table 1).

Table 1.
Comparison of demographics between children with and without NE

	NE (<i>n</i> , %; mean ± <i>SD</i>)	No NE (<i>n</i> , %; mean ± <i>SD</i>)
<i>n</i>	13	48
Age (year)*	8.2 ± 3.2	10.5 ± 3.8
Male:female (% female)	5:8 (38%)	17:31 (35%)
Height z-score	−0.16 ± 1.43	0.13 ± 1.88
Weight z-score	0.39 ± 1.30	0.93 ± 1.55
BMI z-score	0.59 ± 1.34	1.04 ± 1.41
Reason for sleep study		
OSA/SDB	12 (92%)	43 (90%)
Excessive daytime sleepiness	1 (8%)	2 (4%)
Restless leg syndrome	0	3 (6%)
NE on the night of the study*	5 (38%)	0

BMI, body mass index.

**p* < 0.05.

[View Large](#)

Polysomnography was consistent with OSA for 67% of children with no difference in the rate of OSA between children with (77%) and without a history of NE (65%, $\chi^2 = 0.70, p = \text{ns}$). There were no differences in sleep parameters between groups (Table 2); however, children with a history of

NE had more respiratory events and greater variability in respiratory parameters compared to those without NE (Table 2). These included a greater number of central respiratory events accounting for a higher AHI and greater variance in both oxygen and carbon dioxide measures.

Table 2.

Polysomnography characteristics in children with and without NE

	NE (mean \pm SD)	No NE (mean \pm SD)	Mean difference (95% confidence interval)
Total sleep time (min)	458 \pm 49	419 \pm 79	−8.8, 87.8
Sleep efficiency (%)	84.7 \pm 8.7	79.2 \pm 14.2	−2.8, 13.8
Wake after sleep onset (min)	58.3 \pm 38.1	79.5 \pm 67.5	−57.4, 21.0
% SWS	30.5 \pm 7.3	27.0 \pm 9.3	−2.1, 9.1
% REM	19.3 \pm 5.7	18.4 \pm 6.2	−2.9, 4.7
Total arousals (events/h)	12.1 \pm 9.7	9.7 \pm 10.3	−3.9, 8.8
AHI (events/h)	17.8 \pm 31.1	5.7 \pm 13.0	
logAHI*	0.80 \pm 0.61	0.47 \pm 0.44	0.031, 0.64
OMAHI (events/h)	11.2 \pm 18.7	4.4 \pm 11.8	
logOMAHI	0.52 \pm 0.76	0.22 \pm 0.57	−0.94, 0.68
CAHI (events/h)	6.7 \pm 16.2	1.2 \pm 1.5	
logCAHI*	0.29 \pm 0.58	−0.066 \pm 0.47	0.037, 0.67
Mean SpO ₂ (%)**	95.0 \pm 4.8	96.1 \pm 1.4	Non-parametric
Minimum SpO ₂ (%)	88.7 \pm 3.9	88.6 \pm 4.0	−2.5, 2.8
ODI (events/h)	25.4 \pm 41.7	9.3 \pm 14.3	Non-parametric
Mean E _T CO ₂ (mm Hg)**	42.2 \pm 7.1	42.3 \pm 3.7	Non-parametric
Maximum E _T CO ₂ (mm Hg)	51.6 \pm 6.7	49.1 \pm 8.2	−4.4, −1.4
% Time with E _T CO ₂ > 50 mm Hg**	24.7 \pm 6.8	1.6 \pm 0.2	Non-parametric

CAHI, central AHI; E_TCO₂, end-tidal carbon dioxide; OMAHI, obstructive-mixed AHI; REM, rapid eye movement sleep; SpO₂, pulse oxygen saturation; SWS, slow-wave sleep.

* $p < 0.05$.

**Ranges differ between children with and without NE, $p < 0.05$.

[View Large](#)

There were no between-group differences in daily fluid or sodium intake, or urine osmolality. Children with NE had lower overnight urine sodium excretion when compared to children without NE (Table 3); post hoc analysis showed that history of NE was not a significant independent predictor of overnight urine sodium after adjustment for age (history of NE $\beta = 0.022$, $p = \text{ns}$; age $\beta = 0.44$, $p < 0.05$). Average systolic and diastolic blood pressure z -scores, calculated from the average of all measurements within an individual, and rates of elevated blood pressure or hypertension did not differ between groups (Table 3). While the pattern of systolic and diastolic blood pressure across the night did not differ between groups, the heart rate pattern was different (Figure 1). From the start to the end of the study, children with NE showed a 5.4 ± 19.1 beats/min increase in heart rate while children without NE showed a drop in heart rate of 6.0 ± 14.8 beats/min with a mean difference of 11.4 ± 5.6 beats/min ($p < 0.05$) between groups. Post hoc analysis showed that the history of NE remains a significant independent predictor of change in heart rate after adjustment for age

(history of NE: $\beta = -0.35, p < 0.05$; age: $\beta = -0.19, p = \text{ns}$). Change in heart rate was negatively correlated with arousal index (Spearman $-2.88, p < 0.5$), AHI (Spearman $-0.29, p < 0.05$), and ODI (Spearman $-0.31, p < 0.05$) such that as arousals and respiratory event rates increase, change in heart rate across the night decreases.

Table 3.

Urine and blood pressure parameters in children with and without NE

	NE (mean \pm SD; n, %)	No NE (mean \pm SD; n, %)	Mean difference (95% confidence interval)
Fluid intake (L/24 h)	0.999 \pm 0.442	1.325 \pm 0.852	-0.064, 0.716
Sodium intake (mmol)	117 \pm 59	300 \pm 822	-320, 685
Osmolality pre-sleep (mOsm/L)	782 \pm 268	843 \pm 279	-137, 259
Osmolality post-sleep (mOsm/L)	929 \pm 153	860 \pm 234	-185, 47
Change in osmolality (mOsm/L)	130 \pm 217	28 \pm 219	-55, 258
Overnight volume voided (L) ^a	0.409 \pm 0.181	0.591 \pm 293	-0.067, 0.431
Overnight sodium excretion (mmol)*	60.3 \pm 29.8	89.4 \pm 50.9	4.2, 53.9
Average systolic z-score	0.025 \pm 0.83	0.065 \pm 0.90	-0.562, 0.641
Average diastolic z-score	-0.052 \pm 0.66	-0.16 \pm 0.61	-0.530, 0.322
Blood pressure category			
Normal blood pressure	10 (77%)	34 (71%)	
Elevated blood pressure	0	5 (10%)	
Hypertension – stage 1	3 (23%)	7 (15%)	
Hypertension – stage 2	0	2 (4%)	

^aData are included only for children with NE who did not have an episode of NE ($n = 6$) on the study; given the imbalance in subject numbers, statistical comparison may not be valid.

* $p < 0.05$, post hoc multivariate analysis shows that history of NE is not a significant predictor of overnight sodium excretion after adjustment for age.

View Large

Figure 1.

Blood pressure and heart measurements across the night in children with (triangles) and without (circles) a history of NE. Lines represent mean values across the night (solid – NE; dashed – no NE). While blood pressure changes across the night did not differ between groups, there was a mean difference in heart rate across the night of 11.4 ± 5.6 beats/min ($p < 0.05$) between groups.

Discussion

This is the first study to measure multiple physiological parameters, including urine osmolality and blood pressure, in an unselected cohort of children undergoing polysomnography. Our results demonstrate that children with NE, when compared to children without NE, have greater abnormalities in their polysomnographic respiratory parameters, including more central respiratory events, and blunting of normal heart rate depression across the night despite similar incidence of OSA between groups. No differences were seen with respect to measures of sleep disruption, urine osmolality, or hemodynamics between groups. Greater disruption in respiratory parameters with higher variability in oxygen and carbon dioxide parameters as well

as blunting of the decline in heart rate across the night are supportive of impairments in autonomic control in children with NE when compared to those without NE.

There is a clear association between OSA and NE though the mechanism of this association is still debated. One theory is that upper airway obstruction results in increased intrathoracic pressure swings with subsequent myocardial stretch and release of natriuretic peptides resulting in inhibition of the action of ADH. Levels of ADH are lower and 12-hour overnight urine volumes are higher in children with OSA when compared to control children; treatment with adenotonsillectomy normalizes ADH levels and reduces urine volume [18, 19]. The presence of NE with OSA further decreased pre-treatment ADH compared to OSA alone [19]. Brain natriuretic peptide (BNP) is elevated in children with OSA [20], children with OSA and NE [12, 19], and decreases after treatment with adenotonsillectomy [19, 20]. Adenotonsillectomy, however, does not correct NE in all children with approximately half of children continuing to have NE after surgery and changes in post-operative ADH and BNP levels do not differ in those with and without continued post-operative NE [9] suggesting that upper airway obstruction alone cannot explain the association between OSA and NE. In the present study, with similar levels of obstructive respiratory events between children with and without NE, we found no difference in urine osmolality overnight between groups. Looking at prior studies examining the pathway between increased intrathoracic pressure and NE, the majority report an index of total respiratory events without a separate index of obstructive respiratory events. Interestingly, the one study that did report an obstructive respiratory index showed a similar level of airway obstruction between children with and without NE [12] consistent to what we describe here. Together with our study results, the accumulated evidence suggests that airway obstruction leading to alterations in ADH and BNP, ADH inhibition, and subsequent increase in urine volume is insufficient to explain the association between OSA and NE.

Sleep disruption associated with OSA may explain the occurrence of NE for some children and the resolution of NE after adenotonsillectomy. Airway obstruction and the ensuing repetitive intermittent hypoxia lead to disruption of sleep though this is often harder to measure in children [21]. Acute sleep deprivation in healthy children leads to a suppression of ADH and aldosterone levels as well as a substantial increase in nocturnal diuresis, natriuresis, and urine volume [22]. Children with unresolved NE post-adenotonsillectomy have lower amounts of stage 2 sleep and higher amounts of slow-wave, or deep, sleep compared to those whose NE resolved post-operatively [11], a pattern of sleep stage changes that is associated with sleep deprivation, suggesting that resolution of NE is related to resolution of sleep deprivation. In addition to effects on urine parameters, sleep deprivation in healthy children leads to blunting of normal nighttime dipping for heart rate and blood pressure [22]. Not surprisingly, we failed to show a difference in sleep parameters between children with and without NE; that may be because we had similar incidence of OSA between groups or because standard measures of sleep deprivation are insensitive to detecting differences in children.

An alternative mechanism to explain the association between NE and OSA is a common pathway that contributes to the occurrence of both conditions independently with mechanisms related to sleep continuity being clear candidates. NE is a predictor of residual OSA post-adenotonsillectomy [23], supporting a shared factor between OSA and NE that is not improved by adenotonsillectomy. Inherent differences in arousal threshold, or the level of stimuli required to cause arousal from sleep, likely contributes to NE as arousal to bladder stimuli is the main difference between NE and nocturia [24]. Differences in arousal threshold are also seen between children with and without OSA whereby children with OSA, when compared to children without OSA, arouse at a higher inspiratory resistance load and have blunted arousal response to hypercapnia [25, 26]. Corticotropin-releasing factor (CRF), a neuropeptide with a major role in regulating the hypothalamic–pituitary–adrenocortical axis and autonomic function in response to stress, plays a role in regulating sleep and arousal whereby higher levels of CRF promote wake and arousal from sleep [27, 28]. A recent study examining CRF levels in children with and without NE demonstrated lower CRF levels in children with NE in both the evening and morning sample [29], suggesting that lower levels of CRF may contribute to differences in arousal threshold in children with NE. Urocortins, members of the CRF family of peptides, were identified as a strong candidate biomarker for the discrimination of OSA from primary snoring in children [30]. CRF-related peptides are implicated in the regulation of autonomic cardiovascular and respiratory control [31], so could explain the differences in the blunting of nocturnal heart rate dipping and increase in central respiratory events seen in association with NE in our study. While further work is needed to characterize the CRF pathway in NE and OSA, its relationship to sleep continuity, autonomic function, and cardiorespiratory control makes it a highly plausible candidate to explain the association between NE and OSA as well as explaining each disorder independently.

Aspects of our methodology may impact interpretation of our results. We characterized children as having a history of NE but did not differentiate primary from secondary NE; it is possible that mechanisms responsible for NE differ between primary and secondary NE. Our subject sample included a high proportion of children undergoing polysomnography for OSA which may limit the application of our results to a more general population of children with NE. OSA is a known risk factor for NE but mechanisms responsible for NE in the context of OSA may differ from those in children without OSA. Our measurements were taken on a single night in an environment with inherent variability and challenges to a child's normal sleep routine; this alone could alter physiologic parameters unrelated to NE. We did, however, see that a higher proportion of children with NE had enuresis on the study night, suggesting that the laboratory setting maintained relevant differences between the groups. We also limited our measurements to those that could be conducted during the time of the polysomnography. This meant that we used 12-hour urine sodium output as a substitute for a 24-hour collection and only recorded nocturnal blood pressure. To reduce the likelihood of missing important differences between

groups, we included multiple measures and compared both average parameters as well as examining the pattern of change of these parameters across the night.

Our results raise the possibility that autonomic disruption is an important contributor to NE and a plausible link between NE and OSA in children. Other than blunting of normal heart rate dipping, we failed to find defects in urine concentrating ability or hemodynamic parameters between children with and without NE. Further work is needed to understand the mechanisms responsible for NE; the CRF pathway appears to be a good candidate to explain the strong association between NE and OSA. Understanding the link between these conditions will not only shed light on their shared pathophysiology but also has the potential to lead to novel treatment strategies

Acknowledgments

We are grateful to the children and parents/guardians who participated in the study. We acknowledge the contributions of the staff of the Pediatric Sleep Laboratory for their efforts in placing blood pressure monitors and collecting urine samples. We are grateful to Lesley Perez for her assistance in analyzing dietary records. This work was performed at the University of Alberta and the Stollery Children's Hospital.

Funding

This research was supported by a Resident/Fellow Trainee Grant awarded to A.B. through the generosity of the Stollery Children's Hospital Foundation through the Women & Children's Health Research Institute, University of Alberta, Edmonton, Alberta, Canada.

Conflict of interest statement. None declared.

References

1. Firoozi F, et al. Resolution of diurnal incontinence and nocturnal enuresis after adenotonsillectomy in children. *J Urol* . 2006;175(5):1885–1888; discussion 1888.
[Google Scholar](#) [Crossref](#) [PubMed](#)
2. Wolfe-Christensen C, et al. Lower health related quality of life and psychosocial difficulties in children with monosymptomatic nocturnal enuresis—is snoring a marker of severity? *J Urol* . 2013;190 (Suppl. 4):1501–1504.
[Google Scholar](#) [Crossref](#) [PubMed](#)
3. Robson WL. Clinical practice. Evaluation and management of enuresis. *N Engl J Med* . 2009;360(14):1429–1436.
[Google Scholar](#) [Crossref](#) [PubMed](#)
4. Dehoorne JL, et al. Characteristics of a tertiary center enuresis population, with special emphasis on the relation among nocturnal diuresis, functional bladder capacity and desmopressin response. *J Urol* . 2007;177(3):1130–1137.
[Google Scholar](#) [Crossref](#) [PubMed](#)
5. Jeyakumar A, et al. The association between sleep-disordered breathing and enuresis in children. *Laryngoscope* . 2012;122(8):1873–1877.
[Google Scholar](#) [Crossref](#) [PubMed](#)
6. Alexopoulos EI, et al. Nocturnal enuresis is associated with moderate-to-severe obstructive sleep apnea in children with snoring. *Pediatr Res* . 2014;76(6):555–559.
[Google Scholar](#) [Crossref](#) [PubMed](#)
7. Bascom A, et al. High risk of sleep disordered breathing in the enuresis population. *J Urol* . 2011;186 (Suppl. 4):1710–1713.

8. Stone J, et al. Symptoms of sleep-disordered breathing in children with nocturnal enuresis. *J Pediatr Urol* . 2008;4(3):197–202.
[Google Scholar](#) [Crossref](#) [PubMed](#)
9. Kovacevic L, et al. Why does adenotonsillectomy not correct enuresis in all children with sleep disordered breathing? *J Urol* . 2014;191 (Suppl. 5):1592–1596.
[Google Scholar](#) [Crossref](#) [PubMed](#)
10. Kovacevic L, et al. Enuretic children with obstructive sleep apnea syndrome: should they see otolaryngology first? *J Pediatr Urol* . 2013;9(2):145–150.
[Google Scholar](#) [Crossref](#) [PubMed](#)
11. Thottam PJ, et al. Sleep architecture parameters that predict postoperative resolution of nocturnal enuresis in children with obstructive sleep apnea. *Ann Otol Rhinol Laryngol* . 2013;122(11):690–694.
[Google Scholar](#) [Crossref](#) [PubMed](#)
12. Sans Capdevila O, et al. Increased morning brain natriuretic peptide levels in children with nocturnal enuresis and sleep-disordered breathing: a community-based study. *Pediatrics* . 2008;121(5):e1208–e1214.
[Google Scholar](#) [Crossref](#) [PubMed](#)
13. Brockmann PE, et al. Risk factors and consequences of excessive autonomic activation during sleep in children. *Sleep Breath* . 2011;15(3):409–416.
[Google Scholar](#) [Crossref](#) [PubMed](#)
14. Vande Walle J, et al. ; American Academy of Pediatrics; European Society for Paediatric Urology; European Society for Paediatric Nephrology; International Children's Continence Society. Practical consensus guidelines for the management of enuresis. *Eur J Pediatr* . 2012;171(6):971–983.
[Google Scholar](#) [Crossref](#) [PubMed](#)
15. Iber C, et al. *The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and Technical Specifications* . Westchester, IL: American Academy of Sleep Medicine; 2007.
16. Flynn JT, et al. Clinical practice guideline for screening and management of high blood pressure in children and adolescents.[Erratum appears in *Pediatrics*. 2017;140(6)]. *Pediatrics* . 2017;140(3):e20171904.
17. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* . 2004;114 (Suppl. 2, 4th Report):555–576.
[Crossref](#) [PubMed](#)
18. Yue Z, et al. Secretion of antidiuretic hormone in children with obstructive sleep apnea-hypopnea syndrome. *Acta Otolaryngol* . 2009;129(8):867–871.
[Google Scholar](#) [Crossref](#) [PubMed](#)
19. Kovacevic L, et al. Adenotonsillectomy normalizes hormones and urinary electrolytes in children with nocturnal enuresis and sleep-disordered breathing. *Urology* . 2015;86(1):158–161.
[Google Scholar](#) [Crossref](#) [PubMed](#)
20. Goldbart AD, et al. B-type natriuretic peptide and cardiovascular function in young children with obstructive sleep

apnea. *Chest* . 2010;138(3):528–535.

[Google Scholar](#) [Crossref](#) [PubMed](#)

21. Coussens S, et al. Movement distribution: a new measure of sleep fragmentation in children with upper airway obstruction. *Sleep* . 2014;37(12):2025–2034.
[Google Scholar](#) [Crossref](#) [PubMed](#)
22. Mahler B, et al. Sleep deprivation induces excess diuresis and natriuresis in healthy children. *Am J Physiol Renal Physiol* . 2012;302(2):F236–F243.
[Google Scholar](#) [Crossref](#) [PubMed](#)
23. Huang YS, et al. Treatment outcomes of adenotonsillectomy for children with obstructive sleep apnea: a prospective longitudinal study. *Sleep* . 2014;37(1):71–76.
[Google Scholar](#) [Crossref](#) [PubMed](#)
24. Goessaert AS, et al. Nocturnal enuresis and nocturia, differences and similarities—lessons to learn? *Acta Clin Belg* . 2015;70(2):81–86.
[Google Scholar](#) [Crossref](#) [PubMed](#)
25. Marcus CL, et al. Response to inspiratory resistive loading during sleep in normal children and children with obstructive apnea. *J Appl Physiol (1985)* . 1999;87(4):1448–1454.
[Google Scholar](#) [Crossref](#) [PubMed](#)
26. Marcus CL, et al. Arousal and ventilatory responses during sleep in children with obstructive sleep apnea. *J Appl Physiol (1985)* . 1998;84(6):1926–1936.
[Google Scholar](#) [Crossref](#) [PubMed](#)
27. Sanford LD, et al. Mouse strain differences in the effects of corticotropin releasing hormone (CRH) on sleep and wakefulness. *Brain Res* . 2008;1190:94–104.
[Google Scholar](#) [Crossref](#) [PubMed](#)
28. Kimura M, et al. Conditional corticotropin-releasing hormone overexpression in the mouse forebrain enhances rapid eye movement sleep. *Mol Psychiatry* . 2010;15(2):154–165.
[Google Scholar](#) [Crossref](#) [PubMed](#)
29. Motawie AA, et al. Assessment of serum level of corticotropin-releasing factor in primary nocturnal enuresis. *J Pediatr Urol* . 2017;13(1):46.e1–46.e5.
[Google Scholar](#) [Crossref](#)
30. Snow A, et al. Urinary proteins for the diagnosis of obstructive sleep apnea syndrome. *Methods Mol Biol* . 2010;641:223–241.
[Google Scholar](#) [Crossref](#) [PubMed](#)
31. Le Mével JC, et al. Central hyperventilatory action of the stress-related neurohormonal peptides, corticotropin-releasing factor and urotensin-I in the trout *Oncorhynchus mykiss*. *Gen Comp Endocrinol* . 2009;164(1):51–60.
[Google Scholar](#) [Crossref](#) [PubMed](#)

Topic:

[autonomic nervous system](#)

[heart rate](#)

[blood pressure](#)

[child](#)

[polysomnography](#)

[sleep](#)

[urine](#)

[urine osmolality](#)

[nocturnal enuresis](#)

[apnea-hypopnea index procedure](#)

Issue Section: [Sleep Across the Lifespan](#)



[View Metrics](#)

Email alerts

[New issue alert](#)

[Advance article alerts](#)

[Article activity alert](#)

[Subject alert](#)

[Receive exclusive offers and updates from Oxford Academic](#)

More on this topic

Obstructive Sleep Apnea Impairs
Postexercise Sympathovagal Balance in
Patients with Metabolic Syndrome

REM sleep without atonia with REM sleep–
related motor events: broadening the
spectrum of REM sleep behavior disorder

Prevalence and correlates of obstructive sleep
apnea among African Americans: the Jackson
Heart Sleep Study

Associations between polysomnography and actigraphy-based sleep indices and glycemic control among those with and without type 2 diabetes: the Multi-Ethnic Study of Atherosclerosis

Related articles in

Google Scholar

Related articles in PubMed

Pre-eclampsia/Eclampsia.

Cardiovascular Events and Mortality in White Coat Hypertension: A Systematic Review and Meta-analysis.

Blunted sympathoadrenal activation accompanies hemodynamic stability after early ventilation and delayed cord clamping at birth in preterm lambs.

Serum NGF levels may be associated with intrauterine antiepileptic exposure-related developmental problems.

Citing articles via

Google Scholar

CrossRef

Latest | **Most Read** | **Most Cited**

Characterization of the sleep disorder of anti-IgLON5 disease

Actigraphic detection of periodic limb movements: development and validation of a potential device-independent algorithm. A proof of concept study

Simultaneous tonic and phasic REM sleep without atonia best predicts early phenoconversion to neurodegenerative disease in idiopathic REM sleep behavior disorder

Residual symptoms after natural remission of insomnia: associations with relapse over 4 years

Sleep duration and fragmentation in relation to leukocyte DNA methylation in adolescents

Looking for your next opportunity?

Chair of Pain Research
Boston, Massachusetts

PEDIATRIC EMERGENCY PHYSICIAN
Saskatoon Shines, Saskatchewan

Endowed Chair of Occupational
Health/Medicine
Saint John, New Brunswick

CHIEF OF THE DIVISION OF ALLERGY,
IMMUNOLOGY AND INFECTIOUS
DISEASE
New Brunswick, New Jersey

[View all jobs](#)

OXFORD
UNIVERSITY PRESS

[About SLEEP](#)

[Editorial Board](#)

[Author Guidelines](#)

[Facebook](#)

[Twitter](#)

[Contact Us](#)

[Purchase](#)

[Recommend to your Library](#)

[Advertising and Corporate Services](#)

[Journals Career Network](#)

Online ISSN 1550-9109

Print ISSN 0161-8105

Copyright © 2019 Sleep Research Society

[About Us](#)

[Contact Us](#)

[Careers](#)

[Help](#)

[Access & Purchase](#)

[Rights & Permissions](#)

[Open Access](#)

Resources

[Authors](#)

[Librarians](#)

[Societies](#)

[Sponsors & Advertisers](#)

[Press & Media](#)

[Agents](#)

Connect

[Join Our Mailing List](#)

[OUPblog](#)

[Twitter](#)

[Facebook](#)

[YouTube](#)

[Tumblr](#)

Explore

[Shop OUP Academic](#)

[Oxford Dictionaries](#)

[Oxford Index](#)

[Epigeum](#)

[OUP Worldwide](#)

[University of Oxford](#)

*further the University's objective of excellence in research, scholarship,
and education by publishing worldwide*

Copyright © 2019 Oxford University Press
Accessibility

Get Adobe Reader

Cookie Policy

Privacy Policy

Legal Notice

Site Map