

# Hydrocarbons in soil and meltwater stream sediments near Artigas Antarctic Research Station: origin, sources and levels

CAROLINA RODRÍGUEZ<sup>1</sup>, KAREN IGLESIAS<sup>1</sup>, MÁRCIA C. BÍCEGO<sup>2</sup>, SATIE TANIGUCHI<sup>2</sup>,  
SILVIO TAROU SASAKI<sup>2</sup>, NOELIA KANDRATAVICIUS<sup>3</sup>, CAROLINA BUENO<sup>3</sup>, ERNESTO BRUGNOLI<sup>3</sup>  
and NATALIA VENTURINI<sup>1\*</sup>

<sup>1</sup>Laboratorio de Biogeoquímica Marina (LABIM), Instituto de Ecología y Ciencias Ambientales, Facultad de Ciencias, Universidad de la República (UdelaR), Iguá 4225, Montevideo 11400, Uruguay

<sup>2</sup>Laboratório de Química Orgânica Marinha (LabQOM), Instituto Oceanográfico da Universidade de São Paulo, Praça do Oceanográfico 191, Cidade Universitária, 05508-120, São Paulo, Brazil

<sup>3</sup>Oceanografía y Ecología Marina, Instituto de Ecología y Ciencias Ambientales, Facultad de Ciencias, UdelaR, Iguá 4225, Montevideo 11400, Uruguay

\*Corresponding author: rulo@fcien.edu.uy

**Abstract:** Aliphatic and polycyclic aromatic hydrocarbons (AHs and PAHs) were investigated in soil and meltwater stream sediments near the Uruguayan Artigas Research Station (BCAA). Gas chromatography with flame ionization detection and gas chromatography mass spectrometry were used to determine the composition of AHs and PAHs, respectively. Total AH concentrations were in the range 0.57–2333  $\mu\text{g g}^{-1}$ , while total PAH concentrations were in the range 1.36–51 650  $\text{ng g}^{-1}$ . Based on AH and PAH concentrations, sites in the service area and next to the boat storeroom are highly contaminated, while the other sites sampled have moderate to low contamination levels or are not impacted. High unresolved complex mixture concentrations indicate the occurrence of previous petrogenic contamination, but the dominance of low molecular weight and alkyl PAHs indicate recent oil introductions. Anthropogenic hydrocarbons dominate and are related to diesel fuel and organic residue combustion, fuel storage and boat traffic. Petrogenic contamination is relatively high, as at other Antarctic stations, but is restricted to the boat storeroom and the service area where the incinerator, the generator room and fuel tanks are located. Improvements made in fuel management procedures and the dismantling of the old fuel tanks will reduce the risk of diesel leaks and their impact on the environment near the BCAA. This study provides reference information for future environmental monitoring.

Received 21 July 2017, accepted 14 December 2017, first published online 15 February 2018


**Key words:** King George Island, n-alkanes, polycyclic aromatic compounds, unresolved complex mixture, Uruguayan research station

## Introduction

Hydrocarbon contamination of Antarctic soils and sediments has been demonstrated over recent years at a range of sites (e.g. Aislabie *et al.* 1999, Martins *et al.* 2004, Curtosi *et al.* 2007, Bícego *et al.* 2009, Dauner *et al.* 2015, Prus *et al.* 2015), despite the low human presence on the Antarctic continent. Logistical support for scientific operations, growing tourism and fisheries have been reported as the major sources of hydrocarbon pollution in Antarctica (Mazzera *et al.* 1999, Curtosi *et al.* 2007, Tin *et al.* 2009). Hydrocarbons are derived from both natural and anthropogenic sources. Local vegetation such as mosses, lichens and vascular plants represent natural sources of hydrocarbons in Antarctica (e.g. Cabrerizo *et al.* 2016). Anthropogenic sources are associated with fossil fuel combustion, fuel storage, accidental oil spills and sewage discharge (Aislabie *et al.* 1999, Curtosi *et al.* 2007, Bícego *et al.* 2009, Cabrerizo *et al.* 2016). Some polycyclic

aromatic hydrocarbons (PAHs) are teratogenic, mutagenic and carcinogenic, and are considered as priority pollutants because of their high toxicity (USEPA 1985). The impact of hydrocarbon pollution even at a local level on the Antarctic continent is very significant because of its relative pristine environment and sensitive communities. In addition, except in high-energy marine environments where dispersion occurs rapidly, hydrocarbon pollution has proved long-lived in Antarctica (Tin *et al.* 2009). The low temperatures occurring in Antarctica delay the biological degradation of hydrocarbons in contaminated soils, where they are generally highly persistent. However, information about the origin (biogenic or anthropogenic) of these compounds is also preserved under low temperature conditions, so potential sources can be investigated (Prus *et al.* 2015, Cabrerizo *et al.* 2016).

Some significant but mainly localized levels of organic pollutants in Antarctic marine and terrestrial environments have been related to human activities,



**Fig. 1.** Location of the Artigas Research Station (BCAA) on King George Island, South Shetland Islands Archipelago, Antarctica.

especially at coastal research stations, including fuel storage and transportation, sewage release and waste disposal (e.g. Curtosi *et al.* 2007, Bicego *et al.* 2009, Martins *et al.* 2010, Préndez *et al.* 2011, Dauner *et al.* 2015).


In 1991, after the signing of the Protocol on Environmental Protection (Madrid Protocol), Consultative Parties of the Antarctic Treaty recognized the need to improve management practices and mitigating measures, for present and past environmental impacts (Aislabie *et al.* 2004). Uruguay, as a Consultative Party following the Madrid Protocol, has a responsibility both for the preservation and conservation of Antarctica and a duty to assess the environmental impact of its activities in Antarctica including the establishment of mitigation

measures (Aislabie *et al.* 2004). The aim of this study was to assess the origin, main sources, levels and distribution of aliphatic hydrocarbons (AHs) and PAHs in the area close to the Artigas Antarctic Research Station (BCAA) and to evaluate the extent of areas potentially affected by anthropogenic hydrocarbon inputs due to human activities at the BCAA.

## Materials and methods

### *Study site and sampling methods*

The BCAA is located at 62°11'4"S and 58°51'7"W in Collins Bay, a part of Maxwell Bay, on the Fildes



**Fig. 2.** Sketch of the Artigas Research Station (BCAA) showing the location of the sampling sites (based on a Google Earth image). The 11 buildings of the BCAA are: A = radio, B = AINA (scientist staff bedrooms and labs), C = head of station house, D = cavern (water deposit and gym), E = crew housing, F = supply depot, G = nursery, H = kitchen/dining room, I = hangar, J = service area (incinerator, generator room, old and new fuel tanks) and K = boat storeroom. Data from station T12 is not available.

Peninsula, King George Island, South Shetland Islands (Fig. 1). The BCAA with an area of *c.* 30 000 m<sup>2</sup> constitutes the main Uruguayan Antarctic station. It was founded in 1984 and is open throughout the year providing logistical support to activities of the Uruguayan Antarctic Programme. The BCAA consists of 11 buildings including the radio, AINA (scientific staff bedrooms and labs), head of station house, cavern (water deposit and gym), crew housing, supply depot, nursery, kitchen/dining room, hangar, service area (generator room, incinerator, old and new fuel tanks) and boat storeroom (Figs 1 & 2). Three meltwater streams (named for this work: AINA's stream, kitchen's stream and tank's stream) flow across the BCAA into Collins Bay (Fig. 2). The permanent population of the BCAA is composed of nine people, with up to 60 people staying during the summer months. As in most Antarctic stations, diesel generators provide the energy for the BCAA. Residue waste produced in the BCAA is classified, compacted, stored and ultimately removed from Antarctica or incinerated on site following Annex III of the Madrid Protocol. In addition, inspections of the installations and activities developed in the BCAA are regularly performed (e.g. ATIPR 2013).

Soil and surface sediment samples were collected in January and early February 2015 at 26 sites positioned using a GPS (see Table I for co-ordinates). Fourteen sites were located in the ground adjacent to the BCAA buildings ('T' sites), while the other 12 ('C' sites) were positioned throughout the areas of the meltwater streams close to the BCAA (Fig. 2). The top layer (0–2 cm) of undisturbed soil or sediment was collected using a stainless steel scoop or a corer, respectively. Samples were conditioned in pre-muffled (450°C, 4 h) aluminium containers and stored at -20°C. The material was freeze-dried, carefully homogenized in a mortar and kept in clean glass bottles until laboratory analysis. At each site, a second soil or sediment sample was taken for grain size analysis.

#### *Analytical methods*

Grain size analysis was performed by the standard dry sieve method as described in Suguio (1973). Gravel and different sand classes were separated from fine sediments (< 63 µm; silt + clay).

The analytical method for hydrocarbons was adapted from UNEP (1992). Aliquots of 20 g of soil or sediment

**Table 1.** Co-ordinates and percentage of grain size fractions in soil and sediment samples from Artigas Station.


Site	Latitude	Longitude	Gravel	Coarse sand	Medium sand	Fine sand	Silt + clay
<b>Soil</b>							
T0	62°11'0.13"S	58°53'55.28"W	51.10	21.38	8.20	9.61	9.69
T1	62°11'3.60"S	58°54'19.03"W	23.70	53.02	7.92	7.15	8.32
T2	62°11'5.82"S	58°54'16.35"W	80.81	6.58	3.47	4.78	4.58
T3	62°11'6.16"S	58°54'15.70"W	94.10	3.29	0.67	0.71	1.25
T4	62°11'7.58"S	58°54'13.25"W	65.24	30.26	1.87	1.11	1.68
T5	62°11'2.17"S	58°54'14.91"W	44.37	40.83	4.93	4.44	5.48
T6	62°11'3.71"S	58°54'12.61"W	46.53	36.23	5.03	6.02	6.11
T7	62°11'5.53"S	58°54'12.09"W	69.51	19.96	3.33	2.37	4.92
T8	62°11'6.76"S	58°54'10.21"W	67.18	25.35	2.16	2.15	3.15
T9	62°11'1.14"S	58°54'10.58"W	21.57	55.99	4.37	10.78	7.38
T10	62°11'2.36"S	58°54'5.66"W	42.59	41.77	5.64	3.63	6.35
T11	62°11'2.29"S	58°54'4.11"W	11.07	44.81	24.99	12.33	6.16
T13	62°11'2.81"S	58°54'4.15"W	8.95	75.99	5.69	4.81	4.26
T14	62°11'4.91"S	58°54'1.78"W	37.23	22.78	22.48	0.00	17.70
<b>Sediment</b>							
C0	62°11'2.23"S	58°54'24.81"W	55.40	19.48	6.73	5.57	12.76
C1	62°11'4.33"S	58°54'20.84"W	34.80	13.88	3.54	15.00	32.71
C2	62°11'6.12"S	58°54'18.12"W	51.42	10.39	7.61	9.67	18.97
C3	62°11'6.66"S	58°54'17.09"W	n.a.	n.a.	n.a.	n.a.	n.a.
C4	62°11'8.51"S	58°54'15.40"W	33.50	55.94	2.48	1.98	6.06
C5–C9	62°11'1.72"S	58°54'14.13"W	25.42	66.40	4.02	2.15	1.96
C6	62°11'3.54"S	58°54'11.77"W	55.47	37.75	2.63	1.49	2.65
C7	62°11'4.90"S	58°54'9.82"W	73.29	21.77	1.19	0.98	2.71
C8	62°11'6.28"S	58°54'8.09"W	n.a.	n.a.	n.a.	n.a.	n.a.
C10	62°11'3.25"S	58°54'8.63"W	18.80	63.15	10.30	5.24	2.43
C11	62°11'5.49"S	58°54'4.93"W	26.28	57.91	6.64	3.90	5.56
C12	62°11'6.90"S	58°54'3.36"W	n.a.	n.a.	n.a.	n.a.	n.a.

n.a. = not available.

were Soxhlet extracted for 8 h with dichloromethane (DCM)/n-hexane (1:1, v:v). A mixture of surrogates (hexadecene and eicosene to AHs and naphthalene-d<sub>8</sub>, acenaphthene-d<sub>10</sub>, phenanthrene-d<sub>10</sub>, crysene-d<sub>12</sub> and perylene-d<sub>12</sub> to PAHs) was added to each sample before extraction. Extracts were concentrated on a rotary evaporator to 1 ml and fractionated by column chromatography using 5% deactivated alumina and silica. Organic compounds were eluted with 10 ml of n-hexane (F1, AHs) and 15 ml of DCM/n-hexane (3:7) mixture (F2, PAHs). F1 and F2 were concentrated to 1 ml and the internal standards tetradecene (AHs) and benzo [b]fluoranthene-d<sub>12</sub> (PAHs) were respectively added before gas chromatographic (GC) analysis. The AHs were analysed in an Agilent GC (model 6890) with flame ionization detector (GC-FID) and a capillary fused silica column coated with 5% diphenyl dimethylsiloxane (50 m length, 0.32 mm ID and 0.17 µm film thickness). Hydrogen was used as the carrier gas. The oven temperature for AH detection was programmed to ramp from 40°C, held for 2 min, 40–60°C at 20°C per min, then increased to 290°C at 5°C per min, held for 5 min, and finally increased to 300°C at 10°C per min, remaining constant for 10 min. The injector and detector temperature was 300°C and 325°C, respectively. The PAHs were analysed in an Agilent GC (model 6890)

coupled to an Agilent mass spectrometer detector (model 5973N) (GC/MS) and a capillary fused silica column coated with 5% diphenyl dimethylsiloxane (30 m length, 0.25 mm ID and 0.25 µm film thickness). Helium was used as the carrier gas. The oven temperature ramping programme was the same as used for AHs. Data acquisition was undertaken in the SIM mode and the Agilent enhanced chemstation G1701 CA was used to perform the measurements. Individual PAHs were identified and confirmed matching their retention times and their mass to charge ratios with the results of standard mixtures. Quantification was performed by the surrogate peaks area method and a five-point calibration curve for each class of compounds analysed ( $r > 0.995$ ). Procedural blanks were performed for each group of ten samples and no peaks interfere with the analyses of target compounds. Recovery of surrogates ranged from 70–98% for AHs and PAHs. Detection limits (DLs) were obtained as three times the standard deviation of the signal in the same retention time of compounds analysed in the blanks. The DL was 0.001–0.034 µg g<sup>-1</sup> for n-alkanes and 1.00–3.70 ng g<sup>-1</sup> for PAHs. Concentrations of target compounds in reference material provided by the International Atomic Energy Agency (IAEA-408) were also analysed and their concentrations were in agreement with certified values. The 36 PAH compounds detected in the samples





**Fig. 3.** Total aliphatic hydrocarbon (AH) concentrations ( $\mu\text{g g}^{-1}$ ) in soil and meltwater stream sediments from Artigas Station.

were naphthalene, methylnaphthalene, biphenyl, ethylnaphthalene, dimethylantracene, acenaphthylene, acenaphthene, trimethylnaphthalene, fluorene, methylfluorene, dibenzothiophene phenanthrene, anthracene, dimethylfluorene, methyl dibenzothiophene, methylphenanthrene, dimethyldibenzothiophene, dimethylphenanthrene, fluoranthene, pyrene, methylfluoranthene, rethene, methylpyrene, benzo[c]phenanthrene, benzo[a]anthracene,

chrysene, methylchrysene, dimethylchrysene, benzo[b]fluoranthene, benzo[j]fluoranthene, benzo[k]fluoranthene, benzo[e]pyrene, benzo[a]pyrene, perylene, indeno[1,2,3-c,d]pyrene and benzo[g,h,i]perylene.

Since natural samples consist of complex hydrocarbon mixtures, several diagnostic indices, in addition to absolute concentrations, were used to identify the origin (anthropogenic or biogenic) and probable sources of AHs


**Table II.** Aliphatic hydrocarbon (AH) concentrations and related parameters in soil and sediments from Artigas Station.

Site	$\Sigma$ AHs ( $\mu\text{g g}^{-1}$ )	$\Sigma$ n-alkanes ( $\mu\text{g g}^{-1}$ )	Most abundant n-alkane	Short-chain n-alkanes ( $\mu\text{g g}^{-1}$ )	Long-chain n-alkanes ( $\mu\text{g g}^{-1}$ )	UCM ( $\mu\text{g g}^{-1}$ )	% UCM	CPI <sub>1</sub>	CPI <sub>2</sub>	Pri/Phy
<b>Soil</b>										
T0	0.57	0.39	n-C <sub>25</sub>	0.073	0.152	n.d.	n.c.	0.48	7.03	n.c.
T1	9.76	2.86	n-C <sub>14</sub>	1.484	0.884	n.d.	n.c.	0.57	1.76	1.23
T2	3.72	2.01	n-C <sub>29</sub>	0.108	1.647	n.d.	n.c.	1.27	1.89	n.c.
T3	2.15	1.68	n-C <sub>14</sub>	0.954	0.43	n.d.	n.c.	0.20	3.79	n.c.
T4	1.49	1.30	n-C <sub>27</sub>	0.169	0.745	n.d.	n.c.	0.87	3.25	n.c.
T5	11.46	8.30	n-C <sub>16</sub> , n-C <sub>18</sub>	5.493	1.386	n.d.	n.c.	0.76	2.09	1.38
T6	13.82	9.61	n-C <sub>16</sub>	7.955	0.734	n.d.	n.c.	0.71	1.50	1.16
T7	2.29	2.15	n-C <sub>12</sub>	1.267	0.69	n.d.	n.c.	0.18	1.99	0.37
T8	4.71	4.06	n-C <sub>14</sub>	2.61	0.85	n.d.	n.c.	0.22	2.46	0.41
T9	6.09	5.20	n-C <sub>14</sub>	1.94	1.66	n.d.	n.c.	0.23	1.25	0.94
T10	32.28	0.84	n-C <sub>18</sub>	0.71	0.13	27.96	86.63	0.74	5.30	0.75
T11	1009	3.42	n-C <sub>16</sub>	1.23	1.23	855	84.73	0.91	1.26	1.57
T12	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
T13	2332	351	n-C <sub>16</sub> , n-C <sub>17</sub>	220	26.61	1544	66.23	1.07	1.16	1.14
T14	2.20	1.92	n-C <sub>14</sub>	1.25	0.31	n.d.	n.c.	0.26	2.41	0.66
<b>Sediment</b>										
C0	5.48	3.71	n-C <sub>14</sub>	3.29	0.13	n.d.	n.c.	0.30	4.55	1.44
C1	2.57	2.05	n-C <sub>14</sub>	1.5	0.16	n.d.	n.c.	0.24	6.62	0.88
C2	2.80	2.12	n-C <sub>14</sub>	1.82	0.1	n.d.	n.c.	0.11	5.92	0.80
C3	2.48	1.78	n-C <sub>14</sub>	1.33	0.16	n.d.	n.c.	0.20	6.55	1.05
C4	5.56	4.41	n-C <sub>14</sub>	3.08	0.64	n.d.	n.c.	0.13	1.31	0.95
C5–C9	1.73	1.26	n-C <sub>14</sub>	1.15	n.d.	n.d.	n.c.	0.22	n.c.	0.69
C6	31.13	2.15	n-C <sub>14</sub> , n-C <sub>17</sub>	1.34	0.41	n.d.	n.c.	1.10	2.24	1.52
C7	33.23	2.61	n-C <sub>14</sub>	1.95	0.31	26.39	79.42	1.59	2.69	0.79
C8	170	5.56	n-C <sub>16</sub>	5.25	n.d.	136	80.20	0.04	n.c.	n.c.
C10	2.14	1.68	n-C <sub>14</sub>	1.18	0.17	n.d.	n.c.	0.27	3.64	0.80
C11	2.60	1.79	n-C <sub>14</sub>	1.05	0.29	n.d.	n.c.	0.32	2.28	0.73
C12	2.03	1.65	n-C <sub>14</sub>	1.51	2.07	n.d.	n.c.	0.05	2.07	n.c.

CPI<sub>1</sub> =  $1/2 (\Sigma \text{ odd } C_{15-21} / \Sigma \text{ even } C_{14-20}) + \Sigma \text{ odd } C_{15-21} / \Sigma \text{ even } C_{16-22}$ .

CPI<sub>2</sub> =  $1/2 (\Sigma \text{ odd } C_{25-35} / \Sigma \text{ even } C_{24-34}) + \Sigma \text{ odd } C_{25-35} / \Sigma \text{ even } C_{26-36}$ .

CPI = Carbon Preference Index for short-chain (<C<sub>21</sub>) n-alkanes, CPI<sub>2</sub> = Carbon Preference Index for long-chain (>C<sub>25</sub>) n-alkanes, n.a. = not available, n.c. = not calculated, n.d. = not detected, Phy = phytane, Pri = pristane, UCM = unresolved complex mixture.



**Fig. 4.** Total polycyclic aromatic hydrocarbon (PAH) concentrations ( $\text{ng g}^{-1}$ ) in soil and meltwater stream sediments from Artigas Station.

and PAHs in soil and surface sediments. Spearman's correlation analysis was carried out with soil/sediment fractions, AHs and PAHs to explore any relationship

**Table III.** Polycyclic aromatic hydrocarbon (PAH) concentrations and related parameters in soil and sediments from Artigas Station.

Site	$\Sigma$ PAHs ( $\text{ng g}^{-1}$ )	$\Sigma$ 16 PAHs ( $\text{ng g}^{-1}$ )	LMW PAHs ( $\text{ng g}^{-1}$ )	HMW PAHs ( $\text{ng g}^{-1}$ )	Alkylated PAHs ( $\text{ng g}^{-1}$ )	LMW/ HMW
<b>Soil</b>						
T0	1.36	n.d.	1.36	n.d.	1.36	n.c.
T1	268	61.4	260	8.06	184	32.3
T2	93.3	12.0	67.5	25.8	65.0	2.61
T3	19.3	3.91	19.3	n.d.	43.6	n.c.
T4	2.96	n.d.	2.96	n.d.	2.96	n.c.
T5	n.d.	n.d.	n.d.	n.d.	n.d.	n.c.
T6	156	22.2	149	7.35	117	20.3
T7	178	20.5	178	n.d.	157	n.c.
T8	207	140	207	n.d.	65.2	n.c.
T9	204	25.2	201	3.18	144	63.4
T10	2068	766	1202	865	925	1.39
T11	8537	516	7773	763	5291	10.18
T12	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
T13	51 650	2524	49 201	2448	34 577	20.0
T14	159	11.6	157	1.48	14 355	106
<b>Sediment</b>						
C0	423	54.3	420	2.82	344	149
C1	38.30	7.30	38.3	n.d.	49.9	n.c.
C2	24.30	4.28	24.3	n.d.	22.6	n.c.
C3	100	17.0	100	n.d.	71.2	n.c.
C4	143	15.6	142	1.23	122.00	115
C5–C9	69.7	5.54	69.7	n.d.	74.4	n.c.
C6	69.9	16.9	26.1	43.7	46.7	0.60
C7	75.6	15.5	43.6	32.0	54.6	1.36
C8	751.6	70.1	472	279	529	1.69
C10	52.7	8.24	52.7	n.d.	193	n.c.
C11	34.0	5.75	32.6	1.40	26.5	23.3
C12	21.5	5.22	21.5	n.d.	21.6	n.c.

$\Sigma$  16 PAHs = total priori compounds according to USEPA, LMW = low molecular weight (2–3 aromatic rings), HMW = high molecular weight (4–6 aromatic rings), n.a. = not available, n.c. = not calculated, n.d. = not detected.

between grain size and hydrocarbon distribution. A significance level of  $\alpha = 0.05$  was established for Spearman's correlation coefficients. Data analysis was performed using the software package BioEstat 4.0 (Ayres *et al.* 2005).

## Results


### Soil and sediment grain size

Coarse sediment fractions (gravel and coarse sand) dominated at most of the sites. The gravel fraction ranged between 8.95% (site T13) and 94.10% (site T3), and the coarse sand fraction from 3.29% (site T3) to 66.40% (sites C5–C9) (Table I). Mud (silt + clay) varied between 1.25% at site T3 and 32.71% at site C1. The highest mud content was recorded in sites T14, C0, C1 and C2 (Table I).

### Aliphatic hydrocarbons

Total AHs in soil samples close to the BCAA ranged between  $0.57 \mu\text{g g}^{-1}$  (site T0) and  $2332 \mu\text{g g}^{-1}$  (site T13). In sediment samples from the meltwater streams, AHs ranged from  $1.73$ – $170 \mu\text{g g}^{-1}$  (sites C5 and C8, respectively; Fig. 3 and Table II). The highest AH levels were recorded at sites C8, T11 and T13 (Fig. 3 and Table II).

Total n-alkane ( $\text{C}_{12}$ – $\text{C}_{36}$ ) concentrations in soil varied between  $0.39$ – $351 \mu\text{g g}^{-1}$  (sites T0 and T13, respectively; Table II), while in surface sediments they ranged between  $1.78 \mu\text{g g}^{-1}$  (site C5) and  $5.56 \mu\text{g g}^{-1}$  (site C8). Among the short-chain n-alkanes ( $< \text{C}_{21}$ ),  $\text{C}_{14}$ ,  $\text{C}_{16}$  and  $\text{C}_{18}$  were the most abundant, while among the long-chain homologues ( $> \text{C}_{25}$ ),  $\text{C}_{27}$  and  $\text{C}_{29}$  dominated. At most of the sites short-chain n-alkanes were present at higher concentrations compared with long-chain n-alkanes (Table II). The unresolved complex mixture (UCM) was detected only at sites T10, T11, T13, C7 and C8, and



**Fig. 5.** Concentrations of low molecular weight (LMW) and high molecular weight (HMW) polycyclic aromatic hydrocarbons (PAHs) in samples from Artigas Station (sites **a.** T1 and **b.** T13). \*Indicates concentration  $\leq 20.0 \text{ ng g}^{-1}$  or no detected compounds.

presented with the highest concentrations at site T13 ( $1544 \mu\text{g g}^{-1}$ ; Table II). At these sites the UCM corresponded to 66.23–86.63% of total AHs (see example chromatograms found at <http://dx.doi.org/10.1017/S0954102018000019>).

A Carbon Preference Index (CPI) for both short-chain ( $< C_{21}$ ) and long-chain ( $> C_{25}$ ) homologues was calculated. The  $\text{CPI}_1$  calculated for short-chain n-alkanes was 0.18–1.27 for ground sites (T7 and T2, respectively) and 0.04–1.59 for meltwater stream sites (C8 and C7, respectively) (Table II). The  $\text{CPI}_2$  calculated for long-chain n-alkanes was 1.16–7.03 for ground sites (T13 and T0, respectively) and 1.31–6.62 for meltwater stream sites (C4 and C1, respectively) (Table II). The pristane/phytane (Pri/Phy) ratios ranged from 0.37–1.57 (T7 and T11, respectively) and 0.73–1.52 (C11 and C6) for ground and meltwater stream sites, respectively (Table II). For several sites this ratio could not be calculated because of the absence of pristane.


#### Polycyclic aromatic hydrocarbons

Total PAH concentrations (expressed as the sum of the 36 compounds identified) in soil samples ranged from  $1.36 \text{ ng g}^{-1}$  at site T0 to  $51\,650 \text{ ng g}^{-1}$  at site T13 (Fig. 4 and Table III). In surface sediment samples, PAHs ranged from  $21.56$ – $751 \text{ ng g}^{-1}$  at sites C12 and C8, respectively (Fig. 4 and Table III). The highest total PAH

concentrations were recorded at sites T10, T11 and T13 (Fig. 4 and Table III).

Concentrations of the 16 PAHs considered as priority pollutants by the United State Environmental Protection Agency (USEPA) ranged between  $3.91 \text{ ng g}^{-1}$  (site T3) and  $2524 \text{ ng g}^{-1}$  (site T13) (Table III). Sites T10, T11 and T13 showed the highest levels of these compounds (Table III). Low molecular weight (LMW) PAHs (compounds with two to three aromatic rings) showed concentrations between  $1.36 \text{ ng g}^{-1}$  (site T0) and  $49\,201 \text{ ng g}^{-1}$  (site T13) (Table III). High molecular weight (HMW) PAHs (compounds with four to six aromatic rings) ranged from  $1.23$ – $2448 \text{ ng g}^{-1}$  at sites C4 and T13, respectively (Table III). In all the sites, alkylated homologues, such as methylanthracene, dimethylanthracene and trimethylanthracene, dominated (for examples see Fig. 5a & b). The highest concentration of alkylated PAHs was recorded at site T13 (Table III and Fig. 5b).

The LMW/HMW PAHs ratio was  $>1$  for most of the sites, except site C6 (Table III). The ratios anthracene/anthracene + phenanthrene (Ant/178), fluoranthene/fluoranthene + pyrene (Fl/202) and benzo[a]anthracene/benzo[a]anthracene + chrysene (Bz[a]A/228) were only calculated for sites T10, T11, T13 and C8, where these compounds were recorded. These ratios were used to distinguish different sources of PAHs. At sites T10, T11 and T13, the Ant/178 ratio was  $<0.10$  and the Fl/202 ratio showed values  $<0.40$  (Fig. 6a). Site C8 showed



**Fig. 6.** Plots of the polycyclic aromatic hydrocarbon (PAH) isomer pair ratios at sites T10, T11, T13 and C8 based on Yunker *et al.* (2002). **a.** Anthracene/(anthracene + phenanthrene) (Ant/178) vs fluoranthene/(fluoranthene + pyrene) (FI/220) and **b.** Benzo[a]anthracene/(benzo[a]anthracene + chrysene) (Bz[a]A/228) vs fluoranthene/(fluoranthene + pyrene) (FI/220).

Ant/178 > 0.10 and FI/202 > 0.50. The Bz[a]A/228 ratio presented values < 0.20 at sites T11, T13 and C8, and 0.20–0.35 at site T10 (Fig. 6b). Based on these ratios, the main sources of PAHs at these sites are petroleum (crude oil) and combustion.

#### Correlation matrix results

Among the different grain size fractions considered, only fine sand and mud (silt + clay) showed a significantly positive correlation with each other (Table IV). Neither AHs nor PAHs showed significant correlations with any of the grain size fractions. Whereas, total AHs, total n-alkanes, short-chain n-alkanes, long-chain n-alkanes, total PAHs, and LMW, HMW and alkylated compounds all presented significant highly positive correlations among themselves (Table IV). Thus, hydrocarbon deposition seems to be related to the proximity of the source but not connected with soil/sediment grain size. Both AHs and PAHs may derive from the same sources.

#### Discussion

##### Levels and sources of aliphatic hydrocarbons in soil and meltwater stream sediments near Artigas Station

At sites T11, T13 and C8, the concentration of total AHs exceeded the threshold value of  $100 \mu\text{g g}^{-1}$  proposed as an indicator of oil pollution in marine sediments (Volkman *et al.* 1992). Whereas, at the remaining sites AHs levels were below this limit. The high UCM concentration at sites T11, T13 and C8 indicates petroleum residues, supporting cumulative oil inputs and biodegraded oil (Commendatore *et al.* 2012). Sites T11 and T13 are located in the service area, adjacent to the generator room and the fuel tanks (old and new), respectively. Therefore,

**Table IV.** Matrix correlation results among soil/sediment fractions, aliphatic (AHs) and polycyclic aromatic hydrocarbons (PAHs).

	Gravel	Coarse sand	Medium sand	Fine sand	Silt + clay	$\Sigma$ AHs	$\Sigma$ n-alkanes	SC n-alkanes	LC n-alkanes	$\Sigma$ PAHs	LMW PAHs	HMW PAHs	Alkylated PAHs
Gravel	1	–	–	–	–	–	–	–	–	–	–	–	–
Coarse sand	-0.2813	1	–	–	–	–	–	–	–	–	–	–	–
Medium sand	-0.1926	0.2722	1	–	–	–	–	–	–	–	–	–	–
Fine sand	-0.1059	0.1954	0.3787	1	–	–	–	–	–	–	–	–	–
Silt + clay	0.0587	-0.1092	0.3352	<b>0.6581</b>	1	–	–	–	–	–	–	–	–
$\Sigma$ AHs	-0.3352	0.4047	0.2524	0.1555	-0.0843	1	–	–	–	–	–	–	–
$\Sigma$ n-alkanes	-0.2459	0.4036	0.0009	0.0203	-0.0698	<b>0.915</b>	1	–	–	–	–	–	–
SC n-alkanes	-0.2488	0.4001	-0.0033	0.0158	-0.0696	<b>0.9133</b>	<b>0.9999</b>	1	–	–	–	–	–
LC n-alkanes	-0.2464	0.3967	0.0016	0.0263	-0.0935	<b>0.9168</b>	<b>0.9944</b>	<b>0.9932</b>	1	–	–	–	–
$\Sigma$ PAHs	-0.2819	0.4181	0.1087	0.0768	-0.0714	<b>0.9669</b>	<b>0.9859</b>	<b>0.9848</b>	<b>0.9831</b>	1	–	–	–
LMW PAHs	-0.2795	0.4175	0.105	0.0759	-0.07	<b>0.9653</b>	<b>0.9874</b>	<b>0.9864</b>	<b>0.9848</b>	<b>0.9998</b>	1	–	–
HMW PAHs	-0.3111	0.4053	0.1701	0.0889	-0.0935	<b>0.9412</b>	<b>0.8996</b>	<b>0.8988</b>	<b>0.8939</b>	<b>0.9453</b>	<b>0.9395</b>	1	–
Alkylated PAHs	-0.2722	0.3609	0.3167	-0.0165	0.0544	<b>0.8841</b>	<b>0.9126</b>	<b>0.9115</b>	<b>0.9064</b>	<b>0.9219</b>	<b>0.9225</b>	<b>0.8575</b>	1

Significant ( $P < 0.05$ ) Spearman's correlation coefficients are shown in bold.

LC = long-chain ( $> C_{25}$ ), SC = short-chain ( $< C_{21}$ ), LMW = low molecular weight (2–3 aromatic rings), HMW = high molecular weight (4–6 aromatic rings).



oil may be related to accidental spills of fossil fuel during transportation to the generators and storage in the old tanks. The old storage tanks were in poor condition and were replaced. In site C8 located next to the boat storeroom the presence of petroleum residues may be related to boat traffic. In late February 2015, fuel discharge was made directly from the Uruguayan Navy vessel ROU *Vanguardia* to the new fuel storage tanks through a pipe for the first time. Since then, the transport of fuel from the new tanks to the generators has also been through a closed pump system. Furthermore, the old fuel tanks were dismantled to be returned to Uruguay. It is expected that these changes in fuel handling will reduce the risk of leaks and their impact on the Antarctic environment near the BCAA. The occurrence of low concentrations of the UCM at site C7 may be related to occasional wastewater leaks from the septic tank of the kitchen/dining room that drains to the kitchen's stream. The wastewater contains human effluents and associated nutrients (nitrogen, phosphorous), microorganisms (including potential pathogens), organic matter, detergents, heavy metals, disinfection by-products and hydrocarbons (Smith & Riddle 2009). Wastewater contamination has been recently reported at the BCAA (Tort *et al.* 2017).

As expected, site T0 located *c.* 100 m from pollution sources showed the lowest AHs level. AHs levels at sites T11, T13 and C8 are higher than levels recorded in surface marine sediments near other Antarctic research stations with clear evidence of oil pollution (Martins *et al.* 2004, Dauner *et al.* 2015). In contrast, AH concentrations in the other sites are within the range reported for unpolluted ( $< 100 \mu\text{g g}^{-1}$ ) soil and/or marine sediments near other Antarctic research stations (e.g. Dauner *et al.* 2015). Pristane and phytane are dominant in highly weathered crude oil but they may also have biogenic sources (Colombo *et al.* 1989). They derive from the phytol side chain of chlorophyll under reducing (phytane) or oxidizing (pristane) diagenetic conditions. Pristane to phytane ratios (Pri/Phy) of about one have been related to petroleum sources (Commendatore *et al.* 2012). In addition, Pri/Phy ratios considerably lower than one have been linked to the presence of petroleum and/or reducing depositional environments (Moustafa & Morsi 2012). Pri/Phy ratios at the BCAA were close to or lower than one in all the sites measured, which may be related to the prevalence of reducing conditions in soil and sediments, besides petroleum inputs. The absence of pristane in several sites reinforced the likelihood of the occurrence of anoxic hydrocarbon biodegradation processes (Prus *et al.* 2015). Site T11, adjacent to the generator room, and site T13, next to the fuel tanks, presented Pri/Phy ratios close to one, confirming the occurrence of oil pollution and weathered or degraded crude oil in these sites (e.g. Colombo *et al.* 1989).

For most of the sites,  $\text{CPI}_1$  was  $< 1$  indicating even carbon number predominance in short-chain n-alkanes. Even carbon number predominance in short-chain n-alkanes in soil and sediments is related to bacteria, fungi and yeast species, but also to petroleum-derived inputs (Harji *et al.* 2008, Yu *et al.* 2016). In contrast,  $\text{CPI}_2$  was  $> 1$  for most of the sites indicates odd carbon number predominance in long-chain n-alkanes, which has been linked to terrestrial higher plant inputs (Yu *et al.* 2016). Therefore, in soil and sediments from meltwater stream sites near the BCAA that are not impacted by humans the biogenic AHs seem to be derived from microbial and terrestrial inputs. Maximum abundances of  $\text{C}_{27}$  and  $\text{C}_{29}$  have been linked to lichens, mosses and the Antarctic pearlwort (Cabrerizo *et al.* 2016). High abundances of the lichen *Usnea aurantiaco-atra* (Jacq.) Bory and other species have been previously reported in areas around the BCAA (Piñeiro *et al.* 2012). Anthropogenic sources of short-chain n-alkanes may be related to light oils such as diesel (Green & Nichols 1995). In addition, CPI values close to one, as was recorded at site T13 located next to the fuel tanks, are indicative of anthropogenic sources from raw hydrocarbons and combustion of fossil fuels, and have been reported near other Antarctic stations (e.g. Cabrerizo *et al.* 2016). Therefore, anthropogenic inputs of AHs at human-impacted sites near the BCAA seem to derive from petroleum contributions; however, wastewater inputs are another potential source.

#### *Levels and sources of polycyclic aromatic hydrocarbons in soil and meltwater stream sediments near Artigas Station*

Based on the classification proposed by Baumard *et al.* (1998) sites T0, T2, T3, T4 and T5 and most of the sites around the meltwater streams were categorized with low levels of PAH contamination ( $0\text{--}100 \text{ ng g}^{-1}$ ). In addition, sites T1, T6, T7, T8, T9, T14, C0, C3, C4 and C8 were categorized with moderate PAH contamination ( $100\text{--}1000 \text{ ng g}^{-1}$ ). Site T10, T11 and T13 were categorized with high ( $1000\text{--}5000 \text{ ng g}^{-1}$ ) or very high ( $> 5000 \text{ ng g}^{-1}$ ) levels of PAH contamination. High concentrations of both total PAHs and the 16 priority PAHs at sites C8, T10, T11 and T13 confirmed this distributional pattern of PAHs. Major sources of anthropogenic PAHs in environmental samples are combustion processes and the release of uncombusted products (e.g. Yunker *et al.* 2002). Based on isomer ratios, PAHs in sites C8, T10, T11 and T13 most probably originate from high-temperature biomass and petroleum combustion, and from petroleum-derived PAHs (Yunker *et al.* 2002, Xue *et al.* 2016). These findings may be linked with the proximity of sites T10, T11 and T13 to the incinerator, the generator room and the fuel tanks, and with boat traffic at site C8 located next to the boat storeroom. The HMW PAHs (four to six aromatic rings) are generated by diesel vehicles, power


**Table V.** Total aliphatic hydrocarbon (AH), n-alkane and polycyclic aromatic hydrocarbon (PAH) concentrations reported near other Antarctic research stations compared with levels reported in the present study for Artigas Station (adapted from Dauner *et al.* 2015).

Research station		Concentration	Site	Reference
Signy (UK)	n-alkanes	0.04–1.73 $\mu\text{g g}^{-1}$		Cripps (1992)
	PAHs	14–280 $\text{ng g}^{-1}$	m	Cripps (1992)
Davis (AUS)	AHs	0.25–9.0 $\mu\text{g g}^{-1}$		Green & Nichols (1995)
	PAHs	7–7270 $\text{ng g}^{-1}$		Green <i>et al.</i> (1992)
Syowa (JAP)	n-alkanes	0.16–12.2 $\mu\text{g g}^{-1}$		Harada <i>et al.</i> (1995)
Arthur Harbor	PAHs	21–1572 $\text{ng g}^{-1}$	m, i	Kennicutt <i>et al.</i> (1992)
Comandante Ferraz (BRA)	AHs	0.15–13.3 $\mu\text{g g}^{-1}$		Martins <i>et al.</i> (2004)
	n-alkanes	0.10–5.50 $\mu\text{g g}^{-1}$		Bicego <i>et al.</i> (2009)
	PAHs	9.45–270.5 $\text{ng g}^{-1}$	m	Martins <i>et al.</i> (2004)
	PAHs	0.71–234 $\text{ng g}^{-1}$		Bicego <i>et al.</i> (2009)
	PAHs	< 6.0–454.9 $\text{ng g}^{-1}$		Martins <i>et al.</i> (2010)
Presidente Frei (CHL), Bellingshausen (RUS), Artigas (URY), Great Wall (CHN)	n-alkanes	0.01–2.24 $\mu\text{g g}^{-1}$	m	Préndez <i>et al.</i> (2011)
Carlini (ARG)	PAHs	36–1908 $\text{ng g}^{-1}$		Curtosi <i>et al.</i> (2007)
	PAHs	36.5–1908.4 $\text{ng g}^{-1}$		Curtosi <i>et al.</i> (2009)
	PAHs	12.05–210.02 $\text{ng g}^{-1}$	m	Dauner <i>et al.</i> (2015)
	n-alkanes	0.03–0.41 $\mu\text{g g}^{-1}$		Dauner <i>et al.</i> (2015)
	AHs	1.10–3.23 $\mu\text{g g}^{-1}$		Dauner <i>et al.</i> (2015)
Nelson Island	n-alkanes	0.01 $\mu\text{g g}^{-1}$		Dauner <i>et al.</i> (2015)
	AHs	0.52–0.61 $\mu\text{g g}^{-1}$		Dauner <i>et al.</i> (2015)
	PAHs	15.32–33.79 $\text{ng g}^{-1}$	m	Dauner <i>et al.</i> (2015)
Ardley Island	n-alkanes	0.15–0.21 $\mu\text{g g}^{-1}$		Dauner <i>et al.</i> (2015)
	AHs	1.94–4.94 $\mu\text{g g}^{-1}$		Dauner <i>et al.</i> (2015)
	PAHs	7.33–13.64 $\text{ng g}^{-1}$	m	Dauner <i>et al.</i> (2015)
Old Palmer (USA)	PAHs	29–59 478 $\text{ng g}^{-1}$		Kennicutt <i>et al.</i> (1992)
McMurdo (USA)	PAHs	5.3–88 452 $\text{ng g}^{-1}$		Mazzeri <i>et al.</i> (1999)
Mendel (CZ)	PAHs	1.4–205 $\text{ng g}^{-1}$	a	Klánová <i>et al.</i> (2008)
Great Wall (CHN)	PAHs	0.54–228.2 $\text{ng g}^{-1}$	m	Préndez <i>et al.</i> (2011)
Scott Base (NZ)	PAHs	< 30–81 052 $\text{ng g}^{-1}$		Aislabie <i>et al.</i> (1999)
Prydz Bay	PAHs	12.95–30.93 $\text{ng g}^{-1}$	m	Xue <i>et al.</i> (2016)
Artigas (UY)	AHs	0.57–2332 $\mu\text{g g}^{-1}$		This work
	n-alkanes	0.39–351 $\mu\text{g g}^{-1}$		This work
	PAHs	1.36–51 650 $\text{ng g}^{-1}$	s	This work
	PAHs	21.5–751 $\text{ng g}^{-1}$	b	This work

a = before the station was installed, b = meltwater stream sediment, i = intertidal sediment, m = surface marine sediment, s = surface soil.

plant and open-field burning of organic wastes, and are mainly associated with soot and black carbon (Cornelissen *et al.* 2005). High concentrations of HMW PAHs at sites T10, T11, T13 and C8 indicate that at the BCAA pyrogenic sources are restricted. However, the predominance of LMW PAHs and alkylated homologues is generally attributed to the use of a Special

Antarctic Blend diesel as fuel (Kennicutt *et al.* 1992, Green & Nichols 1995) and has been reported in other studies (e.g. Curtosi *et al.* 2007, Dauner *et al.* 2015, Xue *et al.* 2016). The dominance of LMW PAHs has been also linked to natural Antarctic vegetation, such as lichens and mosses (Cabrerizo *et al.* 2012), and to local sources related to low temperature combustion and diesel



**Fig. 7.** Summary of hydrocarbon contamination distribution at the Artigas Station. No or low contamination = green circles, moderate contamination = orange circles, high contamination = red circles.

transport (Xue *et al.* 2016). Furthermore, high levels of alkylated PAHs are derived mainly from petrogenic sources (Cabrerizo *et al.* 2012). Therefore, the dominance of LMW PAHs and alkylated compounds in almost all of the sites near the BCAA seem to be related mainly to petrogenic sources, such as diesel spills, at the contaminated sites or to natural Antarctic vegetation in the sites not impacted by humans.

#### *Comparison between hydrocarbon levels at Artigas Station with other Antarctic research stations*

Total AH and n-alkane concentrations in soil and meltwater stream sediments at the BCAA were considerably higher than the levels reported for pristine or nearly pristine areas in Antarctica (e.g. Nelson Island and Potter Cove) (Table V). AH and n-alkane levels reported here were higher than those previously stated for surface marine sediments near the BCAA and in the vicinity of other research stations including the Presidente Frei, Bellingshausen and Great Wall stations (Préndez *et al.* 2011) (Table V). The levels recorded at sites T11, T13 and C8 were higher than those found in human impacted areas such as the Ardley Island, Great Wall and Davis stations (Green & Nichols, 1995, Préndez *et al.* 2011, Dauner *et al.* 2015) (Table V). Total PAH concentrations recorded in soil samples near the BCAA

were comparable to those reported near other research stations in Antarctica (Table V). The highest concentrations found at the sites T10, T11 and T13 were equivalent to levels reported in surface sediments of highly impacted areas near the old Palmer and McMurdo stations (Kennicutt *et al.* 1992, Mazzer *et al.* 1999) (Table V). In addition, total PAH concentrations recorded in surface sediments from the meltwater streams (except site C8) were similar to those reported in marine surface sediments in areas of low human impact close to other research stations (e.g. Carlini Station in Potter Cove) (Dauner *et al.* 2015) (Table V).

#### **Conclusions**

This work has established the occurrence of local petroleum contamination at the BCAA with the highest levels restricted to sites T10, T11, T13 and C8 (Fig. 7). These sites are exposed to hydrocarbon inputs mainly derived from petrogenic sources but also with certain pyrogenic inputs. Based on the levels of AHs and PAHs detected, sites in the service area, where the incinerator, the generator room and the fuel tanks are located, and next to the boat storeroom are heavily impacted, while the remaining sites showed moderate to low levels or no contamination. Anthropogenic hydrocarbon sources at the BCAA predominate over biogenic sources and are related

to fossil fuel and organic residue combustion, fuel storage and boat traffic. Natural sources seemed to derive from microbial and terrestrial inputs.

Petrogenic contamination in the area close to the BCAA is relatively high as found in impacted areas near other Antarctic stations, but is restricted to the boat storeroom and the service area. It is expected that improvements made in fuel management and the removal of the old fuel tanks will reduce the risk of diesel leaks and their impact on the Antarctic environment near the BCAA. Future monitoring of the degradation process of these contaminants under extreme climatic conditions is recommended, as is investigation of the possibility of applying bioremediation measures. This work provides reference information for future environmental monitoring.

### Acknowledgements

We would like to thank our colleagues from the LabQOM for assistance with laboratory analysis. The Instituto Antártico Uruguayo (IAU) is fully acknowledged for providing support during fieldwork and research execution (Project IAU-P-DCC-15). The Laboratorio Tecnológico del Uruguay (LATU) provided financial support for hydrocarbon analysis (LATU-5000005779). PEDECIBA Geociencias and SNI-ANII supported C. Rodríguez and N. Venturini, respectively. We are also grateful to the Artigas Station crew 'Antarkos XXXI' for their support during sampling surveys. Many thanks to Magdalena Rodríguez who did the sketch of the Artigas Station. We are very grateful for the comments and suggestions of two anonymous reviewers and Professor David W.H. Walton, which helped us to improve an earlier version of this manuscript.

### Author contributions

EB and NV developed and implemented the survey and together with KI, NK and CB participated in the collection of field samples. MCB, ST and STS conducted the hydrocarbon analysis. CB performed the grain size analysis. CR and NV analysed the data and wrote the article. All authors collectively collaborated in the discussion of results, and the editing and organization of the manuscript.

### Data deposit

The depository of hydrocarbon data for this article is the Laboratorio de Biogeoquímica Marina (LABIM), Instituto de Ecología y Ciencias Ambientales, Facultad de Ciencias, Universidad de la República. All data will be available from the corresponding author upon request.

### Supplemental material

Example chromatograms will be found at <https://doi.org/10.1017/S0954102018000019>.

### References

- ASLABIE, J.M., BALKS, M.R., ASTORI, N., STEVENSON, G. & SYMONS, R. 1999. Polycyclic aromatic hydrocarbons in fuel-oil contaminated soils, Antarctica. *Chemosphere*, **39**, 2201–2207.
- ASLABIE, J.M., BALKS, M.R., FOGHT, J.M. & WATERHOUSE, E.J. 2004. Hydrocarbon spills on Antarctic soils: effects and management. *Environmental Science & Technology*, **38**, 1265–1274.
- ATCM (ANTARCTIC TREATY CONSULTATIVE MEETING). 2013. *Antarctic Treaty inspection programme report 2012. Report of Antarctic Treaty inspections undertaken jointly by the United Kingdom, the Netherlands and Spain in accordance with Article VII of the Antarctic Treaty and Article 14 of the Environmental Protocol*. Available at: [https://www.ats.aq/documents/ATCM36/att/ATCM36\\_att108\\_e.pdf](https://www.ats.aq/documents/ATCM36/att/ATCM36_att108_e.pdf).
- AYRES, M., AYRES JR, M., AYRES, D.L. & SANTOS, A.S. 2005. *BioEstat 4.0: aplicações estatísticas nas áreas das ciências biológicas e médicas*. Belém: Sociedade Civil Mamirauá, CNPq, 324 pp.
- BAUMARD, P., BUDZINSKI, H., MICHON, Q., GARRIGUES, P., BURGEOT, T. & BELLOCQ, J. 1998. Origin and bioavailability of PAHs in the Mediterranean Sea from mussel and sediment records. *Estuarine Coastal and Shelf Science*, **47**, 77–90.
- BÍEGO, M.C., ZANARDI-LAMARDO, E., TANIGUCHI, S., MARTINS, C.C., DA SILVA, D.A.M., SASAKI, S.T., ALBERGARIA-BARBOSA, A.C.R., PAOL, F.S., WEBER, R.R. & MONTONE, R.C. 2009. Results from a 15-year study on hydrocarbon concentrations in water and sediment from Admiralty Bay, King George Island, Antarctica. *Antarctic Science*, **21**, 209–220.
- CABRERIZO, A., DACHS, J., BARCELÓ, D. & JONES, K.C. 2012. Influence of organic matter content and human activities on the occurrence of organic pollutants in Antarctic soils, lichens, grass, and mosses. *Environmental Science & Technology*, **46**, 1396–1405.
- CABRERIZO, A., TEJEDO, P., DACHS, J. & BENAYAS, J. 2016. Anthropogenic and biogenic hydrocarbons in soils and vegetation from the South Shetland Islands (Antarctica). *Science of the Total Environment*, **569**, 1500–1509.
- COLOMBO, J.C., PELLETIER, E., BROCHU, C. & KHALIL, M. 1989. Determination of hydrocarbon sources using n-alkane and polyaromatic hydrocarbon distribution indexes. Case study: Rio de la Plata Estuary, Argentina. *Environmental Science & Technology*, **23**, 888–894.
- CORNELISSEN, G., GUSTAFSSON, O., BUCHELI, T.D., JONKER, M.T.O., KOELMANS, A.A. & VAN NOORT, P.C.M. 2005. Extensive sorption of organic compounds to black carbon, coal, kerogen in sediments and soils: mechanisms and consequences for distribution, bioaccumulation, and biodegradation. *Environmental Science & Technology*, **39**, 6881–6895.
- COMMENDATORE, M.G., NIEVAS, M.L., AMIN, O. & ESTEVES, J.L. 2012. Sources and distribution of aliphatic and polyaromatic hydrocarbons in coastal sediments from Ushuaia Bay (Tierra del Fuego, Patagonia, Argentina). *Marine Environmental Research*, **74**, 20–31.
- CRIPPS, G.C. 1992. The extent of hydrocarbon contamination in the marine environment from a research station in the Antarctic. *Marine Pollution Bulletin*, **25**, 288–292.
- CURTOSI, A., PELLETIER, E., VODOPIVEZ, C.L. & MACCORMACK, W.P. 2007. Polycyclic aromatic hydrocarbons in soil and surface marine sediment near Jubany Station (Antarctica). Role of permafrost as a low-permeability barrier. *Science of the Total Environment*, **383**, 193–204.
- CURTOSI, A., PELLETIER, E., VODOPIVEZ, C.L. & MACCORMACK, W.P. 2009. Distribution of PAHs in the water column, sediments and biota of Potter Cove, South Shetland Islands, Antarctica. *Antarctic Science*, **21**, 329–339.
- DAUNER, A.L.L., HERNÁNDEZ, E.A., MACCORMACK, W.P. & MARTINS, C.C. 2015. Molecular characterisation of anthropogenic sources of sedimentary organic matter from Potter Cove, King George Island, Antarctica. *Science of the Total Environment*, **502**, 408–416.
- GREEN, G. & NICHOLS, P.D. 1995. Hydrocarbons and sterols in marine sediments and soils at Davis Station, Antarctica: a survey for human-derived contaminants. *Antarctic Science*, **7**, 137–144.



- GREEN, G., SKERRATT, J.H., LEEMING, R. & NICHOLS, P.D. 1992. Hydrocarbon and coprostanol levels in seawater, sea-ice algae and sediments near Davis-Station in Eastern Antarctica - a regional survey and preliminary-results for a field fuel spill experiment. *Marine Pollution Bulletin*, **25**, 293–302.
- HARADA, N., HANDA, N., FUKUCHI, M. & ISHIWATARI, R. 1995. Source of hydrocarbons in marine-sediments in Lutzow-Holm Bay, Antarctica. *Organic Geochemistry*, 10.1016/0146-6380(94)00124-J.
- HARJI, R.R., YVENAT, A. & BHOSLE, N.B. 2008. Sources of hydrocarbons in sediments of the Mandovi estuary and the Marmugoa harbour, west coast of India. *Environment International*, **34**, 959–965.
- KENNICUTT, M.C., McDONALD, T.J., DENOUX, G.J. & McDONALD, S.J. 1992. Hydrocarbon contamination on the Antarctic Peninsula. 1. Arthur Harbor – subtidal sediments. *Marine Pollution Bulletin*, **24**, 499–506.
- KLÁNOVÁ, J., MATYKIEWICZOVÁ, N., MÁČKA, Z., PROSEK, P., LÁSKA, K. & KLÁN, P. 2008. Persistent organic pollutants in soils and sediments from James Ross Island, Antarctica. *Environmental Pollution*, **152**, 416–423.
- MARTINS, C.C., BÍCEGO, M.C., TANIGUCHI, S. & MONTONE, R.C. 2004. Aliphatic and polycyclic aromatic hydrocarbons in surface sediments in Admiralty Bay, King George Island, Antarctica. *Antarctic Science*, **16**, 117–122.
- MARTINS, C.C., BÍCEGO, M.C., ROSE, N.L., TANIGUCHI, S., LOURENÇO, R.A., FIGUEIRA, R.C.L., MAHIQUES, M.M. & MONTONE, R.C. 2010. Historical record of polycyclic aromatic hydrocarbons (PAHs) and spheroidal carbonaceous particles (SCPs) in marine sediment cores from Admiralty Bay, King George Island, Antarctica. *Environmental Pollution*, **158**, 192–200.
- MAZZERA, D., HAYES, T., LOWENTHAL, D. & ZIELINSKA, B. 1999. Quantification of polycyclic aromatic hydrocarbons in soil at McMurdo Station, Antarctica. *Science of the Total Environment*, **229**, 65–71.
- MOUSTAFA, Y.M. & MORSI, R.E. 2012. Biomarkers. In DHANARASU, S., ed. *Chromatography and its applications*. London: InTech, 210 pp.
- PIÑEIRO, V., EGUREN, G., PEREIRA, I. & ZALDÚA, N. 2012. Líquenes del entorno de la Base Científica Antártica Artigas, B. Collins, Isla Rey Jorge, Antártida. *Polibotánica*, **33**, 105–116.
- PRÉNDEZ, M., BARRA, C., TOLEDO, C. & RICHTER, P. 2011. Alkanes and polycyclic aromatic hydrocarbons in marine surficial sediment near Antarctic stations at Fildes Peninsula, King George Island. *Antarctic Science*, **23**, 578–588.
- PRUS, W., FABIANSKA, M.J. & LABNO, R. 2015. Geochemical markers of soil anthropogenic contaminants in polar scientific stations nearby (Antarctica, King George Island). *Science of the Total Environment*, **518**, 266–279.
- SMITH, J.J. & RIDDLE, M.J. 2009. Sewage disposal and wildlife health in Antarctica. In KERRY, K.R. & RIDDLE, M., eds. *Health of Antarctic wildlife*. Berlin: Springer, 271–315.
- SUGUIO, K. 1973. *Introdução à Sedimentologia*. São Paulo: Edgard Blucher, 317 pp.
- TIN, T., FLEMING, Z.L., HUGHES, K.A., AINLEY, D.G., CONVEY, P., MORENO, C.A., PFEIFFER, S., SCOTT, J. & SNAPE, I. 2009. Impacts of local human activities on the Antarctic environment. *Antarctic Science*, **21**, 3–33.
- TORT, L.F.L., IGLESIAS, K., BUENO, C., LIZASOAIN, A., SALVO, M., CRISTINA, J., KANDRATAVICIUS, N., PÉREZ, L., FIGUEIRA, R., BÍCEGO, M.C., TANIGUCHI, S., VENTURINI, N., BRUGNOLI, E., COLINA, R. & VICTORIA, M. 2017. Wastewater contamination in Antarctic melt-water streams evidenced by virological and organic molecular markers. *Science of the Total Environment*, **609**, 225–231.
- UNEP. 1992. *Determinations of petroleum hydrocarbons in sediments. Reference methods for marine pollution studies 20*. Nairobi: United Nations Environment Programme, 75 pp.
- USEPA. 1985. *Evaluation and estimation of potential carcinogenic risks of polynuclear aromatic hydrocarbons: carcinogen assessment group*. Washington, DC: Office of Health and Environmental Assessment, United States Environmental Protection Agency.
- VOLKMAN, J.K., HOLDSWORTH, D.G., NEILL, G.P. & BYOR JR, H.J. 1992. Identification of natural, anthropogenic and petroleum hydrocarbons in aquatic sediments. *Science of the Total Environment*, **112**, 203–219.
- XUE, R., CHEN, L., LU, Z.B., WANG, J., YANG, H.Z., ZHANG, J. & CAI, M.H. 2016. Spatial distribution and source apportionment of PAHs in marine surface sediments of Prydz Bay, East Antarctica. *Environmental Pollution*, **219**, 528–536.
- YU, Y.L., LI, Y.Y., GUO, Z.G. & ZOU, H. 2016. Distribution and sources of n-alkanes in surface sediments of Taihu Lake, China. *Archives of Environmental Protection*, **42**, 49–55.
- YUNKER, M.B., MACDONALD, R.W., VINGARZAN, R., MITCHELL, R.H., GOYETTE, D. & SYLVESTRE, S. 2002. PAHs in the Fraser River basin: a critical appraisal of PAH ratios as indicators of PAH source and composition. *Organic Geochemistry*, **33**, 489–515.