

Nutritional status and risk for disability in instrumental activities of daily living in older Brazilians

Ligiana Pires Corona^{1,*}, Tábatta Renata Pereira de Brito², Daniella Pires Nunes², Tiago da Silva Alexandre¹, Jair Lício Ferreira Santos³, Yeda Aparecida de Oliveira Duarte² and Maria Lúcia Lebrão¹

¹Department of Epidemiology, Faculty of Public Health, University of São Paulo, Avenida Doutor Arnaldo 715, 1 andar, CEP 01246-904, São Paulo, SP, Brazil; ²Department of Medical Surgical Nursing, School of Nursing, University of São Paulo, São Paulo, SP, Brazil; ³Department of Social Medicine, Faculty of Medicine of Ribeirão Preto, University of São Paulo, Ribeirão Preto, SP, Brazil

Submitted 19 July 2012: Final revision received 4 November 2012: Accepted 12 November 2012: First published online 13 December 2012

Abstract

Objective: The aim of the present study was to examine the association between nutritional status and the incidence of disability regarding instrumental activities of daily living (IADL) among older adults.

Design: The study is part of the longitudinal SABE (Saúde, Bem-Estar e Envelhecimento; Health, Wellbeing and Ageing) Study that began in 2000 (first wave) with a multistage, clustered, probabilistic sample (n 2143) of older adults (≥ 60 years). The second wave was carried out in 2006, when 1115 elders were re-interviewed. The dependent variable was the occurrence of disability in 2006 (report of difficulty on ≥ 1 IADL). Nutritional status (measured at baseline) was classified on the basis of BMI: ≤ 23.0 kg/m² (underweight); > 23.0 and < 28.0 kg/m² (ideal range – reference); ≥ 28.0 and < 30.0 kg/m² (overweight); and ≥ 30.0 kg/m² (obesity).

Setting: São Paulo, Brazil.

Subjects: One thousand and thirty-four individuals without difficulties regarding IADL in 2000 were selected, 611 of whom were re-interviewed in 2006.

Results: In the multiple logistic regression analysis adjusted for baseline variables (gender, age, number of chronic diseases, stroke, osteoarthritis and cognitive status), underweight (OR = 2.03; $P = 0.034$) and obesity (OR = 1.79; $P = 0.022$) remained associated with disability.

Conclusions: Both underweight and obesity are associated with an increased risk of developing disability regarding IADL among older adults, in an independent fashion of other risk factors. Thus, adequate nutritional status is a key point to consider in the establishment of preventive measures.

Keywords
Elderly
Nutritional status
Disability
Instrumental activities of daily living
SABE Study

Disability is defined as difficulty performing activities in an independent fashion due to physical and/or cognitive impairment⁽¹⁾. Disability generally affects older individuals due to the accumulation of chronic diseases; it has a negative effect on quality of life, is a predictor of hospitalization, institutionalization and death, and increases the burden placed on social and health-care systems^(2,3).

Instrumental activities of daily living (IADL) have been widely studied in recent years, as such activities help maintain individuals in the social and cultural realms of life. The loss of function in IADL can lead to a loss of autonomy (especially with regard to household management), confinement and social isolation. Studies have

shown that functional impairment occurs in a hierarchical manner, beginning with IADL and culminating in impaired performance of basic activities of daily living which are related to personal care⁽¹⁾.

Changes in one's nutritional status contribute toward an increase in illness and mortality rates. Excess weight and obesity have been associated with a number of co-morbidities in all phases of life, especially chronic, non-transmittable conditions^(4,5). In recent years, studies have demonstrated that obesity is associated with limitations regarding the performance of IADL, independently of the presence of disease^(6,7). Underweight has also been described as a risk factor for disability due to its strong association with sarcopenia^(8,9).

*Corresponding author: Email lillypires@gmail.com; licorona@usp.br

However, the literature reports divergent findings regarding the association between nutritional status and disability. In some studies, underweight and obesity were not found to be independent risk factors of disability^(10,11). Moreover, this association has not yet been explored in developing countries, in which the process of epidemiological and nutritional transition is ongoing and the coexistence of increasing obesity and a still high proportion of underweight has a considerable impact on health-care services.

The aim of the present study was to examine the association between nutritional status and the incidence of disability regarding the performance of IADL among older adults in the city of São Paulo (Brazil) over a 6-year period.

Methods

Sample and procedures

The data originated from the SABE (Saúde, Bem-Estar e Envelhecimento; Health, Wellbeing and Ageing) Study, which is a longitudinal study that began in 2000 involving a probabilistic sample of older adults (aged ≥ 60 years) residing in the city of São Paulo (n 2143). The baseline sample was obtained with a two-stage stratified sampling method, following the framework of the 1995 National Household Survey based on geographic areas of the city. Individuals aged ≥ 75 years were oversampled to compensate for the greater mortality rate in this age group⁽³⁾. A second wave of the study was carried out in 2006, in which 1115 of the participants from the first wave were interviewed again. Details on the methodology of the study are described elsewhere^(3,12).

The data in the first wave of the study were collected in two stages. The first stage was a household interview conducted by a single interviewer using a standardized questionnaire addressing the living conditions and health status of the older adult. The second stage was a household visit by a pair of interviewers for the measurement of anthropometric data and the determination of physical performance⁽³⁾.

Among the 2143 individuals initially interviewed, 1034 reported no difficulties in performing IADL. However, losses to follow-up occurred (114 individuals had moved to other cities or were not located, 190 had died, four had been institutionalized and 115 individuals refused to participate in the second wave). Thus, 611 individuals were eligible to be followed up.

The study was conducted in compliance with the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects/patients were approved by a human research ethics committee. The first wave of the SABE Study was approved by the National Research Ethics (CONEP), and the second wave was approved by the Research Ethics Committee (COEP) of the Faculty of Public Health, University of São Paulo. Written informed consent was obtained from all participants.

Measures

Disability was the dependent variable and was recorded when the participant reported difficulty on one or more IADL in 2006 for which no difficulty was reported in 2000 (baseline). The following activities were analysed: managing finances, use of transportation within the community, shopping, ability to use the telephone and responsibility for one's own medications. All of the other independent variables and parameters included in the analysis were baseline measures.

BMI was calculated by dividing body mass (in kilograms) by the square of height (in metres; kg/m^2). Nutritional status was classified based on BMI cut-off points adopted by the Pan American Health Organization for the SABE Study⁽¹³⁾: $\leq 23.0 \text{ kg}/\text{m}^2$ = underweight; > 23.0 and $< 28.0 \text{ kg}/\text{m}^2$ = normal range (reference); ≥ 28.0 and $< 30.0 \text{ kg}/\text{m}^2$ = overweight; and $\geq 30.0 \text{ kg}/\text{m}^2$ = obesity.

The following sociodemographic characteristics were included in the analysis: gender; age (grouped into two categories: 60–74 years and ≥ 75 years); household arrangement (living alone or accompanied); income perceived as adequate for expenses; and schooling (in years of formal education).

Health status was assessed based on self-reported diabetes, hypertension, CVD, osteoarthritis, chronic respiratory disease, stroke, osteoporosis, and reported falls and hospitalization in the 12 months preceding the interview.

Cognitive status was evaluated using the modified version of the Mini Mental State Examination (MMSE)⁽¹⁴⁾ validated for the SABE Study, due to the low level of schooling of the South American elderly population. This measure has thirteen items that are less dependent upon schooling and the cut-off point is a score of 12 or less⁽¹⁵⁾. The Brazilian version of the Geriatric Depression Scale was used to assess depressive symptoms, with a score of 6 or more as the cut-off point^(16,17).

Statistical analysis

For the descriptive analysis, means and their standard errors were calculated for the continuous variables and proportions were calculated for the categorical variables. Differences between groups were estimated using the Wald test of mean equality and the Rao–Scott test, which considers sample weights for estimates with population weights⁽¹⁸⁾.

Forward stepwise logistic regression analysis was performed for the assessment of factors associated with the development of disability in the period studied and the dependent variable was positive answer of difficulty in at least one IADL in 2006. Variables that remained significant ($P < 0.05$) or that adjusted the odds ratio by at least 10% were maintained in the model. The inferences considered the design effect. The data analysis was performed using the statistical software package Stata® version 11.

Results

Among the older adults who were independent with regard to IADL in 2000, 611 were interviewed again in 2006. In the comparative analysis between those who were re-interviewed and those lost to follow-up, significant differences were found regarding osteoarthritis and osteoporosis, with higher baseline prevalences in the group lost to follow-up: 35.0% *v.* 20.3% for osteoarthritis ($P < 0.001$) and 18.3% *v.* 12.9% for osteoporosis ($P = 0.031$). No statistically significant differences were found for the other variables analysed (data not shown).

Table 1 displays the baseline characteristics of the participants at the beginning of the study (total sample) and according to functional status at follow-up. At baseline, most of the individuals were between 60 and 74 years of age, had an average of 7.8 years of schooling and a score of 17 on the MMSE. Obesity was identified in ~20% and 73.1% reported at least one chronic disease.

Among the 611 individuals re-evaluated in 2006, 178 reported difficulty on at least one IADL. Those with incident disability differed from those who did not develop disability with regard to some baseline characteristics (Table 1). The incidence of difficulty regarding IADL was greater among women, individuals aged 75 years or older, those with a lower MMSE score, those with reported hypertension, history of stroke or joint disease and

those with a greater number of chronic illnesses. With regard to nutritional status, a greater proportion of either underweight or obesity at baseline was found among those individuals who developed difficulties regarding IADL in comparison to those who did not develop disability.

Table 2 displays the results of the logistic regression analysis for the incidence of disability regarding IADL. Both underweight and obesity were associated with risk of disability in the unadjusted analysis. After controlling for age, MMSE score, report of history of stroke or joint disease and number of reported chronic diseases, both conditions remained independent risk factors for developing functional disability.

Discussion

The results of the present cohort study demonstrate that both underweight and obesity are independent risk factors for functional disability. Underweight was associated with a 100% greater risk of disability in comparison to the ideal weight range (OR = 2.03; $P = 0.034$). This finding may be explained by the loss of muscle mass (sarcopenia), which is known to compromise functionality⁽⁹⁾. Underweight is a multifactor condition among elderly individuals, but has two main causes. The first is inadequate nutrient intake due to factors such as anorexia, oral

Table 1 Sample characteristics at baseline and after follow-up period according to outcome regarding IADL; SABE Study, city of São Paulo, Brazil, 2000 and 2006

Characteristic	Total sample in 2000 (<i>n</i> 1034)	Follow-up		<i>P</i> value*
		Independent in 2006 (<i>n</i> 433)	Dependent in 2006 (<i>n</i> 178)	
Gender (%)				0.024
Male	45.1	44.7	32.8	
Female	54.9	55.3	67.2	
Age (%)				<0.001
60–74 years	87.9	93.9	79.9	
≥75 years	12.1	6.1	20.1	
Years of education (mean)	7.8	9.2	4.1	0.096
Nutritional status (%)				0.027
Ideal weight	48.5	53.6	38.8	
Underweight	17.7	15.4	20.1	
Overweight	13.2	12.6	14.7	
Obesity	20.6	18.4	26.4	
Score on modified version of MMSE (mean)	17.0	17.2	16.3	0.006
Depressive symptoms (%)	12.8	14.0	13.5	0.875
Occurrence of falls in previous 4 months (%)	25.6	24.3	26.9	0.541
Hypertension (%)	49.7	46.0	56.7	0.030
Diabetes (%)	15.3	13.7	17.3	0.235
Chronic respiratory disease (%)	8.6	7.5	6.0	0.602
CVD (%)	14.5	11.9	13.5	0.589
Stroke (%)	3.6	2.3	7.9	0.007
Osteoarticular conditions (%)	25.4	17.3	29.8	0.001
Osteoporosis (%)	13.0	12.6	14.0	0.641
Number of chronic conditions (%)				0.001
No disease	26.9	32.9	23.9	
One disease	34.8	36.1	28.7	
Two or more diseases	38.3	31.0	47.4	

IADL, instrumental activities of daily living; SABE, Saúde, Bem-Estar e Envelhecimento (Health, Wellbeing and Ageing); MMSE, Mini Mental State Examination.

*Rao–Scott and Wald tests were used.

Table 2 Results of the logistic regression model for incidence of disability on IADL in a 6-year period among non-disabled participants at baseline; SABE Study, city of São Paulo, Brazil, 2000 and 2006

Variable	Unadjusted OR	P value	Adjusted OR*	P value
Nutritional status				
Ideal weight	1.00	—	1.00	—
Underweight	1.81	0.040	2.03	0.034
Overweight	1.61	0.122	1.47	0.250
Obesity	1.97	0.005	1.79	0.022
Female	1.66	0.025	1.58	0.074
Age ≥75 years	3.87	0.000	4.40	0.000
Number of chronic conditions				
No disease	1.00	—	1.00	—
One disease	1.09	0.747	0.97	0.917
Two or more diseases	2.11	0.001	1.64	0.112
Osteoarthritis	2.02	0.002	1.48	0.199
Stroke	3.69	0.011	5.41	0.002
MMSE score	0.90	0.025	0.92	0.076

IADL, instrumental activities of daily living; SABE, Saúde, Bem-Estar e Envelhecimento (Health, Wellbeing and Ageing); MMSE, Mini Mental State Examination, modified.

*Hosmer–Lemeshow goodness-of-fit test ($P = 0.468$).

and dental problems, dysphagia and altered sense of smell and taste, leading to weight loss and, if prolonged, malnutrition. Catabolic diseases or associated conditions, such as lung disease, cancer and frailty, constitute the other major cause of underweight^(19–21). Both mechanisms are associated with sarcopenia.

It was not possible to include the adequacy of food intake in the analyses of the present study, as data on food consumption were not addressed on the questionnaires. The same is true regarding frailty and sarcopenia. The other catabolic diseases were not associated with disability in the unadjusted analyses. Although this is suggestive of a relatively greater survival rate (individuals with catabolic diseases and weight loss are believed to have a greater mortality rate), the analysis did not reveal statistically significant differences in these diseases between those re-interviewed and those lost to follow-up or among the deaths having occurred in the sample (data not shown).

Previous studies have also reported underweight to be an independent risk factor for disability, such as a study carried out in the USA⁽¹⁰⁾ which found an association between underweight (BMI ≤ 18.5 kg/m²) and disability; the same association was also found in a study carried out in Canada⁽²²⁾. In a study conducted in France, in which a BMI between 23.0 and 27.0 kg/m² was considered to be the reference category, an effect was found only for elderly individuals with a BMI of <21.0 kg/m², whereas those with a BMI between 21.0 and 23.0 kg/m² were not at increased risk⁽¹¹⁾.

In the present study, the risk of disability was very high among individuals with obesity (79% higher in comparison to those within the ideal weight range, after controlling for other variables: OR = 1.79; $P = 0.022$). This association is less controversial in the literature. It seems to be well established that obesity is associated with a greater incidence of disability in different domains, including the execution of IADL, as reported in a review carried out by Zamboni *et al.*⁽⁵⁾. In other previously cited

studies, obesity was also found to be a risk factor for disability^(11,22). In the American study, however, class I obesity (BMI between 30.0 and 35.0 kg/m²) proved to be a protection factor against disability in men⁽¹⁰⁾. Unlike obesity, overweight appears to play a protective role against diseases and complications among elderly individuals⁽¹⁰⁾. Other studies have not found a significant association between overweight and disability⁽²²⁾. Only the study carried out in France found overweight to be a risk factor for disability regarding IADL – and only among men⁽¹¹⁾. In the present study, overweight was not significantly associated with the incidence of disability.

The analysis of nutritional status was controlled for previously known risk factors of disability. Among the variables that remained in the multivariate model, high odds ratios were found for individuals aged 75 years or older and those who reported a history of stroke – both of which are recognized risk factors for functional disability^(23,24). Osteoarthritis, number of chronic conditions and gender did not achieve statistical significance in the final model, but were maintained due to the fact that these variables adjust the estimates and are commonly reported risk factors in the literature^(24,25).

The findings presented herein contribute to the understanding of the association between nutritional status among elderly individuals and an impaired ability to live an independent life in the community, as measured by difficulties regarding the performance of IADL, especially in developing countries, which are going through a later but faster process of epidemiological and nutritional transition, with a considerable impact on health-care services. In Brazil, the accelerated growth in the prevalence of obesity and the reduction in malnutrition have been following the same patterns as those described for Americans⁽²⁶⁾. In this context, older adults merit special attention due to the increase in the prevalence of obesity, which can contribute towards an increase in illness and disability in this age group, along with the problem of

malnutrition at more advanced ages⁽²⁷⁾, as mentioned above. However, like other developing countries, Brazil does not have a sufficiently organized health-care structure or policies to cope with this rapid process of epidemiological and nutritional transition.

The present study has limitations that should be addressed. Disability was measured using self-reported information. While this may be a source of bias, methodological studies have demonstrated that self-reported data on functional disability have adequate validity and are consistent with medical diagnoses and/or physical tests⁽²⁸⁾. Another limitation is the high proportion of losses, but the analysis demonstrated no statistically significant differences in the baseline characteristics of the individuals interviewed a second time and those lost to follow-up, with the exception of the prevalence of osteoarthritis and osteoporosis. However, this difference tends to underestimate measures of the association of joint disease as a risk factor of disability, as a lower prevalence rate was found among the individuals evaluated a second time in comparison to those lost to follow-up.

Conclusions

Based on the findings of the present study, obesity and underweight are independent risk factors of impairments regarding an autonomous life in the community, as measured by the development of difficulties in the performance of IADL. These findings corroborate data found in the literature and help clarify the role of underweight in the incidence of disability. The understanding of the influence of nutritional status on this process allows the establishment of preventive strategies aimed at the elderly population, contributing to the reformulation of health-care policies and the organization of health services with the aim of diminishing the incidence of disability and favouring quality of life among this portion of the population.

Acknowledgements

Sources of funding: This study was funded by São Paulo Research Foundation (Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP). *Conflicts of interest:* The authors have no conflicts of interests. *Authors' contributions:* Study conception and design: L.P.C.; analysis and/or interpretation of data, drafting of the manuscript: L.P.C., D.P.N., T.R.P.d.B. and T.d.S.A.; revision of the manuscript and approval of the final version of the manuscript: J.L.F.S., Y.A.d.O.D. and M.L.L.; main study (SABE) coordinators: J.L.F.S., Y.A.d.O.D. and M.L.L.

References

- Verbrugge LM & Jette AM (1994) The disablement process. *Soc Sci Med* **38**, 1–14.
- Li LW (2005) Predictors of ADL disability trajectories among low-income frail elders in the community. *Res Aging* **27**, 615–642.
- Andrade FCD, Guevara PE, Lebrão ML *et al.* (2011) Gender differences in life expectancy and disability-free life expectancy among older adults in São Paulo, Brazil. *Womens Health Issues* **21**, 64–70.
- Bray GA (2004) Medical consequences of obesity. *J Clin Endocrinol Metab* **89**, 2583–2589.
- Zamboni M, Mazzali G, Zoico E *et al.* (2005) Health consequences of obesity in the elderly: a review of four unresolved questions. *Int J Obes (Lond)* **29**, 1011–1029.
- Jensen GL & Friedmann JM (2002) Obesity is associated with functional decline in community-dwelling rural older persons. *J Am Geriatr Soc* **50**, 918–923.
- Lang IA, Llewellyn DJ, Alexander K *et al.* (2008) Obesity, physical function, and mortality in older adults. *J Am Geriatr Soc* **56**, 1474–1478.
- Willett WC (1997) Weight loss in the elderly: cause or effect of poor health? *Am J Clin Nutr* **66**, 737–738.
- Lang T, Streeter T, Cawthon P *et al.* (2010) Sarcopenia: etiology, clinical consequences, intervention, and assessment. *Osteoporos Int* **21**, 543–559.
- Imai K, Gregg EW, Chen YJ *et al.* (2008) The association of BMI with functional status and self-rated health in US adults. *Obesity (Silver Spring)* **16**, 402–408.
- Larrieu S, Pérès K, Letenneur L *et al.* (2004) Relationship between body mass index and different domains of disability in older persons: the 3C study. *Int J Obes Relat Metab Disord* **28**, 1555–1560.
- Lebrão ML & Laurenti R (2005) Saúde, bem-estar e envelhecimento: o estudo SABE no Município de São Paulo. *Rev Bras Epidemiol* **8**, 127–141.
- Organización Panamericana de la Salud (2001) XXXVI Reunión del Comité Asesor de Investigaciones en Salud – Encuesta Multicéntrica – Salud Bienestar y Envejecimiento (SABE) en América Latina e el Caribe – Informe preliminar. Washington, DC: División de Promoción y Protección de la Salud, OPAS; available at <http://www.paho.org/Spanish/HDP/HDR/CAIS-01-05.PDF>
- Folstein MF, Folstein SE & McHugh PR (1975) 'Minimal state'. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* **12**, 189–198.
- Icaza MC & Albala C (1999) PROYECTO SABE: Mini Mental State Examination (MMSE) del estudio de demencia en Chile: análisis estadístico. Washington, DC: OPAS.
- Sheikh JI & Yesavage JA (1986) Geriatric Depression Scale (GDS): recent evidence and development of a short version. *Clin Gerontol* **5**, 165–173.
- Paradela EMP, Lourenço RA & Veras RP (2005) Validation of geriatric depression scale in a general outpatient clinic. *Rev Saude Publica* **39**, 918–923.
- Rao JNK & Scott AJ (1984) On chi-squared tests for multiway contingency-tables with cell proportions estimated from survey data. *Ann Stat* **12**, 46–60.
- Hickson M (2006) Malnutrition and ageing. *Postgrad Med J* **82**, 2–8.
- Wannamethee SG, Shaper AG, Whincup PH *et al.* (2000) Characteristics of older men who lose weight intentionally or unintentionally. *Am J Epidemiol* **151**, 667–675.
- Visvanathan R & Chapman IM (2009) Undernutrition and anorexia in the older person. *Gastroenterol Clin North Am* **38**, 393–409.
- Gadalla TM (2010) Relative body weight and disability in older adults: results from a national survey. *J Aging Health* **22**, 403–418.
- Kaplan G, Haan M & Wallace R (1999) Understanding changing risk factor associations with increasing age in adults. *Annu Rev Public Health* **20**, 89–108.

24. Fried LP, Bandeen-Roche K, Kasper JD *et al.* (1999) Association of comorbidity with disability in older women: the Women's Health and Aging Study. *J Clin Epidemiol* **52**, 27–37.
25. Stuck AE, Walthert JM, Nikolaus T *et al.* (1999) Risk factors for functional status decline in community-living elderly people: a systematic literature review. *Soc Sci Med* **48**, 445–469.
26. Popkin BM (1994) The nutrition transition in low-income countries: an emerging crisis. *Nutr Rev* **52**, 285–298.
27. Myers GC, Lamb VL & Agree EM (2003) Patterns of disability change associated with the epidemiologic transition. In *Determining Health Expectancies*, pp. 59–74 [J Robine, C Jagger, CD Mathers *et al.*, editors]. Chichester: Wiley.
28. Reuben DB, Siu AL & Kimpau S (1992) The predictive validity of self-reported and performance-based measures of function and health. *J Gerontol* **47**, M106–M110.