

Short Communication

Geographic measures of retail food outlets and perceived availability of healthy foods in neighbourhoods

Timothy L Barnes^{1,2}, Darcy A Freedman³, Bethany A Bell⁴, Natalie Colabianchi⁵ and Angela D Liese^{2,*}

¹Department of Epidemiology & Community Health, School of Public Health, University of Minnesota, Twin Cities, Minneapolis, MN, USA: ²Department of Epidemiology and Biostatistics and Center for Research in Nutrition and Health Disparities, Arnold School of Public Health, University of South Carolina, Discovery I 461, 915 Greene Street, Columbia, SC 29208, USA: ³Department of Epidemiology and Biostatistics, Case Western Reserve University, Cleveland, OH, USA: ⁴College of Education, University of South Carolina, Columbia, SC, USA: ⁵Institute for Social Research, University of Michigan, Ann Arbor, MI, USA

Submitted 10 April 2015: Final revision received 12 August 2015: Accepted 7 September 2015: First published online 2 October 2015

Abstract

Objective: To examine associations between geographic measures of retail food outlets and perceived availability of healthy foods.

Design: Cross-sectional.

Setting: A predominantly rural, eight-county region of South Carolina, USA.

Subjects: Data from 705 household shoppers were analysed using ordinary least-squares regression to examine relationships between geographic measures (presence and distance) of food outlets obtained via a geographic information system and perceived availability of healthy foods (fresh fruits and vegetables and low-fat foods).

Results: The presence of a supermarket within an 8·05 km (5-mile) buffer area was significantly associated with perceived availability of healthy foods ($\beta=1\cdot09$, $P=0\cdot025$) when controlling for all other food outlet types. However, no other derived geographic presence measures were significant predictors of perceived availability of healthy foods. Distances to the nearest supermarket ($\beta=-0\cdot16$, $P=0\cdot003$), dollar and variety store ($\beta=-0\cdot15$, $P=0\cdot005$) and fast-food restaurant ($\beta=0\cdot11$, $P=0\cdot015$) were all significantly associated with perceptions of healthy food availability.

Conclusions: Our results suggest that distance to food outlets is a significant predictor of healthy food perceptions, although presence is sensitive to boundary size. Our study contributes to the understanding and improvement of techniques that characterize individuals' food options in their community.

Keywords
Food environment
Food outlets
Geographic
Perceptions
Healthy foods

Over the past two decades, the number of studies linking diet to the food environment has increased^(1–4). It is widely believed that the environment can influence eating behaviour and weight status^(1,3,4). Researchers and policy makers have a shared interest in how the food environment contributes to health outcomes⁽⁴⁾.

Geographic information systems (GIS) have been a predominant tool used to characterize the food environment, typically describing the objective, spatial availability of or distance to retail food outlets^(2,5–7). Associations between the food environment and dietary outcomes have varied based on the measurement used in each study⁽³⁾.

Recently, perception measures of the food environment based on individuals' self-report have gained in popularity. Perception-based measures are advantageous, as they allow researchers to capture how people view and utilize their environment⁽³⁾. It has been suggested that perception measures may be more convenient and cheaper, because geographic measures are typically derived from secondary data sources and field validation is often not feasible and is too expensive^(8–10). However, the use of perception-based measures has drawbacks⁽¹¹⁾. Perception measures may be biased because of personal preferences or behaviours, how a person interprets survey question(s) (including

*Corresponding author: Email liese@sc.edu

definitions of food outlets or items) and understanding of the concept of 'neighbourhood'^(11,12). In addition, only a few questionnaires assessing the perceived food environment have been validated by researchers^(13–15).

A handful of studies have examined relationships between geographic measures and perceptions of the food environment^(16–25). However, only a few have focused specifically on the perceived availability of healthy foods, i.e. fresh fruits and vegetables and low-fat foods, compared with direct, objective measures of retail food outlet availability (e.g. presence, count or density) or distance^(17,19,20). Two studies have reported a significant positive association between supermarket availability, assessed by either the presence or density of stores, and perceived availability of healthy foods^(17,19). No study has addressed non-traditional food outlets (e.g. dollar stores or pharmacies) that offer a variety of food products and are attracting an increasing market share^(26,27). Further, many initiatives have been established to improve spatial access to healthier food retailers (e.g. larger grocery stores, food hubs and farmers' markets), such as the Healthy Food Financing Initiative⁽²⁸⁾; however, policy makers are not fully aware whether consumers with access to certain retail outlets are indeed aware of these opportunities and whether positive perceptions of them exist. Moreover, by examining associations between perceptions of healthy foods and food outlets other than supermarkets and supercentres, researchers and policy makers can determine whether current initiatives are sufficient or whether targeted policies and interventions that include non-traditional food outlets and better awareness (including changes) of the environment are warranted.

The objective of the present study was to expand the scope of research and examine the relationship of both GIS-derived presence and distance to various types of retail outlet with participants' perceptions of healthy food availability within their neighbourhoods. In our analyses, distance was defined using the closest food outlet type via street networks, and GIS-derived presence was defined as the existence of retail food outlets (yes or no) within 1.61 and 8.05 km (1- and 5-mile) buffers around each participant's home address. Perceived availability of healthy foods was characterized using a summary score that has been commonly used in the literature to describe a person's food environment^(13–15).

Methods

Study region

The study area consisted of a contiguous geographical area encompassing eight, predominantly rural (e.g. non-urban) counties in the Midlands region of South Carolina, USA and has been described previously^(8,9).

Design and recruitment

The study was approved by the University of South Carolina Institutional Review Board.

Using a cross-sectional design, a geographically based sample of 968 adults serving as the primary food shoppers for their households was recruited in the eight-county study region between April and June 2010 as part of a larger study^(15,25,29). The sample was generated from random selection of landline telephone numbers with listed addresses restricted to sixty-four eligible ZIP codes. To achieve good spatial coverage, research staff aimed to interview approximately fifteen respondents per ZIP code. Recruitment calls were performed by the University of South Carolina Survey Research Laboratory. Respondents were screened with respect to certain eligibility criteria, including being (i) at least 18 years old, (ii) the primary food shopper, (iii) capable of speaking English and (iv) living in the eight-county region. Of 2477 people screened, a total of 968 were eligible and completed the interview. This resulted in a response rate of 47%⁽³⁰⁾, which is comparable to other evaluations among landline households conducted in the USA⁽³¹⁾. For analyses, participants were removed if they were missing data on demographic characteristics (age, n 71; race/ethnicity, n 73; education, n 69; employment status, n 68; household income, n 215; Supplemental Nutrition Assistance Program (SNAP) status, n 69; spouse or partner status, n 64; number of household members, n 74), geographic measures (n 19), perceived availability of healthy foods (n 5) and urbanicity (n 18), resulting in a final sample within non-missing data of 705 participants (73% of completed interviews).

Perceived availability of healthy foods

Perceptions of the food environment were ascertained using previously validated questions^(13–15). Survey participants indicated their agreement on a 5-point Likert scale with the following statements: (i) 'A large selection of fruits and vegetables is available in my neighbourhood'; (ii) 'The fresh fruits and vegetables in my neighbourhood are of high quality'; and (iii) 'A large selection of low-fat products is available in my neighbourhood'. Neighbourhood was defined as a 1.61 km (1-mile) buffer or 20-min walk from home. Responses for each question ranged from 1 to 5 (1 = 'strongly agree', 2 = 'agree', 3 = 'neutral (neither agree nor disagree)', 4 = 'disagree' and 5 = 'strongly disagree'). For ease of interpretability, a summary perceptions score was created^(16,17,22). Responses were reverse coded, summed and then re-scaled to a starting point of zero to reflect a summary score in which 0 indicates the worst availability of healthy foods and 12 indicates the best availability.

Geographic measures of the food environment

All geospatial analyses were conducted using ArcGIS 10.0 (ESRI, Redlands, CA USA; 2010). Geographic measures were calculated using the geocoded participants' home addresses as the point of reference, with both a 1.61 and 8.05 km (1- and 5-mile) street and road network buffer

representing their neighbourhood boundaries. Buffers of 1.61 km (1 mile) and 8.05 km (5 miles) were used to reasonably reflect the range of distances participants may have to travel to the nearest retailer, especially in our predominantly non-urban study region. Furthermore, the nearest retailer ranged from 4.83 to 12.87 km (3 to 8 miles) from home, depending on store type. For urban participants, distance to the nearest retailer ranged from 1.61 to 3.22 km (1 to 2 miles), but for non-urban participants, these distances ranged from 4.83 to 14.48 km (3 to 9 miles). Addresses were then linked with an existing, previously validated geospatial database on food retail outlets^(8,9) and the presence of food outlets within the buffer (availability), defined as yes or no. Distance to the nearest retail food outlet was calculated using the shortest street distance based on StreetMap Premium for ArcGIS (ESRI; 2008). The outlet types included supermarkets, supercentres, warehouse clubs (of note, there is only one warehouse club within the study region), convenience stores, drug and pharmacy stores, dollar and variety stores, and franchised fast-food restaurants. Supermarkets, supercentres and warehouse clubs were aggregated and considered together as supermarkets. This classification has been used previously by the Centers for Disease Control and Prevention in the 2013 State Indicator Report on Fruits and Vegetables⁽³²⁾.

Participant characteristics

Demographic and socio-economic characteristics based on the Behavioral Risk Factor Surveillance Survey (BRFSS)⁽³³⁾ were collected, including age, race/ethnicity, annual household income, education, employment, SNAP use, number of household members and marital/partner status. Each respondent was also classified on level of urbanicity using the 2010 US Census urban classification method via a point-in-polygon operation within ArcGIS⁽³⁴⁾. Specifically, an urban area was defined as any census tract classified as either an 'urbanized area of 50 000 or more people or urban cluster consisting of at least 2500 and less than 50 000 people'. All census tracts not designated as urban were considered to be non-urban.

Statistical analyses

Ordinary least-squares regression was used to assess the relationship between the geographic presence of and distance to specific outlets and the perceived availability of healthy foods in the participants' neighbourhoods. Geographic presence and distance to the nearest food outlet were used as independent variables and perceived availability of healthy foods was the outcome variable for all models. Covariates included age, sex, race/ethnicity, education, employment, household income, SNAP utilization, marital/partner status, household size and urbanicity. No violations of the assumption of independent observations, linearity, homoscedasticity or normality of the residuals were noted and no evidence for

multicollinearity between the geographic measures was found (all variance inflation factors ≤ 10).

First we examined the independent influence of presence of or distance to a specific type of food retail outlet, without controlling for other outlet types, using separate models that adjusted for the listed demographic covariates only. Next we examined models in which presence of or distance to all food outlet types were considered simultaneously. The R^2 of each model was examined to determine how much each model explained the variance in the outcome. The squared semi-partial correlation type II, also known as the unique R^2 , was also used to examine the unique contribution of each specific exposure in explaining variation in the outcome.

Results

The majority of participants in the present study were female (77.7%), non-Hispanic white (65.5%) and lived in non-urban neighbourhoods (79.1%; Table 1). Ninety-four per cent of participants used a personal vehicle for primary shopping trips. Many participants did not have any retail food outlets near their home (i.e. within a 1.61 or 8.05 km (1- or 5-mile) buffer). For example, 88.6% of participants did not have a supermarket in their neighbourhood and 71.5% did not have a convenience store within 1.61 km (1 mile). The mean distance to the nearest supermarket from a participant's home was 9.50 km (5.9 miles) and the distances to other food outlets ranged from 4.67 to 12.55 km (2.9 to 7.8 miles). The mean perceived availability of healthy foods score was 6.2 on a scale of 0 to 12.

Table 2 displays the results for both the separate and full models examining the relationship between the presence of and distance to each type of food outlet and the perceived availability of healthy foods. In separate models, the presence of each type of food outlet was significantly and positively associated with perceived availability of healthy foods. Supermarkets were most strongly associated with perception at the 1.61 km (1-mile) buffer ($\beta = 1.93$, $P < 0.001$), followed by drug and pharmacy outlets ($\beta = 1.77$, $P < 0.001$). Moreover, all food outlet types were found to be significantly associated with perceived availability of healthy foods in separate models using either the 1.61 or 8.05 km (1- or 5-mile) buffer. However, the unique R^2 values were low, accounting for only 1–3% of variation. For each food outlet type, distance to the nearest store was significantly negatively associated with perceived availability of healthy foods. Thus, for every 1.61 km (1-mile) increase in distance to the nearest supermarket, a participant's perceived availability of healthy foods decreased by 0.20 points (unique $R^2 = 4\%$).

When all food outlets were considered jointly, there were no associations between the presence of any type of store and perceived availability of healthy foods using the

Table 1 Descriptive statistics of participants; household shoppers (*n* 705) from a predominantly rural, eight-county region of South Carolina, USA, 2010

Demographic characteristics	
Age (years), mean	57.8
SD	15.1
Sex (%)	
Male	22.3
Female	77.7
Race/ethnicity (%)	
Minority (non-Hispanic black, Hispanic or other)	34.5
Non-Hispanic white	65.5
Education (%)	
Not a high-school graduate	11.4
High-school graduate	35.6
Some college or higher	53.0
Employment (%)	
Not employed	22.6
Retired	31.5
Employed	45.9
Household income per year (%)	
<\$US 20 000	28.9
≥\$US 20 000– <40 000	27.2
≥\$US 40 000– <60 000	17.3
≥\$US 60 000	26.5
SNAP recipient (%)	
Yes	9.9
No	90.1
Spouse or partner (%)	
Yes	64.1
No	35.9
Number of household members, mean	2.5
SD	1.4
Urbanicity (%)	
Urban	20.9
Non-urban	79.1
Transportation for primary shopping (%)	
Personal car/vehicle	93.9
Ride with family/friends	4.7
Public transportation	0.4
Walk, bike or taxi	1.0
Perceived availability of healthy foods	
Availability of healthy foods*, mean	6.2
SD	3.6
Geographic retail food outlet measures	
Presence of supermarkets†	
Presence of a supermarket (%)	11.4
Presence of other food outlets†	
Presence of a convenience store (%)	28.5
Presence of a drug and pharmacy store (%)	13.9
Presence of a dollar and variety store (%)	14.8
Presence of a fast-food restaurant (%)	16.0
Presence of supermarkets‡	
Presence of a supermarket (%)	48.5
Presence of other food outlets‡	
Presence of a convenience store (%)	81.4
Presence of a drug and pharmacy store (%)	40.4
Presence of a dollar and variety store (%)	55.5
Presence of a fast-food restaurant (%)	48.5
Distance to supermarkets (km/miles)	
Distance to nearest supermarket, mean	9.50/5.9
SD	7.24/4.5
Distance to other food outlets (km/miles)	
Distance to nearest convenience store, mean	4.67/2.9
SD	4.18/2.6
Distance to nearest drug and pharmacy store, mean	12.07/7.8
SD	9.50/5.9
Distance to nearest dollar and variety store, mean	8.21/5.1
SD	3.38/4.1
Distance to nearest fast-food restaurant, mean	9.82/6.1
SD	8.21/5.1

SNAP, Supplemental Nutrition Assistance Program.

*Scoring range: 0–12.

†Within a 1.61 km (1-mile) buffer.

‡Within an 8.05 km (5-mile) buffer.

1.61 km (1-mile) buffer. However, the presence of a supermarket within an 8.05 km (5-mile) buffer area was significantly associated with perceived availability of healthy foods ($\beta = 1.09$, $P = 0.025$) when controlling for all other food outlet types. Conversely, distance to the nearest supermarket was significantly inversely associated with perceived availability of healthy foods when accounting for distance to all other outlets ($\beta = -0.16$, $P = 0.003$). Thus, for every 1.61 km (1-mile) increase in distance to the nearest supermarket, the perceived availability healthy foods decreased by 0.16 points. Additionally, distance to the nearest dollar and variety store was also inversely associated with perceived availability of healthy foods, whereas distance to the nearest fast-food restaurant was significantly positively associated with perceived availability of healthy foods. So, as fast-food restaurants get further away, healthy food perceptions go up. For each significant food outlet type, the unique R^2 value was only 1%. In all models, urbanicity explained as much of the variability in perceived availability of healthy foods (unique R^2 values ranging from 2% to 4%) as any of the GIS-derived measures.

Discussion

Our findings show that using GIS-derived presence of retail food outlets may not predict the perceived availability of healthy foods when accounting for all types of food outlet. Previous studies have generally suggested a significant positive association between supermarket availability, assessed by either the presence or number of stores, with perceived availability of healthy foods^(17,19). Our findings are interesting because we examined both presence and distance measures of the food environment and found that only the presence of a supermarket within an 8.05 km (5-mile) buffer and the distance measures to supermarkets, dollar and variety stores, and fast-food restaurants were significant predictors of perceived healthy food options when considering all other food outlets. These findings could reflect that fact that our study took place in a predominantly rural environment and therefore associations are sensitive to boundary selection in the case of GIS-derived presence. Furthermore, our results demonstrate that the selection of study region and geographic measurement may be important factors to consider when assessing geographic- and perception-based food environments together.

Other differences between our study and previous research are in the variety of food outlet types included, i.e. supermarkets ('traditional') and non-traditional retailers. Previous studies have evaluated the relationship between only one geographic measure of a food outlet (typically a supermarket or convenience store) and perceived availability of healthy foods using each outlet type separately. Our study offered a complementary

Table 2 Relationships between geographic retail food outlet measures and perceived availability of healthy foods among household shoppers (*n* 705) from a predominantly rural, eight-county region of South Carolina, USA, 2010

	Separate models†					Full models‡				
	Unique R^{2*}	β	SE	<i>P</i> value	Model R^2	Unique R^{2*}	β	SE	<i>P</i> value	Model R^2
Geographic retail food outlet measures										
Presence of supermarkets§										
Presence of a supermarket	0.03	1.93	0.42	<0.001	0.13	0.002	0.81	0.60	0.176	0.14
Presence of other food outlets§										
Presence of a convenience store	0.02	1.24	0.30	<0.001	0.13	0.002	0.49	0.38	0.201	
Presence of a drug and pharmacy store	0.02	1.77	0.40	<0.001	0.13	0.001	0.54	0.58	0.347	
Presence of a dollar and variety store	0.02	1.53	0.36	<0.001	0.13	0.0005	0.34	0.52	0.509	
Presence of a fast-food restaurant	0.02	1.59	0.37	<0.001	0.13	0.0003	0.28	0.55	0.604	
Presence of supermarkets										
Presence of a supermarket	0.03	1.42	0.30	<0.001	0.15	0.01	1.09	0.49	0.025	0.16
Presence of other food outlets										
Presence of a convenience store	0.01	0.95	0.34	0.005	0.13	0.001	0.39	0.37	0.298	
Presence of a drug and pharmacy store	0.02	1.45	0.33	<0.001	0.14	0.002	0.61	0.47	0.195	
Presence of a dollar and variety store	0.03	1.39	0.29	<0.001	0.15	0.003	0.62	0.42	0.137	
Presence of a fast-food restaurant	0.01	0.85	0.30	0.005	0.13	0.003	-0.75	0.45	0.098	
Distance measures										0.16
Distance to supermarkets (miles)										
Distance to nearest supermarket	0.04	-0.18	0.03	<0.001	0.15	0.01	-0.16	0.05	0.003	
Distance to other food outlets (miles)										
Distance to nearest convenience store	0.02	-0.20	0.05	<0.001	0.13	0.0001	-0.02	0.07	0.764	
Distance to nearest drug and pharmacy store	0.01	-0.07	0.03	0.009	0.12	0.00000003	0.0002	0.04	0.996	
Distance to nearest dollar and variety store	0.04	-0.20	0.03	<0.001	0.16	0.01	-0.15	0.05	0.005	
Distance to nearest fast-food restaurant	0.01	-0.07	0.03	0.018	0.12	0.01	0.11	0.05	0.015	

SNAP, Supplemental Nutrition Assistance Program.

*Squared semi-partial correlation type II for each variable separately.

†Each geographic retail food outlet measure modelled separately.

‡Includes all geographic retail food outlet measures for either presence of or distance to outlets in models. All models adjusted for the covariates age, sex, race/ethnicity, education, employment status, household income, SNAP status, spouse or partner, number of household members and urbanicity.

§Within a 1.61 km (1-mile) buffer.

||Within an 8.05 km (5 mile) buffer.

approach by examining all types of food outlet simultaneously and controlling for possible confounding effects. Thus, it is possible that some significant associations could cancel out in full models because of controlling for both the presence of supermarkets and less healthy options such as fast-food restaurants in the models. This method could be useful in examining future associations involving certain geographic measures of the food environment. However, there is the possibility of multicollinearity⁽³⁵⁾, which would have to be examined. Fortunately, in our study, we found no multicollinearity between measures of food outlets (all variance inflation factors ≤ 10 in the full models).

In our studies, the presence of all food outlet types – including fast-food restaurants – had a significant positive association with the perceived availability of healthy foods in separate models. One possible explanation for this association with fast-food restaurants is spatial co-occurrence of supermarkets and non-traditional food outlets in the food environment⁽³⁶⁾. Thus, other types of food outlet may be located together with supermarkets (e.g. all within the sample shopping area) and are actually markers for the presence of a supermarket that carries fresh fruits and vegetables and low-fat products. However,

when examining whether such co-location of stores existed in our sample, we found that only 9% of participants had a supermarket present within a 1.61 km (1-mile) buffer of their home along with a fast-food restaurant (44% using an 8.05 km (5-mile) buffer), and only 10% of participants had both a supermarket and a convenience store present within a 1.61 km (1-mile) buffer (47% using an 8.05 km (5-mile) buffer). Both fast-food restaurants and convenience stores were present for only 15% of all participants using a 1.61 km (1-mile) buffer (16% using an 8.05 km (5-mile) buffer).

Our study has several limitations. First, our respondents were predominantly female and our landline-based telephone sample yielded a predominantly middle-to-older age distribution. Second, our survey response rate was only 47%; however, this is comparable to the findings from a published evaluation of BRFSS landline response rates (49%)⁽³⁰⁾ and from previous studies⁽¹⁷⁾. Third, the perception questions used may not have adequately characterized the availability of healthy foods in participants' neighbourhoods based on each individual's perspectives and knowledge. It is possible that participants have different perspectives of what constitute fresh or quality foods and/or what food items are

considered fruits and vegetables or low-fat foods. In addition, there was an approximate 1-year interval between the completion of the validation study and the perceptions survey; however, this is a similar or shorter time gap than those reported in other studies⁽¹⁷⁾. Finally, the study operationalized GIS-derived measures centred only on the participants' residences. However, people can purchase food outside the boundaries of their residential neighbourhood (e.g. near or en route to their place of work, school, etc.). Moreover, our study did not examine how food environments are affected by actual shopping patterns or test whether differences in perception of the availability of healthy foods influenced food purchases or consumption. However, the study offers a perspective of the food environment that has been typically characterized throughout the research literature. Future studies should link shopping and dietary habits of individuals with objective and perceived measures of the food environment.

Strengths of our study include the use of GIS-based measures established by validated fieldwork^(8,9), whereas previous researchers have used measures derived from secondary data sources^(17,19). In addition, our study area contained both urban and non-urban communities and may thus be comparable to any new studies examining populations in the south-eastern USA. Our study also included geographic measures of distance in addition to the more traditionally studied presence (availability) of food outlets and included not only supermarkets but also many non-traditional food outlets.

Our study examined the relationship between two types of food environment measure, i.e. geographic and perceived measures. Our results suggest that distance to, not presence of, food outlets is a significant predictor of perceived healthy food options in a person's neighbourhood food environment when considering all types of food outlet. These findings may assist in the development of better measures. Moreover, this research suggests that the perceived availability of healthy foods and GIS-derived food outlet measures may be two different constructs, and researchers should not assume that perception-based measures can be substituted for geographic measures or *vice versa*. In addition, the food environment does not simply comprise presence or proximity to outlets but also includes the selection of products that are offered. Public health professionals and policy makers should consider carefully which type of measure – geographic or perception-based – is most appropriate when assessing the food environment.

Acknowledgements

Financial support: This project was supported by the National Cancer Institute (NCI) (grant number R21CA132133-02S1). The contents of this article are solely

the responsibility of the authors and do not necessarily represent the official views of NCI or the National Institutes of Health. **Conflict of interest:** None. **Authorship:** T.L.B. conceptualized and drafted the manuscript and performed the primary statistical analyses. B.A.B. contributed to the statistical analyses. D.A.F., N.C. and A.D.L. contributed to editing and writing the final manuscript. **Ethics of human subject participation:** This research was approved by the University of South Carolina Institutional Review Board.

References

1. Larson NI & Story MT (2009) A review of environmental influences on food choices. *Ann Behav Med* **38**, Suppl. 1, S56–S73.
2. Larson NI, Story MT & Nelson MC (2009) Neighborhood environments: disparities in access to healthy foods in the US. *Am J Prev Med* **36**, 74–81.
3. Caspi CE, Sorensen G, Subramanian SV *et al.* (2012) The local food environment and diet: a systematic review. *Health Place* **18**, 1172–1187.
4. Minaker LM, Raine KD, Wild TC *et al.* (2013) Objective food environments and health outcomes. *Am J Prev Med* **45**, 289–296.
5. Thornton LE, Pearce JR & Kavanagh AM (2011) Using geographic information systems (GIS) to assess the role of the built environment in influencing obesity: a glossary. *Int J Behav Nutr Phys Act* **8**, 71.
6. Charreire H, Casey R, Salze P *et al.* (2010) Measuring the food environment using geographical information systems: a methodological review. *Public Health Nutr* **13**, 1773–1785.
7. Van Meter EM, Lawson AB, Colabianchi N *et al.* (2010) An evaluation of edge effects in nutritional accessibility and availability measures: a simulation study. *Int J Health Geogr* **9**, 40.
8. Liese AD, Colabianchi N, Lamichhane AP *et al.* (2010) Validation of 3 food outlet databases: completeness and geospatial accuracy in rural and urban food environments. *Am J Epidemiol* **172**, 1324–1333.
9. Liese AD, Barnes TL, Lamichhane AP *et al.* (2013) Characterizing the food retail environment: impact of count, type and geospatial error in two secondary data sources. *J Nutr Educ Behav* **45**, 435–442.
10. Forsyth A, Lytle L & Riper DV (2010) Finding food: issues and challenges in using geographic information systems to measure food access. *J Transp Land Use* **3**, 43–65.
11. Caspi CE, Kawachi I, Subramanian SV *et al.* (2012) The relationship between diet and perceived and objective access to supermarkets among low-income housing residents. *Soc Sci Med* **75**, 1254–1262.
12. Coulton CJ & Jennings MZ (2012) How big is my neighborhood? Individual and contextual effects on perceptions of neighborhood scale. *Am J Community Psychol* **51**, 140–150.
13. Echeverria SE, Diez Roux AV & Link BG (2004) Reliability of self-reported neighborhood characteristics. *J Urban Health* **81**, 682–701.
14. Mujaid MS, Diez Roux AV, Morenoff JD *et al.* (2007) Assessing the measurement properties of neighborhood scales: from psychometrics to econometrics. *Am J Epidemiol* **165**, 858–867.
15. Ma X, Barnes TL, Freedman DA *et al.* (2013) Test–retest reliability of a questionnaire measuring perceptions of neighborhood food environment. *Health Place* **21**, 65–69.

16. Moore LV, Diez Roux AV, Nettleton JA *et al.* (2008) Associations of the local food environment with diet quality – a comparison of assessments based on surveys and geographic information systems: the multi-ethnic study of atherosclerosis. *Am J Epidemiol* **167**, 917–924.
17. Moore LV, Diez Roux AV & Brines S (2008) Comparing perception-based and geographic information system (GIS)-based characterizations of the local food environment. *J Urban Health* **85**, 206–216.
18. Freedman DA & Bell BA (2009) Access to healthful foods among an urban food insecure population: perceptions versus reality. *J Urban Health* **86**, 825–838.
19. Gustafson AA, Sharkey J, Samuel-Hodge CD *et al.* (2011) Perceived and objective measures of the food store environment and the association with weight and diet among low-income women in North Carolina. *Public Health Nutr* **14**, 1032–1038.
20. Zenk SN, Schulz AJ, Lachance LL *et al.* (2009) Multilevel correlates of satisfaction with neighborhood availability of fresh fruits and vegetables. *Ann Behav Med* **38**, 48–59.
21. Williams LK, Thornton L, Ball K *et al.* (2012) Is the objective food environment associated with perceptions of the food environment? *Public Health Nutr* **15**, 291–298.
22. Moore LV, Diez Roux AV & Franco M (2012) Measuring availability of healthy foods: agreement between directly measured and self-reported data. *Am J Epidemiol* **175**, 1037–1044.
23. Giskes K, Van Lenthe FJ, Brug J *et al.* (2007) Socioeconomic inequalities in food purchasing: the contribution of respondent-perceived and actual (objectively measured) price and availability of foods. *Prev Med* **45**, 41–48.
24. Kumar S, Quinn SC, Kriska AM *et al.* (2011) 'Food is directed to the area': African Americans' perceptions of the neighborhood nutrition environment in Pittsburgh. *Health Place* **17**, 370–378.
25. Barnes TL, Bell BA, Freedman DA *et al.* (2015) Do people really know what food retailers exist in their neighborhood? Examining GIS-based and perceived presence of retail food outlets in an eight-county region of South Carolina. *Spat Spatiotemporal Epidemiol* **13**, 31–40.
26. Martinez SW (2007) *The US Food Marketing System: Recent Developments, 1997–2006. Economic Research Report* no. ERR-42. Washington, DC: US Department of Agriculture, Economic Research Service.
27. Gantner LA, Olson CM, Frongillo EA *et al.* (2011) Prevalence of nontraditional food stores and distance to healthy foods in a rural food environment. *J Hunger Environ Nutr* **6**, 279–293.
28. Department of Health and Human Services (2015) *Healthy Food Financing Initiative (HFFI)*. Washington, DC: Department of Health and Human Services; available at <http://www.acf.hhs.gov/programs/ocs/resource/healthy-food-financing-initiative-0>
29. Liese AD, Bell BA, Barnes TL *et al.* (2014) Environmental influences on fruit and vegetable intake: results from a path analytic model. *Public Health Nutr* **17**, 2595–2604.
30. American Association for Public Opinion Research (2011) *Standard Definitions: Final Dispositions of Case Codes and Outcome Rates for Surveys*, 7th ed. https://www.esomar.org/uploads/public/knowledge-and-standards/codes-and-guidelines/ESOMAR_Standard-Definitions-Final-Dispositions-of-Case-Codes-and-Outcome-Rates-for-Surveys.pdf (accessed September 2015).
31. Hu SS, Balluz L, Battaglia MP *et al.* (2011) Improving public health surveillance using a dual-frame survey of landline and cell phone numbers. *Am J Epidemiol* **173**, 703–711.
32. Centers for Disease and Control and Prevention (2013) *State Indicator Report on Fruits and Vegetables 2013*. <http://www.cdc.gov/nutrition/downloads/State-Indicator-Report-Fruits-Vegetables-2013.pdf> (accessed September 2015).
33. Centers for Disease Control and Prevention (2011) *Behavioral Risk Factor Surveillance System Survey Data*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention.
34. US Census Bureau (2010) *2010 Urban and Rural Classification and Urban Area Criteria*. <https://www.census.gov/geo/reference/ua/urban-rural-2010.html> (accessed September 2015).
35. Clary CM, Ramos Y, Shareck M *et al.* (2015) Should we use absolute or relative measures when assessing foodscape exposure in relation to fruit and vegetable intake? Evidence from a wide-scale Canadian study. *Prev Med* **71**, 83–87.
36. Lamichhane AP, Warren J, Puett R *et al.* (2013) Spatial patterning of supermarkets and fast food outlets with respect to neighborhood characteristics. *Health Place* **23**, 157–164.