

Molecular characterization of *Cryptosporidium parvum* detected in Japanese black and Holstein calves in Iwate Prefecture and Tanegashima Island, Kagoshima Prefecture, Japan

Junya AITA¹⁾, Madoka ICHIKAWA-SEKI^{1)*}, Aiko KINAMI²⁾, Seiko YAITA³⁾, Yoshihiro KUMAGAI⁴⁾
Yoshifumi NISHIKAWA⁵⁾ and Tadashi ITAGAKI¹⁾

¹⁾Laboratory of Veterinary Parasitology, Faculty of Agriculture, Iwate University, 3-18-8 Ueda, Morioka, Iwate 020-8550, Japan

²⁾Apartment House Iris B101, 142-12 Shinogikurohata, Takizawa, Iwate 020-0735, Japan

³⁾Kumage Agricultural Mutual Aid Association, Noma-6410-8 Nakatane-cho, Kumage, Kagoshima 891-3604, Japan

⁴⁾Iwate Prefecture Southern Regional Livestock Hygiene Service Center, 41-1 Higashidate, Sakurakawa, Mizusawa-ku, Oushu, Iwate 020-0003, Japan

⁵⁾National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Nishi-2-13 Inada-cho, Obihiro, Hokkaido 080-8555, Japan

(Received 9 February 2015/Accepted 10 March 2015/Published online in J-STAGE 27 March 2015)

ABSTRACT. *Cryptosporidium* oocysts were found in 43 out of 77 calves from two farms in Iwate Prefecture and nine farms on Tanegashima Island, Kagoshima Prefecture, Japan. The DNA fragments of 18S ribosomal RNA (18S rRNA) gene were amplified by a nested PCR from 43 oocyst-positive as well as one oocyst-negative samples. All of them were precisely identified as *C. parvum* by analyzing the nucleotide sequences of the 18S rRNA gene. *C. parvum* oocyst-positive calves ranged in age from 6 to 13 days old and significantly have watery diarrhea ($P < 0.05$). Sequences of the gene encoding the 60-kDa glycoprotein (GP60) in 43 *Cryptosporidium* oocyst-positive samples were identical to that of the zoonotic IIaA15G2R1 subtype. We therefore suggest that calves could be potential sources of *C. parvum* infections in humans.

KEY WORDS: calf, *Cryptosporidium parvum*, GP60, Japan, subtyping

doi: 10.1292/jvms.15-0082; *J. Vet. Med. Sci.* 77(8): 997–999, 2015

Cryptosporidium spp. are protozoan parasites belonging to the phylum Apicomplexa that parasitize the gastrointestinal tract of vertebrates [6, 8, 10]. Infections of *Cryptosporidium parvum*, *C. bovis*, *C. ryanae* and *C. andersoni* have been reported in cattle [5, 8, 9, 13]. The zoonotic subtype *C. parvum* [13] causes severe watery diarrhea in calves [3]. Although *C. parvum* has traditionally been diagnosed by the size and morphology of oocysts detected in fecal samples, the oocysts of *C. parvum* cannot be discriminated from those of *C. bovis* and *C. ryanae*, which are most likely asymptomatic [10]. It is therefore important to accurately distinguish these species from *C. parvum*.

The nucleotide sequences of the 18S ribosomal RNA (18S rRNA) gene have previously been employed for species determination of *Cryptosporidium* spp. [5]. Moreover, intra-specific subtyping of *C. parvum* has been conducted using the sequences of the gene encoding the 60-kDa glycoprotein (GP60) [1, 2, 7, 12, 13], because zoonotic subtypes can be identified by using the GP60 gene. IIa, IIc and III subtype families have been detected in calves worldwide, with the zoonotic subtype IIaA15G2R1 of the IIa subtype family

most frequently reported [13]. Previous studies on *C. parvum* GP60 sequences from Hokkaido [5], Gifu [1], Hyogo [9] and Okinawa [3] Prefectures, Japan, indicated that IIaA15G2R1 was the only subtype distributed in these regions. However, the number of samples and locations analyzed in those studies were insufficient to determine the genetic variation of *C. parvum* throughout Japan. Therefore, this study aimed to characterize the fecal conditions of *C. parvum* infections, as well as to perform molecular characterization of the species from Iwate Prefecture and Tanegashima Island, Kagoshima Prefecture, based on 18S rRNA and GP60 genes, and to compare these results with those of previous studies.

Fecal samples of 46 Holstein calves (under 2 weeks old) were collected from two farms in Iwate Prefecture between March 2013 and September 2014. Fecal samples of 30 Japanese black calves (under 3 months old) were collected from eight farms, and that of one Holstein calf (20 days old) was also collected from a farm in Tanegashima Island, Kagoshima Prefecture between June 2013 and July 2014 (Table 1). The conditions of the fecal samples were recorded, and they were stored at 4°C in the laboratory. The fecal conditions were categorized into watery feces and other fecal conditions. Causation between fecal conditions and *C. parvum* infection was evaluated using Fisher's exact test. The centrifuge sucrose flotation method and microscopic examination were performed to detect *Cryptosporidium* oocysts. After three freeze-thaw cycles at -80°C for 15 min and then at 37°C for 15 min were performed, total DNA was extracted from fecal samples using a QIAamp® DNA Stool Mini Kit (Qiagen,

*CORRESPONDENCE TO: ICHIKAWA-SEKI, M., Laboratory of Veterinary Parasitology, Faculty of Agriculture, Iwate University, 3-18-8 Ueda, Morioka 020-8550, Japan. e-mail: madoka@iwate-u.ac.jp

©2015 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <http://creativecommons.org/licenses/by-nc-nd/3.0/>.

Table 1. Profiles of diarrheal samples and the results of fecal and molecular analyses for *Cryptosporidium parvum*

Location	Farm	Breed of calf	Age (days)	Number of diarrheal samples	Oocyst positive	18S rRNA		GP60	
						Positive	Species identification	Positive	Subtype
Iwate	A	Holstein	6–15	45	32	33	<i>C. parvum</i>	32	IlaA15G2R1
	B	Holstein	8	1	1	1	<i>C. parvum</i>	1	IlaA15G2R1
Subtotal	2			46	33 (72%)	34 (74%)		33 (72%)	
Kagoshima (Tanegashima)	C	Japanese black	6–9	5	4	4	<i>C. parvum</i>	4	IlaA15G2R1
	D	Japanese black	8	1	0	0	<i>C. parvum</i>	0	IlaA15G2R1
	E	Japanese black	8	1	1	1	<i>C. parvum</i>	1	IlaA15G2R1
	F	Holstein	20	1	0	0	<i>C. parvum</i>	0	IlaA15G2R1
	G	Japanese black	10–75	5	1	1	<i>C. parvum</i>	1	IlaA15G2R1
	H	Japanese black	7–49	5	2	2	<i>C. parvum</i>	2	IlaA15G2R1
	I	Japanese black	26–76	5	0	0	<i>C. parvum</i>	0	IlaA15G2R1
	J	Japanese black	4–35	3	0	0	<i>C. parvum</i>	0	IlaA15G2R1
	K	Japanese black	6–33	5	2	2	<i>C. parvum</i>	2	IlaA15G2R1
Subtotal	9			31	10 (32%)	10 (32%)		10 (32%)	
Total	11			77	43 (56%)	44 (57%)		43 (56%)	

Hilden, Germany) according to the manufacturer's instructions. 18S rRNA and *GP60* genes were amplified by nested PCR, as described previously [3, 14]. Second PCR amplicons were sequenced in both directions with the secondary PCR primers, using the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, U.S.A.) and the 3500 Genetic Analyzer (Applied Biosystems).

Small oocysts were detected in 33 out of the 46 calves from Iwate Prefecture and 10 out of the 31 calves from Tanegashima Island. The age of calves discharging oocysts ranged from 6 to 13 days old. 18S rRNA gene fragments were amplified in 34 and 10 samples from Iwate Prefecture and Tanegashima Island, respectively. The nucleotide sequences (786 bp) of these amplicons were identical to that of *C. parvum* deposited in GenBank (accession no. AB746195). The fecal conditions of *C. parvum*-positive samples are summarized in Table 2. The presence of watery feces was causally related with *C. parvum* infection ($P < 0.05$). *GP60* was amplified in 33 and 10 samples from Iwate Prefecture and Tanegashima Island, respectively. The nucleotide sequences (925 bp) of these amplicons were identical to that of the IlaA15G2R1 subtype (AB237131). The representative nucleotide sequences of both genes were deposited in GenBank under accession nos. LC012015–LC012018. In one oocyst-negative fecal sample by the sucrose floatation method and microscopic examination, we were unable to obtain *GP60* amplicons, but successfully amplified the 18S rRNA gene. Because the multi-locus 18S rRNA gene is more efficiently amplified by PCR than the single-locus *GP60* [11], these results suggest that the 18S rRNA nested PCR and sequencing are a superior method capable of accurately detecting *C. parvum*.

Cryptosporidium parvum infection is frequently seen in calves under 2 weeks old [4], so the infected calves in our study (6 to 13 days old) were consistent with the previous report. However, the positive rates of the two locations could not be compared, because the calf ages differed between

Table 2. Causation between *Cryptosporidium parvum* oocyst-positive samples and watery feces

	<i>C. parvum</i> positive	<i>C. parvum</i> negative
Watery feces	27 ^{a)}	14
Other fecal conditions ^{b)}	16	20
Total	43	34

a) *C. parvum* and watery feces were causally related. ($P < 0.05$),
b) Loose feces, mucous feces, muddy feces and formed feces are included.

them. Calves infected with *C. parvum* were previously shown to have watery diarrhea [3]. Watery diarrhea causes severe dehydration in neonatal calves and decreases growth rates of the calves. Therefore, *C. parvum* may lead to a significant economic loss in the cattle industry.

No other subtypes, except for zoonotic IlaA15G2R1 [13], have been reported from calves in Japan [1, 5, 9]. This was also the only subtype detected in the present study. IlaA15G2R1 has previously been identified in a human *Cryptosporidium* infection in Japan [1]. These results indicate that there is a risk of *C. parvum* transmission from calves to humans. Since the subtyping analysis on the basis of *C. parvum* *GP60* amplicons from different areas is as yet insufficient, further work should analyze *C. parvum* from different areas within Japan to investigate the genetic diversity of the species.

ACKNOWLEDGMENT. This study was supported in part by a Joint Research Grant from the National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine (*25-joint-10).

REFERENCES

1. Abe, N., Matsubayashi, M., Kimata, I. and Iseki, M. 2006. Sub-

- genotype analysis of *Cryptosporidium parvum* isolates from humans and animals in Japan using the 60-kDa glycoprotein gene sequences. *Parasitol. Res.* **99**: 303–305. [[Medline](#)] [[CrossRef](#)]
2. Broglia, A., Reckinger, S., Cacció, S. M. and Nöckler, K. 2008. Distribution of *Cryptosporidium parvum* subtypes in calves in Germany. *Vet. Parasitol.* **154**: 8–13. [[Medline](#)] [[CrossRef](#)]
 3. Ichikawa-Seki, M., Aita, J., Masatani, T., Suzuki, M., Nitta, Y., Tamayose, G., Iso, T., Suganuma, K., Fujiwara, T., Matsuyama, K., Niikura, T., Yokoyama, N., Suzuki, H., Yamakawa, K., Inokuma, H., Itagaki, T., Zakimi, S. and Nishikawa, Y. 2015. Molecular characterization of *Cryptosporidium parvum* from two different Japanese prefectures, Okinawa and Hokkaido. *Parasitol. Int.* **64**: 161–166. [[Medline](#)] [[CrossRef](#)]
 4. McNabb, S. J., Hensel, D. M., Welch, D. F., Heijbel, H., McKee, G. L. and Istre, G. R. 1985. Comparison of sedimentation and flotation techniques for identification of *Cryptosporidium* sp. oocysts in a large outbreak of human diarrhea. *J. Clin. Microbiol.* **22**: 587–589. [[Medline](#)]
 5. Murakoshi, F., Tozawa, Y., Inomata, A., Horimoto, T., Wada, Y. and Kato, K. 2013. Molecular characterization of *Cryptosporidium* isolates from calves in Ishikari District, Hokkaido, Japan. *J. Vet. Med. Sci.* **75**: 837–840. [[Medline](#)] [[CrossRef](#)]
 6. Nagano, S., Matsubayashi, M., Kita, T., Narushima, T., Kimata, I., Iseki, M., Hajiri, T., Tani, H., Sasai, K. and Baba, E. 2007. Detection of a mixed infection of a novel *Cryptosporidium andersoni* and its subgenotype in Japanese cattle. *Vet. Parasitol.* **149**: 213–218. [[Medline](#)] [[CrossRef](#)]
 7. O'Brien, E., McInnes, L. and Ryan, U. 2008. *Cryptosporidium* GP60 genotypes from humans and domesticated animals in Australia, North America and Europe. *Exp. Parasitol.* **118**: 118–121. [[Medline](#)] [[CrossRef](#)]
 8. Santin, M., Trout, J. M., Xiao, L., Zhou, L., Greiner, E. and Fayer, R. 2004. Prevalence and age-related variation of *Cryptosporidium* species and genotypes in dairy calves. *Vet. Parasitol.* **122**: 103–117. [[Medline](#)] [[CrossRef](#)]
 9. Satoh, M., Hikosaka, K., Sasaki, T., Suyama, Y., Yanai, T., Ohta, M. and Nakai, Y. 2003. Characteristics of a novel type of bovine *Cryptosporidium andersoni*. *Appl. Environ. Microbiol.* **69**: 691–692. [[Medline](#)] [[CrossRef](#)]
 10. Šlapeta, J. 2013. Cryptosporidiosis and *Cryptosporidium* species in animals and humans: a thirty colour rainbow? *Int. J. Parasitol.* **43**: 957–970. [[Medline](#)] [[CrossRef](#)]
 11. Strong, W. B., Gut, J. and Nelson, R. G. 2000. Cloning and sequence analysis of a highly polymorphic *Cryptosporidium parvum* gene encoding a 60-kilodalton glycoprotein and characterization of its 15- and 45-kilodalton zoite surface antigen products. *Infect. Immun.* **68**: 4117–4134. [[Medline](#)] [[CrossRef](#)]
 12. Wu, Z., Nagano, I., Boonmars, T., Nakada, T. and Takahashi, Y. 2003. Intraspecies polymorphism of *Cryptosporidium parvum* revealed by PCR-restriction fragment length polymorphism (RFLP) and RFLP-single-strand conformational polymorphism analyses. *Appl. Environ. Microbiol.* **69**: 4720–4726. [[Medline](#)] [[CrossRef](#)]
 13. Xiao, L. 2010. Molecular epidemiology of cryptosporidiosis: an update. *Exp. Parasitol.* **124**: 80–89. [[Medline](#)] [[CrossRef](#)]
 14. Xiao, L., Escalante, L., Yang, C., Sulaiman, I., Escalante, A. A., Montali, R. J., Fayer, R. and Lal, A. A. 1999. Phylogenetic analysis of *Cryptosporidium* parasites based on the small-subunit rRNA gene locus. *Appl. Environ. Microbiol.* **65**: 1578–1583. [[Medline](#)]