

The use of indexes evaluating the adherence to the Mediterranean diet in epidemiological studies: a review

Anna Bach^{1,3,*}, Lluís Serra-Majem^{1,2}, Josep L Carrasco³, Blanca Roman¹, Joy Ngo¹, Isabel Bertomeu¹ and Biel Obrador⁴

¹Foundation for the Advancement of the Mediterranean Diet, University of Barcelona Science Park, Spain:

²Department of Clinical Sciences, University of Las Palmas de Gran Canaria, Spain: ³Biostatistics, Department of Public Health, University of Barcelona, Spain: ⁴Faculty of Biology, University of Barcelona, Spain

Abstract

The purpose of this paper is to review some of the methods that several epidemiological studies use to evaluate the adherence of a population to the Mediterranean diet pattern. Among these methods, diet indexes attempt to make a global evaluation of the quality of the diet based on a traditional Mediterranean reference pattern, described as *a priori*, general and qualitative. The Mediterranean diet indexes, hence, summarise the diet by means of a single score that results from a function of different components, such as food, food groups or a combination of foods and nutrients. The reviewed evaluation methods can be classified into three categories depending on the way they are calculated: (1) those based on a positive or negative scoring of the components, (2) those that add or subtract standardised components, and (3) those that are based on a ratio between components.

Dietary scores have been used to explore the multiple associations between the Mediterranean diet, as an integral entity, and health parameters such as life expectancy or the incidence of obesity, cardiovascular diseases and some types of cancers. Moreover, these indexes are also useful tools to measure food consumption trends and to identify the involved factors, as well as to develop comprehensive public health nutrition recommendations.

A more precise and quantitative definition of the Mediterranean diet is required if the adherence to such a dietary pattern is intended to be more accurately measured. Other aspects of the Mediterranean diet indexes should also be taken into account, like the inclusion of typical Mediterranean foods such as nuts and fish and the validation of the dietary pattern approach by using biomarkers.

Keywords

Mediterranean diet
Dietary pattern
Mediterranean diet adherence
Mediterranean diet scores
Mediterranean diet indexes

Historically, in epidemiological studies, the majority being observational, diets consumed by Mediterranean populations have been of interest due to the observation that adults who lived near the Mediterranean Sea had one of the lowest incidences in chronic diseases in the world and one of the highest life expectancies^{1,2}. In the last few decades, numerous associations have been postulated between health status and the Mediterranean diet (MD) and some of its components. For example, it has been suggested that variants of this diet may improve the prognosis in coronary heart disease patients³, that some aspects of the Mediterranean diet pattern (MDP) may protect against the development of diabetes mellitus type II⁴, hypertension⁵, embolisms⁶ and osteoporosis⁷. Additionally, a beneficial effect is suggested with some cancers, such as breast cancer, stomach cancer, colorectal cancer and prostate cancer⁸.

The traditional MD refers to the dietary pattern in the Mediterranean olive grove areas at the beginning of the

1960s, during the post World War II recovery period but before these areas were influenced by fast-food culture⁹. However, the Mediterranean diet is not a homogeneous model within the Mediterranean area. It presents regional variations derived from the same dietary pattern, influenced by various factors, such as socio-cultural, religious and economic determinants, to name a few.

The MDP has been defined in several international scientific meetings^{3,10,11} as varied, not very caloric and based on fresh, local and seasonal products, when possible. This pattern is represented in the Mediterranean Diet Pyramid, a graphic indication that daily intake should be mainly composed of foods of vegetable origin: cereals, fruits, vegetables, legumes and nuts are located at the base of the pyramid. And, with a decreasing intake, in frequency and quantity, in a step up in the pyramid: dairy products, potatoes, poultry, eggs: and on the top, to consume occasionally, sweets, meat and its derivatives.

Other common characteristics are the use of olive oil as the main source of fat, the presence of moderate wine intake at meals and a frequent intake of fish, based on the proximity to the sea.

Most of the scientific knowledge used as evidence for the creation of food guidelines is based on associations between foods or nutrients and the incidence of certain diseases^{12,13}. But it has not been until the 21st century when scientific studies started exploring food patterns in health and disease¹⁴. Until fairly recently there were no systems to evaluate and adequately summarise all the information regarding food patterns¹⁵. To analyse food patterns there are two approximations: developing food indexes, i.e. food scores according to the intake of certain foods; or deriving patterns via multi-variant analysis by means of a factorial analysis, principal component analysis or cluster analysis¹⁶.

The purpose of this paper is to review some of the methods that several epidemiological studies have used to evaluate the adherence of a population to the MDP. The Mediterranean diet indexes attempt to make a global evaluation of the quality of the diet based on a traditional Mediterranean 'reference' pattern, described as '*a priori*', being general and qualitative. The Mediterranean diet indexes, hence, summarise the diet by means of a single score that results from a function of different components, such as food, food groups or a combination of foods and nutrients. These components are previously selected based on prior knowledge or scientific evidence, this approach thus being an '*a priori* approximation'.

Material and methods

An English and Spanish literature search has been done through databases (MEDLINE; NCBI, Bethesda, MD, USA), cited references in related publications, and proceedings of the biannual Barcelona International Congress on the Mediterranean Diet, in order to examine publications on Mediterranean diet adherence indexes. Keywords included were: Mediterranean diet, dietary pattern, Mediterranean diet adherence, Mediterranean diet scores, and Mediterranean diet indexes.

Results

The earlier general diet quality indexes generated the initiative to create the current Mediterranean diet indexes. The reviewed evaluation methods can be classified into three categories depending on the way they are calculated: (1) those based on a positive or negative scoring of the components, (2) those that add or subtract standardized components, and (3) those that are based on a ratio of components. All results are summarised in Table 1.

Index by positive or negative component scoring

The Mediterranean Diet Score

The 'Mediterranean Diet Score' (MDS) was created to measure the adherence gradient to the Greek MDP¹⁷. The MDS is the most extensively used index due to its ease of application, and many variants have been created for the evaluation of multiple diet–health relationships.

The Traditional Greek MD was simplified into eight components to define the MDS-1¹⁷: (1) High ratio of monounsaturated:saturated fat, (2) Moderate alcohol intake, (3) High legume intake, (4) High intake of grains (including bread and potatoes), (5) High fruit intake, (6) High vegetable intake, (7) Low intake of meat and meat products and (8) Moderate intake of milk and dairy products.

The MDS-1 was based on assigning a score from 0 to 1 according to the daily intake of the eight components. In general, the medians of the sample, specific for sex, were used as cut-off points¹⁸ and grams per day were used as the intake measurement¹⁷. A subject received a point if his intake was over the sample median for a protective component (vegetables, fruits, etc.) and below the median for non-protective components (dairy products, meat, etc.). In the case of alcohol (except when specified) 1 point was scored for males if their consumption was within 10 and 50 g/day, and within 5 and 25 g/day for women. If all the characteristics of the diet were incorporated, the highest score was obtained and reflected a greater adherence to the MD. Therefore, the MDS-1 usually ranged from 0 (minimal adherence) to 8 (maximum adherence) if the index had eight components. Generally, a score of 4 or more was associated with satisfactory MDP adherence and better health implications^{17,19}. In most studies, intake was adjusted for calories consumed, 2500 kcal for men and 2000 kcal for women, so the estimations would be independent of the variations present in energy intake.

Greek studies. In the first study of this series, Greek researchers prospectively evaluated the role of the diet in longevity¹⁷. The cohort study included 182 subjects, all of them older than 70, living in three rural towns of Greece. It was observed that the adherence to the Mediterranean diet (MDS ≥ 4) significantly affected elderly life expectancy. Increasing 1 point on the MDS-1 reduced the risk of total mortality by 17% and by 50% with an increase of 4 points.

Subsequently, between 1994 and 1999, another cohort study was carried out in a sample of 22 043 adults²⁰. This showed that a greater adherence to the traditional Mediterranean diet was significantly associated with a reduced total mortality for coronary heart disease and cancer; with an increment of 2 points on the MDS-2 corresponding to a reduction of 25% of the above mentioned mortality. The reduction was stronger in coronary mortality than in cancer mortality. In this study, an MDS-2 of 10 components was used, differing from the

Table 1 Indexes evaluating the adherence to the Mediterranean diet pattern

Reference and index	Objectives	Index components and variables studied	1- Type of study (follow-up period and study's name) 2- Statistics 3- Scoring or cut-off points 4- Dietary assessment	Main results	Country/population (N + age)
Mediterranean diet adherence index by positive or negative component scoring					
Trichopoulos <i>et al.</i>¹⁷ 1995 ‘Mediterranean Diet Score’, original version (MDS-1)	MDP Adh—life expectancy (general and specific mortality)	8-components (g/d): – (+): 1. V/potatoes, Legumes/nuts/seeds, 3. F, 4. Cereals, 5. MRatio, 6. Moderate OH (10–50 g/d ♂; 5–25 g/d ♀) – (–): 7. Dairy products, 8. Meat and poultry Vb: age, sex, smoking, PA, OH	1- Cohort (4–5y) 2- Cox regression models 3- Sex sp medians 4- FFQ	1. ↑ MDP Adh (MDS ≥ 4): ↑ life expectancy + 1p MDS: ↓ 17% RR general mortality (CI: 1, 31%)	Greece/182 elderly (>70y) ♂/♀
Trichopoulos <i>et al.</i>^{20, 22, 30} 2005 Second version of the MDS (MDS-2)	MDP Adh—general mortality	10 ^{20, 30} /g ²² —components (g/d): – (+): 1. V, 2. Legumes, 3. F and nuts ^{20, 30} 4. Cereals/potatoes, 5. Poultry ^{20, 30} 6. MRatio, 7. Moderate OH (10–50 g/d ♂; 5–25 g/d ♀), 8. Fish – (–): 9. Dairy products, 10. Meat products Vb: energy, PA, age, sex, smoking, education, BMI, wt circumf, HTA ²² , chol ²² , DM ²²	1- EPIC Cohort (3.7y) ²⁰ // (3.78y) ²² // (≈ 7y) ³⁰ 2- Cox regression models 3- Sex sp medians 4- FFQ ^{20, 22} //Food diary and 24 h recall (sub sample) ³⁰	↑ MDP Adh (MDS ≥ 4): ↓ general mortality 1. + 2p MDS: ↓ 25% RR of general mortality (CI: 13, 36%) (P < 0.001) ²⁰ 2. + 2p MDS: ↓ 27% RR of general mortality (CI: 7, 42%) ²² 3. + 2p MDS: ↓ 8% RR of general mortality (CI: 3, 12%) ³⁰	1. Greece/22 043 adults ♀/♂ (20–84y) ²⁰ 2. Greece/1 302 ♂/♀ coronary disease patients ²² 3. Europe: Denmark, France, Germany, Greece, Italy, The Netherlands, Spain, Sweden and United Kingdom 174 607 ♂/♀ healthy elderly (≥ 60y) ³⁰
Psaltopoulos <i>et al.</i>²¹ 2004 MDS-2 variant	Olive oil and MDP Adh—HTA	9-components (g/d): – (+): 1. V, 2. Legumes, 3. F, 4. Fish, 5. Moderate OH (10–< 50 g/d ♂; 5–< 25 g/d ♀), 6. MRatio – (–): 7. Cereals, 8. Meat, 9. Dairy products Vb: hypertension, age, city of origin, education, BMI, E, PA, smoking, dietary habits, clinical history	1- Greece-EPIC Cohort EPIC (5y) 2- Regressions 3- Sex sp median 4- FFQ (semi-quantitative and validated)	1. ↑ MDP Adh (+3p MDS): ↓ HTA. The β-coefficient of: -S: -0.8 to -1.0 (CI: -1.1, -0.4) (P < 0.001) -D: -0.2 to -0.4 (CI: -0.5, -0.0) (P = 0.04)	Greece (1994–1999)/ 20 343 individuals without diagnosed HTA: 8685 ♂ and 11 658 ♀ (20–86y)
Osler <i>et al.</i>¹⁹ 1997 MDS-1 variant	MDP Adh—life expectancy	7-components (g/d): – (+): 1. V/legumes/tubers, 2. F, 3. Cereals, 4. MRatio, 5. Moderate OH – (–): 6. Dairy products, 7. Meat Vb: age, sex and smoking, chol	1- Cohort (6y) (SENeca study) 2- Cox regression models 3- Sex sp median 4- Food diary (3d) + FFQ	1. ↑ MDP Adh. (MDS ≥ 4): ↓ general death + 1p MDS: ↓ 21% death RR (CI: 2, 36%)	Denmark/202 ♂/♀ elderly (>70y)

Table 1 Continued

Reference and index	Objectives	Index components and variables studied	1- Type of study (follow-up period and study's name) 2- Statistics 3- Scoring or cut-off points 4- Dietary assessment	Main results	Country/population (N + age)
Kouris-Blazos et al.²³ 1999 MDS-1 variant	MDP Adh—life expectancy	8-components (g/week): – (+): 1. V, 2. Legumes, 3. Cereals, 4. F, 5. MSratio – (–): 6. Milk products, 7. Meat, 8. OH (60 g ♂ and 20 g ♀/d) Vb: age, sex, smoking, ethnic origin	1- Prospective cohort (4y) 2- Cox regression 3- Sex sp medians 4- FFQ	1. 1 MDP Adh (MDS ≥ 4): ↑ survival (Greek-Australians and Anglo-Celts); + 1p MDS: ↓ 17% death RR (CI: – 2, 33%). Anglo-Celts had 40% more mortality vs. Greek-Australians because of dietary differences	Australia/141 Anglo-Celts and 189 Greek-Australians ♀/♂ (≥ 70y)
Woo et al.²⁴ 2001 MDS-1 variant	Chinese dietary pattern. Socio-economic and demographic effects on the index	8-component (g/d): – (+): 1. V + potatoes, 2. F, 3. Legumes/nuts/seeds, 4. Cereals, 5. Fish, 6. MSratio – (–): 7. Milk and dairy products, 8. Poultry and meat Vb: age, sex, smoking, PA, OH	1- 4 Cohorts 2- Multiple comparison; Tukey's methods and chi-squared test 3- Sex sp median ¹⁷ 4- FFQ (semi-quantitative)	1. Between 51% and 96% Chinese had MDS ≥ 4, but vary according to the population and area: – 35–54y group > other age groups (P ≤ 0.001) – ♀ > ♂ (P ≤ 0.001) – Rural areas > urban (P ≤ 0.01)	International/500 ♂ and 510 ♀ Chinese (Hong Kong, Pan Yu, Sydney–Australia and San Francisco)
Lasheras et al.²⁵ 2000 MDS-1 variant	MDP Adh—life expectancy	8 components (g/d): – (+): 1. V, 2. Legumes, 3. Cereals, 4. F, 5. MSratio – (–): 6. Milk and dairy products, 7. Meat, 8. OH (60 g/d ♂ and 20 g/d ♀) Vb: age, sex, BMI, Alb, PA, dieting, health self-assessment	1- Prospective cohort (9.5y) 2- Cox regression models 3- Median 4- FFQ (semi-quantitative)	1. 1 MDP Adh (MDS ≥ 4): ↓ total mortality in < 80y, but not in ≥ 80y + 1p MDS: ↓ 31% death RR (CI: 7, 57%) 2. 1 MDS was related to a ↑ PA and auto-reference to good health	Spain (Asturias)/ 49 ♂ and 112 ♀ elderly institutionalised non-smokers (74 (<80y) and 87 (≥80y))
Gonzalez et al.²⁶ 2002 MDS-1 variant	MDP Adh and sociodemographic factors	9 components (g/d): – (+): 1. V, 2. F, 3. Legumes 4. Cereals, 5. Fish, 6. Olive oil, 7. Wine (40 g/d ♂ and 20 g/d ♀) – (–): 8. Milk and dairy products, 9. Red meat Vb: age, sex, geographic area, education and social class	1- Transverse study (EPIC Study) 2- Multiple linear regression 3- Sex sp quartiles (except wine) 4- Food diary	Variables associated with the MDP Adh: – Age: adults > young adults – Sex: ♂ > ♀ – Social class: low class (22.52) > high class (21.98) – Geography: South (23.5: Murcia) > North (21.64: Asturias)	Spain (Asturias, Navarra, Guipuzcoa, Murcia, Granada) /15634 ♂ and 25812 ♀ (29–69y)

Table 1 Continued

Reference and index	Objectives	Index components and variables studied	1- Type of study (follow-up period and study's name) 2- Statistics 3- Scoring or cut-off points 4- Dietary assessment	Main results	Country/population (N + age)
Schröder <i>et al.</i>²⁷ 2004 MDS-2 variant	Obesity—MDP Adh	10 components (g/d): – (+): 1. V, 2. F, 3. Legumes, 4. Cereals, 5. Fish, 6. Red wine (moderate intake: > 0–20 g/d ♂/♀) 7. Nuts, 8. Olive oil – (–): 9. Milk and dairy (whole), 10. Meat Vb: age, E, PA, education, smoking and OH	1- Transverse study 2- Logistical regression 3- Tertiles 4- FFQ (validated)	1. ↑ MDP Adh: ↓ BMI + 5p MDS: ↓ BMI (coefficient $\beta = -0.043$, $P = 0.03$) in ♂ and (coefficient $\beta = 0.068$, $P = 0.007$) in ♀ 2. Q4 vs. Q1 of the MDS: a 39% less risk of obesity (CI: 0.40, 0.92) in ♂ and (CI: 0.40, 0.93) in ♀	Spain (Girona), 1999–2000/1547 ♂ and 1615 ♀ (25–74y)
Van Staveren <i>et al.</i>²⁸ 2002 MDS-1 ¹⁷ Adapted Mediterranean Diet Score (aMDS) (MDS-1 variant)	MDP Adh—life expectancy	1. MDS-1 ¹⁷ 2. aMDS 7-components (g/d): – (+): 1. V/F, 2. Legumes/nuts/seeds 3. Cereals, 4. MSratio, 5. Moderate OH (percentile50 for ♂ and percentile75 for ♀) – (–): 6. Moderate dairy (percentile25–percentile50), 7. Meat Vb: age, sex and smoking, PA, plasmatic albumin, region	1-Cohort (10y) (SENECA study). 2- Cox regression models 3- Sex sp medians 4- Food diary (3d), and weighted record in sub sample	1. MDS-1: no sig. association 2. aMDS: positive association (not sig.) between the MDP Adh and survival	Europe (12 European countries) / 2586 ♂/♀ elderly (> 70y)
Knoops <i>et al.</i>²⁸ 2004 MDS-2 variant	MDP Adh—life expectancy (general and specific mortality)	8-components (g/d): – (+): 1. V/potatoes, 2. Legumes, nuts, seeds, 3. Fruit, 4. Cereals, 5. Fish, 6. MSratio – (–): 7. Dairy products, 8. Meat and products Vb: age, sex, smoking, PA, OH, nutritional status, morbidity, education, BMI	1- Longitudinal study (10y) (HALE Study) 2- Cox regression models 3- Sex sp medians 4- Food diary	1. ↑ MDP Adh (MDS ≥ 4): ↑ survival 2. The combination of 4 low risk factors (MDS + PA + moderate OH + non-smoking): ↓ 65% RR of TM (CI: 56, 72%)	Europe (11 European countries), 1988–1991 and 1993–1999/ 1507 ♂ and 832 ♀, elderly (70–90y)
Trichopoulos <i>et al.</i>³¹ 2004 MDS-1 variant	MDP Adh—general mortality	7-components (g/d): – (+): 1. V (+legumes), 2. F, 3. Cereals, 4. MS ratio, 5. OH – (–): 6. Dairy products, 7. Meat Vb: age	1- Ecological study 2- Population-weighted averages 3- Mean of the averages of the two groups of countries (Med and non-Med). 4- Food balance sheets	Between the period of 1961–70 and 1990–99: 1. Non-Med countries: ↑ MDS from 2 to 2.5 2. Med countries: ↓ MDS from 4.9 to 4.1 3. MDS difference between Med countries and non-Med countries changed from 2.9 to 1.6	15 EU countries: - Med: Italy/Spain/Greece - Non-Med: rest of EU countries

Table 1 Continued

Reference and index	Objectives	Index components and variables studied	1- Type of study (follow-up period and study's name) 2- Statistics 3- Scoring or cut-off points 4- Dietary assessment	Main results	Country/population (N + age)
Bilenko et al.³² 2005 MDS-1 variant	MDP Adh-CVD	8-components (g/d): - (+): 1. V, 2. Legumes, 3. Cereals, 4. F, 5. MRatio, 6. OH - (-): 7. Milk and dairy products, 8. Meat Vb: age, sex, place of birth, BMI, education, HTA, chol, DM	1- Case-control 2- Fisher's test, logistical regression 3- Sex sp median 4- 24 h recall	1. $\sigma^2 \rightarrow \downarrow$ 1p MDS: \uparrow OR of IM: 1.2 (P = 0.04); of CB: 1.6 (P = 0.01); of A: 1.4 (P = 0.003); and of CVD: 1.3 (P = 0.01). Thus, \uparrow 23–55% the risk of suffering any CVD condition. 2. $\sigma^2 \rightarrow$ non-sig. associations	Israel (Jewish community)/ 1159 adults \geq 35y (520 σ^2 and 639 σ^2)
Bosetti et al.⁸ 2003 MDS-1 variant	MDP-upper aerodigestive tract cancers	8-components (portions/week): - (+): 1. V/potatoes, 2. Legumes/nuts/seeds, 3. F, 4. Cereals, 5. MRatio, 6. Moderate OH (>0–median) - (-): 7. Dairy products, 8. Meat and products Vb: age, sex, education, smoking, BMI and E	1- 3 case-control studies 2- Multiple unconditional models regression 3- Sex sp median 4- FFQ (semi-quantitative and validated)	1. \uparrow MDP Adh: \downarrow aerodigestive tract cancer risk 2. OR MDS \geq 6 vs. MDS < 3: - OC and PHC: 0.40 (CI: 0.26, 0.62) - OEC: 0.26 (CI: 0.13, 0.51) - LC: 0.23 (CI: 0.13, 0.40) 3. + 1p MDS: \downarrow OR OC + PHC: 0.77 (CI: 0.71, 0.83), OEC: 0.72 (CI: 0.65, 0.81), LC: 0.71 (CI: 0.65, 0.78)	Italy 1992–2000 1) 598 cases OC and PHC vs. 14 491 controls 2) 304 OEC vs. 743 controls 3) 460 LC vs. 1088 controls
Martinez-Gonzalez et al.³³ 2002 'A priori' MDP score (1) 'Post hoc' MDP score (2)	MDP Adh-acute MI	8-components (g/d): - (+): 1. V, 2. F, 3. Olive oil, 4. Fibre, 5. Fish, 6. OH - (-): 7. Meat, 8. Elements with 1 GI (white bread, pasta and rice) Vb: smoking, BMI, HTA, chol, DM, PA, CHD family history, aspirin intake, socio-economic status	1- Case-control (MONICA project) 2- Conditional logistical regression 3- (1): Quintiles, and (2): Cut-off according to the dose-response results (components-MI risk) observed in (1) 4- FFQ (semi-quantitative; self-administered)	1. + 1p 'a priori' score: \downarrow OR of MI of 0.92 (CI: 0.86, 0.98) 2. + 1p 'post hoc' score: \downarrow OR of MI of 0.55 (CI: 0.42, 0.73)	Spain/342 adults (<80y): 171 cases (with previous MI) vs. 171 controls
Serra-Majem et al.³⁴ 2004 KIDMED (Mediterranean Diet Quality Index in children and adolescents)	MDP Adh of children and adolescents	16-components: - (+): 1. F o derived/d, 2. 2 pieces F/d, 3. V/d, 4. > 1V/d, 5. Fish 2–3/week, 6. Legumes > 1/week, 7. Pasta or rice \geq 5/week, 8. cereals/breakfast, 9. Nuts 2–3/week, 10. Olive oil at home, 11. Milk products/breakfast, 12. 2 yoghurts or cheese (40 g/d) - (-): 13. > 1 time fast-food, 14. Skip breakfast, 15. Pastries for breakfast, 16. Sweets/day Vb: age, sex, geographical area, social class and rural or urban origin	1- Cross-sectional study of the EnKid Study 2- Descriptive analyses between the three categories (KIDMED: \leq 3 poor MD, 4–7: medium MD and \geq 8: excellent MD) 3- Principles sustaining the MDP and those that undermine it (+1p: positive aspect, -1p: negative connotation) 4- 2 24 h recalls + quantitative FaFQ	KIDMED values of the sample: 1. 4.2%: poor MD, 49.4%: intermediate MD and 46.4%: excellent MD 2. % of excellent MD by areas: 52% North-East vs. 37.5% North 3. % of excellent MD by classes: low (42.8%), medium (47.6%) and high (54.9%) 4. Big cities: \uparrow KIDMED	Spain/3850 children and adolescents (2–24y)

Table 1 Continued

Reference and index	Objectives	Index components and variables studied	1- Type of study (follow-up period and study's name) 2- Statistics 3- Scoring or cut-off points 4- Dietary assessment	Main results	Country/population (N + age)
Goulet <i>et al.</i>³⁵ 2003 Mediterranean Score (MS)	MDP Adh—lipidic profile and body weight changes	11-components (frequency: portions o times/day or week): – (+): 1. Cereals, 2. F, 3. V, 4. Legumes, nuts and seeds, 5. Olive oil, 6. Fish – (In-between foods): 7. Poultry, 8. Dairy products – (–): 9. Eggs, 10. Sweets, 11. Meat Vb: age, abdominal obesity, BMI, lipid levels	1- Nutritional intervention (0.25y) 2- Analysis of variance (Spearman test) 3- Scoring based on the Med Pyramid 4- 4 24 h recalls (telephone interviews) + FFQ	1. MS increase with intervention: 21.1 ± 3.6 (t = 0) $\rightarrow 28.8 \pm 4.5$ (t = 6 and 12 weeks) ($P < 0.0001$). 2. Sig. effect on metabolism (\downarrow apoB, \downarrow chol, and \downarrow BMI ($P < 0.05$)) 3. No sig. changes in HDL, LDL and TG	Canada/77 ♀ adults (30–65y)
Panagiotakos <i>et al.</i>³⁶ 2004 Dietary Score (DS)	MDP Adh—lipidic profile	10-components (frequency: times/month or day): – (+): 1. Whole grain cereals, 2. F, 3. V, 4. Legumes, nuts and olives, 5. Olive oil, 6. Fish, 7. Dairy products 8. Wine (moderately) – (–): 9. Eggs, potatoes and sweets, 10. Meat Vb: sex, age, BMI, smoking, education and PA	1- Cohort (1y) (ATTICA Study) 2- MANCOVA and multiple linear regression 3- Scoring based on the Med Pyramid 4- FFQ (validated)	1. \uparrow 10p DS: \downarrow 22 mg/dl of oxidised-LDL chol (CI: 8,36) ($P = 0.04$) 2. MDP Adh (+statins) vs. Westernised diet: \downarrow 9% of total chol ($P = 0.04$), \downarrow 19% of LDL-chol ($P = 0.02$), \downarrow 32% of oxidised-LDL chol ($P < 0.001$)	Greece (2001–2002) /1128 ♂ and 1154 ♀ (>18y)
Gerber <i>et al.</i>³⁸ 2005 Mediterranean Diet Quality Index (Med-DQI)	MDP Adh—socio-demographic and lifestyle variables	7-components: – (+): 1. Olive oil (ml), 2. Fish (g), 3. Cereals (g), 4. Vegetables + Fruits (g) – (–): 5. Saturated fatty acids (energy %), 6. Cholesterol (mg), 7. Meat (g) Vb: age, BMI, sex, smoking, wine intake, OH, vitamin supplement intake, social class, education, urban or rural origin, biomarkers	1- Transverse study 2- Wilcoxon test rank and Spearman test for Med-DQI—biomarkers correlations 3- Recommended guidelines and when those exist: fertiles 4- FFQ (validated)	- Med-DQI: 1. old > young 2. rural areas > urban areas 3. working class > professional class 4. non-smokers > smokers - \uparrow correlation between Med-DQI and biomarkers (except chol)	France/473 ♂ and 491 ♀ (30–77y)
Ciccarone <i>et al.</i>³⁹ 2003 MDP Score (Med Diet Pattern Score)	MDP Adh—PAD in DM type II patients	14 components (times/week or day): – (+): 1. Cooked V, 2. Raw V, 3. Carrots, 4. F, 5. Fish, 6. Olive oil – (–): 7. Eggs, 8. Meat, 9. Processed meat, 10. Cheese, 11. Vegetable oils, 12. Butter, 13. Milk cream, 14. margarine Vb: smoking, duration of DM, history of hyperlipidemia, hypertension, BMI and PA	1- Cohort study 2- Univariate and multi-variant analysis 3- Food scientific evidence 4- FFQ (semi quantitative)	1. \uparrow MDP score (≥ 11): \downarrow of 56% of PAD risk (CI: 17, 76%), independently associated to the length of DM and the presence of hypertension	Italy/432 adults with DM type II 144 subjects with DM type II and PAD vs. 288 subjects with DM type II

Table 1 Continued

Reference and index	Objectives	Index components and variables studied	1- Type of study (follow-up period and study's name) 2- Statistics 3- Scoring or cut-off points 4- Dietary assessment	Main results	Country/population (N + age)
Mediterranean diet adherence index that adds or subtracts standardised component					
Sanchez-Villegas et al.⁴⁰ 2002 MDP adherence index	MDP Adh—sociodemographic and lifestyles factors	9-components (g/d): – (+): 1. Legumes, 2. Cereals, 3. F. 4. V, 5. Moderate OH (30 g/d ♂ and 20 g/d ♀), 6. MSratio – (–): 7. Trans fatty acids, 8. Meat, 9. Milk and dairy products Vb: age, sex, smoking, PA, nap, education, HDL chol levels self-knowledge	1- Transverse study (SUN Prospective Cohort (2y)) 2- Multi-variant analysis (LOWESS) 3- Sex sp mean 4- FFQ (semi-quantitative)	Associated variables with MDP Adh: – Sex (♀ > ♂) (average difference = 4.1% (CI: 3.2, 4.9) – Age: younger university students > older university students ($P < 0.001$) – Physical activity: active > sedentary ($P = 0.01$ in ♂, $P < 0.001$ in ♀)	Spain (Navarra), 1998/1587 ♂ and 2260 ♀ University of Navarra students
Tur et al.⁴¹ 2004 MDP adherence index	MDP Adh—sociodemographic and lifestyles factors	9-components (g/d): – (+): 1. MSratio, 2. Moderate OH (30 g/d ♂; 20 g/d ♀), 3. Legumes, 4. Cereals and roots, 5. F, 6. V, 7. Fish – (–): 8. Meat, 9. Milk and dairy products Vb: age, sex, PA, marital status, place of origin, education, social-economic status, smoking habit, presence of CVRF	1- Transverse study 2- Logistical regression models 3- Mean 4- 2 24 h recalls + FFQ (semi-quantitative)	1. Mean MDP Adh: 43.1% (SD 5.8%) 2. Variables associated with MDP Adh: – Age: subjects (≥ 46 y) < subjects (< 45y) ($P = 0.01$) – Physical activity: active > sedentary ($P = 0.01$) – Smoking habit: non-smokers > smokers ($P = 0.01$)	Spain (Balearic Islands, 1999–2000)/1200 adult ♂/♀ (16–65y)
Alberti-Fidanza et al. 1999⁴² and 2004⁵⁰ MAI (Mediterranean Adequacy Index)	Adh to a Italian MDP	2 large blocks of products (percentage of total energy) – (+) Med: 1. Carbohydrate group (bread, cereal, legumes (raw-dry), potatoes), 2. Protective food group (vegetables, fruit, legumes (raw-fresh), fish, alcohol (ex. wine), vegetable oils) – (–) Non-Med: 3. Food of animal origin group (milk, cheese, meat, eggs, animal fat and margarines), 4. Sweets food group (sweet beverages, pastries, cookies and sugar). Vb: sex, age and rural or urban origin	1- 3 Italian Cohorts from the 'Seven Countries Study' (31y) 2- Student-t (paired) and Wilcoxon–Mann–Whitney 3- MAI = 1 + 2/3 + 4 4- Food diary (weighted record in a sub sample)	MAI averages from the 1960s to the 1990s: – Crevalcore: 2.9 → 2.2 – Montegiorgio: 5.6 → 3.9 – Nicotera: 7.5 (considered as an Italian MDP reference)	Italy (1960–1991): Crevalcore, Montegiorgio and Nicotera/♂ (40–59y)

Table 1 Continued

Reference and index	Objectives	Index components and variables studied	1- Type of study (follow-up period and study's name) 2- Statistics 3- Scoring or cut-off points 4- Dietary assessment	Main results	Country/population (N + age)
Fuentes-Bol et al.⁴³ 2002 MAI (Mediterranean Adequacy Index)	MDP Adh – sociodemographic differences	2 large blocks of products: – (+) Med: Cereals, fish, fruits, vegetables, legumes, potatoes, vegetable oils, red wine, nuts – (–) Non-Med: eggs, meat, milk, butter, cheese, sugar, pastries, juices and soft drinks Vb: social class, rural or urban origin	1- Transverse study 2- Comparison of averages 3- MAI = (1 + 2)/(3 + 4) 4- Family budget sheets	MAI: - low classes > high classes - towns > cities	Spain/3022 families

Significant results are shown, except those specified, (+): protective component, (–): non-protective component, ↑: increase, ↓ decrease/lower, A: angioplasty, Adh: adherence, Alb: albumin, apoB: apolipoprotein B, BMI: body mass index, CA: cancer, CB: coronary bypass, CHD: coronary heart disease, CH: carbohydrates, CI: 95% confidence interval, chol: cholesterol, CVD: cardiovascular disease, CVRF: cardiovascular risk factor, D: diastolic blood pressure, d: day, DM: diabetes mellitus, DS: disaccharides, E: energy intake, F: fruit, FBS: family budget sheet, FFQ: food frequency questionnaire, GI: glycaemic index, H⁺: hypothesis, HDL: high density lipoprotein, HTA: hypertension, LC: laryngeal cancer, LDL: low density lipoprotein, MDP: Mediterranean diet pattern, MDS: Mediterranean diet score, Med: Mediterranean, MFA: monounsaturated fatty acids, MI: myocardial infarction, MS: monosaccharides, MSratio: monounsaturated–saturated ratio, N-S: non-smoker, OC: oral cancer, OEC: oesophagus cancer, OH: alcohol intake, OR: odds ratio, P: P-value, p: points, PA: physical activity, PAD: peripheral artery disease, PFA: polyunsaturated fatty acids, PHC: pharyngeal cancer, Prot: proteins, Q: quintiles, RR: relative risk, S: systolic blood pressure, SFA: saturated fatty acids, sig.: significant, sp: specific, TM: total mortality, V: vegetables, Vb: study variables, P: polyunsaturated, wt circumf: waist circumference, y: years.

initial 8-point MDS-1, as the second version incorporated fish and moderate poultry intake.

The Greek branch of the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort explored the association between the MDP and olive oil for hypertension, since hypertension can give way to different forms of cardiovascular disease²¹. A nine-component variant of the MDS was applied, finding it inversely related to systolic and diastolic blood pressure, after socio-demographic and anthropometric covariates were controlled. Although cereal intake, a component considered protective, was positively associated with high arterial blood pressure.

Once again, based on the Greek branch of the EPIC cohort, a variant of the MDS was applied to a Greek sample of 1302 coronary disease patients, observing that those who presented a greater MDP adherence had a reduced risk of general mortality²². Specifically, with a 2-point increase in the score the relative risk of general mortality was reduced by 27%, and by 31% in regards to coronary heart disease.

Danish studies. As part of the Euronut Survey in Europe on Nutrition and the Elderly, Olser *et al.*¹⁹ via a Concerted Action (SENECA) examined the influence of the MDS on general survival in a cohort of 202 Danes of advanced ages, with a 6-year follow-up¹⁹. Data regarding food intake was obtained by a 3-day dietary diary and a food frequency questionnaire. The MDS was comprised of seven components, and as a reflection of the Danish food patterns, vegetables and legumes were combined into one component. An increase of 1 point on the score predicted a reduction by 21% of global mortality.

Australian studies. A cohort study of a similar design to the above mentioned studies, with a sample of 141 Anglo-Celts and 189 Greco-Australians, was realised in Melbourne²³. There was a double objective; on one hand to evaluate if the rural Greek results could be replicated in an urban Australian environment and on the other hand, to analyse if the benefits of the Mediterranean diet were applicable to non-Mediterranean populations. The eating habits of the participants were evaluated with extensive and validated frequency questionnaires. The results were compatible with the hypothesis that a high score, i.e. greater or equal to 4, was associated with longer survival. In this study, 153 (81%) Greco-Australians and 39 (28%) Anglo-Celts gathered four or more Mediterranean diet characteristics. An increase of 1 point on the score supposed a reduction of 17% in general mortality.

Chinese studies. The Chinese diet shares some characteristics with the Mediterranean diet, such as a high intake of vegetables and fruit and a low intake of meat. Chinese researchers aimed to determine if dietary habits of some Chinese populations were similar to those of the MDP, with an MDS adapted to the Chinese diet²⁴.

The score was calculated for 1010 Chinese living in four very diverse geographic regions: Hong Kong, a rural town near Pan Yu in southern China, Sydney (Australia) and San Francisco (USA). The results indicated that the majority of the Chinese population, living in China or elsewhere, had dietary patterns similar to the Mediterranean, and achieved in the majority of cases higher MDS scores as compared with Trichopoulou *et al.*'s¹⁷ Greek population. Middle-aged individuals (35–54 years) obtained a higher score than those in other age groups, as did women when compared to men. Moreover, rural populations showed greater adherence due to the ease of maintaining the dietary pattern.

Spanish studies. A prospective cohort study was carried out during 9 years in 161 Spanish nursing home residents, 65 years old or older and non-smokers²⁵. An increase in 1 point on the score was associated with a significant reduction in mortality by 31% in individuals younger than 80 years old. However, no significant associations were found in populations older than 80 years of age.

A transverse study examined the influence of demographic and social variables on the adherence to MDP in 15 634 women and 25 812 men aged 29 to 69, who were volunteers to the EPIC-Spain Study²⁶. A modified version of the initial MDS-1¹⁷ was used. Each of the nine components of the score received 1 to 4 points based on the quartiles of intake (calories adjusted), except in the case of wine, where moderate intake was used as a cut-off point. No variations in adherence to the MDP based on educational level were found, but small differences regarding social status were seen. Less adherence was observed in young adults and in women, and was slightly higher in southern areas as compared to northern Spain.

An MDS variant was applied to study the relationship between the Mediterranean diet and obesity²⁷. The index was calculated with the exception of red wine consumption, according to the tertile distribution of intake. The total score that could be obtained ranged from 9 to 27 points. Increasing the score by 5 units, body mass index (BMI) decreased significantly in men and in women, controlling potentially confounding variables such as sociodemographic and lifestyle factors (i.e. physical activity). Individuals in the upper quartile with respect to those in the lower quartile of the score had 39% less risk of obesity, for both men and women.

European studies. The Healthy Aging: a Longitudinal study in Europe (HALE) project is a prospective study that followed men and women of advanced age during 10 years in 11 countries around Europe to investigate the association between diet and lifestyle factors with mortality due to coronary, cardiovascular and cancer causes²⁸. The HALE project included participants proceeding from two studies: SENECA and the Finland, Italy, the Netherlands, Elderly study (FINE)^{28,29}. The MDS was applied to these data, taking into account some

modifications with respect to the original MDS¹⁷. Potatoes were added to the vegetable group, fish was added as an independent category and alcohol was not included in the score. Sex-specific median intakes were adopted as cut-off points. To evaluate the association between mortality and lifestyle variables, a low risk group was established having the following characteristics: high dietary score, non-smoker, moderate drinker and physically active. The adherence to the MD was associated with 22% less risk of general mortality, being physically active and a non-smoker with 37% and 35% less risk, respectively. The combination of the four protective factors reduced general mortality by 60–64%, supporting the hypothesis that the participants who followed an MD and maintained healthy lifestyle habits had less general and specific mortality, even in ages 70 to 90 years old.

From the final SENECA European study, where 1507 men and 832 women of advanced age from 12 European countries participated, predictive values of dietary patterns in survival during 10 years were evaluated using the original MDS (MDS-1) and an adaptation (aMDS)²⁹. The adaptation consisted of varying the cut-off point of certain components: the optimal intake of dairy products was considered as an interquartile range for men and women, the optimal intake of meat and poultry in women was set as below the 75th percentile and the maximum alcohol intake in women was also set at the 75th percentile. The application of the MDS-1 did not yield a significant positive association between diet and life expectancy. On the other hand, the aMDS, although not contributing significant results, did show a clear tendency that a favourable score was related to higher survival.

Recently, the EPIC-Elderly prospective cohort study evaluated the relationship between the 'modified' Mediterranean diet and the survival of 74 607 individuals aged 60 years or more in nine European countries³⁰. The adherence to the MDP was measured through the MDS-2²⁰, in which a lipid ratio was incorporated where the polyunsaturated fatty acids appear in the numerator, so that the index could be applied to non-Mediterranean European countries. It was found that a greater adherence to the MDP was associated with a significant reduction in general mortality. An increment of 2 points entailed a reduction of 8% of relative risk of mortality and 7% when the exposition factors were calibrated between countries.

Also within the European context, differences in following the MDP were studied between Mediterranean countries and non-Mediterranean countries based on FAO's Food Balance Sheets³¹. During the study period (1961–1970 and 1990–1999) it was observed that non-Mediterranean countries presented an increase of a seven-component mean score from 2 to 2.5, which reflected an increase of fruit, vegetable and vegetable fat consumption. In contrast, the diet of Mediterranean countries was negatively affected by the increased intake of meat and animal fat, as shown by a reduction in MDS mean from 4.9

to 4.1. Therefore, the differences in the index score between the two groups of countries were reduced, this being associated with the reduction in the differences in general mortality observed for the two groups.

Israeli studies. The MDS was applied to 1159 Jewish people³², finding that there were relatively low levels of adherence to the pattern (less than 20%). A strong association between the low MDP adherence (MDS ≤ 4) and cardiovascular disease was found only in men. A reduction of 1 point in the MDS was related to a 23–55% increased risk for cardiovascular disease.

Italian studies. In Italy, it was observed that the Mediterranean diet favourably decreased the risk of suffering aerodigestive cancers (oral, oesophagus, pharynx, and larynx) in three case–control studies, with a total sample of 4684 individuals, where the original MDS-1¹⁷ was applied⁸. Therefore, an increase of 1 point on the score meant a reduced risk of 23% in the case of oral and pharynx cancer, 28% in oesophageal cancer and 29% in larynx cancer.

'Post hoc' Mediterranean diet pattern score

To understand if the MDP plays a protective role in re-infarction and cardiovascular death, a '*post hoc*' index was created based on '*a priori*' defined MDP³³. The '*a priori*' index was made up of eight components, where apart from foods, such as those with high glycaemic index, nutrients could also be found. According to the quintile distribution by the intake of each component (adjusted by energy) a score was assigned from 1 to 5 for each element. In the case of protective components 1 point was assigned to the lower quintile and 5 points to the upper quintile, and for the non-protective components, the scoring was inverted. This way, each participant had a total score that ranged from 8 to 40. In the '*post hoc*' index a single cut-off point was used for each component based on the 'previous' results of the dose–response associations between the intake of each component and the risk of suffering a second myocardial infarction. For the majority of index components, individuals in the second quintile had a major reduction of risk in comparison to those in the first quintile, but no significant differences between Q2 and Q5 were observed. Therefore, with the '*post hoc*' index if the participant consumed more than this value (Q2) a point was assigned for each component, thus obtaining a score between 0 and 8. The results of both indexes indicated that when the score increased, the odds ratio for myocardial infarctions was significantly reduced.

KIDMED Index

The KIDMED Index was a Mediterranean diet quality index constructed to evaluate the food habits of a population of 3850 Spanish children and adolescents aged between 2–24 years in the Enkid study³⁴. The index contains 16 elements and is composed of a scale from 0 to

12 points. A point was added if a series of Mediterranean characteristics were met, and subtracting a point with 'Westernised' or harmful food behaviours such as frequently consuming 'fast foods', pastries and sweets and not having breakfast. 4.2% of the sample presented a poor MDP, 49.4% had an intermediate pattern and 46.4% an excellent MDP. In high social classes there was a greater proportion of children and adolescents with excellent Mediterranean diets (54.9%) as compared to lower (42.8%) and medium (47.6%) classes.

Indexes based on the Mediterranean Diet Pyramid

Mediterranean score. Canadian researchers studied 77 women to examine food habits, plasma lipoprotein profiles and body weight modifications based on a 12-week nutritional intervention promoting the MD³⁵. Scoring based on 11 components of the Mediterranean pyramid by Oldways Preservation Trust was designed to evaluate MDP adherence. A partial score of 0 to 4 was attributed to each component. Food found at the base of the pyramid received a high score when consumed frequently. However, food found at the peak of the pyramid (meat, sweets and eggs) was given a high score when consumed less frequently. From the nutritional intervention, it was observed that the 'Mediterranean score' sample mean increased significantly from 21.1 points to 28 points, and resulted in a slight but significant improvement of the metabolic profile (total cholesterol, apoB levels and BMI), a significant decrease in energy coming from lipids and a significant reduction in weight and waist circumference.

Dietary score. Greek researchers constructed a dietary score also based on the Mediterranean Diet Pyramid¹¹ to study the differences in plasma lipids according to MDP adherence³⁶. The index was constructed based on higher scoring from 0 to 5 points according to intake frequency of typical Mediterranean products, and for those components far from the MDP a decreased scoring. The adherence to the Mediterranean diet resulted in significant reductions only for levels of oxidised LDL-cholesterol.

Mediterranean Diet Quality Index

The Mediterranean Diet Quality Index (Med-DQI) was an adaptation of the 'Diet Quality Index'³⁷ to evaluate the MDP. Olive oil, fish, and alternative meat substitutes were added³⁸. The objective of the study was to evaluate the adherence to the MDP in a French population, to study the socio demographic and lifestyle associated factors, and to validate and correlate the Med-DQI with biological markers (carotene, vitamin E, EPA and DHA). In the Med-DQI a score from 0 to 2 was assigned to each food group according to the recommendations when existing, or otherwise using the population intake tertiles to assign cut-off points. The total score of the index was from 0 to 14 points. The lower the Med-DQI value, the healthier the diet. Elder individuals and those living in rural areas,

working class individuals and non-smokers presented a greater Med-DQI. There was a high correlation between Med-DQI and studied biomarkers, with the exception of cholesterol.

Mediterranean diet pattern score

In an Italian cohort study, a score was created to be able to study the association between the MDP and the risk of developing peripheral vascular disease in type II diabetic patients³⁹ as it was postulated that more than a specific food alone, a high score may play a protective role. A score was developed where a point was added according to food intake with sufficient evidence of its beneficial effect on coronary artery disease, and 0 for those foods that have potentially harmful effects. Specifically, the study found that individuals who obtained 11 points or more had 56% less risk of suffering this illness.

Adherence index that adds and subtracts standardised components

The SUN study is a prospective cohort study based on a sample of 4259 University of Navarra students with the objective of assessing the protective role of the MDP on coronary diseases and evaluating variables associated with this pattern⁴⁰. An index was built that calculated a value of adherence as a percentage, standardising the food intake values to add and subtract the components depending on its nature. The results suggest that there is a progressive abandonment of the traditional MDP in younger individuals and those individuals who led an active lifestyle had greater adherence to the MDP.

In a transverse study of similar characteristics carried out in the Balearic Islands⁴¹, a variant of the Sanchez-Villegas index was applied. It also had the objective of analysing the prevalence of the MDP in a sample of 1200 individuals and the sociodemographic and lifestyle factors related to this pattern. The adherence to the MDP was defined through 9 points or characteristics, with small component modifications as compared to Sanchez-Villegas *et al.*, but with the same calculation methodology. The MDP adherence in the Balearic population was 43.1%, similar for all socio-demographic groups and lifestyles but with differences in age, sex, physical activity and smoking habits.

Mediterranean Adequacy Index: quotient between components

The adherence to the Italian MDP reference was measured in two Italian cohorts of the Seven Countries Study (Crevalcore and Montegiorgio)⁴². The 'Mediterranean Adequacy Index' (MAI) was based on a quotient between the sum of energy proceeding from Mediterranean products (carbohydrate and protective food groups) and the sum of energy from non-Mediterranean products (animal origin foods and sweets). High index values indicated a greater MDP adherence. During the 31-year follow-up, a progressive abandonment of the MDP was

observed, taking as a reference an Italian MDP from the town of Nicotera.

The MAI was also applied to Spanish Family Food Balance Sheets from the Ministry of Food and Agriculture⁴³. A total of 3022 Spanish households participated in the study to evaluate the influence of sociodemographic variables towards the adherence to the MDP. To apply MAI to Spanish data and to define Spanish MDP products a classification based on the Mediterranean Diet Pyramid was created, placing in the numerator those foods at the base of the pyramid and in the denominator those found at the vertex. Generally, high MAI values were observed in lower classes than in higher classes. Also, less adherence to the MDP was found in cities as compared to towns.

Discussion

Food pattern studies summarise the complexity of a diet taking into account the synergic effects or interactions between nutrients or foods that comprise the diet^{44,45}.

Currently, indexes to measure the MDP are being explored in epidemiological studies, but not so much in experimental studies⁴⁶. Simultaneously, there have been some attempts for outlining and specifying the definition of this food pattern^{3,10,11}. According to the experts, there is still the need for a more precise and quantified definition of the pattern, which could be obtained, for example, by establishing inferior and/or superior component limits⁴⁷. However, reaching a consensus is complex since the pattern may be related to a specific Mediterranean region and to a selected period that are adopted as references. Moreover, apart from using the traditional MDP as the baseline, current debates include the admission of a 'modern' MDP definition that incorporates current scientific knowledge on the relationship between diet and chronic diseases.

These problems with the definition of the pattern indirectly affect pattern evaluation methods and components used. For instance, there is a debate on the type of fat to be included in the definition (monounsaturated versus polyunsaturated)²², on how to include dairy products (their proportion and composition)²¹, the importance of different types of meat³³, the classification of refined cereals as protective or 'non-protective' components²¹, the establishment of a definition for moderate alcohol intake, and the presence of nuts and fish as independent components²⁷.

On the other hand, Mediterranean diet evaluation methods that utilise scorings are limited by subjectivity in the selection of scoring components, mostly conditioned by available data and by the study's own objectives in assigning cut-off points and by the interpretation of the diet-disease relationship¹⁶. The variability in choosing cut-off points in the score, and distributing the population into different intake groups according to the grade of

adherence, may influence the interpretation of results. Also, the use of indexes and cut-off points by other authors is a limitation since the population in which the index is applied may differ from the population for which the index was originally designed.

Furthermore, the majority of indexes do not really measure the adherence to a universal MDP but rather to a specific pattern, based on the distribution of selected food groups in the same population. This makes international comparisons more difficult. Some indexes postulate monotonic relations and do not take into account possible tolerance limits⁴⁸. Another component-related problem is that the same importance is usually arbitrarily given to all index components, independently of the components' proportions in the diet and of the scientific evidence on the diet–disease relationship²⁶.

Even though it is difficult to quantify the adherence to a diet and despite the fact that indexes require operational definitions, grouping foods to obtain complex scores is a very useful method to evaluate epidemiological associations³⁹. It is a simple and intuitive approximation to estimate attributable risk to a dietary pattern, although it does not result in such apparently strong associations as those obtained with *a posteriori* analytical methods⁴⁹. Earlier studies have focused on the Mediterranean diet and life expectancy relationships within elderly populations: a 4-point increase in the MDS was associated with a significant reduction of total mortality in elderly in studies conducted in Greece, Denmark, Australia, Spain and throughout Europe^{15,23,28}. In the last few years, however, age ranges of the studied populations have been amplified and the studied MDP–health associations expanded²⁰. For instance, a 5-point increase in the MDS led to a reduction in BMI²⁷. On the other hand, a 1-point reduction of the MDS was related to a significant increase in cardiovascular risk³². A greater adherence to MDP led to a significant reduction in blood pressure²¹ and in the risk of aerodigestive tract cancers⁸, of suffering an acute myocardial infarction³³ and of peripheral vascular disease in type II diabetes patients³⁹. A simple nutritional intervention to promote the MDP may be effective in modifying nutritional habits in people in free-living conditions and may result in significant effects on apoB and BMI reduction³⁵.

Some positive aspects of the reviewed studies can be found in the use of samples of extensive populations with a wide variety of patterns and lifestyles, and the use of valid frequency questionnaires^{22,28}. Other elements that allow a critical evaluation of the MD–health hypothesis are the prospectiveness of the study, the evaluation of many confounding variables, the participation of cohorts from different European regions and the adjustment of dietary exposure factors between countries²². The Mediterranean diet adherence indexes are especially useful when the MD is compared with other food patterns, such as the Western pattern.

Some of the revised studies show problems related to dietary assessment methods. The majority of indexes are based on food groups, which require food categorisation, with the associated difficulties when evaluating mixed dishes.

The identified co-variables to be controlled when studying the MDP–health relations are also diverse in the different studies, basically influenced by their objectives. Age, sex, smoking and physical activity are the most frequently controlled variables because of their strong association to the MDP. To control for dieting is also considered to be necessary, since modifying food habits due to presence of disease may bring on increased MDP adherence²⁵.

Other commonly controlled demographic variables are the geographic origin of the population, educational level, ethnic origin or social class. Among clinical and anthropometric variables are cholesterol, self-assessment of health status, mobility, blood pressure, diabetes and BMI.

In the different studies the MDP is analysed in several ways. Dietary indexes are being used to evaluate the degree of adherence to the MDP, to study the socio-economic and lifestyle variables that influence the adherence to the pattern and to explore the multiple associations between the Mediterranean diet, as an integral entity, and health parameters such as life expectancy or the incidence of obesity, cardiovascular diseases and certain types of cancers. The evaluation of these associations is based on the risk reduction for chronic diseases, or on the modification of a clinical parameter of nutritional status (biochemical, anthropometric and clinical). Moreover, these indexes are also useful tools to measure food consumption trends and to identify the involved factors, as well as to develop comprehensive public health nutrition recommendations.

To sum up, a prudent dietary pattern for health promotion apparently coincides with the Mediterranean diet. MD indexes are useful tools to study this pattern and its association with health. A more precise and quantitative definition of the Mediterranean diet, however, is required if the adherence to such a dietary pattern is intended to be more accurately measured. Other aspects of the Mediterranean diet indexes should also be taken into account, such as the inclusion of typical Mediterranean foods like nuts and fish and the validation of the dietary pattern approach via the use of biomarkers.

References

- 1 Keys A, Menotti A, Karvonen M. The diet and 15-year death rate in the Seven Countries study. *American Journal of Epidemiology* 1986; **124**: 903–15.
- 2 Nestle M. Mediterranean diets: historical and research overview. *American Journal of Clinical Nutrition* 1995; **61**: 1313S–20S.

- 3 De Lorgeril M, Salen P, Martin J, Monjaud I, Delaye J, Mamelle N. Mediterranean diet, traditional risk factors and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999; **99**: 779–85.
- 4 Ryan M, McInerney D, Owens D, Collins P, Johnson A, Tomkin G. Diabetes and the Mediterranean diet: beneficial effects of oleic acid. *Quarterly Journal of Medicine* 2000; **93**: 85–91.
- 5 Ferrara L, Raimondi S, D'Episcopo L, Guida L, Dello Russo A, Marotta T. Olive oil and reduced need for antihypertensive medications. *Archives of Internal Medicine* 2000; **160**: 837–42.
- 6 Joshipura KJ, Ascherio A, Manson JE, Stampfer MJ, Rimm EB, Speizer FE, *et al.* Fruit and vegetable intake in relation to risk of ischemic stroke. *Journal of the American Medical Association* 1999; **282**: 1233–9.
- 7 Trichopoulou A, Georgiou E, Bassiakos Y, Lipworth L, Lagiou P, Proukakis C, *et al.* Energy intake and mono-unsaturated fat in relation to bone mineral density among women and men in Greece. *Preventive Medicine* 1997; **26**: 395–400.
- 8 Bosetti C, Gallus S, Trichopoulou A, Talamini R, Franceschi S, Negri E, *et al.* Influence of the Mediterranean diet on the risk of cancers of the upper aerodigestive tract. *Cancer Epidemiology, Biomarkers & Prevention* 2003; **12**: 1091–4.
- 9 Matelas A, Zampelas A, Stavrinou V, Wolinsky I. *The Mediterranean Diet: Constituents and Health Promotion*. Boca Raton, FL: CRC Press, 2000.
- 10 Serra-Majem L, Helsing E. Changing patterns of fat intake in Mediterranean countries. *European Journal of Clinical Nutrition* 1993; **47**: 1S–100S.
- 11 Willett W, Sacks F, Trichopoulou A, Drescher G, Ferro-Luzzi A, Helsing E, *et al.* Mediterranean diet pyramid: a cultural model for healthy eating. *American Journal of Clinical Nutrition* 1995; **61**: 1402S–6S.
- 12 WCRF. *Nutrition and the Prevention of Cancer: A Global Perspective*. Washington, DC: World Cancer Research Fund and American Institute for Cancer Prevention, 1997.
- 13 WHO. *Diet, Nutrition and the Prevention of Chronic Diseases*. Geneva: World Health Organization, 2003.
- 14 Hu F, Rimm E, Stampfer M. Prospective study of major dietary patterns and risk of coronary heart disease in men. *American Journal of Clinical Nutrition* 2000; **72**: 912–21.
- 15 Trichopoulos D, Lagiou P. Dietary patterns and mortality. *British Journal of Nutrition* 2001; **85**: 133–4.
- 16 Hu F. Dietary pattern analysis: a new direction in nutritional epidemiology. *Current Opinion in Lipidology* 2002; **13**: 3–9.
- 17 Trichopoulou A, Kouris-Blazos A, Wahlqvist M, Gnardellis D, Lagiou P, Polychronopoulos E, *et al.* Diet and overall survival in elderly people. *British Medical Journal* 1995; **311**: 1457–60.
- 18 Costacou T, Bamia C, Ferrari P, Riboli E, Trichopoulos D, Trichopoulou A. Tracing the Mediterranean diet through principal components and cluster analyses in the Greek population. *European Journal of Clinical Nutrition* 2003; **57**: 1378–85.
- 19 Osler M, Schroll M. Diet and mortality in a cohort of elderly people in a north European community. *International Journal of Epidemiology* 1997; **26**: 155–9.
- 20 Trichopoulou A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. *New England Journal of Medicine* 2003; **348**: 2599–608.
- 21 Psaltopoulou T, Naska A, Orfanos P, Trichopoulos D, Mountokalakis T, Trichopoulou A. Olive oil, the Mediterranean diet, and arterial blood pressure: the Greek European Prospective Investigation into Cancer and Nutrition (EPIC) Study. *American Journal of Clinical Nutrition* 2004; **80**: 1012–18.
- 22 Trichopoulou A, Bamia C, Trichopoulos D. Mediterranean diet and survival among patients with coronary heart disease in Greece. *Archives of Internal Medicine* 2005; **165**: 929–35.
- 23 Kouris-Blazos A, Gnardellis C, Wahlqvist M, Trichopoulos D, Lukito W, Trichopoulou A. Are the advantages of the Mediterranean diet transferable to other populations? A cohort study in Melbourne, Australia. *British Journal of Nutrition* 1999; **82**: 57–61.
- 24 Woo J, Woo K, Leung S. The Mediterranean score of dietary habits in Chinese populations in four different geographical areas. *European Journal of Clinical Nutrition* 2001; **55**: 215–20.
- 25 Lasheras C, Fernandez S, Patterson. Mediterranean diet and age with respect to overall survival in institutionalised, non-smoking elderly people. *American Journal of Clinical Nutrition* 2000; **71**: 987–92.
- 26 González C, Argilaga S, Agudo A, Amiano P, Barricarte A, Beguiristain JM, *et al.* Sociodemographic differences in adherence to the Mediterranean dietary pattern in Spanish populations. *Gaceta Sanitaria* 2002; **16**: 214–21.
- 27 Schröder H, Marrugat J, Vila J, Covas M, Elosua R. Adherence to the traditional mediterranean diet is inversely associated with body mass index and obesity in Spanish population. *Journal of Nutrition* 2004; **134**: 3355–61.
- 28 Knuops K, De Groot L, Kromhout D, Perrin A, Moreiras-Varela O, Menotti A, *et al.* Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women. *Journal of the American Medical Association* 2004; **292**: 1433–9.
- 29 Van Staveren W, De Groot L, Haveman-Nies A. The SENECA study: potentials and problems in relating diet to survival over 10 years. *Public Health Nutrition* 2002; **5**: 901–5.
- 30 Trichopoulou A, Orfanos P, Norat T, Bueno-de-Mesquita B, Oake MC, Peeters PH, *et al.* Modified Mediterranean diet and survival: EPIC-elderly prospective cohort study. *British Medical Journal* 2005; **330**: 991–7.
- 31 Trichopoulos D, Lagiou P. Mediterranean diet and overall mortality differences in the European Union. *Public Health Nutrition* 2004; **7**: 949–51.
- 32 Bilenko N, Fraser D, Vardi H, Shai I, Shahar D. Mediterranean diet and cardiovascular diseases in an Israeli population. *Preventive Medicine* 2005; **40**: 299–305.
- 33 Martínez-González M, Fernández-Jarne E, Serrano-Martínez M, Martí A, Martínez J, Martín-Moreno J. Mediterranean diet and reduction in the risk of a first acute myocardial infarction: an operational healthy dietary score. *European Journal of Nutrition* 2002; **41**: 153–60.
- 34 Serra-Majem L, Ribas L, Ngo J, Ortega R, García A, Pérez-Rodrigo C, *et al.* Food youth and the Mediterranean diet in Spain. Development of KIDMED, Mediterranean Diet Quality Index in children and adolescents. *Public Health Nutrition* 2004; **7**: 931–5.
- 35 Goulet J, Lamarche B, Nadeau G, Lemieux S. Effect of a nutritional intervention promoting the Mediterranean food pattern on plasma lipids, lipoproteins and body weight in healthy French-Canadian women. *Atherosclerosis* 2003; **170**: 115–24.
- 36 Panagiotakos D, Pitsavos C, Polychronopoulos E, Chrysohoo C, Zampelas A, Trichopoulou A. Can a Mediterranean diet moderate the development and clinical progression of coronary heart disease? A systematic review. *Medical Science Monitor* 2004; **10**: 193–8.
- 37 Patterson RE, Haines PS, Popkin BM. Diet quality index: capturing a multidimensional behaviour. *Journal of the American Dietetic Association* 1994; **94**: 57–64.
- 38 Gerber M. Qualitative methods to evaluate Mediterranean diet in adults. *Public Health Nutrition* 2006; **9** (1A): 147–51.
- 39 Ciccarone E, Di Castelnuovo A, Salcuni M, Siani A, Giacco A, Donati M, *et al.* A high-score Mediterranean dietary pattern is associated with a reduced risk of peripheral arterial disease

- in Italian patients with type 2 diabetes. *Journal of Thrombosis and Haemostasis* 2003; **1**: 1744–52.
- 40 Sanchez-Villegas A, Martinez J, de Irala J, Martínez-González M. Determinants of the adherence to an 'a priori' defined Mediterranean dietary pattern. *European Journal of Nutrition* 2002; **41**: 249–57.
 - 41 Tur J, Romaguera D, Pons A. Adherence to the Mediterranean dietary pattern among the population of the Balearic Islands. *British Journal of Nutrition* 2004; **92**: 341–6.
 - 42 Alberti-Fidanza A, Fidanza F, Chiuchiu M, Verducci G, Fruttini D. Dietary studies on two rural Italian population groups of the Seven Countries Study. 3. Trend of food and nutrient intake from 1960 to 1991. *European Journal of Clinical Nutrition* 1999; **53**: 854–60.
 - 43 Fuentes-Bol M. La dieta mediterránea: su evolución. In: Serra-Majem L, Ngo J, eds. *¿Qué es la Dieta Mediterránea? Fundación para el desarrollo de la Dieta Mediterránea*. Barcelona: Nexus Editions, 2002; 60–73.
 - 44 Kant A. Dietary patterns and health outcomes. *Journal of the American Dietetic Association* 2004; **104**: 615–36.
 - 45 Hu F. The Mediterranean diet and mortality – olive oil and beyond. *New England Journal of Medicine* 2003; **348**: 2595–6.
 - 46 Serra-Majem L, Roman B, Estruch R. Scientific evidence of interventions using the Mediterranean diet: a systematic review. *Nutrition Reviews* 2006; **64**: 531–52.
 - 47 Serra-Majem L, Trichopoulou A, Ngo J, Cervera P, García A, La Vecchia C, *et al.* Does the definition of the Mediterranean diet need to be updated? *Public Health Nutrition* 2004; **7**: 927–9.
 - 48 Trichopoulou A, Naska A, Antoniou A, Friel S, Trygg K, Turrini A. Vegetable and fruit: the evidence in their favour and the public health perspective. *International Journal for Vitamin and Nutrition Research* 2003; **73**: 63–9.
 - 49 Jacques P, Tucker K. Are dietary patterns useful for understanding the role of diet in chronic disease? *European Journal of Clinical Nutrition* 2001; **73**: 1–2.
 - 50 Alberti-Fidanza A, Findanza F. Mediterranean Adequacy Index of Italian diets. *Public Health Nutrition* 2004; **7**: 937–41.