

Effects of revision of Japanese food composition tables on estimation of nutrient intakes, with reference to age-dependent differences

Naoko Matsuda-Inoguchi¹, Shinichiro Shimbo¹, Haruo Nakatsuka², Takao Watanabe³, Kae Higashikawa⁴ and Masayuki Ikeda^{4,*}

¹Department of Food and Nutrition, Kyoto Women's University, Kyoto, Japan: ²Miyagi University, Taiwa-cho, Japan:

³Miyagi University of Education, Sendai, Japan: ⁴Kyoto Industrial Health Association, 67 Nishinokyo-Kitatsuboicho, Nakagyo-ku, Kyoto 604-8472, Japan

Submitted 1 October 2003: Accepted 3 February 2004

Abstract

Objective: To identify effects of revision of the Japanese food composition tables from the fourth version to the fifth version on nutrient intake estimation.

Design: A database on 783 samples of 24-hour food duplicate portions was re-visited. Nutrients in the duplicate portions were estimated by use of the fourth and fifth versions of the Japanese food composition tables in parallel, together with supplemental use of other databases. The two sets of estimates were subjected to comparison.

Setting: The sample collection was conducted at 31 sites all over Japan.

Subjects: The sample donors were 783 women aged 20–78 years.

Results: Compared with the estimates by use of the fourth version of the tables, the estimates by the fifth version were substantially higher for intakes of energy, carbohydrate, dietary fibre, vitamin A and niacin, and lower for iron intake. The increase in carbohydrate intake estimates was more evident in older women than in young women, whereas the decrease in the intake estimation of iron and the increase in that of dietary fibre were more marked in young women than in older women.

Conclusion: The recent revision of food composition tables in Japan induced substantial changes in the estimation of nutrient intakes, i.e. an increase in energy, carbohydrate, dietary fibre, vitamin A and niacin, and a decrease in iron. The extent of the changes varied depending on age.

Keywords
Adult women
Age
Carbohydrate
Comparison between versions
Energy
Food composition tables
Iron
Nutrient intakes

Adequate intake of nutrients is one of the essentials for healthy life. For the estimation of nutrient intakes, food composition tables are an important tool having wide application in both clinical and public health practice¹. In addition, food composition tables are often a common tool when several methods of nutrient intake estimation are compared^{2–6}. As the tables should be reliable in estimating nutrient intakes, the tables themselves are also subject to validation studies^{7–9}.

In Japan, the food composition tables were revised recently to produce the fifth version of the tables¹⁰ almost 20 years after publication of the previous version (the fourth version¹¹) in 1982. Revision of the tables has been practised several times in the history of the National Nutritional Survey^{12,13} in Japan: the first version of the food composition tables was used from 1948 to 1954, the second version from 1955 to 1963, and the third version from 1964 to 1970. After a period of atypical use of categorised food group tables, the fourth version with several supplemental publications enjoyed a long period

of application from 1988 to 2001. Regardless of these changes in the tables used, the differences between the estimates by the incoming new version and those by the previous version of the tables were generally small, i.e. less than 4%¹⁴.

The latest (i.e. fifth) version¹⁰ covers more food items (1882 items, compared with 1621 items in the fourth version¹¹) with more detailed classification. It was thought possible, therefore, that the revision might influence the estimation of nutrient intakes. Accordingly, this group conducted a pilot study on 71 girl students to compare their nutrient intakes as estimated by use of the fourth version of the food composition tables with that estimated by the fifth version. The study indicated that estimation by use of the fifth version gave increases in energy (7%), carbohydrate (8%) and vitamin A (20%) intakes, and a decrease in iron intake (–13%)¹⁴.

It is known, however, that the food habits of girl students may be biased and not typical of those

of women^{15–19}. The present analyses were initiated, therefore, to examine if the conclusions obtained in that study of girl students are applicable also to women in general, by expanding the study population both in number and age range of the subjects.

Methods

Study population

In 1991 to 1998, 783 adult women at 31 sites in eight regions of Japan (for locations, see Fig. 1) offered food duplicate portion samples on a voluntary basis^{14,20}. The duplicate portion database thus established^{14,20} was re-visited for the present analysis. The ages of the participants (mostly housewives, except for young women) were distributed over a wide range from 20 to 78 years, with a mean \pm standard deviation (SD) of 49.8 ± 12.8 years. The majority of women were in their 50s (264 cases), followed by those in their 40s (198 cases) and women their 60s (157 cases)²⁰.

Collection of food duplicate portion samples

Each participant provided her informed consent. She was carefully instructed to cook an ordinary everyday meal (i.e. no special dishes)¹⁴. Subsequently, she offered 24-hour food duplicate portion samples²¹ together with food menus (the names of each dish) of the day. The procedures for recording the food items have been described previously in detail¹⁴; in short, each food item in the sample was separated manually under supervision of a veteran nutritionist, and the weight of each food item was recorded with reference to the menu.

Estimation of nutrient intakes

For the fourth version-based estimation of daily nutrient intakes, the nutritionist coded all items by use of the fourth

version of the standard tables of food composition¹¹, the standard tables of dietary fibre²², the standard tables of fatty acids, cholesterol and vitamin E²³, and the food composition tables on commercial pre-cooked foods²⁴, in combination. In the case that a code for a food item of concern was not listed in the tables, an available item of close botanical or zoological relevance was taken as a surrogate²⁵. Nutrient intake was first estimated from the food weight record and the code number for each food item, and the estimates were then summed up for daily intake on an individual basis as well as on a group basis.

In the next step, to make the fifth version-based estimation of daily nutrient intakes, the same nutritionist coded all items again, but with reference to the fifth version of the tables¹⁰ (as amended after corrections given on the Resources Council home page on 28 March 2001), together with the food composition tables on commercial pre-cooked foods²⁴. Although more food codes were available in the fifth version of the tables as described above, use of the pre-cooked food tables²⁴ still remained necessary.

Data analysis

Comparisons were made for energy, three macronutrients (including carbohydrate), dietary fibre, five minerals, five vitamins, fatty acids and cholesterol, as listed in Table 1. A normal distribution was assumed for all nutrient intakes, so that the distribution was expressed as mean (arithmetic) \pm SD (arithmetic). Vitamin A contents were expressed in terms of international units (IU) in both the fourth version of the tables¹¹ and the pre-cooked food tables²⁴, whereas it was in retinol equivalents (RE) in the fifth version¹⁰. For conversion of the value in IU to that in RE, 10 IU was thought to be equivalent to 3 μ g RE²⁶.




Fig. 1 Locations of survey sites, grouped by administrative region. Each solid circle indicates a survey site where food duplicate portion samples were collected

Table 1 Recommended dietary allowances for Japanese women by age range

Nutrient (units day ⁻¹)	Age range (years)			
	18–29	30–49	50–69	70 and over
Energy (kcal)	2050	2000	1900	1700
Protein (g)	55	55	55	55
Lipid (g)	38–57	38–57	38–57	38–57
Dietary fibre (g)	17–20.5	17–20.5	17–20.5	17–20.5
Minerals				
Ca (mg)	600	600	600	600
P (mg)	700	700	700	700
Fe (mg)	12	12†	12†	10
K (mg)	2000	2000	2000	2000
NaCl* (g)	< 10	< 10	< 10	< 10
Vitamins				
Vitamin A (µg RE†)	540–1500	540–1500	540–1500	540–1500
Vitamin B ₁ (mg)	0.8	0.8	0.8	0.8
Vitamin B ₂ (mg)	1.0	1.0	1.0	1.0
Vitamin C (mg)	100	100	100	100
Niacin (mg)	13–30	13–30	13–30	13–30

The sixth recommended daily allowances (RDAs) for women with moderate physical exercise²⁶.

* Assuming that all Na is attributable to sodium chloride (NaCl); the value is not an RDA but the recommended upper limit of acceptance.

† Retinol equivalent; 300 µg RE = 1000 IU.

‡ 10 mg day⁻¹ after the menopause.

Table 2 Intakes of macro- and micronutrients

Nutrient (units day ⁻¹)	Intake (mean ± standard deviation) as estimated with		Difference in mean†	P-value‡	% change§
	Fourth version	Fifth version			
Energy (kcal)	1818 ± 422	1945 ± 447	127	**	7.1
Protein (g)	67.0 ± 18.4	66.8 ± 18.6	-0.2	**	-0.4
Carbohydrate (g)	277 ± 73	306 ± 79	29	**	10.4
Lipid (g)	47.1 ± 17.3	47.9 ± 18.3	0.8	**	1.6
Dietary fibre (g)	13.0 ± 6.0	17.1 ± 7.0	4.1	**	38.1
Water-soluble (g)	2.58 ± 1.50	3.48 ± 1.46	0.90	**	62.7
Water-insoluble (g)	10 ± 4.45	12.15 ± 5.05	2.11	**	25.6
Minerals					
Ca (mg)	600 ± 249	610 ± 253	10	**	2.8
P (mg)	951 ± 286	1020 ± 304	69	**	7.5
Fe (mg)	10.0 ± 3.2	8.9 ± 3.0	-1.1	**	-10.8
K (mg)	2583 ± 959	2666 ± 961	83	**	3.4
NaCl¶ (g)	10.8 ± 4.0	10.5 ± 3.8	-0.3	**	-0.6
Vitamins					
Vitamin A (µg RE)	830 ± 826	1066 ± 852	236	**	40.0
Vitamin B ₁ (mg)	0.83 ± 0.31	0.80 ± 0.32	-0.03	**	-3.3
Vitamin B ₂ (mg)	1.23 ± 0.43	1.21 ± 0.44	-0.02	**	-0.9
Vitamin C (mg)	127 ± 73.3	126.2 ± 71.0	-0.5	NS	1.4
Niacin (mg)	12.3 ± 6.1	13.7 ± 7.6	1.5	**	11.6
Fatty acids					
Saturated (g)	9.9 ± 5.0	12.4 ± 6.1	2.5	**	34.0
Monounsaturated (g)	12.7 ± 5.9	14.8 ± 6.7	2.0	**	20.9
Polyunsaturated (g)	10.4 ± 4.8	11.7 ± 5.1	1.3	**	14.7
Cholesterol (mg)	230 ± 161	281 ± 161	51.6	**	73.2

† The difference is calculated as the value obtained from the 5th version minus the value obtained from the fourth version.

‡ The statistical significance of the difference: **, $P \leq 0.01$; *, $P \leq 0.05$; NS, $P > 0.05$, as examined by Student's paired *t*-test ($n = 783$).

§ The average % change for each item is shown; the percentage change – i.e. (the difference divided by the value obtained from the fourth version) × 100 – was first calculated on an individual basis, and then the average of the percentages was calculated.

¶ Assuming that all Na is attributable to sodium chloride (NaCl).

|| Retinol equivalent; 300 µg RE = 1000 IU.

All data were analysed using SPSS version 10 (SPSS Inc., Chicago, IL, USA). Student's paired *t*-test was employed for detection of a significant difference between the two estimates of nutrient intakes, one based on the fourth version of the tables and the other on the fifth. One-way analysis of variance and the multiple comparison test

(Tukey–Kramer) were employed to detect possible significant differences among means of various age groups. Nationally recommended dietary allowances (RDA)²⁶ were employed for evaluation of the estimated intakes; the RDA values for adult non-pregnant and non-lactating women with moderate physical exercise are cited in Table 1.

+++, $F \leq 0.01$; +, $F \leq 0.05$; for difference from the value of the 40–49-year-old group, by multiple comparison test (Tukey–Kramer).
 ++, $F \leq 0.01$; +, $F \leq 0.05$; for difference from the value of the 30–39-year-old group, by multiple comparison test (Tukey–Kramer).
 +, $F \leq 0.05$; for difference from the value of the 20–29-year-old group, by multiple comparison test (Tukey–Kramer).
 ., $F \leq 0.05$; for difference from the value of the 10–19-year-old group, by multiple comparison test (Tukey–Kramer).

Table 4 Differences in the intakes of macro- and micronutrients by food group

Food group*	Energy (kcal)		Carbohydrate (g)		Dietary fibre (g)		Iron (µg)		NaCl (g)		Niacin (mg)		Vitamin A (µg)†	
	5th‡	Diff.§	5th‡	Diff.§	5th‡	Diff.§	5th‡	Diff.§	5th‡	Diff.§	5th‡	Diff.§	5th‡	Diff.§
1. Cereals	860.7	85.6	185.3	22.7	2.65	0.02	847	-53	0.45	0.05	1.38	-0.28	0.2	-0.1
2. Potatoes, etc.	46.7	-0.4	11.0	0.4	1.24	0.34	280	-32	0.00	-0.01	0.37	-0.03	0.5	0.5
3. Sugars, etc.	43.8	0.1	11.3	0.0	0.00	0.00	28	-10	0.00	0.00	0.00	0.00	0.0	0.0
4. Confectioneries	92.8	-5.7	16.9	-0.7	0.52	0.52	266	10	0.11	-0.01	0.15	0.05	8.1	2.6
5. Oils and fats	71.9	7.1	0.0	0.0	0.00	0.00	1	1	0.02	0.01	0.00	0.00	16.0	3.5
6. Nuts and seeds	15.2	0.1	0.9	0.0	0.26	0.21	125	-1	0.00	0.00	0.24	0.00	0.1	0.0
7. Pulses	86.2	-31.9	3.9	-3.0	1.55	-0.46	1161	-758	0.02	-1.80	0.17	-0.23	0.0	0.0
8. Fish and shellfish	118.1	6.2	1.8	0.1	0.00	0.00	923	-263	0.86	-0.46	4.38	0.59	32.9	5.5
9. Meats	93.4	2.3	0.2	0.0	0.00	0.00	416	-143	0.18	-0.03	1.93	0.11	105.7	-12.5
10. Eggs	51.3	-2.2	0.1	-0.2	0.00	0.00	618	27	0.12	0.02	0.04	0.00	50.4	-13.2
11. Milks	106.5	7.4	8.1	0.2	0.00	0.00	34	-113	0.21	-0.03	0.15	0.01	55.9	3.5
12. Vegetables	70.0	0.5	15.2	2.2	5.80	1.58	1241	-468	0.61	-0.28	1.17	0.13	596.4	128.3
13. Fruits	89.2	3.2	23.4	1.6	2.21	-0.31	272	-1	0.25	0.02	0.42	0.01	128.2	110.1
14. Fungi	2.5	2.5	0.8	0.2	0.51	0.24	57	-19	0.00	0.00	0.45	-0.02	0.0	0.0
15. Algae	6.3	6.3	2.2	-0.2	1.39	0.99	507	-65	0.38	0.02	0.18	0.02	56.0	11.8
16. Beverages	31.8	11.0	4.4	0.5	0.01	0.01	717	312	0.02	-0.01	1.97	0.73	0.2	-0.1
17. Seasonings, etc.	81.1	37.9	8.2	5.1	0.87	0.85	1065	463	6.88	2.26	0.64	0.37	2.6	0.7
18. Pre-cooked foods	9.9	0.8	0.9	-0.1	0.00	0.00	52	-4	0.06	-0.02	0.06	-0.01	2.2	-0.7
19. Commercial pre-cooked foods	67.5	-4.0	10.9	-0.2	0.07	0.07	280	32	0.37	0.02	0.03	0.02	10.1	-4.3

* Classification after Resources Council¹⁰, except for Group 19 which is a new addition.

† In retinol equivalents.

‡ The estimates by use of the fifth version of the tables.

§ The difference between estimates from the fifth version of the tables and the fourth version.

Food groups that induced an increase or a decrease in estimations

To identify the food groups that played major roles in inducing these differences, food items from each woman were classified into 19 groups in accordance with the 18 groups in the food composition tables¹⁰ together with a supplemental group (Group 19) for commercially available pre-cooked foods. The average (for women of all ages) estimates from the fifth version, the difference between the two estimates (i.e. the value from the fifth version minus the value from the fourth version) and the percentage change (the difference divided by the value from the fourth version in %) of each food group, respectively, were calculated for energy, carbohydrate, dietary fibre (water-soluble and -insoluble), iron, NaCl and niacin (Table 4).

The calculation for energy showed (Table 4) that the majority of the difference, i.e. 85.6 kcal out of 127 kcal (or 67%), was attributable to cereals. The same also applied to carbohydrate, as cereals were the major sources of carbohydrates. Comparison of the two composition tables, the fourth¹¹ and the fifth versions¹⁰, made it clear that this was due to re-evaluation of the energy per unit weight of boiled rice^{10,11} in reflection of a reduced estimation of the water content in boiled rice after shifts in the liking of people. The reduction of energy, iron and NaCl in pulses and the increase in seasonings were due to classification changes of *miso* paste (a fermentation product of soy beans, salt and rice (or wheat)²⁷) from the pulses group in the fourth version¹¹ to the seasoning group in the fifth version¹⁰. The increase in dietary fibre was primarily due to re-evaluation (an increase) of the water-insoluble fibre in some vegetables (e.g. pumpkin) in the fifth version¹⁰.

In the case of iron intake estimation, both the major sources and the food groups that induced the differences were various. Nevertheless, it is worthy to note that the major reduction came from a re-evaluation of iron losses from vegetables during the cooking process. For example, iron in spinach after boiling was revised from 2.0 mg/100 g in the fourth version¹¹ to 0.9 mg/100 g in the fifth version¹⁰. The increase in niacin was mostly due to increased evaluation of the niacin content in some types of sea fish and several types of soft drink such as Japanese green tea¹⁰. With regard to vitamin A, the increase was attributable to re-evaluation of the content in plant foods such as pumpkin, spinach and mandarins¹⁰.

Discussion

The present analysis makes it clear that revision of the food composition tables from the fourth to the fifth version induced substantial increases in the estimated intakes of energy, carbohydrate and vitamin A, and a decrease in the estimation of iron intake. The observation is in close

agreement with the results of a previous study on girl students¹⁴, suggesting that the applicability of the conclusion is not limited to young women, but can be extended to adult Japanese women in general. Additional findings from the present analysis are increases in the estimates of dietary fibre, niacin, fatty acids and cholesterol, possibly due to the 10-fold increase in the number of women studied, from 71 cases in the previous study¹⁴ to 783 cases in the present analysis.

The meaning of these observations should be evaluated at least from two viewpoints. One is the evaluation with reference to the RDA²⁵ as listed in Table 1. The comparison on energy suggests that the estimates (1818 or 1945 kcal day⁻¹; Table 2) are most probably not in excess of (although very close to) the RDA for energy intake (1700–2050 kcal day⁻¹; Table 1), irrespective of the version of the food composition tables. Nevertheless, elevated risk of obesity-associated adverse health effects such as diabetes, hyperlipidaemia and hypertension has been observed for Japanese populations at an even lower grade of obesity than for American and European populations²⁸. Thus, more attention is necessary for the prevention of excess weight gain. A simple calculation suggests that an excess intake of 127 kcal day⁻¹ (Table 2) may result in weight gain of 15.88 g day⁻¹ [= (127 kcal day⁻¹)/(8 kcal/g fat)] or about 5.8 kg (= 15.88 g day⁻¹ × 365 days) per year, assuming that the energy is converted to fat and stored in the body. Thus, an accurate estimation of total energy intake is apparently desirable.

Intakes of vitamin A and niacin (Table 2) are either sufficient or barely so when compared with the corresponding RDA values (Table 1). In the case of iron intake, however, the 10% reduction in estimated intake may need attention because the current level of intake, 8.9 mg day⁻¹ (the estimation based on the fifth version; Table 2), does not meet the recommendation (12 mg day⁻¹; Table 1). It was further made clear (Table 3) through the present analysis that the effects of re-evaluation of iron intake were greater in young women than in older women. Insufficient iron intake especially among young women has been a problem elsewhere in the world as well as in Japan^{14,18,19,29–34}. In this connection, it has been noticed that estimations based on food composition tables tend to give a larger iron intake than the results of inductively coupled plasma–mass spectrometry analysis⁸ of food duplicate portions, in support of the present observation.

In contrast, results for the estimation of NaCl intake by use of the two versions agreed well with each other (Table 2). It was previously observed that the estimates of salt intake based on the tables agree well with the results of instrumental analysis⁷. Nevertheless, NaCl intake by young women (and not of older women) as estimated by use of the fifth version of the tables was greater by 6% than the estimate by the fourth version of

the tables (Table 3). The observation suggests the possibility that the NaCl intake of younger people has been underestimated for years, which is apparently not favourable for the prevention of NaCl-induced hypertension in later life.

With regard to fatty acids, the ratio of S:M:P = 3:4:3 has been recommended for intakes of saturated (S), mono-unsaturated (M) and polyunsaturated (P) fatty acids, but no numerical values were given²⁶ for quantitative evaluation. Thus, no further evaluation was made on fatty acids in the present study, as available information on recommended intake is still insufficient. It should be added that the current S:M:P ratio of 10:13:10 or 12:15:12 (Table 2) is close to the recommendation described above²⁶. With regard to cholesterol, a reduction in intake to less than 300 mg day⁻¹ is recommended for those with a tendency to hypercholesterolaemia²⁶, but this recommendation may not necessarily be applicable to general healthy populations as was the survey population of the present study.

The extent of the effects of the revision should also be evaluated from a chronological viewpoint. The last five annual (i.e. from 1996 to 2000) reports from the National Nutritional Survey in Japan gave nutrient intake values for women separately from the values for men¹³. The estimations were based on the fourth version of the food composition tables. Linear regression analyses of the reported values, taking the survey years from 1996 to 2000

(Fig. 2) on the horizontal axis and the nutrient intakes on the vertical axis, gives:

$$Y_1 = 31565 - 14.9X \quad (r = -0.981, \quad P < 0.01) \text{ for energy,}$$

$$Y_2 = 4644 - 2.2X \quad (r = -0.996, \quad P < 0.01) \text{ for carbohydrate,}$$

$$Y_3 = 260.7 - 0.10X \quad (r = -0.870, \quad P > 0.10) \text{ for iron, and}$$

$$Y_4 = 351.7 - 0.17X \quad (r = -0.981, \quad P < 0.01) \text{ for NaCl,}$$

where X is the year of the survey, and Y_1 , Y_2 , Y_3 and Y_4 are daily intake of energy (kcal day⁻¹), carbohydrate (g day⁻¹), iron (mg day⁻¹) and NaCl (g day⁻¹), respectively. Division of the fifth version minus the fourth version difference in the estimates (Table 2) by the corresponding slope of the regression line gives -8.5, -4.8, 10.9 and 1.8 years for the four items, respectively, as the time in years necessary to induce the changes based on the current chronological trends in nutrient intakes; the plus and minus signs indicate the future and the past, respectively. The longest time is about 11 years for iron followed by -9 years for energy, which suggests that revision of the food composition tables may induce a substantially large artificial gap in the time-dependent trends of energy and iron (as the major source of energy is carbohydrate, the implication of the values for energy and carbohydrate should be the same). The estimates for the year 2001¹³, which were based on the fifth version of the




Fig. 2 Chronological trends observed in the Japanese National Nutritional Survey for (a) energy, (b) carbohydrate, (c) iron and (d) NaCl. Solid circles show the values for the years 1996 to 2000, which were based on the fourth version of the tables, and open squares show the values for the year 2001, which were based on the fifth version of the tables¹³. The line is a calculated regression line based on the data shown by the five solid circles

tables¹⁰, are also plotted in Fig. 2 to show the deviation from the continuity observed in the National Nutritional Survey results. Whereas food composition tables should be revised to keep up with the latest developments in science¹, it is also important to pay attention to maintain continuity in the evaluation of time trends.

Conclusions

The revision of the food composition tables from the fourth version to the fifth version induced substantial increases in intake estimation for energy, carbohydrate, dietary fibre, vitamin A and niacin, and decreases in iron. The changes were age-dependent, such that the increase in carbohydrate intake estimation was more marked in older women, whereas the increase in intake of dietary fibre and the decrease in intake of iron were more evident in young women. The importance of continuity in food intake evaluation was stressed in chronological analysis.

Acknowledgements

This work was supported in part by the 1999 Kyoto City Research Fund for Promotion of Health of Citizens to S.S., T.W. and M.I., and by sponsorship by the Ministry of Education, Science, Sports and Culture, the Government of Japan. The authors are grateful to Miss F Ohashi of Kyoto Industrial Health Association for her skilful PC operation.

References

- 1 Dwyer JT. Future directions in food composition studies. *Journal of Nutrition* 1994; **124**(Suppl. 9): 1783S–8S.
- 2 Clark AJ, Mossholder S. Sodium and potassium intake measurements; dietary methodology problems. *American Journal of Clinical Nutrition* 1986; **43**: 470–6.
- 3 Knuiman JT, Rasanen L, Ahola M, West CE, van den Snoek L. The relative validity of reports of food intake of Dutch and Finnish boys aged 8 and 9 years. *Journal of the American Dietetic Association* 1987; **87**: 303–7.
- 4 Nagata Y, Suzue R. Comparison of young women's salt intake between Japan and England: consideration on different measurement methods in national nutrition surveys. *Journal of Nutritional Science and Vitaminology* 1996; **42**: 19–26.
- 5 Tsubono Y, Sasaki S, Kobayashi M, Akabane M, Tsugane S. Food composition and empirical weight methods in predicting nutrient intakes from food frequency questionnaire. *Annals of Epidemiology* 2001; **11**: 213–8.
- 6 Sasaki S, Matsumura Y, Ishihara J, Tsugane S. Validity of a self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study Cohort I to assess dietary fiber intake; comparison with dietary records. *Journal of Epidemiology* 2003; **13**(Suppl. 1): S106–14.
- 7 Shimbo S, Zhang Z-W, Miyake K, Watanabe T, Nakatsuka H, Matsuda-Inoguchi N, *et al.* Estimates of mineral intakes using food composition tables vs measures by inductively-coupled plasma mass spectrometry: Part 2. Sodium, potassium, magnesium, copper and zinc. *European Journal of Clinical Nutrition* 1999; **53**: 233–8.
- 8 Zhang Z-W, Shimbo S, Miyake K, Watanabe T, Nakatsuka H, Matsuda-Inoguchi N, *et al.* Estimates of mineral intakes using food composition tables vs measures by inductively-coupled plasma mass spectrometry: Part 1. Calcium, phosphorus and iron. *European Journal of Clinical Nutrition* 1999; **53**: 226–32.
- 9 Kim E-S, Ko Y-S, Kim J, Matsuda-Inoguchi N, Nakatsuka H, Watanabe T, *et al.* Food composition table-based estimation of energy and major nutrient intake in comparison with chemical analysis: a validation study in Korea. *Toboku Journal of Experimental Medicine* 2003; **200**: 7–15.
- 10 Resource Council, Science and Technology Agency, the Government of Japan. *Standard Tables of Food Composition in Japan*, 5th revised version. Tokyo: Ministry of Finance Printing Bureau, 2000 [in Japanese with English translation].
- 11 Resource Council, Science and Technology Agency, the Government of Japan. *Standard Tables of Food Composition in Japan*, 4th revised version. Tokyo: Ministry of Finance Printing Bureau, 1982 [in Japanese with English translation].
- 12 Ministry of Health and Welfare, the Government of Japan. *Review on Public Nutrition in Japan in 1955 to 1988*. Tokyo: Dai-ichi Shuppan Press, 2000 [in Japanese].
- 13 Ministry of Health and Welfare, the Government of Japan. *National Nutritional Survey in Japan, 1989–2001* [annual reports]. Tokyo: Dai-ichi Shuppan Press, 1991–2003 [in Japanese].
- 14 Matsuda-Inoguchi N, Nakatsuka H, Watanabe T, Shimbo S, Higashikawa K, Ikeda M. Estimation of nutrient intake by the new version of Japanese food composition tables in comparison with that by the previous version. *Toboku Journal of Experimental Medicine* 2001; **194**: 229–39.
- 15 Ikeda J, Asano H, Nagata H. Dietary life of female college students (Part 1). Survey on nutrition awareness and their mode of living affecting their nutrient intake [in Japanese with English abstract]. *Japanese Journal of Nutrition* 1983; **41**: 103–16.
- 16 Shiraki M, Iwasaki N. Effects of a habit of skipping meals on dietary life of college students [in Japanese with English abstract]. *Japanese Journal of Nutrition* 1986; **44**: 257–65.
- 17 Someya R, Negishi Y, Mizuno K, Muto S. Dietary life of female college students [in Japanese with English abstract]. *Japanese Journal of Nutrition* 1989; **47**: 251–8.
- 18 Cai MQ, Yan WY. Study on iron nutritional status in adolescence. *Biomedical and Environmental Sciences* 1990; **3**: 113–9.
- 19 Hendricks KM, Herbold NH. Diet, activity, and other health-related behaviors in college women. *Nutrition Reviews* 1998; **56**: 65–75.
- 20 Watanabe T, Zhang Z-W, Moon C-S, Shimbo S, Nakatsuka H, Matsuda-Inoguchi N, *et al.* Cadmium exposure of women in general populations in Japan during 1991–1997 compared with 1977–1981. *International Archives of Occupational and Environmental Health* 2000; **73**: 26–34.
- 21 World Health Organization (WHO). *Guidelines for the Study of Dietary Intakes of Chemical Contaminants*. WHO Offset Publication No. 87. Geneva: WHO, 1985.
- 22 Resource Council, Science and Technology Agency, the Government of Japan. *Standard Tables of Food Composition in Japan; Dietary Fibers*. Tokyo: Ministry of Finance Printing Bureau, 1992 [in Japanese with English translation].
- 23 Resource Council, Science and Technology Agency, the Government of Japan. *Standard Tables of Food Composition in Japan; Fatty Acids, Cholesterol and Vitamin E*. Tokyo: Ministry of Finance Printing Bureau, 1989 [in Japanese with English translation].
- 24 Kagawa Y. *Food Composition Tables for Commercial Precooked Food*. Tokyo: Kagawa University of Nutrition Press, 1995 [in Japanese].
- 25 Shimbo S, Hayase A, Murakami M, Hatai I, Higashikawa K,

- Moon C-S, *et al.* Use of a food composition database to estimate daily dietary intake of nutrient or trace elements in Japan, with reference to its limitation. *Food Additives and Contaminants* 1996; **13**: 775–86.
- 26 Ministry of Health and Welfare, the Government of Japan. *Recommended Dietary Allowances*, 6th version. Tokyo: Dai-ichi Shuppan Press, 1999 [in Japanese].
- 27 Watanabe T, Miyasaka M, Koizumi A, Ikeda M. Regional difference in sodium chloride content in home-made and store-bought preparations of miso paste. *Toboku Journal of Experimental Medicine* 1982; **137**: 305–13.
- 28 Obesity Criteria Committee, Japan Society for Study on Obesity. New concept of obesity and diagnostic criteria [in Japanese]. *Journal of the Japanese Society for the Study of Obesity* 2000; **6**: 18–28.
- 29 Shimbo S, Imai Y, Tominaga N, Gotoh T, Yokota M, Inoguchi N, *et al.* Insufficient calcium and iron intakes among general female population in Japan, with special reference to inter-regional differences. *Journal of Trace Elements in Medicine and Biology* 1996; **10**: 133–8.
- 30 Shimbo S, Moon C-S, Zhang Z-W, Watanabe T, Noor Hassim I, Romzi MA, *et al.* Nutritional evaluation of working Malay women in Kuala Lumpur as studied by total food duplicate method. *Toboku Journal of Experimental Medicine* 1996; **180**: 99–114.
- 31 Shimbo S, Moon C-S, Zhang Z-W, Watanabe T, Guo LL, Ma W-C, *et al.* Nutritional evaluation of Chinese working women in the city of Tainan, Taiwan. *Toboku Journal of Experimental Medicine* 1997; **181**: 339–52.
- 32 Moon C-M, Zhang Z-W, Imai M, Shimbo S, Watanabe T, Moon D-H, *et al.* Nutritional status of women in urban and rural areas in Korea as assayed by total food duplicate method. *Toboku Journal of Experimental Medicine* 1997; **181**: 245–65.
- 33 Nakatsuka H, Zhang Z-W, Agetano MG, Subida RD, Inoguchi N, Watanabe T, *et al.* Total food duplicate study on nutrient intake of working women in Manila, the Philippines. *Toboku Journal of Experimental Medicine* 1998; **184**: 189–205.
- 34 Matsuda-Inoguchi N, Shimbo S, Zhang Z-W, Srianujata S, Banjong O, Chitchumroonchokchai C, *et al.* Nutrient intake of working women in Bangkok, Thailand, as studied by total food duplicate method. *European Journal of Clinical Nutrition* 2000; **54**: 187–94.