

Received 5 September 2014
Accepted 22 September 2014

Edited by M. Zeller, Youngstown State University, USA

Keywords: crystal structure; coordination compound; ruthenium; stacking

CCDC reference: 1025408

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structure of (2,11-diaza[3.3](2,6)pyridinophane- $\kappa^4N,N',N'',N''')(1,6,7,12-tetraazaperylene- $\kappa^2N^1,N^{12})\text{ruthenium(II)}\text{ bis(hexafluoridophosphate)}\text{ acetonitrile 1.422-solvate}$$


Thomas Brietzke, Falko Otto Rottke, Alexandra Kelling, Uwe Schilde* and Hans-Jürgen Holdt

Universität Potsdam, Institut für Chemie, Anorganische Chemie, Karl-Liebknecht-Str. 24-25, D-14476 Potsdam, Germany. *Correspondence e-mail: us@chem.uni-potsdam.de

In the title compound, $[\text{Ru}(\text{C}_{14}\text{H}_{16}\text{N}_4)(\text{C}_{16}\text{H}_8\text{N}_4)](\text{PF}_6)_2 \cdot 1.422\text{CH}_3\text{CN}$, discrete dimers of complex cations, $[\text{Ru}(\text{L}-\text{N}_4\text{H}_2)\text{tape}]^{2+}$ are formed ($\text{L}-\text{N}_4\text{H}_2 = 2,11$ -diaza[3.3](2,6)pyridinophane; tape = 1,6,7,12-tetraazaperylene), held together by $\pi-\pi$ stacking interactions *via* the tape ligand moieties with a centroid-centroid distance of 3.49 (2) Å, assisted by hydrogen bonds between the non-coordinating tape ligand α,α' -diimine unit and the amine proton of a 2,11-diaza[3.3](2,6)-pyridinophane ligand of the opposite complex cation. The combination of these interactions leads to an unusual nearly face-to-face $\pi-\pi$ stacking mode. Additional weak C—H···N, C—H···F, N—H···F and P—F···π-ring (tape, py) (with F···centroid distances of 2.925–3.984 Å) interactions are found, leading to a three-dimensional architecture. The Ru^{II} atom is coordinated in a distorted octahedral geometry, particularly manifested by the $\text{N}_{\text{amine}}-\text{Ru}-\text{N}_{\text{amine}}$ angle of 153.79 (10)°. The counter-charge is provided by two hexafluoridophosphate anions and the asymmetric unit is completed by acetonitrile solvent molecules of crystallization. Disorder was observed for both the hexafluoridophosphate anions as well as the acetonitrile solvate molecules, with occupancies for the major moieties of 0.801 (6) for one of the PF₆ anions, and a shared occupancy of 0.9215 (17) for the second PF₆ anion and a partially occupied acetonitrile molecule. A second CH₃CN molecule is fully occupied, but 1:1 disordered across a crystallographic inversion center.


1. Chemical context

Heteroaromatic ligands with more than three fused rings are commonly called large-surface ligands. Such ligands have attracted attention due to their use as connecting building blocks for supramolecular assemblies. If large-surface ligands feature more than one ligand donor site, connection between neighboring complexes can be realized through normal metal coordination (Ishow *et al.*, 1998), but the large π system also allows for strong $\pi-\pi$ stacking interactions. (Kammer *et al.*, 2006; Gut *et al.*, 2002). In order to study the properties of ruthenium complexes containing large-surface ligands, we have recently reported an easy entry to such complexes (Brietzke, Mickler, Kelling, Schilde *et al.*, 2012). Therein, we formulated the advantages of the 2,11-dimethyl-2,11-diaza[3.3](2,6)-pyridinophane (L-N₄Me₂) macrocycle over bi-pyridine (bpy)-type ligands in saturating the coordination sphere of an octahedral ruthenium complex containing the large-surface ligand of interest. The microwave-assisted synthesis of the precursor [Ru(L-N₄Me₂)]²⁺, starting from [Ru(DMSO)₄Cl₂] and L-N₄Me₂, in an ethanolic solution finished within 30 min. It is not only fast, but also reproducible



OPEN ACCESS

with only few byproducts, and hence requires no labor-intensive workup. Moreover, using the C_{2v} symmetric macrocycle rather than bipyridine-type ligands avoids the formation of mono- and dinuclear complexes with multiple stereoisomeric forms (Brietzke, Mickler, Kelling & Holdt, 2012; Brietzke *et al.*, 2014). To test the applicability of our microwave-assisted synthetic strategy for use with other related macrocyclic ligands, we choose the unmethylated parent compound of L-N₄Me₂, 2,11-diaza[3.3](2,6)-pyridinophane (L-N₄H₂) as a new ligand for Ru^{II}. Herein, we present the structure of the complex [Ru(L-N₄H₂)tape](PF₆)₂ (tape = 1,6,7,12-tetraazaperylene), obtained as its acetonitrile solvate.



2. Structural commentary

Fig. 1 illustrates the molecular structure of the complex [Ru(L-N₄H₂)tape]²⁺ in [Ru(C₁₄H₁₆N₄)(C₁₆H₈N₄)](PF₆)₂·1.422CH₃CN. The Ru—N bond lengths formed by the tape ligand (Table 1) are very close to those reported earlier for [Ru(L-N₄Me₂)tape]²⁺ (Brietzke, Mickler, Kelling, Schilde *et al.*, 2012). The deviation of the N_{amine}—Ru—N_{amine} angle [153.79 (10) $^{\circ}$] from the idealized value of 180 $^{\circ}$ is slightly larger than for analogous ruthenium L-N₄Me₂ complexes




Figure 1

The molecular structure of [Ru(L-N₄H₂)tape]²⁺ in [Ru(L-N₄H₂)tape](PF₆)₂·1.422CH₃CN with the atomic numbering scheme and 30% probability displacement ellipsoids. Anions and solvent molecules are omitted for clarity.

Table 1
Selected bond lengths (\AA).

Ru1—N6	2.005 (2)	Ru1—N2	2.055 (2)
Ru1—N4	2.018 (2)	Ru1—N5	2.132 (3)
Ru1—N1	2.045 (2)	Ru1—N3	2.135 (3)

[155.46 (9)–155.93 (17) $^{\circ}$; Brietzke, Mickler, Kelling, Schilde *et al.*, 2012].

3. Supramolecular features

In the crystal structure, the cations form discrete centrosymmetric dimers, similar to those seen previously in mononuclear ruthenium–tape complexes. The dimers are held together by π – π stacking interactions *via* the planar tetraazaperylene units, with a typical interplanar distance of 3.39 \AA . For the





Figure 2

(a) Illustration of the asymmetric unit rendering the disorder of the hexafluoridophosphate anions and acetonitrile solvate molecules (see *Refinement* section for details). An additional π – π stacked [Ru(L-N₄H₂)tape]²⁺ cation demonstrates, due to the view along the normal of the tape ligand's r.m.s. plane, the nearly face-to-face π – π stacking motif between the tape ligand moieties. The atomic numbering is shown for the anions and solvent molecules as well as for the ruthenium atoms. Hydrogen atoms are omitted for clarity. [Symmetry codes: (ii) 1 – x , – y , 1 – z , (ix) 1 – x , 1 – y , – z .] (b) A side view of the dimer formed by two [Ru(L-N₄H₂)tape]²⁺ in [Ru(L-N₄H₂)tape](PF₆)₂·1.422CH₃CN, featuring the stacking interactions *via* planar tape ligand moieties. Only H atoms essential for illustration of the hydrogen bonds, shown as orange dashed lines, are included.

**Figure 3**

A packing diagram of the title compound is displayed along the c axis, illustrating the herringbone-type motif formed by two $[\text{Ru}(\text{L}-\text{N}_4\text{H}_2)\text{tape}]^{2+}$ dimers. The disordered minor atoms are omitted for clarity.

tape ligand, the root-mean-square deviation from planarity was calculated to be 0.0211 \AA . However, in the case of $[\text{Ru}(\text{L}-\text{N}_4\text{H}_2)\text{tape}]^{2+}$, the dimers are also connected through bifurcated hydrogen bonds between one of the two $\text{L}-\text{N}_4\text{H}_2$ ligand amine protons and both nitrogen atoms of the non-coordinating tape ligand α,α' -diimine unit of the second complex cation of the dimer. In the crystal structure, these additional hydrogen bonds result in a short $\text{Ru}\cdots\text{Ru}$ distance of $8.8306(2) \text{ \AA}$, a tape ligand centroid–centroid distance of $3.49(2) \text{ \AA}$ and an angle of $13.7(1.4)^\circ$ between the ring normal and the centroid-to-centroid vector. Therefore, the $\pi\cdots\pi$ stacking motif can be described as parallel-displaced, but near to face-to-face (Fig. 2). In metal complexes, a near face-to-face alignment of the polycyclic units is extremely rare (Janiak, 2000). Furthermore, a large number of weak hydrogen bonds connect cations, anions and solvent molecules, stabilizing the crystal packing (Table 2), supported by $\text{P}\cdots\pi$ -ring (tape, py) interactions with $\text{F}\cdots\text{centroid}$ distances from 2.925 to 3.984 \AA . In the packing, the dimers are oriented in a herringbone-like motif, surrounded by hexafluoridophosphate anions. The solvent acetonitrile molecules fill the space between complex moieties (Fig. 3). For a description of the disorder of the anions and solvent molecules, see the *Refinement* section.

4. Database survey

For related Ru^{II} complexes with 2,11-dimethyl-2,11-diazab-[3.3](2,6)-pyridinophane, see Brietzke, Mickler, Kelling, Schilde *et al.* (2012). For Ru^{II} tetraazaperylene complexes containing bipyridine-type ligands, see: Brietzke, Mickler, Kelling & Holdt (2012); Brietzke *et al.* (2014).

5. Synthesis and crystallization

The syntheses of the ligands $\text{L}-\text{N}_4\text{H}_2$ (Bottino *et al.*, 1988) and tape (Brietzke, Mickler, Kelling & Holdt, 2012) have been reported previously. $[\text{Ru}(\text{L}-\text{N}_4\text{H}_2)\text{tape}](\text{PF}_6)_2$ was synthesized as reported for $[\text{Ru}(\text{L}-\text{N}_4\text{Me}_2)\text{tape}](\text{PF}_6)_2$ (Brietzke, Mickler, Kelling, Schilde *et al.*, 2012), using $\text{L}-\text{N}_4\text{H}_2$ (73.5 mg ,

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H3N \cdots F2A _a ⁱ	0.91	2.22	3.037 (4)	150
N3—H3N \cdots F4A _d ⁱ	0.91	2.55	3.236 (4)	133
N5—H5N \cdots N7 ⁱⁱ	0.88	2.20	3.006 (3)	153
N5—H5N \cdots N8 ⁱⁱ	0.88	2.69	3.373 (4)	135
C1—H1 \cdots F11A _a ⁱⁱⁱ	0.95	2.48	3.376 (5)	157
C1—H1 \cdots F11B _b ⁱⁱⁱ	0.95	2.28	3.019 (11)	134
C3—H3 \cdots F8A _a ^{iv}	0.95	2.51	3.301 (8)	141
C3—H3 \cdots F12A _a ^{iv}	0.95	2.60	3.414 (5)	144
C3—H3 \cdots F8B _b ^{iv}	0.95	2.50	3.28 (3)	140
C3—H3 \cdots F12B _b ^{iv}	0.95	2.37	3.283 (10)	161
C5—H5 \cdots F7A _a ^v	0.95	2.50	3.404 (5)	159
C8—H8 \cdots F1A _a	0.95	2.53	3.211 (4)	128
C8—H8 \cdots F4A _a	0.95	2.40	3.085 (4)	129
C8—H8 \cdots F1B _b	0.95	2.50	3.42 (2)	163
C17—H17A \cdots N9 _a ^{vi}	0.99	2.65	3.500 (8)	145
C17—H17B \cdots F4B _b	0.99	2.34	3.15 (2)	138
C19—H19 \cdots F9B _b ^{vi}	0.95	2.61	3.287 (12)	128
C21—H21 \cdots F11A _a ^{vii}	0.95	2.48	3.166 (4)	129
C23—H23B \cdots F6A _a ⁱⁱⁱ	0.99	2.51	3.409 (4)	151
C24—H24A \cdots F7A _a ⁱⁱⁱ	0.99	2.53	3.224 (5)	127
C24—H24B \cdots F2B _b ⁱⁱⁱ	0.99	2.09	3.00 (2)	152
C24—H24B \cdots F5B _b ⁱⁱⁱ	0.99	2.51	3.41 (3)	150
C26—H26 \cdots F1A _a ⁱⁱⁱ	0.95	2.55	3.328 (5)	140
C26—H26 \cdots F4B _b ⁱⁱⁱ	0.95	2.36	3.26 (3)	156
C28—H28 \cdots N10 ^{viii}	0.95	2.23	3.169 (12)	169
C30—H30A \cdots N9 _a ^{vi}	0.99	2.59	3.484 (7)	151
C30—H30B \cdots F4A _a ⁱ	0.99	2.54	3.182 (5)	122
C32 _a \cdots H32A _a \cdots F12A _a ^{vi}	0.98	2.36	3.273 (8)	155
C32 _a \cdots H32B _a \cdots F7A _a	0.98	2.54	3.150 (7)	121
C32 _a \cdots H32C _a \cdots F1A _a	0.98	2.58	3.446 (8)	148

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x + 1, -y, -z + 1$; (iii) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (iv) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (v) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (vi) $-x + 1, -y + 1, -z + 1$; (vii) $x - 1, y, z$; (viii) $x, y, z + 1$.

$306 \mu\text{mol}$) instead of $\text{L}-\text{N}_4\text{Me}_2$. A yield of 44% (120.0 mg , $135 \mu\text{mol}$) was obtained; m.p. $> 573 \text{ K}$. ^1H NMR = (MeCN- d_3): $\delta = 8.69$ ($d, J = 5.5 \text{ Hz}$, 2H, C^d—H), 8.56 ($d, J = 6.6 \text{ Hz}$, 2H, C^a—H), 8.01 ($t, J = 8.0 \text{ Hz}$, 2H, C^d—H), 7.77 ($d, J = 5.5 \text{ Hz}$, 2H, C^c—H), 7.72 ($d, J = 6.6 \text{ Hz}$, 2H, C^b—H), 7.65 ($d, J = 8.0 \text{ Hz}$, 4H, C³—H + C⁵—H), 5.6 (bs , 2H, N—H), 4.83 ($bd, J = 14.0 \text{ Hz}$, 4H, CH₂), 4.47 ($d, J = 17.4 \text{ Hz}$, 4H, CH₂) p.p.m. ^{13}C NMR = (MeCN- d_3): $\delta = 160.0$ (C² + C⁶), 152.4 (C^e), 150.28 (C^d), 150.24 (C^a), 145.7 (C^f), 138.9 (C⁴), 136.7 (C^{b'}), 123.5 (C^b), 122.7 (C³ + C⁵), 122.0 (C^c), 119.3 (C^e), 64.8 (CH₂) p.p.m. ESI=MS: calculated for $[\text{M}-\text{PF}_6]^+$ 743.0809 ; found 743.0778 .

Crystals suitable for X-ray structure analysis were obtained by vapor diffusion of diethyl ether into a saturated acetonitrile solution of $[\text{Ru}(\text{L}-\text{N}_4\text{H}_2)\text{tape}](\text{PF}_6)_2$. The solution was filled into a test tube, which was placed into a diethyl ether-containing bottle. Dark-green crystals began to form at ambient temperature within a few days.

6. Refinement

Disorder was observed for both the hexafluoridophosphate anions as well as the acetonitrile solvate molecules. Both PF₆ anions were refined as disordered over one major and one minor moiety each. The geometry of the minor moieties were each restrained to be similar to that of the major moieties (within an estimated standard deviation of 0.02 \AA). The minor

Table 3
Experimental details.

Crystal data	
Chemical formula	[Ru(C ₁₄ H ₁₆ N ₄)(C ₁₆ H ₈ N ₄)](PF ₆) ₂ ·1.422C ₂ H ₃ N
M _r	946.11
Crystal system, space group	Monoclinic, P2 ₁ /n
Temperature (K)	150
a, b, c (Å)	11.7027 (3), 21.7157 (7), 13.9377 (4)
β (°)	97.938 (2)
V (Å ³)	3508.08 (18)
Z	4
Radiation type	Mo Kα
μ (mm ⁻¹)	0.65
Crystal size (mm)	1.30 × 0.65 × 0.31
Data collection	
Diffractometer	STOE IPDS 2
Absorption correction	Integration (X-RED; Stoe & Cie, 2011)
T _{min} , T _{max}	0.613, 0.843
No. of measured, independent and observed [I > 2σ(I)] reflections	60767, 9454, 7744
R _{int}	0.087
(sin θ/λ) _{max} (Å ⁻¹)	0.689
Refinement	
R[F ² > 2σ(F ²)], wR(F ²), S	0.048, 0.146, 1.09
No. of reflections	9454
No. of parameters	651
No. of restraints	183
H-atom treatment	H-atom parameters constrained
Δρ _{max} , Δρ _{min} (e Å ⁻³)	1.84, -1.22

Computer programs: X-AREA and X-RED (Stoe & Cie, 2011), SHELXS97 and SHELXL2014 (Sheldrick, 2008), Mercury (Macrae *et al.*, 2006), ORTEP-3 for Windows (Farrugia, 2012), SHELXLE (Hübschle *et al.*, 2011) and publCIF (Westrip, 2010).

moieties were subjected to a rigid bond restraint (RIGU command of SHELX2014, estimated standard deviation 0.004 Å²), and the anisotropic displacement parameters of the major and minor phosphorus atoms were each constrained to be identical. Associated with the major moiety of the PF₆ anion of P1 is an acetonitrile molecule that is absent for the minor moiety. Subject to the restraints and constraints used, the occupancy ratios refined to 0.9215 (17) to 0.0785 (17) for the PF₆ units of P1A and P1B, and to 0.801 (6) and 0.199 (6) for those of P2A and P2B.

A second acetonitrile molecule is disordered across a crystallographic inversion center, with substantial overlap for the two carbon atoms of symmetry-related molecules. The

geometry of the molecule was restrained to be similar to that of the first acetonitrile molecule, and the ADPs of its C and N atoms were restrained to be have similar U_{ij} components to their neighbors closer than 2 Å, including those of symmetry-related atoms (SIMU restraint in SHELX2014, estimated standard deviation 0.01 Å²).

All hydrogen atoms connected to C and N atoms were placed in their expected calculated positions and refined as riding with C—H = 0.98 (CH₃), 0.99 (CH₂), 0.95 (C_{arom}), N—H = 1.0 Å, and with U_{iso}(H) = 1.2U_{eq}(C) with the exception of methyl hydrogen atoms, which were refined with U_{iso}(H) = 1.5U_{eq}(C).

Crystal data, data collection and structure refinement details are summarized in Table 3.

Acknowledgements

We gratefully acknowledge financial assistance provided by the University of Potsdam.

References

- Bottino, F., Di Grazia, M., Finocchiaro, P., Fronczek, F. R., Mamo, A. & Pappalardo, S. (1988). *J. Org. Chem.* **53**, 3521–3529.
- Brietzke, T., Kässler, D., Kelling, A., Schilde, U. & Holdt, H.-J. (2014). *Acta Cryst. E70*, m238–m239.
- Brietzke, T., Mickler, W., Kelling, A. & Holdt, H.-J. (2012). *Dalton Trans.* **41**, 2788–2797.
- Brietzke, T., Mickler, W., Kelling, A., Schilde, U., Krüger, H.-J. & Holdt, H.-J. (2012). *Eur. J. Inorg. Chem.* pp. 4632–4643.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Gut, D., Rudi, A., Kopilov, J., Goldberg, I. & Kol, M. (2002). *J. Am. Chem. Soc.* **124**, 5449–5456.
- Hübschle, C. B., Sheldrick, G. M. & Dittrich, B. (2011). *J. Appl. Cryst.* **44**, 1281–1284.
- Ishow, E., Gourdon, A., Launay, J.-P., Lecante, P., Verelst, M., Chiorboli, C., Scandola, F. & Bignozzi, C.-A. (1998). *Inorg. Chem.* **37**, 3603–3609.
- Janiak, C. (2000). *J. Chem. Soc. Dalton Trans.* pp. 3885–3896.
- Kammer, S., Müller, H., Grunwald, N., Bellin, A., Kelling, A., Schilde, U., Mickler, W., Dosche, C. & Holdt, H.-J. (2006). *Eur. J. Inorg. Chem.* pp. 1547–1551.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Stoe & Cie (2011). X-AREA and X-RED. Stoe & Cie, Darmstadt, Germany.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2014). E70, 265-268 [doi:10.1107/S1600536814021060]

Crystal structure of (2,11-diaza[3.3](2,6)pyridinophane- κ^4N,N',N'',N''')(1,6,7,12-tetraazaperylene- κ^2N^1,N^{12})ruthenium(II) bis(hexafluoridophosphate) acetonitrile 1.422-solvate


Thomas Brietzke, Falko Otto Rottke, Alexandra Kelling, Uwe Schilde and Hans-Jürgen Holdt

Computing details

Data collection: *X-AREA* (Stoe & Cie, 2011); cell refinement: *X-AREA* (Stoe & Cie, 2011); data reduction: *X-RED* (Stoe & Cie, 2011); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006), *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2008), *SHELXLE* (Hübschle *et al.*, 2011), *publCIF* (Westrip, 2010).

(2,11-Diaza[3.3](2,6)pyridinophane- κ^4N,N',N'',N''')(1,6,7,12-tetraazaperylene- κ^2N^1,N^{12})ruthenium(II) bis(hexafluoridophosphate) acetonitrile 1.422-solvate

Crystal data



$M_r = 946.11$

Monoclinic, $P2_1/n$

$a = 11.7027 (3)$ Å

$b = 21.7157 (7)$ Å

$c = 13.9377 (4)$ Å

$\beta = 97.938 (2)^\circ$

$V = 3508.08 (18)$ Å³

$Z = 4$

$F(000) = 1893.4$

$D_x = 1.791$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 63561 reflections

$\theta = 1.5\text{--}29.6^\circ$

$\mu = 0.65$ mm⁻¹

$T = 150$ K

Prism, dark green

1.30 × 0.65 × 0.31 mm

Data collection

STOE IPDS 2

diffractometer

Radiation source: sealed X-ray tube

Plane graphite monochromator

Detector resolution: 6.67 pixels mm⁻¹

rotation method scans

Absorption correction: integration

(*X-RED*; Stoe & Cie, 2011)

$T_{\min} = 0.613$, $T_{\max} = 0.843$

60767 measured reflections

9454 independent reflections

7744 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.087$

$\theta_{\max} = 29.3^\circ$, $\theta_{\min} = 1.8^\circ$

$h = -16 \rightarrow 14$

$k = -29 \rightarrow 29$

$l = -19 \rightarrow 19$

Refinement

Refinement on F^2

$S = 1.09$

Least-squares matrix: full

9454 reflections

$R[F^2 > 2\sigma(F^2)] = 0.048$

651 parameters

$wR(F^2) = 0.146$

183 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: mixed
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0787P)^2 + 2.127P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.002$$

$$\Delta\rho_{\max} = 1.84 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -1.22 \text{ e \AA}^{-3}$$

$$\text{Extinction correction: } SHELXL,$$

$$Fc^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$$

$$\text{Extinction coefficient: } 0.0041 (4)$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ru1	0.40042 (2)	0.19157 (2)	0.58944 (2)	0.03723 (9)	
N1	0.49654 (19)	0.11825 (11)	0.64626 (15)	0.0367 (5)	
N2	0.46354 (19)	0.15956 (11)	0.46830 (16)	0.0368 (4)	
N3	0.5182 (2)	0.26652 (13)	0.6188 (2)	0.0537 (7)	
H3N	0.5905	0.2585	0.6061	0.064*	
N4	0.3005 (2)	0.26143 (11)	0.53082 (16)	0.0398 (5)	
N5	0.2354 (2)	0.14885 (11)	0.58215 (17)	0.0411 (5)	
H5N	0.2357	0.1112	0.5582	0.049*	
N6	0.3522 (2)	0.22164 (12)	0.71383 (17)	0.0458 (6)	
N7	0.7375 (2)	-0.04526 (12)	0.55594 (17)	0.0410 (5)	
N8	0.7102 (2)	0.00109 (12)	0.36755 (17)	0.0431 (5)	
C1	0.5101 (2)	0.09624 (14)	0.73954 (19)	0.0419 (6)	
H1	0.4729	0.1176	0.7861	0.050*	
C2	0.5738 (2)	0.04565 (14)	0.76894 (18)	0.0403 (6)	
H2	0.5793	0.0321	0.8342	0.048*	
C3	0.7009 (3)	-0.03870 (14)	0.72236 (19)	0.0416 (6)	
H3	0.7137	-0.0553	0.7860	0.050*	
C4	0.7498 (3)	-0.06536 (15)	0.6482 (2)	0.0464 (7)	
H4	0.7961	-0.1009	0.6634	0.056*	
C5	0.6950 (3)	0.02554 (16)	0.2771 (2)	0.0489 (7)	
H5	0.7336	0.0061	0.2296	0.059*	
C6	0.6285 (3)	0.07603 (15)	0.2488 (2)	0.0465 (7)	
H6	0.6210	0.0904	0.1839	0.056*	
C7	0.4978 (3)	0.15793 (14)	0.3018 (2)	0.0435 (6)	
H7	0.4836	0.1756	0.2390	0.052*	
C8	0.4473 (3)	0.18253 (14)	0.3751 (2)	0.0433 (6)	
H8	0.3983	0.2173	0.3615	0.052*	
C9	0.5501 (2)	0.08765 (12)	0.58204 (17)	0.0330 (5)	
C10	0.6189 (2)	0.03545 (12)	0.60524 (17)	0.0327 (5)	
C11	0.6734 (2)	0.00476 (12)	0.53437 (18)	0.0339 (5)	
C12	0.6582 (2)	0.02915 (12)	0.43437 (18)	0.0349 (5)	

C13	0.5882 (2)	0.08220 (13)	0.41366 (17)	0.0341 (5)
C14	0.5334 (2)	0.11075 (12)	0.48493 (18)	0.0335 (5)
C15	0.6323 (2)	0.01304 (13)	0.70162 (18)	0.0357 (5)
C16	0.5714 (2)	0.10613 (14)	0.31837 (19)	0.0392 (6)
C17	0.4761 (3)	0.31805 (17)	0.5503 (3)	0.0596 (9)
H17A	0.5040	0.3580	0.5787	0.072*
H17B	0.5074	0.3126	0.4884	0.072*
C18	0.3465 (3)	0.31816 (15)	0.5318 (3)	0.0486 (7)
C19	0.2756 (3)	0.36940 (16)	0.5145 (2)	0.0534 (7)
H19	0.3077	0.4096	0.5146	0.064*
C20	0.1575 (3)	0.36070 (16)	0.4970 (2)	0.0499 (7)
H20	0.1076	0.3952	0.4850	0.060*
C21	0.1115 (3)	0.30167 (15)	0.4970 (2)	0.0467 (7)
H21	0.0304	0.2953	0.4854	0.056*
C22	0.1864 (2)	0.25246 (14)	0.51419 (18)	0.0406 (6)
C23	0.1520 (3)	0.18614 (14)	0.5144 (2)	0.0447 (6)
H23A	0.1483	0.1693	0.4480	0.054*
H23B	0.0741	0.1828	0.5340	0.054*
C24	0.2034 (3)	0.14619 (16)	0.6821 (2)	0.0505 (7)
H24A	0.2333	0.1076	0.7143	0.061*
H24B	0.1183	0.1462	0.6788	0.061*
C25	0.2531 (3)	0.20074 (15)	0.7401 (2)	0.0486 (7)
C26	0.2081 (4)	0.22746 (19)	0.8173 (2)	0.0605 (10)
H26	0.1382	0.2129	0.8367	0.073*
C27	0.2687 (4)	0.2763 (2)	0.8653 (2)	0.0725 (13)
H27	0.2399	0.2952	0.9186	0.087*
C28	0.3680 (4)	0.2973 (2)	0.8372 (3)	0.0711 (13)
H28	0.4082	0.3308	0.8703	0.085*
C29	0.4110 (3)	0.26922 (17)	0.7592 (2)	0.0578 (9)
C30	0.5223 (3)	0.2836 (2)	0.7230 (3)	0.0664 (11)
H30A	0.5388	0.3281	0.7309	0.080*
H30B	0.5856	0.2607	0.7620	0.080*
P2A_a	0.8068 (3)	0.42105 (13)	0.4826 (2)	0.0447 (4) 0.801 (6)
F7A_a	0.7169 (3)	0.4256 (2)	0.3859 (2)	0.0686 (11) 0.801 (6)
F8A_a	0.8952 (6)	0.4132 (4)	0.5788 (4)	0.073 (2) 0.801 (6)
F9A_a	0.8867 (5)	0.4699 (2)	0.4427 (3)	0.119 (2) 0.801 (6)
F10A_a	0.7290 (3)	0.3697 (2)	0.5217 (3)	0.1071 (18) 0.801 (6)
F11A_a	0.8748 (3)	0.3704 (2)	0.4319 (2)	0.0715 (11) 0.801 (6)
F12A_a	0.7436 (5)	0.4740 (2)	0.5315 (3)	0.122 (2) 0.801 (6)
P2B_b	0.7944 (11)	0.4130 (6)	0.4882 (10)	0.0447 (4) 0.199 (6)
F7B_b	0.6911 (14)	0.4026 (9)	0.4033 (11)	0.076 (4) 0.199 (6)
F8B_b	0.8948 (19)	0.4222 (13)	0.5756 (15)	0.053 (4) 0.199 (6)
F9B_b	0.7949 (14)	0.4848 (5)	0.4693 (10)	0.075 (4) 0.199 (6)
F10B_b	0.8082 (15)	0.3414 (5)	0.4931 (10)	0.090 (4) 0.199 (6)
F11B_b	0.8886 (12)	0.4117 (9)	0.4167 (9)	0.081 (4) 0.199 (6)
F12B_b	0.6988 (9)	0.4239 (8)	0.5536 (8)	0.068 (3) 0.199 (6)
P1A_a	0.33081 (8)	0.31111 (5)	0.17529 (8)	0.0538 (3) 0.9215 (17)
F1A_a	0.4320 (3)	0.31248 (12)	0.2642 (3)	0.0900 (11) 0.9215 (17)

F2A_a	0.2279 (2)	0.30908 (12)	0.08663 (17)	0.0644 (6)	0.9215 (17)
F3A_a	0.3730 (4)	0.37384 (18)	0.1391 (4)	0.1430 (19)	0.9215 (17)
F4A_a	0.2860 (2)	0.24711 (14)	0.2119 (2)	0.0810 (8)	0.9215 (17)
F5A_a	0.2477 (3)	0.3457 (2)	0.2363 (2)	0.1186 (15)	0.9215 (17)
F6A_a	0.4133 (3)	0.2749 (2)	0.1160 (3)	0.1041 (11)	0.9215 (17)
P1B_b	0.3791 (8)	0.3534 (5)	0.2328 (8)	0.0538 (3)	0.0785 (17)
F1B_b	0.2932 (17)	0.3068 (10)	0.2780 (18)	0.061 (5)	0.0785 (17)
F2B_b	0.4610 (19)	0.3981 (11)	0.185 (2)	0.082 (6)	0.0785 (17)
F3B_b	0.2821 (18)	0.3626 (12)	0.1461 (17)	0.073 (5)	0.0785 (17)
F4B_b	0.4705 (19)	0.3486 (12)	0.3286 (16)	0.076 (5)	0.0785 (17)
F5B_b	0.433 (2)	0.2970 (9)	0.187 (2)	0.065 (5)	0.0785 (17)
F6B_b	0.324 (2)	0.4083 (11)	0.288 (2)	0.085 (6)	0.0785 (17)
N9_a	0.5253 (5)	0.5581 (3)	0.2892 (6)	0.135 (3)	0.9215 (17)
C31_a	0.4952 (4)	0.5135 (3)	0.3184 (5)	0.0832 (15)	0.9215 (17)
C32_a	0.4527 (5)	0.4596 (3)	0.3578 (6)	0.105 (2)	0.9215 (17)
H32A_a	0.3799	0.4689	0.3821	0.158*	0.9215 (17)
H32B_a	0.5092	0.4446	0.4111	0.158*	0.9215 (17)
H32C_a	0.4394	0.4279	0.3074	0.158*	0.9215 (17)
N10	0.4690 (12)	0.4178 (5)	-0.0541 (15)	0.189 (6)	0.5
C33	0.482 (2)	0.4656 (7)	-0.023 (2)	0.151 (6)	0.5
C34	0.5079 (14)	0.5265 (5)	0.0086 (15)	0.088 (4)	0.5
H34A	0.4426	0.5534	-0.0144	0.132*	0.5
H34B	0.5769	0.5410	-0.0172	0.132*	0.5
H34C	0.5219	0.5273	0.0796	0.132*	0.5

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ru1	0.03631 (13)	0.04128 (15)	0.03304 (13)	0.00739 (8)	0.00104 (8)	-0.00181 (8)
N1	0.0365 (11)	0.0423 (12)	0.0302 (10)	0.0077 (9)	0.0011 (8)	-0.0006 (8)
N2	0.0355 (10)	0.0411 (12)	0.0333 (10)	0.0032 (9)	0.0029 (8)	0.0002 (9)
N3	0.0371 (12)	0.0515 (15)	0.0704 (18)	0.0019 (11)	-0.0005 (12)	-0.0090 (13)
N4	0.0406 (12)	0.0421 (12)	0.0352 (10)	0.0084 (10)	-0.0001 (9)	0.0012 (9)
N5	0.0420 (12)	0.0401 (12)	0.0417 (12)	0.0047 (10)	0.0078 (9)	-0.0039 (9)
N6	0.0511 (14)	0.0498 (14)	0.0344 (11)	0.0186 (11)	-0.0018 (10)	-0.0052 (10)
N7	0.0402 (12)	0.0447 (13)	0.0363 (11)	0.0072 (10)	-0.0011 (9)	-0.0051 (9)
N8	0.0434 (12)	0.0516 (14)	0.0348 (11)	0.0036 (10)	0.0076 (9)	-0.0061 (10)
C1	0.0432 (14)	0.0517 (16)	0.0308 (12)	0.0084 (12)	0.0055 (10)	-0.0032 (11)
C2	0.0434 (14)	0.0486 (15)	0.0282 (11)	0.0044 (12)	0.0024 (10)	0.0005 (10)
C3	0.0447 (14)	0.0463 (15)	0.0318 (12)	0.0063 (12)	-0.0018 (10)	-0.0003 (10)
C4	0.0505 (16)	0.0466 (16)	0.0394 (14)	0.0125 (13)	-0.0029 (12)	-0.0025 (12)
C5	0.0502 (16)	0.0620 (19)	0.0362 (13)	0.0018 (14)	0.0122 (12)	-0.0040 (13)
C6	0.0510 (16)	0.0565 (18)	0.0329 (12)	-0.0012 (14)	0.0093 (11)	0.0022 (12)
C7	0.0464 (15)	0.0485 (16)	0.0349 (12)	-0.0006 (12)	0.0029 (11)	0.0080 (11)
C8	0.0422 (14)	0.0469 (15)	0.0394 (14)	0.0027 (12)	0.0010 (11)	0.0070 (11)
C9	0.0286 (10)	0.0381 (13)	0.0316 (11)	0.0003 (9)	0.0015 (9)	-0.0022 (9)
C10	0.0302 (11)	0.0356 (12)	0.0311 (11)	-0.0008 (9)	0.0000 (9)	-0.0022 (9)
C11	0.0300 (11)	0.0376 (12)	0.0331 (11)	-0.0031 (9)	0.0006 (9)	-0.0044 (9)

C12	0.0321 (11)	0.0410 (13)	0.0312 (11)	-0.0022 (10)	0.0033 (9)	-0.0039 (10)
C13	0.0305 (11)	0.0400 (13)	0.0314 (11)	-0.0034 (9)	0.0026 (9)	-0.0015 (9)
C14	0.0296 (11)	0.0380 (12)	0.0321 (11)	-0.0008 (9)	0.0015 (9)	-0.0012 (9)
C15	0.0353 (12)	0.0402 (13)	0.0303 (11)	0.0002 (10)	0.0006 (9)	-0.0018 (10)
C16	0.0389 (13)	0.0461 (15)	0.0327 (12)	-0.0044 (11)	0.0053 (10)	0.0006 (10)
C17	0.0472 (17)	0.0492 (18)	0.081 (3)	-0.0014 (14)	0.0055 (17)	0.0021 (17)
C18	0.0466 (16)	0.0450 (15)	0.0534 (17)	0.0032 (13)	0.0043 (13)	0.0038 (13)
C19	0.0590 (19)	0.0432 (16)	0.0572 (18)	0.0068 (14)	0.0060 (15)	0.0073 (14)
C20	0.0555 (18)	0.0518 (17)	0.0414 (14)	0.0147 (14)	0.0025 (13)	0.0042 (13)
C21	0.0440 (15)	0.0586 (18)	0.0354 (13)	0.0153 (13)	-0.0018 (11)	-0.0015 (12)
C22	0.0407 (13)	0.0492 (15)	0.0303 (11)	0.0089 (12)	-0.0002 (10)	-0.0032 (10)
C23	0.0376 (14)	0.0517 (17)	0.0433 (15)	0.0064 (12)	0.0005 (11)	-0.0073 (12)
C24	0.0557 (18)	0.0502 (17)	0.0493 (16)	0.0112 (14)	0.0208 (14)	0.0039 (13)
C25	0.0599 (18)	0.0540 (17)	0.0324 (13)	0.0250 (14)	0.0081 (12)	0.0026 (11)
C26	0.074 (2)	0.072 (2)	0.0370 (14)	0.0348 (19)	0.0137 (14)	0.0020 (14)
C27	0.088 (3)	0.088 (3)	0.0377 (15)	0.049 (2)	-0.0052 (17)	-0.0161 (17)
C28	0.072 (3)	0.079 (3)	0.0531 (19)	0.037 (2)	-0.0241 (18)	-0.0289 (18)
C29	0.0607 (19)	0.0594 (19)	0.0465 (16)	0.0259 (16)	-0.0171 (14)	-0.0149 (14)
C30	0.0500 (18)	0.068 (2)	0.074 (2)	0.0122 (17)	-0.0178 (17)	-0.0264 (19)
P2A_a	0.0495 (8)	0.0536 (9)	0.0305 (5)	0.0068 (6)	0.0035 (5)	-0.0021 (6)
F7A_a	0.0612 (19)	0.095 (3)	0.0446 (14)	0.0162 (18)	-0.0104 (13)	0.0000 (16)
F8A_a	0.074 (3)	0.103 (5)	0.038 (2)	0.017 (3)	-0.0093 (19)	-0.001 (2)
F9A_a	0.144 (4)	0.119 (4)	0.087 (3)	-0.066 (3)	-0.012 (3)	0.035 (2)
F10A_a	0.065 (2)	0.146 (4)	0.111 (3)	-0.030 (2)	0.017 (2)	0.056 (3)
F11A_a	0.0552 (16)	0.095 (3)	0.0623 (18)	0.0215 (17)	-0.0006 (13)	-0.0264 (18)
F12A_a	0.166 (5)	0.118 (4)	0.070 (2)	0.093 (4)	-0.020 (3)	-0.035 (2)
P2B_b	0.0495 (8)	0.0536 (9)	0.0305 (5)	0.0068 (6)	0.0035 (5)	-0.0021 (6)
F7B_b	0.075 (7)	0.082 (9)	0.067 (7)	0.024 (5)	-0.008 (5)	-0.032 (6)
F8B_b	0.056 (7)	0.062 (8)	0.042 (7)	-0.006 (5)	0.008 (5)	-0.015 (5)
F9B_b	0.101 (8)	0.057 (5)	0.070 (7)	0.003 (4)	0.015 (6)	-0.008 (4)
F10B_b	0.115 (10)	0.067 (5)	0.080 (7)	0.016 (5)	-0.020 (6)	-0.016 (4)
F11B_b	0.085 (7)	0.110 (10)	0.050 (5)	0.029 (6)	0.021 (5)	-0.011 (6)
F12B_b	0.052 (5)	0.097 (9)	0.056 (5)	-0.012 (5)	0.011 (4)	-0.038 (5)
P1A_a	0.0385 (4)	0.0567 (6)	0.0637 (6)	0.0047 (4)	-0.0018 (4)	0.0080 (4)
F1A_a	0.0674 (17)	0.0674 (17)	0.119 (3)	-0.0021 (13)	-0.0435 (18)	0.0102 (16)
F2A_a	0.0545 (13)	0.0872 (18)	0.0500 (12)	0.0095 (11)	0.0013 (10)	0.0056 (10)
F3A_a	0.107 (3)	0.093 (2)	0.210 (5)	-0.032 (2)	-0.046 (3)	0.080 (3)
F4A_a	0.0538 (13)	0.0899 (19)	0.0937 (19)	-0.0141 (13)	-0.0099 (13)	0.0333 (16)
F5A_a	0.104 (2)	0.160 (4)	0.082 (2)	0.069 (3)	-0.0224 (18)	-0.049 (2)
F6A_a	0.0612 (17)	0.136 (3)	0.121 (3)	0.0272 (19)	0.0349 (17)	0.007 (2)
P1B_b	0.0385 (4)	0.0567 (6)	0.0637 (6)	0.0047 (4)	-0.0018 (4)	0.0080 (4)
F1B_b	0.040 (7)	0.061 (9)	0.078 (10)	0.011 (6)	-0.008 (7)	0.022 (7)
F2B_b	0.054 (9)	0.066 (9)	0.123 (12)	0.009 (7)	0.004 (8)	0.029 (8)
F3B_b	0.050 (8)	0.063 (11)	0.103 (8)	0.001 (7)	-0.004 (6)	0.033 (7)
F4B_b	0.052 (8)	0.066 (11)	0.103 (8)	0.006 (7)	-0.014 (6)	0.008 (7)
F5B_b	0.047 (9)	0.058 (7)	0.087 (10)	0.002 (6)	-0.001 (8)	0.020 (7)
F6B_b	0.061 (10)	0.073 (8)	0.119 (11)	0.012 (7)	0.002 (8)	0.005 (7)
N9_a	0.069 (3)	0.072 (3)	0.264 (9)	0.004 (3)	0.021 (4)	-0.011 (4)

C31_a	0.047 (2)	0.073 (3)	0.127 (5)	0.007 (2)	0.005 (3)	-0.018 (3)
C32_a	0.064 (3)	0.099 (4)	0.152 (6)	0.001 (3)	0.010 (3)	0.007 (4)
N10	0.114 (9)	0.095 (8)	0.335 (18)	-0.005 (7)	-0.052 (11)	-0.012 (11)
C33	0.079 (9)	0.113 (10)	0.247 (15)	-0.004 (9)	-0.025 (10)	-0.005 (11)
C34	0.039 (5)	0.085 (6)	0.142 (10)	0.010 (5)	0.018 (6)	0.025 (7)

Geometric parameters (\AA , $^{\circ}$)

Ru1—N6	2.005 (2)	C20—C21	1.390 (5)
Ru1—N4	2.018 (2)	C20—H20	0.9500
Ru1—N1	2.045 (2)	C21—C22	1.382 (4)
Ru1—N2	2.055 (2)	C21—H21	0.9500
Ru1—N5	2.132 (3)	C22—C23	1.495 (4)
Ru1—N3	2.135 (3)	C23—H23A	0.9900
N1—C9	1.338 (3)	C23—H23B	0.9900
N1—C1	1.374 (3)	C24—C25	1.505 (5)
N2—C14	1.339 (3)	C24—H24A	0.9900
N2—C8	1.380 (4)	C24—H24B	0.9900
N3—C30	1.492 (5)	C25—C26	1.388 (4)
N3—C17	1.509 (5)	C26—C27	1.395 (6)
N3—H3N	0.9051	C26—H26	0.9500
N4—C22	1.338 (4)	C27—C28	1.356 (7)
N4—C18	1.343 (4)	C27—H27	0.9500
N5—C24	1.493 (4)	C28—C29	1.400 (5)
N5—C23	1.499 (4)	C28—H28	0.9500
N5—H5N	0.8825	C29—C30	1.494 (6)
N6—C25	1.342 (5)	C30—H30A	0.9900
N6—C29	1.349 (5)	C30—H30B	0.9900
N7—C11	1.330 (4)	P2A_a—F9A_a	1.566 (4)
N7—C4	1.347 (4)	P2A_a—F12A_a	1.573 (4)
N8—C12	1.328 (3)	P2A_a—F11A_a	1.581 (4)
N8—C5	1.356 (4)	P2A_a—F10A_a	1.583 (4)
C1—C2	1.359 (4)	P2A_a—F8A_a	1.585 (4)
C1—H1	0.9500	P2A_a—F7A_a	1.594 (4)
C2—C15	1.424 (4)	P2B_b—F12B_b	1.556 (13)
C2—H2	0.9500	P2B_b—F10B_b	1.565 (14)
C3—C4	1.376 (4)	P2B_b—F9B_b	1.581 (14)
C3—C15	1.388 (4)	P2B_b—F8B_b	1.584 (14)
C3—H3	0.9500	P2B_b—F11B_b	1.585 (13)
C4—H4	0.9500	P2B_b—F7B_b	1.586 (14)
C5—C6	1.371 (5)	P1A_a—F3A_a	1.556 (3)
C5—H5	0.9500	P1A_a—F6A_a	1.566 (3)
C6—C16	1.411 (4)	P1A_a—F5A_a	1.569 (3)
C6—H6	0.9500	P1A_a—F1A_a	1.592 (3)
C7—C8	1.358 (4)	P1A_a—F4A_a	1.594 (3)
C7—C16	1.416 (4)	P1A_a—F2A_a	1.603 (3)
C7—H7	0.9500	P1B_b—F3B_b	1.552 (15)
C8—H8	0.9500	P1B_b—F5B_b	1.555 (16)

C9—C10	1.402 (4)	P1B_b—F2B_b	1.576 (15)
C9—C14	1.431 (3)	P1B_b—F4B_b	1.594 (15)
C10—C11	1.415 (3)	P1B_b—F6B_b	1.601 (16)
C10—C15	1.417 (3)	P1B_b—F1B_b	1.614 (15)
C11—C12	1.479 (4)	N9_a—C31_a	1.125 (8)
C12—C13	1.420 (4)	C31_a—C32_a	1.413 (8)
C13—C14	1.400 (3)	C32_a—H32A_a	0.9800
C13—C16	1.414 (4)	C32_a—H32B_a	0.9800
C17—C18	1.504 (5)	C32_a—H32C_a	0.9800
C17—H17A	0.9900	N10—C33	1.125 (8)
C17—H17B	0.9900	C33—C34	1.413 (8)
C18—C19	1.389 (5)	C34—H34A	0.9800
C19—C20	1.383 (5)	C34—H34B	0.9800
C19—H19	0.9500	C34—H34C	0.9800
N6—Ru1—N4	83.61 (9)	C21—C22—C23	125.5 (3)
N6—Ru1—N1	97.14 (9)	C22—C23—N5	111.7 (2)
N4—Ru1—N1	177.60 (10)	C22—C23—H23A	109.3
N6—Ru1—N2	175.16 (9)	N5—C23—H23A	109.3
N4—Ru1—N2	100.11 (9)	C22—C23—H23B	109.3
N1—Ru1—N2	79.27 (9)	N5—C23—H23B	109.3
N6—Ru1—N5	79.72 (10)	H23A—C23—H23B	107.9
N4—Ru1—N5	80.65 (10)	N5—C24—C25	110.0 (3)
N1—Ru1—N5	97.22 (9)	N5—C24—H24A	109.7
N2—Ru1—N5	103.86 (9)	C25—C24—H24A	109.7
N6—Ru1—N3	80.65 (12)	N5—C24—H24B	109.7
N4—Ru1—N3	80.06 (11)	C25—C24—H24B	109.7
N1—Ru1—N3	102.31 (11)	H24A—C24—H24B	108.2
N2—Ru1—N3	96.87 (11)	N6—C25—C26	120.3 (3)
N5—Ru1—N3	153.79 (10)	N6—C25—C24	113.8 (3)
C9—N1—C1	117.2 (2)	C26—C25—C24	125.8 (4)
C9—N1—Ru1	114.26 (17)	C25—C26—C27	117.8 (4)
C1—N1—Ru1	128.52 (18)	C25—C26—H26	121.1
C14—N2—C8	116.8 (2)	C27—C26—H26	121.1
C14—N2—Ru1	113.92 (17)	C28—C27—C26	121.2 (3)
C8—N2—Ru1	129.2 (2)	C28—C27—H27	119.4
C30—N3—C17	113.3 (3)	C26—C27—H27	119.4
C30—N3—Ru1	108.0 (2)	C27—C28—C29	119.4 (4)
C17—N3—Ru1	107.4 (2)	C27—C28—H28	120.3
C30—N3—H3N	109.4	C29—C28—H28	120.3
C17—N3—H3N	104.6	N6—C29—C28	118.9 (4)
Ru1—N3—H3N	114.3	N6—C29—C30	114.4 (3)
C22—N4—C18	121.6 (3)	C28—C29—C30	126.5 (4)
C22—N4—Ru1	118.0 (2)	N3—C30—C29	111.4 (3)
C18—N4—Ru1	118.4 (2)	N3—C30—H30A	109.3
C24—N5—C23	112.5 (2)	C29—C30—H30A	109.3
C24—N5—Ru1	108.35 (19)	N3—C30—H30B	109.3
C23—N5—Ru1	107.56 (18)	C29—C30—H30B	109.3

C24—N5—H5N	109.4	H30A—C30—H30B	108.0
C23—N5—H5N	107.3	F9A_a—P2A_a—F12A_a	89.9 (4)
Ru1—N5—H5N	111.7	F9A_a—P2A_a—F11A_a	87.2 (3)
C25—N6—C29	122.3 (3)	F12A_a—P2A_a—F11A_a	177.0 (4)
C25—N6—Ru1	118.9 (2)	F9A_a—P2A_a—F10A_a	177.8 (4)
C29—N6—Ru1	117.9 (2)	F12A_a—P2A_a—F10A_a	92.3 (4)
C11—N7—C4	117.6 (2)	F11A_a—P2A_a—F10A_a	90.7 (3)
C12—N8—C5	117.3 (3)	F9A_a—P2A_a—F8A_a	91.2 (4)
C2—C1—N1	123.6 (2)	F12A_a—P2A_a—F8A_a	90.2 (4)
C2—C1—H1	118.2	F11A_a—P2A_a—F8A_a	89.6 (3)
N1—C1—H1	118.2	F10A_a—P2A_a—F8A_a	88.4 (4)
C1—C2—C15	119.8 (2)	F9A_a—P2A_a—F7A_a	90.8 (3)
C1—C2—H2	120.1	F12A_a—P2A_a—F7A_a	91.5 (3)
C15—C2—H2	120.1	F11A_a—P2A_a—F7A_a	88.8 (2)
C4—C3—C15	118.2 (3)	F10A_a—P2A_a—F7A_a	89.5 (2)
C4—C3—H3	120.9	F8A_a—P2A_a—F7A_a	177.3 (4)
C15—C3—H3	120.9	F12B_b—P2B_b—F10B_b	101.7 (11)
N7—C4—C3	125.4 (3)	F12B_b—P2B_b—F9B_b	88.2 (10)
N7—C4—H4	117.3	F10B_b—P2B_b—F9B_b	170.0 (12)
C3—C4—H4	117.3	F12B_b—P2B_b—F8B_b	92.7 (12)
N8—C5—C6	125.2 (3)	F10B_b—P2B_b—F8B_b	91.7 (13)
N8—C5—H5	117.4	F9B_b—P2B_b—F8B_b	89.2 (12)
C6—C5—H5	117.4	F12B_b—P2B_b—F11B_b	171.9 (13)
C5—C6—C16	118.5 (3)	F10B_b—P2B_b—F11B_b	86.2 (11)
C5—C6—H6	120.8	F9B_b—P2B_b—F11B_b	83.9 (10)
C16—C6—H6	120.8	F8B_b—P2B_b—F11B_b	88.8 (12)
C8—C7—C16	120.6 (3)	F12B_b—P2B_b—F7B_b	85.6 (9)
C8—C7—H7	119.7	F10B_b—P2B_b—F7B_b	87.3 (10)
C16—C7—H7	119.7	F9B_b—P2B_b—F7B_b	92.1 (10)
C7—C8—N2	123.2 (3)	F8B_b—P2B_b—F7B_b	177.8 (14)
C7—C8—H8	118.4	F11B_b—P2B_b—F7B_b	93.0 (11)
N2—C8—H8	118.4	F3A_a—P1A_a—F6A_a	91.3 (3)
N1—C9—C10	123.7 (2)	F3A_a—P1A_a—F5A_a	90.3 (3)
N1—C9—C14	116.3 (2)	F6A_a—P1A_a—F5A_a	178.4 (3)
C10—C9—C14	120.0 (2)	F3A_a—P1A_a—F1A_a	90.06 (19)
C9—C10—C11	121.3 (2)	F6A_a—P1A_a—F1A_a	88.5 (2)
C9—C10—C15	118.8 (2)	F5A_a—P1A_a—F1A_a	91.1 (2)
C11—C10—C15	119.9 (2)	F3A_a—P1A_a—F4A_a	179.29 (18)
N7—C11—C10	121.4 (2)	F6A_a—P1A_a—F4A_a	89.1 (2)
N7—C11—C12	119.8 (2)	F5A_a—P1A_a—F4A_a	89.3 (2)
C10—C11—C12	118.8 (2)	F1A_a—P1A_a—F4A_a	90.53 (15)
N8—C12—C13	122.5 (2)	F3A_a—P1A_a—F2A_a	90.70 (18)
N8—C12—C11	119.3 (2)	F6A_a—P1A_a—F2A_a	91.83 (18)
C13—C12—C11	118.2 (2)	F5A_a—P1A_a—F2A_a	88.55 (15)
C14—C13—C16	119.1 (2)	F1A_a—P1A_a—F2A_a	179.18 (18)
C14—C13—C12	121.6 (2)	F4A_a—P1A_a—F2A_a	88.72 (14)
C16—C13—C12	119.3 (2)	F3B_b—P1B_b—F5B_b	94.2 (14)
N2—C14—C13	123.8 (2)	F3B_b—P1B_b—F2B_b	90.7 (13)

N2—C14—C9	116.2 (2)	F5B_b—P1B_b—F2B_b	90.3 (13)
C13—C14—C9	120.0 (2)	F3B_b—P1B_b—F4B_b	173.7 (15)
C3—C15—C10	117.4 (2)	F5B_b—P1B_b—F4B_b	91.9 (13)
C3—C15—C2	125.6 (2)	F2B_b—P1B_b—F4B_b	91.0 (13)
C10—C15—C2	117.0 (2)	F3B_b—P1B_b—F6B_b	88.9 (13)
C6—C16—C13	117.2 (3)	F5B_b—P1B_b—F6B_b	175.5 (15)
C6—C16—C7	126.2 (3)	F2B_b—P1B_b—F6B_b	92.9 (14)
C13—C16—C7	116.5 (2)	F4B_b—P1B_b—F6B_b	85.0 (13)
C18—C17—N3	110.1 (3)	F3B_b—P1B_b—F1B_b	87.5 (12)
C18—C17—H17A	109.6	F5B_b—P1B_b—F1B_b	88.8 (12)
N3—C17—H17A	109.6	F2B_b—P1B_b—F1B_b	178.0 (15)
C18—C17—H17B	109.6	F4B_b—P1B_b—F1B_b	90.9 (12)
N3—C17—H17B	109.6	F6B_b—P1B_b—F1B_b	88.1 (13)
H17A—C17—H17B	108.2	N9_a—C31_a—C32_a	176.6 (7)
N4—C18—C19	120.3 (3)	C31_a—C32_a—H32A_a	109.5
N4—C18—C17	113.2 (3)	C31_a—C32_a—H32B_a	109.5
C19—C18—C17	126.5 (3)	H32A_a—C32_a—H32B_a	109.5
C20—C19—C18	118.6 (3)	C31_a—C32_a—H32C_a	109.5
C20—C19—H19	120.7	H32A_a—C32_a—H32C_a	109.5
C18—C19—H19	120.7	H32B_a—C32_a—H32C_a	109.5
C19—C20—C21	120.3 (3)	N10—C33—C34	174.0 (19)
C19—C20—H20	119.9	C33—C34—H34A	109.5
C21—C20—H20	119.9	C33—C34—H34B	109.5
C22—C21—C20	118.4 (3)	H34A—C34—H34B	109.5
C22—C21—H21	120.8	C33—C34—H34C	109.5
C20—C21—H21	120.8	H34A—C34—H34C	109.5
N4—C22—C21	120.8 (3)	H34B—C34—H34C	109.5
N4—C22—C23	113.7 (2)		
C9—N1—C1—C2	-0.3 (4)	C1—C2—C15—C10	-0.9 (4)
Ru1—N1—C1—C2	177.9 (2)	C5—C6—C16—C13	0.7 (4)
N1—C1—C2—C15	1.0 (5)	C5—C6—C16—C7	-178.7 (3)
C11—N7—C4—C3	0.5 (5)	C14—C13—C16—C6	179.9 (3)
C15—C3—C4—N7	0.4 (5)	C12—C13—C16—C6	-1.4 (4)
C12—N8—C5—C6	-1.5 (5)	C14—C13—C16—C7	-0.7 (4)
N8—C5—C6—C16	0.8 (5)	C12—C13—C16—C7	178.1 (2)
C16—C7—C8—N2	0.0 (5)	C8—C7—C16—C6	179.6 (3)
C14—N2—C8—C7	0.3 (4)	C8—C7—C16—C13	0.2 (4)
Ru1—N2—C8—C7	177.6 (2)	C30—N3—C17—C18	-85.1 (4)
C1—N1—C9—C10	-0.5 (4)	Ru1—N3—C17—C18	34.0 (4)
Ru1—N1—C9—C10	-178.98 (19)	C22—N4—C18—C19	-0.6 (5)
C1—N1—C9—C14	179.0 (2)	Ru1—N4—C18—C19	-164.4 (3)
Ru1—N1—C9—C14	0.6 (3)	C22—N4—C18—C17	-179.1 (3)
N1—C9—C10—C11	179.9 (2)	Ru1—N4—C18—C17	17.1 (4)
C14—C9—C10—C11	0.3 (4)	N3—C17—C18—N4	-34.3 (4)
N1—C9—C10—C15	0.6 (4)	N3—C17—C18—C19	147.3 (3)
C14—C9—C10—C15	-179.0 (2)	N4—C18—C19—C20	0.5 (5)
C4—N7—C11—C10	-0.8 (4)	C17—C18—C19—C20	178.8 (3)

C4—N7—C11—C12	179.3 (3)	C18—C19—C20—C21	0.0 (5)
C9—C10—C11—N7	−179.1 (2)	C19—C20—C21—C22	−0.4 (5)
C15—C10—C11—N7	0.2 (4)	C18—N4—C22—C21	0.3 (4)
C9—C10—C11—C12	0.8 (4)	Ru1—N4—C22—C21	164.1 (2)
C15—C10—C11—C12	−179.9 (2)	C18—N4—C22—C23	179.1 (3)
C5—N8—C12—C13	0.7 (4)	Ru1—N4—C22—C23	−17.1 (3)
C5—N8—C12—C11	−179.1 (3)	C20—C21—C22—N4	0.2 (4)
N7—C11—C12—N8	−0.9 (4)	C20—C21—C22—C23	−178.4 (3)
C10—C11—C12—N8	179.2 (2)	N4—C22—C23—N5	31.3 (3)
N7—C11—C12—C13	179.3 (2)	C21—C22—C23—N5	−150.0 (3)
C10—C11—C12—C13	−0.6 (4)	C24—N5—C23—C22	90.0 (3)
N8—C12—C13—C14	179.4 (2)	Ru1—N5—C23—C22	−29.3 (3)
C11—C12—C13—C14	−0.7 (4)	C23—N5—C24—C25	−86.8 (3)
N8—C12—C13—C16	0.7 (4)	Ru1—N5—C24—C25	32.0 (3)
C11—C12—C13—C16	−179.5 (2)	C29—N6—C25—C26	−0.9 (4)
C8—N2—C14—C13	−0.8 (4)	Ru1—N6—C25—C26	−169.9 (2)
Ru1—N2—C14—C13	−178.5 (2)	C29—N6—C25—C24	−178.3 (3)
C8—N2—C14—C9	179.6 (2)	Ru1—N6—C25—C24	12.7 (3)
Ru1—N2—C14—C9	1.9 (3)	N5—C24—C25—N6	−30.1 (4)
C16—C13—C14—N2	1.0 (4)	N5—C24—C25—C26	152.7 (3)
C12—C13—C14—N2	−177.7 (2)	N6—C25—C26—C27	0.3 (5)
C16—C13—C14—C9	−179.4 (2)	C24—C25—C26—C27	177.3 (3)
C12—C13—C14—C9	1.9 (4)	C25—C26—C27—C28	0.4 (5)
N1—C9—C14—N2	−1.7 (3)	C26—C27—C28—C29	−0.5 (6)
C10—C9—C14—N2	177.9 (2)	C25—N6—C29—C28	0.8 (5)
N1—C9—C14—C13	178.7 (2)	Ru1—N6—C29—C28	170.0 (2)
C10—C9—C14—C13	−1.7 (4)	C25—N6—C29—C30	176.9 (3)
C4—C3—C15—C10	−0.9 (4)	Ru1—N6—C29—C30	−13.9 (4)
C4—C3—C15—C2	178.9 (3)	C27—C28—C29—N6	−0.1 (5)
C9—C10—C15—C3	180.0 (2)	C27—C28—C29—C30	−175.7 (4)
C11—C10—C15—C3	0.7 (4)	C17—N3—C30—C29	90.1 (4)
C9—C10—C15—C2	0.1 (4)	Ru1—N3—C30—C29	−28.7 (4)
C11—C10—C15—C2	−179.2 (2)	N6—C29—C30—N3	28.9 (4)
C1—C2—C15—C3	179.3 (3)	C28—C29—C30—N3	−155.4 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N3—H3N \cdots F2A $_a^i$	0.91	2.22	3.037 (4)	150
N3—H3N \cdots F4A $_a^i$	0.91	2.55	3.236 (4)	133
N5—H5N \cdots N7 ii	0.88	2.20	3.006 (3)	153
N5—H5N \cdots N8 ii	0.88	2.69	3.373 (4)	135
C1—H1 \cdots F11A $_a^{iii}$	0.95	2.48	3.376 (5)	157
C1—H1 \cdots F11B $_b^{iii}$	0.95	2.28	3.019 (11)	134
C3—H3 \cdots F8A $_a^{iv}$	0.95	2.51	3.301 (8)	141
C3—H3 \cdots F12A $_a^{iv}$	0.95	2.60	3.414 (5)	144
C3—H3 \cdots F8B $_b^{iv}$	0.95	2.50	3.28 (3)	140
C3—H3 \cdots F12B $_b^{iv}$	0.95	2.37	3.283 (10)	161

C5—H5···F7A_ <i>a</i> ^v	0.95	2.50	3.404 (5)	159
C8—H8···F1A_ <i>a</i>	0.95	2.53	3.211 (4)	128
C8—H8···F4A_ <i>a</i>	0.95	2.40	3.085 (4)	129
C8—H8···F1B_ <i>b</i>	0.95	2.50	3.42 (2)	163
C17—H17A···N9_ <i>a</i> ^{vi}	0.99	2.65	3.500 (8)	145
C17—H17B···F4B_ <i>b</i>	0.99	2.34	3.15 (2)	138
C19—H19···F9B_ <i>b</i> ^{vi}	0.95	2.61	3.287 (12)	128
C21—H21···F11A_ <i>a</i> ^{vii}	0.95	2.48	3.166 (4)	129
C23—H23B···F6A_ <i>a</i> ⁱⁱⁱ	0.99	2.51	3.409 (4)	151
C24—H24A···F7A_ <i>a</i> ⁱⁱⁱ	0.99	2.53	3.224 (5)	127
C24—H24B···F2B_ <i>b</i> ⁱⁱⁱ	0.99	2.09	3.00 (2)	152
C24—H24B···F5B_ <i>b</i> ⁱⁱⁱ	0.99	2.51	3.41 (3)	150
C26—H26···F1A_ <i>a</i> ⁱⁱⁱ	0.95	2.55	3.328 (5)	140
C26—H26···F4B_ <i>b</i> ⁱⁱⁱ	0.95	2.36	3.26 (3)	156
C28—H28···N10 ^{viii}	0.95	2.23	3.169 (12)	169
C30—H30A···N9_ <i>a</i> ^{vi}	0.99	2.59	3.484 (7)	151
C30—H30B···F4A_ <i>a</i> ⁱ	0.99	2.54	3.182 (5)	122
C32_ <i>a</i> —H32A_ <i>a</i> ···F12A_ <i>a</i> ^{vi}	0.98	2.36	3.273 (8)	155
C32_ <i>a</i> —H32B_ <i>a</i> ···F7A_ <i>a</i>	0.98	2.54	3.150 (7)	121
C32_ <i>a</i> —H32C_ <i>a</i> ···F1A_ <i>a</i>	0.98	2.58	3.446 (8)	148

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $-x+1, -y, -z+1$; (iii) $x-1/2, -y+1/2, z+1/2$; (iv) $-x+3/2, y-1/2, -z+3/2$; (v) $-x+3/2, y-1/2, -z+1/2$; (vi) $-x+1, -y+1, -z+1$; (vii) $x-1, y, z$; (viii) $x, y, z+1$.