

4-Methyl-N-[2-(pyridin-2-yl)ethyl]carbamothioyl]benzamide

Farook Adam,^{a*} Nadiah Ameram^a and Naser Eltaher Eltayeb^{b†}

^aSchool of Chemical Sciences, 11800, USM Pulau Pinang, Malaysia, and ^bCollege of Sciences and Arts, Rabigh, King Abdulaziz University, Saudi Arabia. Correspondence e-mail: farook@usm.my

Received 15 June 2014; accepted 15 July 2014


Edited by G. Smith, Queensland University of Technology, Australia

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.068; wR factor = 0.182; data-to-parameter ratio = 17.1.

In the title compound, $\text{C}_{16}\text{H}_{17}\text{N}_3\text{OS}$, the dihedral angle between the planes of the benzene and pyridine rings is $71.33(15)^\circ$. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond is present. In the crystal, weak aromatic $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into chains extending along a .

Related literature

For related structures, see: Saeed & Flörke (2007); Yusof *et al.* (2008, 2011); Shoukat *et al.* (2007); Hassan *et al.* (2008a,b,c). For standard bond lengths, see: Allen *et al.* (1987). For graph-set analysis, see Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{17}\text{N}_3\text{OS}$	$V = 1478.7(2)\text{ \AA}^3$
$M_r = 299.39$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 16.0467(12)\text{ \AA}$	$\mu = 0.22\text{ mm}^{-1}$
$b = 4.8824(4)\text{ \AA}$	$T = 100\text{ K}$
$c = 23.0403(18)\text{ \AA}$	$0.47 \times 0.20 \times 0.14\text{ mm}$
$\beta = 124.997(5)^\circ$	

Data collection

Bruker APEXII CCD diffractometer	12777 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	3409 independent reflections
$T_{\min} = 0.903$, $T_{\max} = 0.980$	2221 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.082$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.068$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.182$	$\Delta\rho_{\text{max}} = 0.66\text{ e \AA}^{-3}$
$S = 1.04$	$\Delta\rho_{\text{min}} = -0.44\text{ e \AA}^{-3}$
3409 reflections	
199 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H1N2 \cdots O2	0.87 (4)	1.90 (3)	2.645 (3)	143 (3)
C14—H14A \cdots O2 ⁱ	0.95	2.51	3.421 (4)	161

Symmetry code: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

The authors thank Universiti Sains Malaysia for a research grant (No. PKIMIA846017) which partially supported this work.

Supporting information for this paper is available from the IUCr electronic archives (Reference: ZS2306).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hassan, I. N., Yamin, B. M. & Kassim, M. B. (2008a). *Acta Cryst. E64*, o1727.
- Hassan, I. N., Yamin, B. M. & Kassim, M. B. (2008b). *Acta Cryst. E64*, o2083.
- Hassan, I. N., Yamin, B. M. & Kassim, M. B. (2008c). *Acta Cryst. E64*, o2167.
- Saeed, A. & Flörke, U. (2007). *Acta Cryst. E63*, o4259.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Shoukat, N., Rauf, M. K., Bolte, M. & Badshah, A. (2007). *Acta Cryst. E63*, o3207.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Yusof, M. S. M., Muhamram, S. H., Kassim, M. B. & Yamin, B. M. (2008). *Acta Cryst. E64*, o1137.
- Yusof, M. S. M., Wong, S. T. & Yamin, B. M. (2011). *Acta Cryst. E67*, o2483.

† Additional address: Department of Chemistry, International University of Africa, Sudan

supporting information

Acta Cryst. (2014). E70, o885 [doi:10.1107/S1600536814016377]

4-Methyl-N-[2-(pyridin-2-yl)ethylcarbamothioyl]benzamide

Farook Adam, Nadiah Ameram and Naser Eltaher Eltayeb

S1. Comment

In the title compound, $C_{16}H_{17}N_3SO$ (Fig. 1), the bond lengths and angles are generally normal compared to those in *N*-alkyl-*N*-benzoylthiourea compounds (Allen *et al.*, 1987). The bond lengths of the carbonyl and thiocarbonyl groups [$C7—O2 = 1.229$ (5) Å and $C8—S1 = 1.677$ (4) Å, respectively] have typical $C=O$ and $C=S$ double-bond character (Yusof *et al.* 2011). However, the thiocarbonyl is longer compared to the typical $C=S$ bond which is 1.660 (2) Å. The $C—N$ bond lengths for the title compound [$C7—N1 = 1.375$ (4) Å, $C8—N1 = 1.397$ (4) Å, $C9—N2 = 1.460$ (4) Å, $C11—N3 = 1.335$ (4), $C15—N3 = 1.351$ (4) Å] are all shorter than the average $C—N$ single bond length [1.472 (5) Å], thus showing varying degrees of single bond character (Yusof *et al.* 2008). These bond features in the structure are presumed as a result of the intramolecular H-bonding interactions "locking" the molecule into a planar six-membered ring structure and are consistent with the expected delocalization in the title compound, confirmed by the $C9—N2—C8$ and $C8—N1—C7$ bond angles [125.0 (3) and 128.8 (3)°, respectively], showing sp^2 hybridization on the N2 and N1 atoms. The molecule maintains its *cis-trans* configuration with respect to the position of the methyl benzene and ethyl pyridine groups relative to the thiocarbonyl sulfur atom across the $N1—C7$ and $N2—C8$ bonds, respectively (Hassan *et al.* (2008b,2008c)). The conformation of the molecule with respect to the thiocarbonyl and carbonyl moieties is twisted, as reflected by the torsion angles [$C8—N1—C7—O2$, $C7—N1—C8—N2$ and $C7—N1—C8—S1$: 2.1 (5), -4.4 (4) and 175.9 (2)°, respectively. The angle between the benzene and pyridine rings is 71.33 (15)°. The N2 H-atom forms bifurcated intramolecular interactions with both a carbonyl O-atom and the pyridine N-atom (Table 1): a hydrogen bond with $O2$ ($N2—H···O2$) and an interaction with $N3$ ($N2—H···N3$), giving cyclic motifs [graph sets $S6$ (Bernstein *et al.*, 1995)]. Present also are weak intramolecular $C1—H···O2$ and $C9—H···S1$ interactions [graph set $S(5)$]. In the crystal, molecules are connected through weak intermolecular $C14—H···O2$ hydrogen-bonding interactions, giving one-dimensional chain structures extending along x (Fig. 2). The $N1—H1N1$ group has no acceptor in the crystal.


S2. Experimental

S2.1. Synthesis and crystallization


Freshly prepared substituted *p*-benzoyl chloride (13 mmol) was added dropwise to a stirred acetone solution (30 ml) of ammonium thiocyanate (13 mmol). The mixture was stirred for 10 min. A solution of 2-(2-aminethylpyridine) in acetone was added and the reaction mixture was refluxed for 3 h., after which the solution was poured into a beaker containing some ice cubes. The resulting precipitate was collected by filtration, washed several times with a cold ethanol/water mixture and purified by recrystallization from ethanol (Hassan *et al.*, 2008a). Yield 65%; white transparent crystals, m.p. 126.3 °C. Anal Calc. for $C_{16}H_{17}N_3OS$: C, 64.9; H, 5.6; N, 15.9; S, 8.2%. Found: C, 64.8; H, 5.7; N, 14.8; S, 8.7%.

S2.2. Refinement

The H-atoms on the N atoms were located in a difference-Fourier and were fully refined. All other H-atoms were positioned geometrically and refined using a riding model with C—H = 0.95–0.99 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{aromatic C})$, $1.5U_{\text{eq}}(\text{methyl C})$ and $1.2U_{\text{eq}}(\text{methylene C})$.

**Figure 1**

The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme. Intramolecular interactions are shown as dashed lines.

**Figure 2**

The crystal packing of the title compound viewed down the *c* axis. Hydrogen bonds are shown as dashed lines.

(I)

Crystal data

$C_{16}H_{17}N_3OS$
 $M_r = 299.39$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 16.0467 (12) \text{ \AA}$
 $b = 4.8824 (4) \text{ \AA}$
 $c = 23.0403 (18) \text{ \AA}$
 $\beta = 124.997 (5)^\circ$
 $V = 1478.7 (2) \text{ \AA}^3$
 $Z = 4$

$F(000) = 632$
 $D_x = 1.345 \text{ Mg m}^{-3}$
Melting point: 399.3 K
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 $\theta = 2.6\text{--}25.5^\circ$
 $\mu = 0.22 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
Block, colourless
 $0.47 \times 0.20 \times 0.14 \text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)
 $T_{\min} = 0.903$, $T_{\max} = 0.980$

12777 measured reflections
3409 independent reflections
2221 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.082$
 $\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -17 \rightarrow 20$
 $k = -6 \rightarrow 6$
 $l = -30 \rightarrow 24$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.068$
 $wR(F^2) = 0.182$
 $S = 1.04$
3409 reflections
199 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0899P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.66 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.44 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.72652 (6)	0.04528 (17)	0.87044 (4)	0.0286 (3)
O2	0.78657 (14)	0.5226 (4)	0.72802 (11)	0.0272 (5)
N1	0.69712 (19)	0.3898 (5)	0.77218 (13)	0.0227 (6)
N2	0.84392 (17)	0.1337 (5)	0.82384 (13)	0.0211 (5)
N3	1.02858 (17)	0.3126 (5)	0.84401 (12)	0.0229 (6)
C1	0.6442 (2)	0.9043 (6)	0.63777 (15)	0.0236 (7)
H1A	0.7034	0.8759	0.6386	0.028*
C2	0.5737 (2)	1.0965 (6)	0.59246 (15)	0.0250 (7)
H2A	0.5853	1.1996	0.5627	0.030*
C3	0.4852 (2)	1.1422 (6)	0.58956 (15)	0.0227 (6)
C4	0.4718 (2)	0.9901 (6)	0.63468 (16)	0.0246 (7)
H4A	0.4128	1.0199	0.6341	0.030*
C5	0.5426 (2)	0.7954 (6)	0.68073 (15)	0.0245 (7)
H5A	0.5315	0.6933	0.7108	0.029*
C6	0.6301 (2)	0.7508 (6)	0.68249 (14)	0.0200 (6)

C7	0.7113 (2)	0.5479 (6)	0.72911 (14)	0.0203 (6)
C8	0.7611 (2)	0.1903 (6)	0.82127 (15)	0.0218 (6)
C9	0.9261 (2)	-0.0462 (6)	0.87572 (15)	0.0240 (7)
H9A	0.9508	-0.1556	0.8523	0.029*
H9B	0.9001	-0.1743	0.8952	0.029*
C10	1.0135 (2)	0.1207 (6)	0.93572 (15)	0.0242 (7)
H10A	0.9874	0.2330	0.9578	0.029*
H10B	1.0650	-0.0066	0.9723	0.029*
C11	1.0649 (2)	0.3085 (6)	0.91261 (15)	0.0224 (6)
C12	1.1454 (2)	0.4739 (6)	0.96253 (16)	0.0282 (7)
H12A	1.1707	0.4633	1.0112	0.034*
C13	1.1880 (2)	0.6550 (7)	0.93981 (18)	0.0315 (8)
H13A	1.2426	0.7710	0.9728	0.038*
C14	1.1503 (2)	0.6643 (6)	0.86918 (18)	0.0323 (8)
H14A	1.1775	0.7882	0.8523	0.039*
C15	1.0716 (2)	0.4885 (6)	0.82314 (17)	0.0269 (7)
H15A	1.0466	0.4919	0.7745	0.032*
C16	0.4073 (2)	1.3480 (6)	0.53901 (16)	0.0279 (7)
H16C	0.3579	1.3783	0.5505	0.042*
H16D	0.3721	1.2787	0.4904	0.042*
H16A	0.4412	1.5211	0.5431	0.042*
H1N2	0.852 (2)	0.242 (7)	0.7974 (17)	0.039 (10)*
H1N1	0.651 (3)	0.424 (9)	0.775 (2)	0.078 (15)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0251 (4)	0.0384 (5)	0.0310 (5)	0.0068 (3)	0.0211 (4)	0.0085 (3)
O2	0.0175 (11)	0.0343 (12)	0.0359 (12)	0.0052 (9)	0.0189 (10)	0.0070 (9)
N1	0.0166 (13)	0.0302 (14)	0.0268 (14)	0.0042 (10)	0.0157 (12)	0.0023 (10)
N2	0.0151 (12)	0.0249 (14)	0.0253 (14)	0.0030 (10)	0.0127 (11)	0.0040 (10)
N3	0.0185 (13)	0.0269 (14)	0.0266 (14)	0.0036 (10)	0.0149 (11)	0.0030 (10)
C1	0.0173 (14)	0.0322 (17)	0.0246 (16)	-0.0003 (12)	0.0140 (13)	0.0001 (12)
C2	0.0212 (15)	0.0317 (17)	0.0258 (16)	0.0013 (12)	0.0156 (13)	0.0055 (13)
C3	0.0184 (15)	0.0252 (16)	0.0234 (16)	0.0012 (12)	0.0114 (13)	-0.0001 (12)
C4	0.0180 (15)	0.0286 (16)	0.0311 (17)	0.0010 (12)	0.0163 (13)	0.0005 (13)
C5	0.0235 (16)	0.0284 (17)	0.0273 (16)	0.0017 (12)	0.0180 (14)	0.0038 (13)
C6	0.0154 (14)	0.0229 (15)	0.0214 (15)	-0.0028 (11)	0.0104 (12)	-0.0031 (11)
C7	0.0165 (14)	0.0234 (15)	0.0212 (15)	-0.0030 (11)	0.0109 (12)	-0.0029 (12)
C8	0.0173 (14)	0.0265 (16)	0.0222 (15)	0.0005 (12)	0.0116 (12)	0.0002 (12)
C9	0.0208 (15)	0.0247 (16)	0.0303 (17)	0.0046 (12)	0.0169 (14)	0.0056 (12)
C10	0.0180 (15)	0.0315 (17)	0.0238 (16)	0.0037 (12)	0.0124 (13)	0.0037 (12)
C11	0.0156 (15)	0.0248 (16)	0.0285 (16)	0.0051 (12)	0.0137 (13)	0.0030 (12)
C12	0.0216 (16)	0.0360 (18)	0.0268 (17)	-0.0014 (13)	0.0137 (14)	-0.0002 (13)
C13	0.0214 (16)	0.0293 (17)	0.044 (2)	-0.0020 (13)	0.0185 (15)	-0.0035 (14)
C14	0.0265 (17)	0.0298 (18)	0.052 (2)	0.0030 (14)	0.0292 (17)	0.0059 (15)
C15	0.0246 (16)	0.0330 (18)	0.0327 (17)	0.0063 (13)	0.0220 (14)	0.0052 (13)
C16	0.0229 (16)	0.0327 (18)	0.0286 (17)	0.0048 (13)	0.0150 (14)	0.0027 (13)

Geometric parameters (\AA , $\text{^{\circ}}$)

S1—C8	1.678 (3)	C5—H5A	0.9500
O2—C7	1.228 (3)	C6—C7	1.493 (4)
N1—C7	1.376 (4)	C9—C10	1.523 (4)
N1—C8	1.397 (4)	C9—H9A	0.9900
N1—H1N1	0.80 (5)	C9—H9B	0.9900
N2—C8	1.325 (3)	C10—C11	1.519 (4)
N2—C9	1.460 (3)	C10—H10A	0.9900
N2—H1N2	0.87 (3)	C10—H10B	0.9900
N3—C11	1.335 (4)	C11—C12	1.392 (4)
N3—C15	1.351 (4)	C12—C13	1.391 (4)
C1—C2	1.376 (4)	C12—H12A	0.9500
C1—C6	1.393 (4)	C13—C14	1.374 (4)
C1—H1A	0.9500	C13—H13A	0.9500
C2—C3	1.402 (4)	C14—C15	1.386 (4)
C2—H2A	0.9500	C14—H14A	0.9500
C3—C4	1.390 (4)	C15—H15A	0.9500
C3—C16	1.502 (4)	C16—H16C	0.9800
C4—C5	1.391 (4)	C16—H16D	0.9800
C4—H4A	0.9500	C16—H16A	0.9800
C5—C6	1.399 (4)		
C7—N1—C8	128.8 (2)	N2—C9—H9A	109.5
C7—N1—H1N1	119 (3)	C10—C9—H9A	109.5
C8—N1—H1N1	111 (3)	N2—C9—H9B	109.5
C8—N2—C9	125.0 (2)	C10—C9—H9B	109.5
C8—N2—H1N2	113 (2)	H9A—C9—H9B	108.1
C9—N2—H1N2	121 (2)	C11—C10—C9	114.1 (2)
C11—N3—C15	117.7 (3)	C11—C10—H10A	108.7
C2—C1—C6	121.1 (3)	C9—C10—H10A	108.7
C2—C1—H1A	119.4	C11—C10—H10B	108.7
C6—C1—H1A	119.4	C9—C10—H10B	108.7
C1—C2—C3	120.9 (3)	H10A—C10—H10B	107.6
C1—C2—H2A	119.5	N3—C11—C12	122.6 (3)
C3—C2—H2A	119.5	N3—C11—C10	117.9 (3)
C4—C3—C2	117.8 (3)	C12—C11—C10	119.5 (3)
C4—C3—C16	121.4 (3)	C13—C12—C11	118.7 (3)
C2—C3—C16	120.8 (3)	C13—C12—H12A	120.6
C3—C4—C5	121.8 (3)	C11—C12—H12A	120.6
C3—C4—H4A	119.1	C14—C13—C12	119.3 (3)
C5—C4—H4A	119.1	C14—C13—H13A	120.3
C4—C5—C6	119.7 (3)	C12—C13—H13A	120.3
C4—C5—H5A	120.2	C13—C14—C15	118.4 (3)
C6—C5—H5A	120.2	C13—C14—H14A	120.8
C1—C6—C5	118.7 (3)	C15—C14—H14A	120.8
C1—C6—C7	116.4 (2)	N3—C15—C14	123.3 (3)
C5—C6—C7	124.9 (2)	N3—C15—H15A	118.4

O2—C7—N1	122.0 (3)	C14—C15—H15A	118.4
O2—C7—C6	121.0 (3)	C3—C16—H16C	109.5
N1—C7—C6	117.0 (2)	C3—C16—H16D	109.5
N2—C8—N1	115.7 (2)	H16C—C16—H16D	109.5
N2—C8—S1	126.4 (2)	C3—C16—H16A	109.5
N1—C8—S1	117.8 (2)	H16C—C16—H16A	109.5
N2—C9—C10	110.5 (2)	H16D—C16—H16A	109.5
C6—C1—C2—C3	−0.5 (5)	C9—N2—C8—N1	173.5 (3)
C1—C2—C3—C4	1.0 (4)	C9—N2—C8—S1	−6.9 (4)
C1—C2—C3—C16	−178.7 (3)	C7—N1—C8—N2	−4.4 (4)
C2—C3—C4—C5	−0.9 (4)	C7—N1—C8—S1	175.9 (2)
C16—C3—C4—C5	178.8 (3)	C8—N2—C9—C10	−98.0 (3)
C3—C4—C5—C6	0.4 (4)	N2—C9—C10—C11	−63.9 (3)
C2—C1—C6—C5	−0.1 (4)	C15—N3—C11—C12	−1.0 (4)
C2—C1—C6—C7	179.8 (3)	C15—N3—C11—C10	177.2 (3)
C4—C5—C6—C1	0.1 (4)	C9—C10—C11—N3	1.1 (4)
C4—C5—C6—C7	−179.7 (3)	C9—C10—C11—C12	179.4 (3)
C8—N1—C7—O2	2.1 (5)	N3—C11—C12—C13	1.5 (4)
C8—N1—C7—C6	−178.7 (3)	C10—C11—C12—C13	−176.6 (3)
C1—C6—C7—O2	0.8 (4)	C11—C12—C13—C14	−0.5 (4)
C5—C6—C7—O2	−179.3 (3)	C12—C13—C14—C15	−0.9 (4)
C1—C6—C7—N1	−178.3 (3)	C11—N3—C15—C14	−0.5 (4)
C5—C6—C7—N1	1.5 (4)	C13—C14—C15—N3	1.5 (4)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H1N2···O2	0.87 (4)	1.90 (3)	2.645 (3)	143 (3)
N2—H1N2···N3	0.87 (4)	2.41 (4)	2.860 (4)	113 (3)
C1—H1A···O2	0.95	2.42	2.751 (4)	100
C9—H9B···S1	0.99	2.72	3.166 (4)	108
C14—H14A···O2 ⁱ	0.95	2.51	3.421 (4)	161

Symmetry code: (i) $-x+2, y+1/2, -z+3/2$.