

Fluorescence and nonphotochemical quenching responses to simulated vertical mixing in the marine diatom *Thalassiosira weissflogii*

Allen J. Milligan*, Uriel A. Aparicio, Michael J. Behrenfeld

Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon 97331, USA

ABSTRACT: The fluorescence signal emitted from phytoplankton exposed to natural sunlight has been considered a potentially useful tool to examine phytoplankton physiology from *in situ* radiometers and satellites, but variability in the fluorescence signal is confounded by non-photochemical quenching (NPQ). This pathway dissipates excitation energy as heat rather than fluorescence. It is necessary to first correct for this NPQ signal in order to extract physiological information, such as rates or nutrient stresses. Here we examine the fluorescence–irradiance response and NPQ in a marine diatom grown under simulated vertical mixing. Contrary to previous studies, we find that low-light-acclimated cultures express greater NPQ than high-light-acclimated cultures. Our results differ from previous studies because we employ realistic vertical mixing, rather than constant light conditions. We also find that fluorescence yields do not saturate at high-light levels. This observation impacts how we might apply NPQ corrections to radiometer and satellite data.

KEY WORDS: Fluorescence · Non-photochemical quenching · Fluctuating light · Diatom

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Oxygenic photosynthesis involves light harvesting to form chemical energy through the action of photosystem II (PSII) and photosystem I (PSI). A small fraction (~2%) of the solar energy absorbed by phytoplankton for photosynthesis is re-radiated by chlorophyll pigments as fluorescence. Both photosystems contribute to total cellular chlorophyll concentration, but *in vivo* fluorescence is dominated by emissions from antennae pigments at the core of PSII (Krause & Weis 1991). Under natural sunlight conditions, chlorophyll fluorescence can be detected in the subsurface- and above-surface upwelling radiance spectrum (Morel & Prieur 1977, Neville & Gower 1977). This sun-induced natural fluorescence feature has been proposed as a tool for deriving phytoplankton photosynthetic rates (Kiefer 1973, Falkowski & Kiefer 1985, Kiefer et al. 1989) and physiological vari-

ability (Letelier et al. 1997, Morrison 2003, Schallenberg et al. 2008).

A wide range of physiological factors influence fluorescence, but the most important of these are photochemical quenching (qP) and non-photochemical quenching (NPQ). Photochemical quenching is associated with charge separation in the PSII reaction center, while NPQ is a protective response to high-light exposure that is broadly expressed in phytoplankton (Niyogi et al. 1997, Lavaud et al. 2007, Miloslavina et al. 2009). NPQ reduces fluorescence and encompasses a variety of physiological mechanisms that dissipate excess absorbed light energy as heat before it reaches the core of PSII. One mechanism of NPQ is a rapid response (seconds) to super-saturating light that is dependent on a thylakoid proton gradient and photoprotective pigments (xanthophylls) (Holt et al. 2004). This form of NPQ is referred to as qE. In diatoms, qE is likely achieved through conformational changes in

*Email: allen.milligan@science.oregonstate.edu

PSII and conversion of diadinoxanthin to diatoxanthin (a fluorescence quencher) in a single de-epoxidation reaction (Lavaud et al. 2002, Lavaud & Kroth 2006, Grouneva et al. 2009).

A second form of NPQ is a rebalancing of light energy between the 2 photosystems and involves the physical movement of light harvesting antennae complexes from PSII to PSI. This process, termed a state transition or qT, functions on longer time scales (minutes) than qE and allows movement of a pool of chlorophyll from fluorescence-emitting PSII to weakly-fluorescent PSI. There is no direct evidence for state transitions in diatoms (Owens 1986, Ting & Owens 1994, Miloslavina et al. 2009). A final form of NPQ involves quenching at the PSII reaction center and can be associated with either protein damage or reversible down-regulation of PSII. We distinguish the former form as qI (I for inhibition; Holt et al. 2004) and the later, reversible form as qRC (RC for reaction center). qRC is thought to involve a loss of excitation energy when PSII becomes inactive and functions as a quencher rather than a photochemical trap (Öquist et al. 1992, Lee et al. 2001). There is conflicting evidence for (Eisenstadt et al. 2008) and against (Goss et al. 2006) qRC in diatoms. Both qI and qRC function independently of qE and qT.

Relaxation of NPQ occurs as light is reduced or in darkness and the recovery time is a key characteristic often used to distinguish different quenching processes (Horton & Hague 1988). In vascular plants and green algae, qE relaxes on time scales of seconds to minutes, qT relaxes on the order of 10 min, qRC relaxes on time scales of minutes to an hour, and qI relaxes on the order of hours due to the requirement for protein synthesis to rebuild PSII reaction centers. In diatoms by contrast, qE relaxes on several time scales, from seconds to several 10s of min and relaxation can require low light (Goss et al. 2006, Grouneva et al. 2007, 2008).

Since the emergence of fluorescence studies in oceanographic settings, it has been recognized that NPQ is an important variable affecting the fluorescence quantum yield (i.e. fluorescence emitted per unit light absorbed). Accordingly, NPQ impacts the relationship between chlorophyll concentration and fluorescence in the water column (Berman 1972) and total measured fluorescence between clear and cloudy days (Neville and Gower 1977). Furthermore, removing the effects of NPQ on fluorescence allows additional information to be retrieved on phytoplankton physiology and photosynthesis from the remaining variability in the corrected data (e.g. Falkowski & Kiefer 1985, Morrison 2003, Westberry

& Siegel 2003). Cullen & Lewis (1995) have called for studies that focus on NPQ and fluorescence responses of phytoplankton grown in high light. However, few studies to date have measured fluorescence in cultures grown under high-light intensities delivered as diurnal or fluctuating light (Laney et al. 2005, Dimier et al. 2009, van de Poll et al. 2010). Understanding fluorescence variability under these conditions is important because they better represent the variable light conditions of natural systems. Proper representation of NPQ for the natural light environment is particularly important when evaluating satellite fluorescence data because these measurements register fluorescence properties of phytoplankton in the highly dynamic light environment of the surface mixed layer and are collected near noon and only under clear skies when incident irradiance is maximal (Behrenfeld et al. 2009).

Previous laboratory studies using constant light conditions have suggested that high-light-acclimated phytoplankton exhibit a strong NPQ response, while low-light-acclimated phytoplankton have reduced NPQ capacity (Niyogi et al. 1997, Cruz & Serodio 2008, Ragni et al. 2008). Here we examine NPQ responses of *Thalassiosira weissflogii* grown under light conditions simulating natural surface layers of shallow and deep vertical mixing. Importantly, both conditions encompassed exposures to the same maximal light levels (i.e. the treatments simulated 2 mixing regimes with identical surface light levels). We find that low-light-acclimated phytoplankton (i.e. deep mixing regime) exhibit higher NPQ than high-light-acclimated phytoplankton (i.e. shallow mixing). This finding thus directly contradicts previous conclusions regarding NPQ capacity and likely reflects the fact that earlier experiments have rarely employed growth conditions of fluctuating light with saturation of photochemistry well below the maximum light encountered by the cells. We also find that the fluorescence-versus-incident irradiance (iPAR) response curves do not saturate, suggesting the presence of a chl a fluorescence source that is insensitive to NPQ. This behavior of the fluorescence-iPAR relationship impacts the form of NPQ corrections applied to natural fluorescence data.

MATERIALS AND METHODS

Culture conditions

A culture of the marine diatom *Thalassiosira weissflogii* (clone Actin, CCMP 1336) was obtained

from the Center for the Culture of Marine Protozoa. Cultures (1L) were grown under simulated shallow and deep mixing using natural seawater (from coastal Oregon) amended with f/2 nutrients. The medium was sterilized by autoclaving. Temperature was maintained at 20°C. Light (12 h d⁻¹) was provided by high intensity fluorescent tubes (55 W Dulux, Sylvania) with variable ballasts that were controlled by computer using Labview (ver. 6.0, National Instruments). Specifically, the diurnal light cycle (i.e. the light cycle representing daily changes in incident light at the surface ocean) ranged from 42 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ to a maximum of 1550 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ for both treatments (Fig. 1). Superimposed on this sinusoidal function were higher-frequency variations in light simulating shallow or deep vertical mixing with a realistic period of 1.5 h (D'Asaro 2003) (Fig. 1). In addition to these variable light conditions, we also examined the light dependent recovery of NPQ in cultures grown in continuous light of 500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. Scalar irradiance in all culture vessels was continuously monitored with a 4 π sensor (Biospherical). Cultures were acclimated to each light condition for up to 6 mo and maintained at steady state growth for at least 2 wk prior to sampling. All sampling occurred early in the early exponential growth phase (cell abundance < 3 $\times 10^4 \text{ ml}^{-1}$).

Physiological properties

Coincident with NPQ measurements, we also collected samples for determination of chl *a* concentration, pigment spectral absorption, cell concentration, growth rate, and photosynthesis-irradiance relationships. Samples (20 to 30 ml) for chl *a* concentration were collected on glass fiber filters (GFF Whatman). Chlorophyll was extracted overnight with 90% acetone and determined spectrophotometrically following the methods of Jeffrey & Humphrey (1975). Cell concentrations were measured using a Coulter (Multisizer 3) particle counter fitted with a 70 μm aperture. Daily growth rates were calculated from sequential cell counts using the formula:

$$\mu = \frac{\ln(C_1/C_0)}{t_1 - t_0} \quad (1)$$

where μ is the intrinsic rate of growth in units of d⁻¹, C_0 is initial (t_0) cell concentration, and C_1 is the cell concentration 1 d later (t_1).

Phytoplankton pigment absorption spectra were determined for each culture and light level using the

glass fiber filter pad technique on a dual beam spectrophotometer (Shimadzu, UV2401 PC) and corrected for pathlength amplification following NASA protocols (Mitchell et al. 2003). The spectrally-averaged chl *a*-specific absorption cross-section (\bar{a}^*) was calculated following Falkowski et al. (1985) for both the fluorescent tube and fluorometer blue light LED (see below) spectra:

$$\bar{a}^* = \int_{400}^{750} a^*(\lambda) E(\lambda) d\lambda \cdot \left[\int_{400}^{750} [E(\lambda) d\lambda] \right]^{-1} \quad (2)$$

where $E(\lambda)$ is the spectral irradiance of the light source and $a^*(\lambda)$ is the chl *a*-specific spectral absorption cross-section calculated from:

$$a^*(\lambda) = 2.3 \cdot A(\lambda) / (\text{Chl}) \quad (3)$$

where $A(\lambda)$ is the sample spectral absorbance, 2.3 is the conversion factor from log₁₀ to ln, and Chl is the chl *a* concentration (mg m^{-3}).

Photosynthesis-irradiance (PE) relationships for samples collected at noon were determined from 20 min incubations with ¹⁴C bicarbonate following the photosynthetron protocol of Lewis & Smith (1983). During each incubation, 10 irradiances were used, with the resultant data fit to the equation of Jassby & Platt (1976):

$$P^B = P_{\text{max}}^B \tanh(\alpha^B E / P_{\text{max}}^B) \quad (4)$$

where P^B is the photosynthetic rate normalized to chl *a*, P_{max}^B is the maximum chl *a* specific photosynthetic rate, α^B is the chl *a* specific initial slope of the PE relationship, and E is the photosynthetically active radiation (PAR: 400 to 700 nm).




Fig. 1. Photoperiod (12 h) light regimes for simulated deep (dashed line) and shallow (solid line) mixing. Light was controlled by a variable-output ballast recreating both the sinusoidal function of daylight and a vertical mixing function with a period of 1.5 h (D'Asaro 2003). Maximum and minimum light = 1550 and 42 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, respectively

Fluorescence and NPQ

Fluorescence emission was determined in a temperature controlled (20°C) pulse amplitude modulated (PAM) fluorometer (DUALPAM-100, Heinz Walz) fitted with a photomultiplier detector (DUAL-DPM). For fluorescence yield determinations, a sample was removed from the culture vessel, placed in the fluorometer, and dark acclimated for 25 min. Initial fluorescence (F_0) was then measured using a low-intensity blue LED source (DUAL-DB) running at a fixed frequency of 20 Hz and an intensity of 24 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. The sample was then exposed to a sequence of 20 blue actinic light levels ranging from 6 to 1952 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, with steady-state fluorescence (F_s) monitored continuously. At the end of each 7 min actinic light exposure, F_0 was re-measured during a brief exposure to darkness and then maximum fluorescence (F_m) determined with a saturating red LED flash of intensity of 3000 to 5000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and duration from 400 to 600 ms. The specific intensity and duration of the saturating flash was determined prior to running the full fluorescence yield protocol to ensure saturation. From these measurements, variable fluorescence (F_v/F_m) for each actinic light level was calculated as (Govindjee 2004):

$$F_v/F_m = (F_m - F_0)/F_m \quad (5)$$

where F_m and F_0 are determined at the end of actinic light exposure. Fluorescence yields (F_v) were then calculated for each light level as (Govindjee 2004):

$$F_v = (F_s/F_0) - 1 \quad (6)$$

where F_0 is the F_0 value measured when F_v/F_m was maximal (i.e. when NPQ was fully relaxed; see 'Results') and functions to account for differences in total pigment absorption between samples.

NPQ kinetics were determined using a PAM protocol entailing an initial dark acclimation period of 25 min, followed by an actinic light exposure of 10 min, and finally a second dark period of 35 min. During the first dark period, F_0 was monitored continuously using the low-intensity blue LED described above and F_m measured periodically (to determine F_v/F_m) with a saturating red LED flash (as above). During the subsequent actinic light exposure, the sample was exposed to a single background saturating light level of 1100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ to induce NPQ. Throughout this period, F_s was continuously monitored and maximum fluorescence under actinic light exposure (F_m') periodically determined. NPQ was calculated as (Govindjee 2004):

$$\text{NPQ} = (F_m/F_m') - 1 \quad (7)$$

During the final dark period, F_0 was continuously monitored and the kinetics of fluorescence recovery and NPQ relaxation determined through frequent determinations of F_m .

One of the motivations for these experiments was to characterize phytoplankton fluorescence properties relevant to the interpretation of satellite fluorescence data. Accordingly, samples for fluorescence yield and NPQ kinetics measurements were collected at midday. This sampling strategy ensured that all data were consistent with respect to their position within the photoperiod; however, they differed with respect to the incident light level at the time of sampling (see 'Results' section),

To further examine details of NPQ relaxation, samples from the continuous light cultures (500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) were collected and fluorescence measured during a modified protocol of 3 h darkness, 15 min actinic light (140 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$), and then 20 min darkness. Throughout this protocol, F_0 was monitored continuously and F_m and F_m' measured periodically (as above). Five replicate experiments were conducted.

Finally, we evaluated variability in the magnitude of NPQ over the diurnal cycle in the 2 simulated mixing conditions by conducting fluorescence measurements on samples collected at 2 h intervals over the daytime cycle (5 samples d^{-1} for 3 d). Samples were dark acclimated for 30 min and exposed to 10 actinic light levels for 5 min, with intervening 5 min dark periods. During the dark phase of this protocol, we monitored F_0 continuously and measured F_m (to determine F_v/F_m) just prior to the actinic light exposure. F_m' was measured at the end of each actinic light period to determine NPQ.

RESULTS

Photoacclimation state

Thalassiosira weissflogii grew at equivalent rates (0.95 to 0.96 d^{-1}) in the 2 light conditions simulating shallow and deep mixing. This result implies that cells in the deep-mixing regime adjusted their light harvesting capacity to fully compensate for their lower daily light exposure. This acclimation was observed as a 3-fold increase in cellular chl *a* concentration in the low-light (deep mixing) treatment (Table 1), which consequently decreased the chl *a* normalized absorption cross section ($\bar{\alpha}^*$) in a manner

Table 1. *Thalassiosira weissflogii*. Growth and photosynthetic parameters for cultures grown under simulated shallow- and deep-mixing light regimes. Absorption cross sections are calculated using spectral distribution for fluorescent lights (\bar{a}^*_{Fluor}) under growth conditions and blue light emitting diodes (\bar{a}^*_{LED}) used as light source in Dual-PAM. Values in parentheses = SD for triplicates

Mixing regime light treatment	Mean light intensity ($\mu\text{mol quanta m}^{-2} \text{s}^{-1}$)	Integrated flux (mol quanta $\text{m}^{-2} \text{d}^{-1}$)	Specific growth rate (d^{-1})	Chl <i>a</i> quota (pg chl <i>a</i> cell $^{-1}$)	Spectrally weighted, chl <i>a</i> normalized absorption cross section ($\text{m}^2 \text{mg chl a}^{-1}$)	
					\bar{a}^*_{Fluor}	\bar{a}^*_{LED}
Shallow mixing						
High light	487	20.8	0.96 (0.26)	1.51 (0.19)	2.44×10^{-3} (0.91)	9.8×10^{-3} (0.19)
Deep mixing						
Low light	138	5.9	0.95 (0.08)	4.68 (0.10)	2.02×10^{-3} (0.37)	6.92×10^{-3} (0.14)




Fig. 2. *Thalassiosira weissflogii*. Photosynthesis vs. irradiance (PE) response curves for low- and high-light-acclimated cultures. Data are fit to Eq. (4) (Jassby & Platt 1976). Low light: $\alpha^B = 2.29$ ($\mu\text{mol C mg chl a h}^{-1}$ [$\mu\text{mol photons m}^{-2} \text{s}^{-1}$] $^{-1}$), photosynthetic rate $P_{\text{max}} = 943$ ($\mu\text{mol C mg chl a h}^{-1}$), saturation irradiance $E_k = 411$ ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$). High light: $\alpha^B = 2.43$ ($\mu\text{mol C mg chl a h}^{-1}$ [$\mu\text{mol photons m}^{-2} \text{s}^{-1}$] $^{-1}$), $P_{\text{max}}^B = 2185$ ($\mu\text{mol C mg chl a h}^{-1}$), $E_k = 899$ ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$). If we assume that P_{max}^B cannot exceed measured values, and hold P_{max}^B at 1720 and 910 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ for high- and low-light respectively, E_k becomes 707 and 398 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, respectively

consistent with increased self-shading (Falkowski et al. 1985). Photosynthesis versus irradiance curves yielded similar initial slopes (α^B) for the 2 mixing regimes, but maximum rates (P_{max}^B) that were nearly 2-fold greater in the high-light-acclimated (shallow mixing) cells (Fig. 2). As growth rates were equivalent between the 2 treatments (i.e. similar carbon fixing capacity), this difference in P_{max}^B is driven in part by the normalization of P_{max} to the lower cellular chl *a* in high light (Table 1).




Fig. 3. *Thalassiosira weissflogii*. (A) Fluorescence yields in low- and high-light-acclimated cultures. Note: F_0 used in the calculation of F_Y is determined at maximum F_V/F_m . (B) Dark measured variable fluorescence (F_V/F_m) for same treatments as (A)

Fluorescence yield

Noon-time PAM-derived fluorescence yields (F_Y) were similar in both light treatments (Fig. 3A). This finding was not expected because total light absorption differs between high- and low-light-acclimated cultures (Table 1). Specifically, taking the product of

cellular chl *a* and \bar{a}^* for the blue light LED spectrum (Table 1), we obtained cellular absorption cross section values of 14.8×10^{-12} and $32.4 \times 10^{-12} \text{ m}^2 \text{ cell}^{-1}$ for the high-light and low-light treatments respectively. This 2.2 fold higher light absorption in low-light cells (1) is driven primarily through changes in pigment concentration, (2) should result in higher fluorescence yields for low-light-acclimated cultures, and (3) suggests that there are significant differences in NPQ capacity in these treatments.

As noted in the 'Materials and methods' section above, the calculation of F_Y requires assessment of F_o when NPQ is fully relaxed. In *Thalassiosira weissflogii*, NPQ had a greater impact on F_m than F_o . Thus, observed changes in F_v/F_m (Eq. 5) can be used to evaluate the NPQ relaxation state. During our fluorescence yield experiments, we observed that F_v/F_m rose to a maximum (0.63 to 0.66) at 70 and 126 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ for low and high light, respectively, and then declined at higher light levels (Fig. 3B). Although our F_Y results are calculated using F_o values at maximum F_v/F_m , this initial rise in F_v/F_m indicates that NPQ was still relaxing at these lowest light levels. In other words, the assumption of negligible NPQ after 25 min of dark acclimation prior to the actinic light sequence is violated.

NPQ dynamics

Fluorescence properties for low-light and high-light cultures for our NPQ induction-relaxation protocol are shown in Fig. 4A & B, respectively. At the time of sampling (noon), the high-light cultures were receiving near maximum iPAR ($1100 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$) while the low-light cultures were receiving only $42 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$. Given this difference, we would anticipate that the low-light cells would be expressing little NPQ at the time of sampling, while the high-light cells would express high NPQ. Comparison of fluorescence emissions between treatments during the initial 25 min dark period shows that the low-light-acclimated cells had much higher variable fluorescence (F_v/F_m) than the high-light cells (Fig. 4). This finding is consistent with anticipated differences in NPQ, as variable fluorescence values can be high only when NPQ is minimal. Furthermore, the persistently low F_v/F_m values for the high-light cells indicates that NPQ relaxed little during the initial dark period of the PAM protocol.

A common assumption applied in PAM measurements of NPQ is that pre-exposure of phytoplankton samples to a period of darkness before the NPQ-




Fig. 4. *Thalassiosira weissflogii*. Example fluorescence traces of (A) low- and (B) high-light-acclimated cultures sampled at noon (14:10 h). Light intensities at the time of sampling were 42 and $1100 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ for low- and high-light-acclimated cultures, respectively. Samples were first exposed to a dark period of 25 min (black bar on x-axes), during which initial (F_o) and maximum (F_m) fluorescence were monitored. Fluorescence (F_s) and maximum fluorescence (F_m') were then monitored during a 15 min exposure to super-saturating background light ($1100 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$). Finally, fluorescence recovery was observed over a subsequent 20 min dark period (black bar on x-axes). Variable fluorescence (F_v/F_m) values are given for 3 points in each time course (at the beginning of each protocol, after exposure to saturating light, and at the end of the second dark period). Maximum F_v/F_m for *T. weissflogii* under our growth conditions (when NPQ is fully relaxed) = 0.6 to 0.65 (horizontal dashed line)

induction protocol (i.e. exposure to high actinic light) is adequate for full relaxation of pre-existing NPQ. Results in Fig. 4B show that this assumption was not valid for *Thalassiosira weissflogii* under our conditions. This retention of NPQ through the initial dark period is clear from the difference between the F_m values and observed F_m associated with maximal F_v/F_m (horizontal dashed lines in Fig. 4).

In both treatments, the rise in F_m during the initial dark period is small. However, during the dark period following actinic light exposure, a more dramatic rise in F_m is observed (Fig. 4), suggesting that




Fig. 5. *Thalassiosira weissflogii*. Expanded time-course of dark relaxation of non-photochemical quenching (NPQ) in cultures grown under continuous light of $500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$. Maximum fluorescence rises only gradually for the first 3 h of darkness (black bar on x-axis), but then quickly rises with a background illumination of $140 \mu\text{mol photons m}^{-2} \text{s}^{-1}$. Values: slopes of each regression line. Each point is average of 5 replicates. Shaded area: 95% CI

NPQ relaxation in *Thalassiosira weissflogii* was enhanced by previous light exposure. This recovery is greatest during the first 6 to 8 min following actinic light for both treatments, a time scale consistent with qE relaxation (Holt et al. 2004). It is important to note that the incomplete NPQ relaxation in the initial dark period has consequences for derived fluorescence properties. Of particular interest, it can result in underestimates, or even negative values, of calculated NPQ (i.e. $F_m < F_m'$; Eq. 7).

We further examined the light dependency of NPQ relaxation using samples from our constant light condition. Despite a prolonged 3 h initial dark period, maximum fluorescence exhibited little recovery (increasing from ~ 0.7 to 0.8) (Fig. 5). However, a marked increase in NPQ recovery was observed upon exposure to low-level actinic irradiance. Recovery, slowed again during subsequent exposure to darkness (Fig. 5). This result demonstrates that complete relaxation of NPQ in *Thalassiosira weissflogii* requires light energy when cells are harvested from growth conditions that induce NPQ.

NPQ capacity

Fluorescence properties illustrated in Figs. 3 & 4 are all for samples collected at midday. We additionally investigated NPQ capacities over the diurnal

light cycle. As described in the 'Materials and methods' section, these assessments entailed sequential light-dark exposures that generally ensured NPQ relaxation and high F_v/F_m within 40 to 50 min into the protocol. NPQ capacities were calculated using F_m' values observed at each actinic light level and the F_m value associated with the maximum F_v/F_m value exceeding 0.6 (Eq. 7). We found NPQ capacity to be independent of diurnal sampling time (thus recent light history) for the low light cultures. The relationship between NPQ and iPAR for the pooled data is shown by the black symbols in Fig. 6. For the high light treatment, full NPQ relaxation ($F_v/F_m > 0.6$) was achieved on 3 occasions. Pooling these data yields the NPQ-iPAR relationship shown by open symbols in Fig. 6. The 2 growth conditions both give relationships that begin with declining NPQ at low-light levels because of incomplete relaxation early in the protocol, as discussed above. NPQ values then begin to rise above 70 and $240 \mu\text{mol m}^{-2} \text{s}^{-1}$ for low- and high-light *Thalassiosira weissflogii*, respectively (Fig. 6). The 2.3-fold higher NPQ for the low-light-acclimated cells is consistent with the 2.2-fold higher cellular absorption cross section calculated for low-light-acclimated cells. This finding suggests that the lack of difference in F_y relationships between treatments (Fig. 3A) was due to enhanced NPQ in the low-light treatment.




Fig. 6. *Thalassiosira weissflogii*. Comparison of non-photochemical quenching (NPQ) in low- and high-light-acclimated cultures plotted against incident irradiance (iPAR). Low-light represents 5 light curves (1 every 2 h) collected over a period of 3 d (each point is average of 15 values) under semicontinuous culture conditions. For high-light data only 3 light curves (3 out of 15 data sets) met the criteria of $F_v/F_m > 0.6$. Error bars = SD

DISCUSSION

Field and satellite measurements of fluorescence are a potentially powerful tool for optically quantifying chl *a* concentrations and investigating variability in phytoplankton physiology. These applications, however, require a thorough understanding of the fluorescence–iPAR relationship because of the strong influence of NPQ on the quantum yield of fluorescence. This problem is particularly acute when interpreting satellite-based retrievals of fluorescence, as these data are acquired under clear-sky, near-noon conditions when incident light is maximal. We examined fluorescence–iPAR relationships with a culture system that simulated the vertically-mixing, upper water column to test whether low-light-acclimated phytoplankton (i.e. deep mixing) have greater NPQ than those that are high-light-acclimated (i.e. shallow mixing). This approach differs from many previous NPQ investigations that have employed the unnatural conditions of constant low and/or high light. Importantly, both light environments used in the current study saturated growth of *Thalassiosira weissflogii*, but the different simulated mixing regimes yielded 2 distinctly different photoacclimation states, as evidenced by changes in chl *a* cell quotas and PE relationships.

Fluorescence response to light acclimation

Close inspection of data in Fig. 2 shows a separation in PE curves between the 2 simulated mixing conditions beginning around $180 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, well below the light level for saturation of carbon fixation (E_{k-C}). Measurements of F_v/F_m for these treatments likewise revealed a progressive decrease above $\sim 150 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, indicating increasing NPQ (Fig. 3B). This early induction of NPQ below the E_{k-C} has also been reported by Laney et al. (2005) and Ting & Owens (1993). These findings have 2 important implications. First, the saturation irradiance for fluorescence (E_{k-F}) is not a quantitative measure of E_{k-C} (Laney et al. 2005). Second, modeling of NPQ for corrections to satellite fluorescence data (e.g. Behrenfeld et al. 2009) should describe NPQ induction as a function of E_{k-F} , rather than E_{k-C} .

NPQ dynamics

Results from our F_v experiments also demonstrate an increase in F_v/F_m at the very lowest light levels

(Fig. 3B). This result would not be expected if NPQ was fully relaxed prior to the actinic light exposures (Ting & Owens 1994). Thus our initial 25 min dark period was not sufficient for complete relaxation. This finding is noteworthy because dark acclimation times of 5 to 30 min are often used to relax NPQ in diatom studies (Lavaud et al. 2007, Miloslavina et al. 2009). Our additional experiments with cells acclimated to constant light show that low-light pretreatment is required to completely relax NPQ (Fig. 5), in agreement with several recent diatom-specific NPQ studies (Ting & Owens 1993, Grouneva et al. 2008, 2009, Miloslavina et al. 2009). The failure of F_m values to rise to maximal levels during dark pretreatment (Fig. 4) is thought to be due to a light requirement for qE relaxation in diatoms. More specifically, NADPH is required to convert the quenching xanthophyll, diatoxanthin, to its epoxidated non-quenching state, diadinoxanthin (Grouneva et al. 2009). This reductant is supplied through photosynthesis, creating a direct link between NPQ relaxation and availability of light energy (Jakob et al. 2001, Grouneva et al. 2009). There is also evidence that diatoms induce NPQ in darkness through chlororespiratory reduction of the plastoquinone pool and establishment of a weak transthylakoid pH gradient (Ting & Owens 1993, Jakob et al. 2001, Dijkman & Kroon 2002). This electron flow is much slower than PSI turnover, such that exposure to low light reoxidizes the plastoquinone pool and relaxes NPQ (Grouneva et al. 2009).

NPQ capacity

One issue of particular interest during this study was the relationship between NPQ and growth irradiance conditions. Behrenfeld et al. (2009) proposed that the fluorescence quantum yield at a given iPAR value will be dependent on the photoacclimation state of the phytoplankton population. They envisioned low-light-acclimated cells (i.e. low E_{k-F}) as showing earlier induction of NPQ and thus lower fluorescence emission at a given iPAR than high-light-acclimated cells (Fig. 7). Our findings demonstrate a greater capacity and earlier induction of NPQ in low-light-acclimated cells than high-light, in agreement with the predictions of Behrenfeld et al. (2009). Greater NPQ capacity in low-light-acclimated cells may at first appear to contradict results from previous algal NPQ studies, where NPQ capacity was greatest in high-light treatments (Niyogi et al. 1997, Cruz & Serodio 2008, Ragni et al. 2008). However, these




Fig. 7. *Thalassiosira weissflogii*. Fluorescence response of low- and high-light-acclimated cultures normalized to iPAR and hypothesized curves of quantum yield for 2 different photoacclimation states (Behrenfeld et al. 2009). With increasing iPAR, there is a rise in fluorescence as reaction centers close. At E_{k-F} , light reactions are saturated and fluorescence declines as an inverse function of incident light ($iPAR^{-1}$) as each additional photon is lost as heat through NPQ. Data points are from light levels $>E_{k-F}$. Data $<E_{k-F}$ do not follow a power function. Low-light fit (dashed line): $Y = 0.037 \times iPAR^{-0.31}$; High-light fit: $Y = 0.155 \times iPAR^{-0.54}$. 95% CI for exponents = 0.1 and 0.04 for low- and high-light, respectively

earlier studies used constant high- and low-light conditions and were focused on resolving NPQ mechanisms, rather than characterizing the fluorescence–iPAR relationship. Of the few studies that have employed fluctuating light, the focus has generally been on examining short-term responses to increased light flux (Lavaud et al. 2007, van de Poll et al. 2007, 2010), rather than steady-state fluorescence characteristics of fully acclimated cultures (Dimier et al. 2009). We therefore propose that the apparent discrepancies between our results and earlier findings largely reflect differences in the culturing conditions employed. Specifically, phytoplankton maintained at a constant low-light level have little need to protect PSII from photodamage. In contrast, growth under constant high-light requires a much larger NPQ capacity to safely dissipate the excess absorbed light energy. In a variable light regime ranging from sub- to super-saturating intensities (as experienced by a vertically mixed population), phytoplankton must simultaneously have attributes of both high- and low-light acclimated cells. In particular and as observed here for *Thalassiosira weissflogii*, such conditions require both enhanced light absorption capacity (i.e. increased cellular pigment levels) for periods

of low-light exposure near the bottom of the mixing layer and high NPQ capacity to dissipate the great excess of absorbed light energy experienced near noon at the surface of the mixing layer.

We find that the magnitude of NPQ (maximum of 0.5) in *Thalassiosira weissflogii* is lower than previous studies but within the general range for diatoms (0.5 to 2.5) (Lavaud et al. 2007). Variability in NPQ magnitude within a species can be large and is related to growth conditions. For instance, NPQ in *Phaeodactylum tricornutum* ranges from a value of 2.3 for cultures grown under a 1:1 h light:dark cycle at $45 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ (Lavaud et al. 2007) to a value of 6.0 for cultures grown under a 5:55 min light:dark cycle at $40 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ (Ruban et al. 2004). These growth conditions are, of course, far from natural and highlight the fact that many NPQ studies focus on resolving NPQ mechanisms rather than NPQ behavior in natural settings.

Application to fluorescence studies

Our results show that the F_v –iPAR relationship does not saturate in *Thalassiosira weissflogii* (Fig. 3). This finding agrees with the laboratory and field observations of Laney et al. (2005) and Morrison (2003), but contrasts with field observations of Schalenberg et al. (2008). In this latter study, saturation of *in situ* fluorescence was observed in all profiles but one. However, it appears that non-saturating fluorescence is the more common behavior and it implies that there is an NPQ-insensitive source of fluorescence that increases with iPAR. One likely candidate for this residual fluorescence is PSI. PSI has no variable fluorescence and exciton trapping is more efficient than in PSII, but it still gives a low, constant background fluorescence signal (but see Joly & Carpentier 2007). To evaluate the PSI contribution necessary to explain our nonsaturating F_v –iPAR relationships (Fig. 3), we employed the fluorescence model of Behrenfeld et al. (2009). We find that PSI contributions of 8.5 and 11.9% give best fits to our data for high- and low-light cultures, respectively. These values are in close agreement with reports that PSI fluorescence is $\sim 10\%$ of total cellular fluorescence (Franck et al. 2002). Furthermore, the slightly higher PSI contribution for the low-light condition is consistent with earlier findings suggesting that low-light acclimation can cause a decrease in PSII:PSI (Strzepek & Harrison 2004).

It is important to note that the contribution of PSI to cellular fluorescence is wavelength dependent and

therefore dependant on the type of instrument and filter sets used. Peak fluorescence emission of PSI *in vivo* is ~722 nm and is minimal around the *in vivo* fluorescence peak of PSII (684 nm) (Franck et al. 2002, 2003). The fluorometer used in this study is equipped with a 665 nm cut-off filter, so all PSI fluorescence (>665 nm) is included in the fluorescence measurement. For fluorescence studies in marine systems, radiometers are often employed with narrow band (683 with 10 nm half-maximum band-width) fluorescence detection (Morrison 2003, Schallenberg et al. 2008). The Moderate-resolution Imaging Spectroradiometer (MODIS) ocean color sensor uses a similar narrow band method although it is centered at 678 nm to avoid an atmospheric oxygen absorption band (Behrenfeld et al. 2009). Using narrow band-pass detection will minimize the fluorescence signal from PSI, but even at 685 nm the PSI contribution can be as high as 10% at low iPAR (Franck et al. 2002). The contribution of PSI to total fluorescence under brief high iPAR exposure is very low (~0.2%), but under sustained high iPAR it increases due to NPQ of PSII fluorescence (Franck et al. 2002, Govindjee 2004). Given that field-measured quantum yields of fluorescence decline roughly 5-fold with NPQ, we might expect the contribution of PSI fluorescence to be an important contributor to total fluorescence (Morrison 2003). A better understanding of the contribution of PSI to fluorescence in phytoplankton is clearly needed.

Satellite-derived natural fluorescence is by necessity measured in the presence of high photon flux and, as such, is influenced by the multiple NPQ pathways. In order to use these data as a physiological indicator, it is necessary to first correct for the confounding NPQ signal. Behrenfeld et al. (2009) applied an inverse incident light function ($1/iPAR$) to MODIS fluorescence data as a first-order NPQ correction, but recognized that this function assumed NPQ behaved as a perfect protective quencher. They also noted that the NPQ correction should also account for photoacclimation and that neglecting this component likely impacted derived fluorescence quantum yields at high latitudes and under deep mixing conditions. Findings of the current study provide insight on the concerns raised by Behrenfeld et al. (2009) regarding the inverse-light function applied for NPQ correction. Our results demonstrate enhanced NPQ capacity in low-light-acclimated cells and illustrate the non-saturable nature of F_Y with increasing light. This latter phenomenon necessitates the employment of an NPQ correction that deviates from a $1/iPAR$ relationship.

Normalizing our F_Y data to iPAR and fitting this ratio to a power function at $iPAR > E_{k-F}$ gives best-fit exponents for our low- and high-light treatments of -0.31 (95% CI = 0.1) and -0.54 (95% CI = 0.04), respectively (Fig. 7). These exponents are significantly different than the exponent of -1 associated with the inverse light function of Behrenfeld et al. (2009). Similarly, we obtain exponents of -0.43 and -0.45 using data from the Laney et al. (2005) study. These findings are consistent with an unquenched PSI contribution to fluorescence that becomes more significant as PSII fluorescence is increasingly quenched via NPQ. Our results further suggest that a variable model for NPQ (e.g. corrections ranging from $iPAR^{-0.3}$ to $iPAR^{-0.6}$) may be more appropriate than the single fixed relationship of Behrenfeld et al. (2009). To put these corrections in perspective, the change in the quantum yield of fluorescence due to iron limitation, the largest nutrient-related signal in global data, is about 4-fold (Behrenfeld et al. 2009). If we apply an $iPAR^{-0.4}$ correction for NPQ (rather than $iPAR^{-1}$), fluorescence quantum yields change by up to 2-fold (data not shown). While this result illustrates sensitivity to the choice of NPQ correction exponents, it is not yet clear what range of exponent values should be applied to satellite data. As detailed above, instruments that employ long-pass cutoff filters collect a greater fraction of unquenched PSI fluorescence than instruments measuring a narrow spectral band centered around the PSII fluorescence peak (e.g. MODIS). Accordingly, data from narrow-band measurements require an NPQ correction with an exponent closer to -1 than those derived from long-pass instruments. Employing radiometer based measures of F_Y in laboratory studies would help to determine appropriate corrections (Peperzak et al. 2011).

A notable limitation of the current study is its restriction to a single diatom species. Additional work is still needed to evaluate whether our results are broadly applicable or if taxonomic composition is yet another important confounding factor that must be considered when characterizing NPQ effects. In particular, it remains unclear how changes in PSII:PSI associated with variations in the composition of phytoplankton assemblages influence the contribution of PSI to solar-induced fluorescence emissions.

Acknowledgements. We thank T. Westberry for helpful discussions and comments on early versions of this manuscript. Funding was provided by NASA (NNX08AF72G, NNX07AL80G) and NSF, Biological Oceanography (NSF-OCE 0726116). We also thank the Howard Hughes Medical Institute and K. G. Ahern for advisory support for U.A.A.

LITERATURE CITED

- Behrenfeld MJ, Westberry TK, Boss ES, O'Malley RT and others (2009) Satellite-detected fluorescence reveals global physiology of ocean phytoplankton. *Biogeochemistry* 6:779–794
- Berman T (1972) Profiles of chlorophyll concentrations by *in vivo* fluorescence: some limnological applications. *Limnol Oceanogr* 17:616–618
- Cruz S, Serodio J (2008) Relationship of rapid light curves of variable fluorescence to photoacclimation and non-photochemical quenching in a benthic diatom. *Aquat Bot* 88:256–264
- Cullen JJ, Lewis MR (1995) Biological processes and optical measurements near the sea-surface: some issues relevant to remote-sensing. *J Geophys Res* 100:13255–13266
- D'Asaro EA (2003) Performance of autonomous Lagrangian floats. *J Atmos Ocean Technol* 20:896–911
- Dijkman NA, Kroon BMA (2002) Indications for chlororespiration in relation to light regime in the marine diatom *Thalassiosira weissflogii*. *J Photochem Photobiol B* 66:179–187
- Dimier C, Brunet C, Geider R, Raven J (2009) Growth and photoregulation dynamics of the picoeukaryote *Pelagomonas calceolata* in fluctuating light. *Limnol Oceanogr* 54:823–836
- Eisenstadt D, Ohad I, Keren N, Kaplan A (2008) Changes in the photosynthetic reaction centre II in the diatom *Phaeodactylum tricornerutum* result in non-photochemical fluorescence quenching. *Environ Microbiol* 10:1997–2007
- Falkowski P, Kiefer DA (1985) Chlorophyll-a fluorescence in phytoplankton: relationship to photosynthesis and biomass. *J Plankton Res* 7:715–731
- Falkowski PG, Dubinsky Z, Wyman K (1985) Growth-irradiance relationships in phytoplankton. *Limnol Oceanogr* 30:311–321
- Franck F, Juneau P, Popovic R (2002) Resolution of the Photosystem I and Photosystem II contributions to chlorophyll fluorescence of intact leaves at room temperature. *Biochim Biophys Acta* 1556:239–246
- Franck VM, Bruland KW, Hutchins DA, Brzezinski MA (2003) Iron and zinc effects on silicic acid and nitrate uptake kinetics in three high-nutrient, low-chlorophyll (HNLC) regions. *Mar Ecol Prog Ser* 252:15–33
- Goss R, Pinto EA, Wilhelm C, Richter M (2006) The importance of a highly active and Δ pH-regulated diatoxanthin epoxidase for the regulation of the PSII antenna function in diatoxanthin cycle containing algae. *J Plant Physiol* 163:1008–1021
- Govindjee (2004) Chlorophyll a fluorescence: a bit of basics and history. In: Papageorgiou GC, Govindjee (eds) *Chlorophyll a fluorescence: a signature of photosynthesis*, Book 19. Springer, Dordrecht, p 1–42
- Grouneva I, Jakob T, Wilhelm C, Goss R (2007) Evidence for the existence of a fast xanthophyll cycle-independent NPQ component in diatoms. *Photosynth Res* 91:PS1522
- Grouneva I, Jakob T, Wilhelm C, Goss R (2008) A new multicomponent NPQ mechanism in the diatom *Cyclotella meneghiniana*. *Plant Cell Physiol* 49:1217–1225
- Grouneva I, Jakob T, Wilhelm C, Goss R (2009) The regulation of xanthophyll cycle activity and of non-photochemical fluorescence quenching by two alternative electron flows in the diatoms *Phaeodactylum tricornerutum* and *Cyclotella meneghiniana*. *Biochim Biophys Acta* 1787:929–938
- Holt NE, Fleming GR, Niyogi KK (2004) Toward an understanding of the mechanism of nonphotochemical quenching in green plants. *Biochemistry* 43:8281–8289
- Horton P, Hague A (1988) Studies on the induction of chlorophyll fluorescence in isolated barley protoplasts. 4. Resolution of non-photochemical quenching. *Biochim Biophys Acta* 932:107–115
- Jakob T, Goss R, Wilhelm C (2001) Unusual pH-dependence of diadinoxanthin de-epoxidase activation causes chlororespiratory induced accumulation of diatoxanthin in the diatom *Phaeodactylum tricornerutum*. *J Plant Physiol* 158:383–390
- Jassby AD, Platt T (1976) Mathematical formulation of relationship between photosynthesis and light for phytoplankton. *Limnol Oceanogr* 21:540–547
- Jeffrey SW, Humphrey GF (1975) New spectrophotometric equations for determining chlorophylls *a*, *b*, *c*₁ and *c*₂ in higher plants, algae, and natural phytoplankton. *Biochem Physiol Pflanz* 167:191–194
- Joly D, Carpentier R (2007) Regulation of energy dissipation in photosystem I by the redox state of the plastoquinone pool. *Biochemistry* 46:5534–5541
- Kiefer DA (1973) Chlorophyll *a* fluorescence in marine centric diatoms: responses of chloroplasts to light and nutrient stress. *Mar Biol* 23:39–46
- Kiefer DA, Chamberlin WS, Booth CR (1989) Natural fluorescence of chlorophyll *a*: relationship to photosynthesis and chlorophyll concentration in the western South-Pacific Gyre. *Limnol Oceanogr* 34:868–881
- Krause GH, Weis E (1991) Chlorophyll fluorescence and photosynthesis: the basics. *Annu Rev Plant Physiol Plant Mol Biol* 42:313–349
- Laney SR, Letelier RM, Abbott MR (2005) Parameterizing the natural fluorescence kinetics of *Thalassiosira weissflogii*. *Limnol Oceanogr* 50:1499–1510
- Lavaud J, Kroth PG (2006) In diatoms, the transthylakoid proton gradient regulates the photoprotective non-photochemical fluorescence quenching beyond its control on the xanthophyll cycle. *Plant Cell Physiol* 47:1010–1016
- Lavaud J, Rousseau B, Etienne AL (2002) In diatoms, a transthylakoid proton gradient alone is not sufficient to induce a non-photochemical fluorescence quenching. *FEBS Lett* 523:163–166
- Lavaud J, Strzepak RF, Kroth PG (2007) Photoprotection capacity differs among diatoms: possible consequences on the spatial distribution of diatoms related to fluctuations in the underwater light climate. *Limnol Oceanogr* 52:1188–1194
- Lee HY, Hong YN, Chow WS (2001) Photoinactivation of photosystem II complexes and photoprotection by non-functional neighbours in *Capsicum annum* L. leaves. *Planta* 212:332–342
- Letelier RM, Abbott MR, Karl DM (1997) Chlorophyll natural fluorescence response to upwelling events in the Southern Ocean. *Geophys Res Lett* 24:409–412
- Lewis MR, Smith JC (1983) A small volume, short-incubation-time method for measurement of photosynthesis as a function of incident irradiance. *Mar Ecol Prog Ser* 13:99–102
- Miloslavina Y, Grouneva I, Lambrev PH, Lepetit B, Goss R, Wilhelm C, Holzwarth AR (2009) Ultrafast fluorescence study on the location and mechanism of non-photochemical quenching in diatoms. *Biochim Biophys Acta* 1787:1189–1197

- Mitchell BG, Kahru M, Wieland J, Stramska M (2003) Determination of spectral absorption coefficients of particles, dissolved material and phytoplankton for discrete water samples. In: Mueller JL, Fargoin GS, McClain CR (eds) Ocean optics protocols for satellite ocean color sensor validation, Vol. 4. NASA/TM-2003-211621/Rev4-Vol4. NASA Goddard Space Flight Center, Greenbelt, MD, p 39–56
- Morel A, Prieur L (1977) Analysis of variations in ocean color. *Limnol Oceanogr* 22:709–722
- Morrison JR (2003) *In situ* determination of the quantum yield of phytoplankton chlorophyll *a* fluorescence: a simple algorithm, observations, and a model. *Limnol Oceanogr* 48:618–631
- Neville RA, Gower JFR (1977) Passive remote-sensing of phytoplankton via chlorophyll *a* fluorescence. *J Geophys Res* 82:3487–3493
- Niyogi KK, Bjorkman O, Grossman AR (1997) *Chlamydomonas* xanthophyll cycle mutants identified by video imaging of chlorophyll fluorescence quenching. *Plant Cell* 9:1369–1380
- Öquist G, Chow WS, Anderson JM (1992) Photoinhibition of photosynthesis represents a mechanism for the long-term regulation of photosystem-II. *Planta* 186:450–460
- Owens TG (1986) Light-harvesting function in the diatom *Phaeodactylum tricorutum*. 2. Distribution of excitation-energy between the photosystems. *Plant Physiol* 80: 739–746
- Peperzak L, Timmermans KR, Wernand MR, Oosterhuis S, van der Woerd HJ (2011) A mesocosm tool to optically study phytoplankton dynamics. *Limnol Oceanogr Methods* 9:232–244
- Ragni M, Ainsworth RL, Leonardos N, Geider RJ (2008) Photoinhibition of PSII in *Emiliania huxleyi* (Haptophyta) under high light stress: the roles of photoacclimation, photoprotection, and photorepair. *J Phycol* 44:670–683
- Ruban AV, Lavaud J, Rousseau B, Guglielmi G, Horton P, Etienne AL (2004) The super-excess energy dissipation in diatom algae: comparative analysis with higher plants. *Photosynth Res* 82:165–175
- Schallenberg C, Lewis MR, Kelley DE, Cullen JJ (2008) Inferred influence of nutrient availability on the relationship between sun-induced chlorophyll fluorescence and incident irradiance in the Bering Sea. *J Geophys Res* 113:C07046, doi:10.1029/2007JC004355
- Strzepek RF, Harrison PJ (2004) Photosynthetic architecture differs in coastal and oceanic diatoms. *Nature* 431: 689–692
- Ting CS, Owens TG (1993) Photochemical and nonphotochemical fluorescence quenching processes in the diatom *Phaeodactylum tricorutum*. *Plant Physiol* 101: 1323–1330
- Ting CS, Owens TG (1994) The effects of excess irradiance on photosynthesis in the marine diatom *Phaeodactylum tricorutum*. *Plant Physiol* 106:763–770
- van de Poll WH, Visser RJW, Buma AGJ (2007) Acclimation to a dynamic irradiance regime changes excessive irradiance sensitivity of *Emiliania huxleyi* and *Thalassiosira weissflogii*. *Limnol Oceanogr* 52:1430–1438
- van de Poll WH, Buma AGJ, Visser RJW, Janknegt PJ, Villafane VE, Helbling EW (2010) Xanthophyll cycle activity and photosynthesis of *Dunaliella tertiolecta* (Chlorophyceae) and *Thalassiosira weissflogii* (Bacillariophyceae) during fluctuating solar radiation. *Phycologia* 49:249–259
- Westberry TK, Siegel DA (2003) Phytoplankton natural fluorescence variability in the Sargasso Sea. *Deep-Sea Res I* 50:417–434

Editorial responsibility: Matthias Seaman,
Oldendorf/Luhe, Germany

Submitted: March 9, 2011; Accepted: December 5, 2011
Proofs received from author(s): February 17, 2012