

False-Positive “Necklace Sign” on Whole-Body Thyroid Cancer Survey—A Case Report

Sylvain Beaulieu, MD; David Djang, MD; and Satoshi Minoshima, MD, PhD

Division of Nuclear Medicine, University of Washington, Seattle, Washington

A woman with papillary thyroid carcinoma treated with surgery and postoperative ^{131}I returned 4 y later for a whole-body survey. The imaging results initially suggested disease recurrence but were later found to be false positive and due to ^{131}I uptake in a necklace that had been contaminated by the patient's saliva. This case stresses the importance of having the patient remove all jewelry before undergoing a radioiodine survey.

Key Words: endocrinology; oncology; quality assurance; false positive; ^{131}I ; thyroid carcinoma

J Nucl Med Technol 2005; 33:42–43

Whole-body radioiodine cancer surveys are an integral part of the diagnostic algorithm for patients with thyroid cancer. Radioiodine has several regions of physiologic uptake, including the salivary glands, the gastrointestinal tract, and the urinary tract, that can have a variable appearance (1–2). In uncharacteristic regions of uptake, contamination must be distinguished from actual disease (3–5).

CASE REPORT

A 25-y-old Russian woman with a history of papillary thyroid carcinoma, which was resected in 1998, underwent a postoperative ^{131}I cancer survey revealing only a small amount of tissue in the thyroid bed, likely remnant thyroid tissue. The patient was treated postoperatively with 2,849 MBq of ^{131}I . The patient returned for a recombinant thyrotropin whole-body survey December 2002 while having near-undetectable levels of thyroglobulin. At first glance, the images (Fig. 1) were suggestive of recurrence of disease at the base of the neck on the left side. After careful consideration, the initial finding was suspected of being false positive. The patient was called at home and agreed to return to the nuclear medicine clinic the following day. Images were taken (Fig. 2) without the necklace. The neck-

lace was then removed and imaged alone and showed activity (Fig. 3). The signature spectrum on a multichannel analyzer confirmed that the necklace activity was from ^{131}I , thus excluding the less likely possibility of an unidentified contaminant, such as a heavy isotope uncommonly contained in Eastern European jewelry. After further questioning, the patient described a habit of placing the necklace in her mouth and confirmed doing so after taking her radioiodine dose. Figure 4 shows the necklace on the patient.




FIGURE 1. Recombinant thyrotropin whole-body survey that suggests recurrence of disease at base of neck on left side. Shown are $^{99\text{m}}\text{Tc}$ image with radioactive markers (A) and ^{131}I image (B). Both are anterior views.




FIGURE 2. $^{99\text{m}}\text{Tc}$ image (A) and ^{131}I image (B) taken without necklace. Both are anterior views.

DISCUSSION

In this patient, a whole-body survey showed a finding that initially created concern about recurrent thyroid carcinoma in the base of the neck but later was revealed to be ^{131}I

For correspondence or reprints contact: Sylvain Beaulieu, MD, 1511 Rue Lajoie, Outremont, Quebec, Canada, H2V 1R2.
E-mail: sylvain.beaulieu@bigfoot.com




FIGURE 3. ^{131}I image of necklace alone.



FIGURE 4. Photograph of necklace on patient.

uptake in a necklace that had been contaminated by the patient's saliva. This case stresses the importance of having the patient remove all jewelry before undergoing a radioiodine survey.

REFERENCES

1. Carlisle MR, Lu C, McDougall IR. The interpretation of ^{131}I scans in the evaluation of thyroid cancer, with an emphasis on false positive findings. *Nucl Med Commun*. 2003;24:715–735.
2. Leitha T, Staudenherz A. Frequency of diagnostic dilemmas in ^{131}I whole body scanning. *Nuklearmedizin*. 2003;42:55–62.
3. Pochis WT, Krasnow AZ, Isitman AT, et al. The radioactive handkerchief sign: a contamination artifact in I-131 imaging for metastatic thyroid carcinoma. *Clin Nucl Med*. 1990;15:491–494.
4. Dick C, Mudun A, Alazraki NP. False-positive I-131 images mimicking thyroid cancer metastasis: the nose ring sign. *Clin Nucl Med*. 1995;20:876–877.
5. Joyce WT, Cowan RJ. A potential false-positive posttherapy radioiodine scan secondary to I-131 excretion in perspiration. *Clin Nucl Med*. 1995;20:368–369.





False-Positive "Necklace Sign" on Whole-Body Thyroid Cancer Survey—A Case Report

Sylvain Beaulieu, David Djang and Satoshi Minoshima

JNMT 2005;33:42-43.

This article and updated information are available at:
<http://tech.snmjournals.org/content/33/1/42>

Information about reproducing figures, tables, or other portions of this article can be found online at:
<http://tech.snmjournals.org/site/misc/permission.xhtml>

Information about subscriptions to can be found at:
<http://tech.snmjournals.org/site/subscriptions/online.xhtml>

Journal of Nuclear Medicine Technology is published quarterly.
SNMMI | Society of Nuclear Medicine and Molecular Imaging
1850 Samuel Morse Drive, Reston, VA 20190.
(Print ISSN: 0091-4916, Online ISSN: 1535-5675)

© Copyright 2005 SNMMI; all rights reserved.